

DEX-0142

EL

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12P 21/04, C12N 15/00, C07H 21/02		A1	(11) International Publication Number: WO 00/55351
		(43) International Publication Date: 21 September 2000 (21.09.00)	
(21) International Application Number: PCT/US00/05883 (22) International Filing Date: 8 March 2000 (08.03.00) (30) Priority Data: 60/124,270 12 March 1999 (12.03.99) US (71) Applicant (for all designated States except US): HUMAN GENOME SCIENCES, INC. [US/US]; 9410 Key West Avenue, Rockville, MD 20850 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): ROSEN, Craig, A. [US/US]; 22400 Rolling Hill Road, Laytonsville, MD 20882 (US). RUBEN, Steven, M. [US/US]; 18528 Heritage Hills Drive, Laytonsville, MD 20882 (US). (74) Agents: WALES, Michele, M. et al.; Human Genome Sciences, Inc., 9410 Key West Avenue, Rockville, MD 20850 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	
(54) Title: HUMAN COLON CANCER ASSOCIATED GENE SEQUENCES AND POLYPEPTIDES			
(57) Abstract			
<p>This invention relates to newly identified colon or colon cancer related polynucleotides and the polypeptides encoded by these polynucleotides herein collectively known as "colon cancer antigens", and to the complete gene sequences associated therewith and to the expression products thereof, as well as the use of such colon cancer antigens for detection, prevention and treatment of disorders of the colon, particularly the presence of colon cancer. This invention relates to the colon cancer antigens as well as vectors, host cells, antibodies directed to colon cancer antigens and recombinant and synthetic methods for producing the same. Also provided are diagnostic methods for diagnosing and treating, preventing and/or prognosing disorders related to the colon, including colon cancer, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying agonists and antagonists of colon cancer antigens of the invention. The present invention further relates to methods and/or compositions for inhibiting the production and/or function of the polypeptides of the present invention.</p>			

53

5110-X3G



THIS PAGE BLANK (USPTO)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

THIS PAGE BLANK (USPTO)

Human Colon Cancer Associated Gene Sequences and Polypeptides

5

Field of the Invention

This invention relates to newly identified colon or colon cancer related polynucleotides and the polypeptides encoded by these polynucleotides herein collectively known as "colon cancer antigens," and to the complete gene sequences
10 associated therewith and to the expression products thereof, as well as the use of such colon cancer antigens for detection, prevention and treatment of disorders of the colon, particularly the presence of colon cancer. This invention relates to the colon cancer antigens as well as vectors, host cells, antibodies directed to colon cancer antigens and recombinant and synthetic methods for producing the same. Also
15 provided are diagnostic methods for diagnosing and treating, preventing and/or prognosing disorders related to the colon, including colon cancer, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying agonists and antagonists of colon cancer antigens of the invention. The present invention further relates to methods and/or compositions for
20 inhibiting the production and/or function of the polypeptides of the present invention.

Background of the Invention

Colorectal cancers are among the most common cancers in men and women in the U.S. and are one of the leading causes of death. Other than surgical resection no
25 other systemic or adjuvant therapy is available. Vogelstein and colleagues have described the sequence of genetic events that appear to be associated with the multistep process of colon cancer development in humans (Trends Genet 9(4):138-41 (1993)). An understanding of the molecular genetics of carcinogenesis, however, has not led to preventative or therapeutic measures. It can be expected that advances in
30 molecular genetics will lead to better risk assessment and early diagnosis but colorectal cancers will remain a deadly disease for a majority of patients due to the

lack of an adjuvant therapy. Adjuvant or systemic treatments are likely to arise from a better understanding of the autocrine factors responsible for the continued proliferation of cancer cells.

Colorectal carcinoma is a malignant neoplastic disease. There is a high incidence of colorectal carcinoma in the Western world, particularly in the United States. Tumors of this type often metastasize through lymphatic and vascular channels. Many patients with colorectal carcinoma eventually die from this disease. In fact, it is estimated that 62,000 persons in the United States alone die of colorectal carcinoma annually.

At the present time the only systemic treatment available for colon cancer is chemotherapy. However, chemotherapy has not proven to be very effective for the treatment of colon cancers for several reasons, the most important of which is the fact that colon cancers express high levels of the MDR gene (that codes for multi-drug resistance gene products). The MDR gene products actively transport the toxic substances out of the cell before the chemotherapeutic agents can damage the DNA machinery of the cell. These toxic substances harm the normal cell populations more than they harm the colon cancer cells for the above reasons.

There is no effective systemic treatment for treating colon cancers other than surgically removing the cancers. In the case of several other cancers, including breast cancers, the knowledge of growth promoting factors (such as EGF, estradiol, IGF-11) that appear to be expressed or effect the growth of the cancer cells, has been translated for treatment purposes. But in the case of colon cancers this knowledge has not been applied and therefore the treatment outcome for colon cancers remains bleak.

There is a need, therefore, for identification and characterization of such factors that modulate activation and differentiation of colon cells, both normally and in disease states. In particular, there is a need to isolate and characterize additional molecules that mediate apoptosis, DNA repair, tumor-mediated angiogenesis, genetic imprinting, immune responses to tumors and tumor antigens and, among other things, that can play a role in detecting, preventing, ameliorating or correcting dysfunctions or diseases of the colon.

Summary of the Invention

The present invention includes isolated nucleic acid molecules comprising, or alternatively, consisting of, a colon and/or colon cancer associated polynucleotide sequence disclosed in the sequence listing (as SEQ ID Nos: 1 to 773) and/or contained in a human cDNA clone described in Tables 1, 2 and 5 and deposited with the American Type Culture Collection ("ATCC"). Fragments, variant, and derivatives of these nucleic acid molecules are also encompassed by the invention. The present invention also includes isolated nucleic acid molecules comprising, or alternatively consisting of, a polynucleotide encoding a colon or colon cancer polypeptide. The present invention further includes colon and/or colon cancer polypeptides encoded by these polynucleotides. Further provided for are amino acid sequences comprising, or alternatively consisting of, colon and/or colon cancer polypeptides as disclosed in the sequence listing (as SEQ ID Nos: 774 to 1546) and/or encoded by a human cDNA clone described in Tables 1, 2 and 5 and deposited with the ATCC. Antibodies that bind these polypeptides are also encompassed by the invention. Polypeptide fragments, variants, and derivatives of these amino acid sequences are also encompassed by the invention, as are polynucleotides encoding these polypeptides and antibodies that bind these polypeptides. Also provided are diagnostic methods for diagnosing and treating, preventing, and/or prognosing disorders related to the colon, including colon cancer, and therapeutic methods for treating such disorders. The invention further relates to screening methods for identifying agonists and antagonists of colon cancer antigens of the invention.

Detailed Description

25

Tables

Table 1 summarizes some of the colon cancer antigens encompassed by the invention (including contig sequences (SEQ ID NO:X) and the cDNA clone related to the contig sequence) and further summarizes certain characteristics of the colon cancer polynucleotides and the polypeptides encoded thereby. The first column shows the "SEQ ID NO:" for each of the 773 colon cancer antigen polynucleotide sequences of the invention. The second column provides a unique "Sequence/Contig ID"

30

identification for each colon and/or colon cancer associated sequence. The third column, "Gene Name," and the fourth column, "Overlap," provide a putative identification of the gene based on the sequence similarity of its translation product to an amino acid sequence found in a publicly accessible gene database and the database
5 accession no. for the database sequence having similarity, respectively. The fifth and sixth columns provide the location (nucleotide position nos. within the contig), "Start" and "End", in the polynucleotide sequence "SEQ ID NO:X" that delineate the preferred ORF shown in the sequence listing as SEQ ID NO:Y. The seventh and eighth columns provide the "% Identity" (percent identity) and "% Similarity"
10 (percent similarity), respectively, observed between the aligned sequence segments of the translation product of SEQ ID NO:X and the database sequence. The ninth column provides a unique "Clone ID" for a cDNA clone related to each contig sequence.

Table 2 summarizes ATCC Deposits, Deposit dates, and ATCC designation numbers of deposits made with the ATCC in connection with the present application.

15 Table 3 indicates public ESTs, of which at least one, two, three, four, five, ten, fifteen or more of any one or more of these public EST sequences are optionally excluded from certain embodiments of the invention.

Table 4 lists residues comprising antigenic epitopes of antigenic epitope-bearing fragments present in most of the colon or colon cancer associated
20 polynucleotides described in Table 1 as predicted by the inventors using the algorithm of Jameson and Wolf, (1988) Comp. Appl. Biosci. 4:181-186. The Jameson-Wolf antigenic analysis was performed using the computer program PROTEAN (Version 3.11 for the Power MacIntosh, DNASTAR, Inc., 1228 South Park Street Madison, WI). Colon and colon cancer associated polypeptides shown in Table 1 may possess
25 one or more antigenic epitopes comprising residues described in Table 4. It will be appreciated that depending on the analytical criteria used to predict antigenic determinants, the exact address of the determinant may vary slightly. The residues and locations shown in Table 4 correspond to the amino acid sequences for most colon and colon cancer associated polypeptide sequence shown in the Sequence Listing.

30 Table 5 shows the cDNA libraries sequenced, and ATCC designation numbers and vector information relating to these cDNA libraries.

Definitions

The following definitions are provided to facilitate understanding of certain terms used throughout this specification.

5 In the present invention, "isolated" refers to material removed from its original environment (e.g., the natural environment if it is naturally occurring), and thus is altered "by the hand of man" from its natural state. For example, an isolated polynucleotide could be part of a vector or a composition of matter, or could be contained within a cell, and still be "isolated" because that vector, composition of
10 matter, or particular cell is not the original environment of the polynucleotide. The term "isolated" does not refer to genomic or cDNA libraries, whole cell total or mRNA preparations, genomic DNA preparations (including those separated by electrophoresis and transferred onto blots), sheared whole cell genomic DNA preparations or other compositions where the art demonstrates no distinguishing
15 features of the polynucleotide/sequences of the present invention.

As used herein, a "polynucleotide" refers to a molecule having a nucleic acid sequence contained in SEQ ID NO:X (as described in column 1 of Table 1) or the related cDNA clone (as described in column 9 of Table 1 and contained within a library deposited with the ATCC). For example, the polynucleotide can contain the
20 nucleotide sequence of the full length cDNA sequence, including the 5' and 3' untranslated sequences, the coding region, as well as fragments, epitopes, domains, and variants of the nucleic acid sequence. Moreover, as used herein, a "polypeptide" refers to a molecule having an amino acid sequence encoded by a polynucleotide of the invention as broadly defined (obviously excluding poly-Phenylalanine or poly-
25 Lysine peptide sequences which result from translation of a polyA tail of a sequence corresponding to a cDNA).

In the present invention, "SEQ ID NO:X" was often generated by overlapping sequences contained in multiple clones (contig analysis). A representative clone containing all or most of the sequence for SEQ ID NO:X is deposited at Human
30 Genome Sciences, Inc. (HGS) in a catalogued and archived library. As shown in column 9 of Table 1, each clone is identified by a cDNA Clone ID. Each Clone ID is unique to an individual clone and the Clone ID is all the information needed to

retrieve a given clone from the HGS library. In addition to the individual cDNA clone deposits, most of the cDNA libraries from which the clones were derived were deposited at the American Type Culture Collection (hereinafter "ATCC"). Table 5 provides a list of the deposited cDNA libraries. One can use the Clone ID to
5 determine the library source by reference to Tables 2 and 5. Table 5 lists the deposited cDNA libraries by name and links each library to an ATCC Deposit. Library names contain four characters, for example, "HTWE." The name of a cDNA clone ("Clone ID") isolated from that library begins with the same four characters, for example "HTWEP07". As mentioned below, Table 1 correlates the Clone ID names
10 with SEQ ID NOs. Thus, starting with a SEQ ID NO, one can use Tables 1, 2 and 5 to determine the corresponding Clone ID, from which library it came and in which ATCC deposit the library is contained. Furthermore, it is possible to retrieve a given cDNA clone from the source library by techniques known in the art and described elsewhere herein. The ATCC is located at 10801 University Boulevard, Manassas,
15 Virginia 20110-2209, USA. The ATCC deposits were made pursuant to the terms of the Budapest Treaty on the international recognition of the deposit of microorganisms for the purposes of patent procedure.

A "polynucleotide" of the present invention also includes those polynucleotides capable of hybridizing, under stringent hybridization conditions, to
20 sequences contained in SEQ ID NO:X, or the complement thereof (e.g., the complement of any one, two, three, four, or more of the polynucleotide fragments described herein), and/or sequences contained in the related cDNA clone within a library deposited with the ATCC. "Stringent hybridization conditions" refers to an overnight incubation at 42 degree C in a solution comprising 50% formamide, 5x SSC
25 (750 mM NaCl, 75 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared salmon sperm DNA, followed by washing the filters in 0.1x SSC at about 65 degree C.

Also included within "polynucleotides" of the present invention are nucleic acid molecules that hybridize to the polynucleotides of the present invention at lower
30 stringency hybridization conditions. Changes in the stringency of hybridization and signal detection are primarily accomplished through the manipulation of formamide concentration (lower percentages of formamide result in lowered stringency); salt

conditions, or temperature. For example, lower stringency conditions include an overnight incubation at 37 degree C in a solution comprising 6X SSPE (20X SSPE = 3M NaCl; 0.2M NaH₂PO₄; 0.02M EDTA, pH 7.4), 0.5% SDS, 30% formamide, 100 ug/ml salmon sperm blocking DNA; followed by washes at 50 degree C with
5 1XSSPE, 0.1% SDS. In addition, to achieve even lower stringency, washes performed following stringent hybridization can be done at higher salt concentrations (e.g. 5X SSC).

Note that variations in the above conditions may be accomplished through the inclusion and/or substitution of alternate blocking reagents used to suppress
10 background in hybridization experiments. Typical blocking reagents include Denhardt's reagent, BLOTTO, heparin, denatured salmon sperm DNA, and commercially available proprietary formulations. The inclusion of specific blocking reagents may require modification of the hybridization conditions described above, due to problems with compatibility.

15 Of course, a polynucleotide which hybridizes only to polyA+ sequences (such as any 3' terminal polyA+ tract of a cDNA shown in the sequence listing), or to a complementary stretch of T (or U) residues, would not be included in the definition of "polynucleotide," since such a polynucleotide would hybridize to any nucleic acid molecule containing a poly (A) stretch or the complement thereof (e.g., practically
20 any double-stranded cDNA clone generated using oligo dT as a primer).

The polynucleotides of the present invention can be composed of any polyribonucleotide or polydeoxribonucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. For example, polynucleotides can be composed of single- and double-stranded DNA, DNA that is a mixture of single- and double-
25 stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, the polynucleotide can be composed of triple-stranded regions comprising RNA or DNA or both RNA and DNA. A
30 polynucleotide may also contain one or more modified bases or DNA or RNA backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of

modifications can be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically, or metabolically modified forms.

In specific embodiments, the polynucleotides of the invention are at least 15, at least 30, at least 50, at least 100, at least 125, at least 500, or at least 1000 continuous nucleotides but are less than or equal to 300 kb, 200 kb, 100 kb, 50 kb, 15 kb, 10 kb, 7.5kb, 5 kb, 2.5 kb, 2.0 kb, or 1 kb, in length. In a further embodiment, polynucleotides of the invention comprise a portion of the coding sequences, as disclosed herein, but do not comprise all or a portion of any intron. In another embodiment, the polynucleotides comprising coding sequences do not contain coding sequences of a genomic flanking gene (i.e., 5' or 3' to the gene of interest in the genome). In other embodiments, the polynucleotides of the invention do not contain the coding sequence of more than 1000, 500, 250, 100, 50, 25, 20, 15, 10, 5, 4, 3, 2, or 1 genomic flanking gene(s).

"SEQ ID NO:X" refers to a colon cancer antigen polynucleotide sequence described in Table 1. SEQ ID NO:X is identified by an integer specified in column 1 of Table 1. The polypeptide sequence SEQ ID NO:Y is a translated open reading frame (ORF) encoded by polynucleotide SEQ ID NO:X. There are 773 colon cancer antigen polynucleotide sequences described in Table 1 and shown in the sequence listing (SEQ ID NO:1 through SEQ ID NO:773). Likewise there are 773 polypeptide sequences shown in the sequence listing, one polypeptide sequence for each of the polynucleotide sequences (SEQ ID NO:774 through SEQ ID NO:1546). The polynucleotide sequences are shown in the sequence listing immediately followed by all of the polypeptide sequences. Thus, a polypeptide sequence corresponding to polynucleotide sequence SEQ ID NO:1 is the first polypeptide sequence shown in the sequence listing. The second polypeptide sequence corresponds to the polynucleotide sequence shown as SEQ ID NO:2, and so on. In other words, since there are 773 polynucleotide sequences, for any polynucleotide sequence SEQ ID NO:X, a corresponding polypeptide SEQ ID NO:Y can be determined by the formula $X + 773 = Y$. In addition, any of the unique "Sequence/Contig ID" defined in column two of Table 1, can be linked to the corresponding polypeptide SEQ ID NO:Y by reference to Table 4.

The polypeptides of the present invention can be composed of amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres, and may contain amino acids other than the 20 gene-encoded amino acids. The polypeptides may be modified by either natural processes, such as posttranslational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched, for example, as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched, and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. (See, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York (1993); POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, pgs. 1-12 (1983); Seifter et al., Meth Enzymol 182:626-646 (1990); Rattan et al., Ann NY Acad Sci 663:48-62 (1992).)

The colon and colon cancer polypeptides of the invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced

polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

The polypeptides may be in the form of the secreted protein, including the mature form, or may be a part of a larger protein, such as a fusion protein (see below).

5 It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification, such as multiple histidine residues, or an additional sequence for stability during recombinant production.

10 The colon and colon cancer polypeptides of the present invention are preferably provided in an isolated form, and preferably are substantially purified. A recombinantly produced version of a polypeptide, including the secreted polypeptide, can be substantially purified using techniques described herein or otherwise known in the art, such as, for example, by the one-step method described in Smith and Johnson, Gene 67:31-40 (1988). Polypeptides of the invention also can be purified from
15 natural, synthetic or recombinant sources using techniques described herein or otherwise known in the art, such as, for example, antibodies of the invention raised against the polypeptides of the present invention in methods which are well known in the art.

By a polypeptide demonstrating a "functional activity" is meant, a polypeptide
20 capable of displaying one or more known functional activities associated with a full-length (complete) protein of the invention. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a polypeptide for binding) to an anti-polypeptide antibody], immunogenicity (ability to generate antibody which binds to a specific polypeptide of the invention), ability to
25 form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide.

"A polypeptide having functional activity" refers to polypeptides exhibiting activity similar, but not necessarily identical to, an activity of a polypeptide of the present invention, including mature forms, as measured in a particular assay, such as,
30 for example, a biological assay, with or without dose dependency. In the case where dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to

the polypeptide of the present invention (i.e., the candidate polypeptide will exhibit greater activity or not more than about 25-fold less and, preferably, not more than about tenfold less activity, and most preferably, not more than about three-fold less activity relative to the polypeptide of the present invention).

5 The functional activity of the colon cancer antigen polypeptides, and fragments, variants derivatives, and analogs thereof, can be assayed by various methods.

For example, in one embodiment where one is assaying for the ability to bind or compete with full-length polypeptide of the present invention for binding to an
10 antibody to the full length polypeptide antibody, various immunoassays known in the art can be used, including but not limited to, competitive and non-competitive assay systems using techniques such as radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoradiometric assays, gel diffusion precipitation reactions, immunodiffusion assays, in situ immunoassays
15 (using colloidal gold, enzyme or radioisotope labels, for example), western blots, precipitation reactions, agglutination assays (e.g., gel agglutination assays, hemagglutination assays), complement fixation assays, immunofluorescence assays, protein A assays, and immunoelectrophoresis assays, etc. In one embodiment, antibody binding is detected by detecting a label on the primary antibody. In another
20 embodiment, the primary antibody is detected by detecting binding of a secondary antibody or reagent to the primary antibody. In a further embodiment, the secondary antibody is labeled. Many means are known in the art for detecting binding in an immunoassay and are within the scope of the present invention.

In another embodiment, where a ligand is identified, or the ability of a
25 polypeptide fragment, variant or derivative of the invention to multimerize is being evaluated, binding can be assayed, e.g., by means well-known in the art, such as, for example, reducing and non-reducing gel chromatography, protein affinity chromatography, and affinity blotting. See generally, Phizicky, E., et al., Microbiol. Rev. 59:94-123 (1995). In another embodiment, physiological correlates polypeptide
30 of the present invention binding to its substrates (signal transduction) can be assayed.

In addition, assays described herein (see Examples) and otherwise known in the art may routinely be applied to measure the ability of polypeptides of the present

invention and fragments, variants derivatives and analogs thereof to elicit polypeptide related biological activity (either in vitro or in vivo). Other methods will be known to the skilled artisan and are within the scope of the invention.

5

Colon and Colon Cancer Associated Polynucleotides and Polypeptides of the Invention

It has been discovered herein that the polynucleotides described in Table 1 are expressed at significantly enhanced levels in human colon and/or colon cancer tissues.

10 Accordingly, such polynucleotides, polypeptides encoded by such polynucleotides, and antibodies specific for such polypeptides find use in the prediction, diagnosis, prevention and treatment of colon related disorders, including colon cancer as more fully described below.

15 Table 1 summarizes some of the polynucleotides encompassed by the invention (including contig sequences (SEQ ID NO:X) and the related cDNA clones) and further summarizes certain characteristics of these colon and/or colon cancer associated polynucleotides and the polypeptides encoded thereby.

Table 1

Seq ID No.	Sequence/ Contig ID	Gene Name	Overlap	HGS Nucleotide Start End	% Identity	% Similarity	Clone ID
1	500802	Immunoglobulin kappa light chain variable region L25 [Homo sapiens] >pir S41816 S41816 Ig kappa chain V region L25 - human Length = 119 (AB008790) Grb7V protein [Homo sapiens] >sp D103000 D103000 GRB7V PROTEIN. >gi 1526535 Grb7 protein [Homo sapiens] {SUB 130-343} Length = 447	gi 415381	2 304	73	86	HGBA183
2	531091			2 292			HUKDY21
3	553147			3 440			HCASG85
4	558860	(AF039700) antigen NY-CO-38 [Homo sapiens] >sp G3170200 G3170200 ANTIGEN NY-CO-38. >gi 3170198 (AF039699) antigen NY-CO-37 [Homo sapiens] {SUB 1-403} Length = 652 MDA-7 [Homo sapiens] >sp Q13007 MDA7_HUMAN MDA-7 PROTEIN PRECURSOR (MELANOMA DIFFERENTIATION ASSOCIATED PROTEIN 7). Length = 206 disintegrin-protease [Homo sapiens] >sp O15204 O15204 DISINTEGRIN- PROTEASE. Length = 470	gn PID d1030000	33 635	97	98	HCEGY28
5	561730	(AF039700) antigen NY-CO-38 [Homo sapiens] >sp G3170200 G3170200 ANTIGEN NY-CO-38. >gi 3170198 (AF039699) antigen NY-CO-37 [Homo sapiens] {SUB 1-403} Length = 652 MDA-7 [Homo sapiens] >sp Q13007 MDA7_HUMAN MDA-7 PROTEIN PRECURSOR (MELANOMA DIFFERENTIATION ASSOCIATED PROTEIN 7). Length = 206 disintegrin-protease [Homo sapiens] >sp O15204 O15204 DISINTEGRIN- PROTEASE. Length = 470	gi 3170200	34 393	98	98	HSDFA48
6	585938	(AF039700) antigen NY-CO-38 [Homo sapiens] >sp G3170200 G3170200 ANTIGEN NY-CO-38. >gi 3170198 (AF039699) antigen NY-CO-37 [Homo sapiens] {SUB 1-403} Length = 652 MDA-7 [Homo sapiens] >sp Q13007 MDA7_HUMAN MDA-7 PROTEIN PRECURSOR (MELANOMA DIFFERENTIATION ASSOCIATED PROTEIN 7). Length = 206 disintegrin-protease [Homo sapiens] >sp O15204 O15204 DISINTEGRIN- PROTEASE. Length = 470	gi 1141751	206 538	81	81	HMQBR31
7	587785	(AF039700) antigen NY-CO-38 [Homo sapiens] >sp G3170200 G3170200 ANTIGEN NY-CO-38. >gi 3170198 (AF039699) antigen NY-CO-37 [Homo sapiens] {SUB 1-403} Length = 652 MDA-7 [Homo sapiens] >sp Q13007 MDA7_HUMAN MDA-7 PROTEIN PRECURSOR (MELANOMA DIFFERENTIATION ASSOCIATED PROTEIN 7). Length = 206 disintegrin-protease [Homo sapiens] >sp O15204 O15204 DISINTEGRIN- PROTEASE. Length = 470	gn PID e332729	2 331	100	100	HIOB086

8	588916	Human apoC-II gene for preapoplipoprotein C-II [Homo sapiens] >gi 757915 apoCII protein [Homo sapiens] >gi 178836 apolipoprotein C-II [Homo sapiens] >pir A24238 LPHUC2 apolipoprotein C- II precursor - human	gi 296636	5	376	100	100	100	HLIDQU56
9	613825	KHS1 [Homo sapiens] >sp G185733 G185733.1 KHS1. Length = 846 protein kinase Dyrk2 [Homo sapiens] >sp Q92630 Q92630 PROTEIN KINASE DYRK2 (PROTEIN KINASE, DYRK2). >gnl PIDe280618 Dyrk2 [Homo sapiens] {SUB 320-528} Length = 528	gi 1857331	3	260				HMSHB03
10	639090			254	559				HCRME22
11	651644			63	194				HCFO73
12	659544			109	249				HJMBU15
13	659739			238	1140	94			HSYAM68
14	661057	(AJ003061) most expressed alternative spliced form [Homo sapiens] >sp O60852 O60852 PROTEIN ENCODED BY SACCHAROMYCES CEREVISIAE SPC98 HOMOLOGUE. Length = 907	gnl PIDe321513	3	425	100			HCDBX83
15	661313			894	1118				HHEMN11
16	666316			193	369				HCDC84
17	669229			430	762				HOHDD51
18	670471			203	937	92	93		HAGGX21
19	676611		gnl PIDe1293754	207	530				HCE5C73
20	691240			2	385				HISAN54

21	702977	26-kDa cell surface protein TAPA-1 [Homo sapiens] >pir A35649 A35649 cell surface protein TAPA-1 - human >sp P18582 CD81_HUMAN CD81 ANTIGEN (26 KD CELL SURFACE PROTEIN TAPA-1). Length = 236	gi 338678	34	819	80	80	HGCMV09
22	709517	(AF062476) retinoic acid-responsive protein; STRA6 [Mus musculus] >sp O70491 O70491 RETINOIC ACID- RESPONSIVE PROTEIN. Length = 670	gi 3126975	344	478	75	88	HWLJX38
23	714730			1	534			HCRND05
24	714834			1	657			IIA'PTL75
25	715016			1	411	62	64	HCEOQ15
26	719584	(AF076856) small espin [Rattus norvegicus] >sp G33818569 G33818569 SMALL ESPIN. Length = 253	gi 3818569	530	886			HWLFA47
27	724637	muskelin [Mus musculus] >sp O89050 O89050 MUSKELIN. Length = 735	gi 3493462	1	444	92	95	HUSXP30
28	728392	(AB015318) gamma2-adaptin [Homo sapiens] >sp O75843 O75843 GAMMA2-ADAPTIN. Length = 785	gn PI D d1034356	160	801	100	100	HBJIG25
29	738716			137	289			HCRMQ71
30	739056	similar to ADP-ribosylation factor; (AF054179) H beta 58 homolog [Homo sapiens] >sp O75436 O75436 H BETA 58 HOMOLOG. Length = 327	gn PI D e1350748	2	502	87	97	HSDZB27
31	739143		gi 3342000	1	1083	100	100	HKABV36
32	742329			2	277			HSRDE23
33	742557			2	814			HWHGD94
34	745481			1229	1396			HPMFL67
35	746035			414	959			HCHCJ20
36	753731			1	357			HPTTL69

54	784446	(AB002086) p47 [Rattus norvegicus]	19	282		1H3JFL85
55	784832	>gnl PID e294068 XY40 protein [Rattus norvegicus]	134	751		HCGMI84
56	786813	>sp O35987 O35987 P47, COMPLETE CDS. Length = 370	114	347		HE2OI55
57	792139		32	334	83	H6EEC65
58	793987	steroidogenic acute regulatory protein [Mus musculus]	100	564		HCIAEI8
59	805715	>sp A55455 A55455 steroidogenic acute regulatory protein precursor, mitochondrial - mouse Length = 284	513	1226		HDPKI64
60	811111		1	438		HCEDF72
61	811113		2	718	28	HWBEX78
62	823902	(AF028722) fetal globin inducing factor [Mus musculus]	36	497	87	HDTBD43
63	826518	>sp G4103857 G4103857 FETAL GLOBIN INDUCING FACTOR. Length = 238				
		RNase 4 [Homo sapiens]	1	231	100	HLQCQ62
		>pir I52489 I52489 ribonuclease 4 (EC 3.1.-.-) precursor - human Length = 147				
64	826704	(AB013456) aquaporin 8 [Homo sapiens]	475	726		HCQB118
65	827720	>gnl PID d1035202 (AB013456) aquaporin 8 [Homo sapiens]	789	1076		HFICY86
66	828102	>sp D1035202 D1035202 AQUAPORIN 8. Length = 261	106	297		HSRFC02
67	828180		20	883	83	IIWLFM26

68	828386	(AF093821) RRM RNA binding protein GRY-RBP [Mus musculus] >sp O8899 O88991 RRM RNA BINDING PROTEIN GRY-RBP. Length = 625	gij3694986	3	650	100	100	100	110HAD26
69	828658	protein-tyrosine-phosphatase [Homo sapiens] >gnl PID d1032930 (AB013601) DUSP6 [Homo sapiens] >gnl PID d1035350 (AB013382) DUSP6 [Homo sapiens] >gnl PID d1032930 (AB013601) DUSP6 [Homo sapiens] >sp Q16828 DUS6_HUMAN DUAL SPECIFICITY PROTEIN PHOSPHATASE 6	gnl PID e218263	2	568	100	100	100	HLHCO24
70	828919	RNA helicase [Homo sapiens] >pir S71758 S71758 DEAD box protein MrDb, Myc-regulated - human >sp Q92732 Q92732 RNA HELICASE. Length = 610	gnl PID e254454	2	661	99	100	100	HFOYL30
71	829572	similar to Glyoxalase [Caenorhabditis elegans] Length = 281		163	411				HSVAK51
72	830138	UbcHSB [Homo sapiens] >gij 595668 ubiquitin conjugating enzyme [Rattus norvegicus] >gij 1480742 ubiquitin conjugating enzyme [Mus musculus] >pir S53359 S53359 ubiquitin conjugating enzyme (E217kB) - rat Length = 147	gnl PID e1344082	134	475	53	67	67	HYAAH90
73	830208		gij 145689	2	205	92	95	95	HIBCN46
74	830248	A33 antigen precursor [Homo sapiens] >sp Q99795 A33_HUMAN CELL SURFACE A33 ANTIGEN	gij 1814277	3	1097	30	39	39	HWLHJ13

PRECURSOR. Length = 319

75	830275	Similar to D.melanogaster parallel sister chromatids protein [Homo sapiens] >sp Q92549 Q92549 MYELOBLAST KIAA0261 (FRAGMENT). Length = 1287	gnl PID d1014081	3	647	100	100	HWLFO28
76	830286	interferon-related putative protein [Homo sapiens] >sp Q12894 Q12894 HYPOTHETICAL_48.0 KD PROTEIN. >gij 209022 interferon-related putative protein [Homo sapiens] {SUB 2-442} Length = 442	gij 2880033	385	1488	91	91	HWLFE46
77	830347	(AF039401) calcium-dependent chloride channel-1 [Homo sapiens] >sp G4009460 G4009460 CALCIUM-DEPENDENT CHLORIDE CHANNEL-1. Length = 914	gij 4009460	3	656	63	76	HWLEL81
78	830348	inorganic pyrophosphatase (EC 3.6.1.1) - bovine >sp P37980 PYR_BOVIN	3	911				HWLQR45
79	830364	INORGANIC PYROPHOSPHATASE (EC 3.6.1.1) (PYROPHOSPHATE PHOSPHO- HYDROLASE) (PPASE). Length = 289	pir A45153 A45153 3	3	1022	67	85	HWLEI47
80	830394			1	951			HDPVF62
81	830398			526	627			HWBCR84

82	830412	SDF2 [Homo sapiens] >pir JC5106 JC5106 stromal cell-derived factor 2 - human >sp Q99470 Q99470 SDF2. Length = 211	gnl PID d1009953	233	928	91	92	HWHHQ57
83	830436	(AJ005821) X-like 1 protein [Homo sapiens] >sp E1291794 E1291794 X- LIKE 1 PROTEIN. Length = 3027	gnl PID e1291794	83	523	65	78	HWABR83
84	830464	CLP36 [Rattus norvegicus] >pir JC4385 JC4385 LIM protein - rat >sp P52944 CL36_RAT LIM PROTEIN CLP36. Length = 327	gil1020151	2	289	72	81	HUSGB72
85	830471	(AF011794) cell cycle progression restoration 8 protein [Homo sapiens] >sp O14712 O14712 CELL CYCLE PROGRESSION RESTORATION 8 PROTEIN. Length = 375	gil2352906	95	229			HUSIK51
86	830477			116	2389	95	96	HULAT84
87	830500	ORF YGR036c [Saccharomyces cerevisiae] >pir S64327 S64327 probable membrane protein YGR036c - yeast (Saccharomyces cerevisiae) Length = 239	gnl PID e243385	185	736	38	54	HJPCP29
88	830509	(AL021813) phenylalanyl-tRNA synthetase alpha chain [Schizosaccharomyces pombe] >sp O42849 O42849 PHENYLALANYL-tRNA SYNTHETASE ALPHA CHAIN. Length = 589	gnl PID e1250585	2	1081	40	63	HUFAU68

89	830528	hepatoma-derived growth factor [Mus musculus] >pir JC5662 JC5662 hepatoma-derived growth factor-related protein 2 - mouse >sp O35540 O35540 HEPATOMA-DERIVED GROWTH FACTOR, RELATED PROTEIN 2 Length = 669	dbj ID63850_1	38	1591	78	87	HUFBF32
90	830542	mitochondrial 3-oxoacyl-CoA thiolase [Homo sapiens] >pir S43440 S43440 3-oxoacyl-CoA thiolase - human Length = 397	gnl PID d1004316	324	1637	92	92	HTTDO45
91	830564	IgM heavy chain VH1 region precursor [Homo sapiens] Length = 146		702	1343			HTPBU79
92	830611		gi 2344934	1	495	78	79	HTJMB28
93	830618			655	915			HDIM121
94	830620			452	754			HTGDM95
95	830630	mitochondrial benzodiazepine receptor [Homo sapiens] >pir 38724 38724 mitochondrial benzodiazepine receptor - human >gi 3411163 (AF075589) peripheral-type benzodiazepine receptor [Homo sapiens] {SUB 27-169} Length = 169	gi 529946	14	259	100	100	HTGFS43
96	830654	RNA-binding protein [Saccharomyces cerevisiae] Length = 497	gi 295631	2	1687	40	51	HSYBQ96
97	830660			122	694			HSYDW13
98	830661			555	779			HSXDG80
99	830704	pp21 [Homo sapiens] >pir I53785 I53785 gene pp21 protein - human >sp Q15170 Q15170 (PP21). Length = 157	gi 521207	1	609	51	76	HSUSF13
100	830765			39	236			HSKES11

101	830778	methionine aminopeptidase [Homo sapiens] >gil687243 eIF-2-associated p67 homolog [Homo sapiens] >pirS52112 DPHUM2 methionyl aminopeptidase (EC 3.4.11.18) 2 - human >sp P50579 AMP2_HUMAN METHIONINE AMINOPEPTIDASE 2 (EC 3.4.11.18) (METAP 2) (PEPTIDASE M 2)	gil903982	26	718	99	100	HSPAX18
102	830784			595	858			HSIFY77
103	830800	(AF039918) CD39L4 [Homo sapiens] >sp O75356 O75356 CD39L4. Length = 428	gil3335102	1	990	94	94	HHPDD94
104	830821			449	754			HAQND53
105	830849			464	868			HHHEAA48
106	830903			1	525			HPJCT75
107	830913	tumor necrosis factor type I receptor associated protein [Homo sapiens] >pir A55877 A55877 tumor necrosis factor type I receptor associated protein TRAP-1 - human microsomal glutathione S-transferase 2 [Homo sapiens] >sp Q99735 GST2_HUMAN MICROSOMAL GLUTATHIONE S-TRANSFERASE II (EC 2.5.1.18) (MICROSOMAL GST- II). Length = 147	gil687237	3	1193	99	99	HPIBH48
108	830920	peroxisome proliferator activated receptor gamma 2 [Homo sapiens] >gil171117 ligand activated transcription factor PPARgamma2 [Homo sapiens]	gil1747521	90	650	87	87	HIPHA84
109	830938		gil1432177	227	610	98	98	HONAE45

110	830980	beta COP [Rattus norvegicus] >pir S13520 S13520 beta-COP protein - rat >sp P23514 COPB_RAT COATOMER BETA SUBUNIT (BETA- COAT PROTEIN) (BETA-COP). >pir S13636 S13636 110K protein - rabbit {SUB 451-500} Length = 953 (AF016687) similar to alpha-actinin [Caenorhabditis elegans] >sp O16785 O16785 T21D12.4 PROTEIN. Length = 375	gil55819	47	289	95	98	HCEG53
111	831014		gil2315828	310	1188	53	73	HOEBV08
112	831026			340	687			HOBAE30
113	831031			526	765			HTXOK56
114	831055	(AF091395) Trio isoform [Homo sapiens] >sp O75962 O75962 TRIO ISOFORM. Length = 3038	gil3644048	674	1921	93	94	HNTAT24
115	831057			3	1106			HNTCW73
116	831062			3	821			HNTBD04
117	831117			400	579			HMWBR70
118	831122	cell surface glycoprotein [Homo sapiens] >gil567110 [Human CD79b/Ig beta/B29 gene, complete coding sequence.]. gene product [Homo sapiens] >bbs 122035 membrane immunoglobulin beta chain, Ig-beta=Ag receptor complex [human, B cells, Peptide, 229 aa] [Homo	gil179312	2	772	91	92	HMWCV70
119	831125			868	1023			HMWFH12
120	831132			36	185			HMUAR55

121	831152	(AC004668) similar to murine cell cycle regulator MIDA1; similar to A57591 (PID:g2137417) [Homo sapiens] >sp O60414 O60414 WUGSC:H_RG276O03.1A PROTEIN (FRAGMENT). Length = 635 (AF030109) regulator of G protein signaling 12 [Homo sapiens] >gj 2766633 (AF030152) regulator of G protein signaling 12 [Homo sapiens] Length = 799	gj 3115346	111	875	90	91	HMVA157
122	831157		gj 2605780	664	1110	100	100	HMVA24
123	831160	ezrin (AA 1-586) [Homo sapiens] >pir A34400 A34400 ezrin - human >sp P15311 EZRL_HUMAN EZRIN (P81) (CYTOVILLIN) (VILLIN-2). {SUB 2-586} >gj 340217 cyto villin 2 [Homo sapiens] {SUB 12-586} Length = 586	gj 31283	3	1907	100	100	HCRPE60
124	831193			256	378			IIM1AG77
125	831197			884	1267			HMELO02
126	831217			152	427			HITAAN07
127	831239			420	638			HAKBB67
128	831248			84	443			HCFLLO8
129	831313	c-fos protein [Homo sapiens] >gj 29904 c-fos gene product [Homo sapiens] >gj 4063509 (AF111167) cfos [Homo sapiens] >pir A01342 TVHUF1 transforming protein fos - human >sp P01100 FOS_HUMAN P55-C-FOS PROTO-ONCOGENE PROTEIN (G0S7 PROTEIN). >sp G4063509 G406	gj 182735	1182	1670	83	88	HAGDZ30
130	831369			31	1464			HDFQB94

131	831371	cytochrome P450j [Homo sapiens] >gil181356 cytochrome P450IIE1 [Homo sapiens] >pir A31949 A31949 cytochrome P450 2E1 - human >sp P05181 CPE1_HUMAN CYTOCHROME P450 2E1 (EC 1.14.14.1) (CYP1IIE1) (P450-J). >gnl PID d1001366 cytochrome P450IIE1 [Homo sapiens] hydroxymethylglutaryl-CoA synthase [Homo sapiens] >gil2463646 3-hydroxy-3-methylglutaryl CoA synthase [Homo sapiens] >pir S71623 S71623 hydroxymethylglutaryl-CoA synthase (EC 4.1.3.5) precursor, mitochondrial - human >sp P54868 HMCN_HUMAN HYDROXYMETHYLGLU	gil181360	81	344	94	94	94	HLADA28 HWADP47
132	831373			221	1744				
133	831387	mucin 2 precursor, intestinal - human (fragments) >gil186396 mucin [Homo sapiens] {SUB 626-1895} >gil186398 MUC2 [Homo sapiens] {SUB 2037-3020} >gil188874 intestinal mucin [Homo sapiens] {SUB 1916-2193} >gil188615 mucin-like protein [Homo sapiens] {SUB 23 calcium-modulated protein S100-beta [artificial sequence] >pir A91254 BCBO1B S-100 protein beta chain - bovine {SUB 2-92} Length = 92	pir A49963 A4393 2	2	727	95	96	96	HCQDM23
134	831410								
135	831448		gil554574	126	482	32	60		HKACO81
136	831450			807	1319				HKABK55

137	831472	(AF020043) chromosome-associated	gi 3089368	1	138		HJMBH59
138	831473	polypeptide [Homo sapiens] >sp O60464 O60464 CHROMOSOME-ASSOCIATED POLYPEPTIDE (BAMACAN PROTEIN). >gnl PID e1285055 (AJ005015) bamacan protein [Homo sapiens] {SUB 827-1217} Length = 1217		40	3765	89	HKACE68
139	831474	excision repair protein [Homo sapiens]		1231	1746		HWHPX60
140	831494	>gi 182174 excision repair protein [Homo sapiens] >gi 2583146 (AF001925) excision repair protein [Homo sapiens] >pir A32875 A24781 excision repair protein - human	gi 182177	2	616		HISES08
141	831506	>sp P07992 ERC1_HUMAN DNA EXCISION REPAIR PROTEIN ERC similar to yeast adenylate cyclase (S56776) [Homo sapiens] >sp Q92627 Q92627 MYELOBLAST KIAA0231 (FRAGMENT). Length = 476		3	596	100	HIICAF79
142	831533	growth and transformation dependent protein [Rattus norvegicus] >pir A26882 A26882 p11.2 hypothetical protein - rat (fragment) >sp Q63571 Q63571 RAT GROWTH AND TRANSFORMATION-DEPENDENT (FRAGMENT). Length = 175	gnl PID d1013909	1	900	51	HCRPH87
143	831539		gi 207250	102	572	81	HDTIT02
144	831556			395	625		HDTLJ87

145	831594	protein serine/threonine kinase [Homo sapiens] >gi 468789 CDK activating kinase [Homo sapiens] >gi 485909 MO15/CDK-activating kinase (CAK) [Homo sapiens] >gnl PID e257806 Cdk-activating kinase [Homo sapiens] >pir A54820 A54820 CDK-activating protein kinase	gi 348243	117	677			HHECU01
146	831598			23	802	99	99	HHEDO14
147	831608	translational initiation factor beta subunit [Homo sapiens] >pir A31226 A31226 translation initiation factor eIF-2 beta chain - human >pir S13147 S13147 protein synthesis factor - rabbit >sp P20042 F2B_HUMAN EUKARYOTIC TRANSLATION INITIATION FACTOR 2 BET	gi 182067	120	1154	87	87	HHEFB46
148	831613	Human giant larvae homologue [Homo sapiens] >pir S55474 S55474 Human giant larvae homolog - human >sp Q14521 Q14521 GIANT LARVAE HOMOLOGUE. Length = 1015	gi 854124	3	104	96	100	HISAU33
149	831622	alpha l-acid glycoprotein [Homo sapiens] >gi 1340138 alpha l-acid glycoprotein [Homo sapiens] {SUB 39-86} Length = 201	gnl PID e222211	46	690	99	99	HGBI1256

150	831631	aldose reductase-like peptide [Homo sapiens] >sp O60218 O60218 ALDOSE REDUCTASE-LIKE PEPTIDE (ALDOSE REDUCTASE-RELATED PROTEIN). >gij3098514 (AF044961) aldose reductase-related protein [Homo sapiens] {SUB 232-316; Length = 316	gij3150035	100	1173	100	100	HGBAX75
151	831632	lambda-crystallin precursor [Oryctolagus cuniculus] >pir A31992 A31992 lambda-crystallin - rabbit		2	226			HGBCC19
152	831653	>sp P14755 CRYL_RABIT LAMBDA-CRYSTALLIN. {SUB 2-320} Length = 320	gij164905	172	927	85	90	HTJN173
153	831655	weak similarity to TPR domains [Caenorhabditis elegans] >sp Q23049 Q23049 SIMILARITY TO TPR DOMAINS. Length = 458	gij1465826	3	662	32	54	HFVHF47
154	831708	vascular endothelial growth factor [Homo sapiens] >sp Q16889 Q16889 VASCULAR ENDOTHELIAL GROWTH FACTOR (FRAGMENT). >pir A41551 A41551 vascular endothelial growth factor 206 precursor - human {SUB 23-254} >bbs 85194 vascular endothelial growth factor; VEGF	gij3712671	96	410	98	100	HFIUT25
155	831738			313	573			HFCAL79

156	831741	myelodysplasia/myeloid leukemia factor 2 [Homo sapiens] >gi13387897 (AF070539) myelodysplasia/myeloid leukemia factor 2 [Homo sapiens] >sp Q15773 Q15773	gi1399745	186	974	77	77	HEBT03
		MYELODYSPLASIA/MYELOID LEUKEMIA FACTOR 2. Length = 248						
157	831754	multidrug resistance protein 3 [Homo sapiens] >gnl PID e1288198 multidrug resistance protein 3 [Homo sapiens] >sp O60922 O60922 MULTIDRUG RESISTANCE PROTEIN 3. Length = 1526	gnl PID e1288198	1	924	92	92	HWMEZ67
158	831760			373	510			HETEH76
159	831780			2	1003			HELGH58
160	831796			892	1158			HE9RY54
161	831800	nuclear protein SA-2 [Homo sapiens] >sp O00540 O00540 NUCLEAR PROTEIN SA-2. Length = 1162	gnl PID e250094	600	1541	93	93	HFIAU59
162	831807			1015	1341			HE9QD17
163	831812			520	765			HE9OY91
164	831813			83	793			HEAIIA84
165	831830	isoleucyl-tRNA synthetase [Homo sapiens] >pir 59314 59314 isoleucine--tRNA ligase (EC 6.1.1.5) - human Length = 1266	gnl PID d1006382	52	2307	98	99	HE8TV13
166	831860	Similarity to S. Pombe BEM1/BUD5 suppressor;	gnl PID e1347870	465	776	69	84	HE8OT93
167	831872			1	1671			HE8CL14
168	831896			1	2121			HDTDX05

169	831928	(AF061795) dynamin-like protein Dymple isoform [Homo sapiens] >sp O60709 O60709 DYNAMIN-LIKE PROTEIN DYMPLE ISOFORM. Length = 699	gi 3126874	2	778	77	77	HSYBO86
170	831949	carbonic anhydrase II [Homo sapiens] >gi 179780 carbonic anhydrase II [Homo sapiens] >gi 179795 carbonic anhydrase II [Homo sapiens] >gi 29587 carbonic anhydrase II (AA 1-260) [Homo sapiens]	gi 179772	3	1109			HE8TX12
171	831950			48	521			HAPQS51
172	831953			106	987	100	100	H'WLHA60
173	831975	human phosphotyrosine phosphatase kappa [Homo sapiens] Length = 1439	gn P1D e234080	555	761			HDTBO06
174	832036			2	490	82	82	HCYAC13
175	832047			877	1137			HCWKS85
176	832078	Ca2+ ATPase of fast-twitch skeletal muscle sarcoplasmic reticulum, adult isoform [Homo sapiens] >sp O14983 O14983 CA2+ ATPASE OF FAST-TWITCH SKELETAL MUSCLE SARCOPLASMIC RETICULUM, ADULT ISOFORM. Length = 1001	gi 2052522	751	1014			HASAB14
177	832100			687	917			HCRNM09
178	832104			95	220			HCRMU71
179	832268			18	191			HTXOU56
180	832270			622	1290	90	91	HBKDW03

181	832279	acetyl-CoA synthetase [Drosophila melanogaster] >pir[S52154]S52154 acetyl-CoA synthetase - fruit fly (Drosophila melanogaster) >sp Q24226 Q24226 ACETYL- COENZYME A SYNTHETASE (EC 6.2.1.1) (ACETATE--COA LIGASE) (ACYL-ACTIVATING ENZYME). Length = 581	gi 608694	2	1237	65	77	HIBKDN33
182	832317	11kD protein [Homo sapiens] Length = 111	gi 897917	270	719	100	100	HBIAX17
183	832354			1	408			HBBB52
184	832364			3	1385			HDPQA93
185	832378	sialidase [Homo sapiens] >gi 2773339 (AF040958) lysosomal neuraminidase precursor [Homo sapiens] >gi 4099141 lysosomal sialidase [Homo sapiens] >sp Q99519 Q99519 SIALIDASE PRECURSOR. >sp G4099141 G4099141 LYSOSOMAL SIALIDASE PRECURSOR (EC 3.2.1.18). Lengt (AF048700) gastrointestinal peptide [Homo sapiens] >sp O60575 O60575 GASTROINTESTINAL PEPTIDE. Length = 86	gn PID e303801	3	746	96	96	HATCO72
186	832385		gi 2935440	2	316	90	90	HARAG42
187	832428	APO-1 ANTIGEN, FAS ANTIGEN. Length = 335	sp G249613 G249613	136	846	97	97	HAMGD53
188	832485			202	597			HAGHC54

189	832494	Ku protein subunit [Homo sapiens] >gi 78650 p70 autoantigen [Homo sapiens] >gi 339667 thyroid autoantigen [Homo sapiens] >bbs 107206 Ku autoantigen p70 subunit [human, Peptide, 609 aa] [Homo sapiens] >pir A30299 A30894 70K thyroid autoantigen - human >sp	gi 307095	80	1918	90	90	90	HAIBY70
190	832512	Similar to Human C219-reactive peptide (L34688) [Homo sapiens] >sp Q92580 Q92580 MYELOBLAST KIA0268 (FRAGMENT). >gi 511639 C219-reactive peptide [Homo sapiens] {SUB 592-727; Length = 1193 integrin alpha6 subunit [Homo sapiens] Length = 1067 nuclear factor RIP140 [Homo sapiens] >pir S57348 S57348 nuclear factor RIP140 - human Length = 1158 protein tyrosine kinase [Homo sapiens] >pir A55922 A55922 tyrosine kinase A6 - human >sp Q12792 Q12792 PROTEIN TYROSINE KINASE. Length = 350	gnl PID d1014138	3	1058	87	87	87	HDPTT16
191	832515		gi 33942	2	1660	96	96	96	HCRPH70
192	832526		gi 940539	34	693	95	95	95	HADCX04
193	832575		gi 451482	49	1203	99	99	99	H2LAJ21
194	832576	BTG1 gene product [Homo sapiens] >gi 293306 BTG1 [Mus musculus] >gi 50188 big1 [Mus musculus] >pir S20947 S20947 BTG1 protein - human >pir 48272 48272 big1 protein - mouse >sp P31607 BTG1_HUMAN BTG1 PROTEIN (B-CELL TRANSLOCATION GENE 1 PROTEIN). Length	gi 29509	388	1050	100	100	100	HKGAJ67

195	832588	mitochondrial ATP synthase subunit 9 precursor [Homo sapiens] >pir 38612 38612 ATP synthase chain 9 precursor, mitochondrial - human >sp P48201 A193_HUMAN ATP SYNTHASE LIPID-BINDING PROTEIN P3 PRECURSOR (EC 3.6.1.34) (ATPASE PROTEIN 9) (SUBUNIT C). Leng	gi 511450	2	637	85	85	H2LAD51
196	832634			253	924			HCRMZ25
197	832728			3	542			HKAIL83
198	833094	immunoglobulin from VH4 family [Homo sapiens] >pir S13519 S13519 Ig heavy chain V region precursor - human >gi 553385 immunoglobulin heavy chain [Homo sapiens] {SUB 24-125} Length = 147	gi 37725	2	391	77	81	HRADC46
199	833395	novel stromal cell protein [Mus musculus] >pir JC4761 JC4761 recombination activating gene 1 inducing protein - human >sp Q62275 Q62275 RECOMBINATION ACTIVATING PROTEIN 1 PROTEIN ACTIVATION (NOVEL STROMAL CELL PROTEIN). Length = 221	gnl PID e229590	224	853			HHENV68
200	834326			1	744	69	76	HWLEQ41
201	834583	(AF073957) CXCL chemokine BRAX [Homo sapiens] Length = 99	gi 4140394	2	607	98	100	HHGDE66
202	834944	(AF061443) G protein-coupled receptor LGR4 [Rattus norvegicus] >sp G3885470 G3885470 G PROTEIN-COUPLED RECEPTOR 1.GR4. Length	gi 3885470	2	781	85	86	HE8QE56

= 951

203	835012	(AB017169) Slit-3 protein [Homo sapiens] >sp D1036172 D1036172 SLIT-3 PROTEIN. >gnl PID d1033429	gnl PID d1036172	3	344	92	92	HCCMD55
204	835104	(AB011538) MEGF5 [Homo sapiens] {SUB 785-1523} Length = 1523		580	1818			HLHTJ57
205	835332	(AF065389) tetraspan NET-4 [Homo sapiens] >sp O60746 O60746 TETRASPAN NET-4. Length = 268	gi 3152703	268	1080	100	100	HCROP84
206	835487	(AC002528) alpha2(I) collagen [Homo sapiens] >sp G238855 G238855 ALPHA2(I) COLLAGEN (FRAGMENT). Length = 1186	gi 2388555	2218	4239	100	100	HTSGZ29
207	836182			39	398			HFLUE31
208	836522			1819	2046			HSLFO17
209	836655			1	624			HTPCU04
210	836787	calmodulin-dependent protein kinase II-delta (EC 2.7.1.37) [Rattus norvegicus] >pir A34366 A34366 Ca2+/calmodulin-dependent protein kinase (EC 2.7.1.123) II delta chain - rat	gi 203267	767	1549	92	94	HAIED73
		>sp I5791 KCCD_RAT CALCIUM/CALMODULIN-DEPENDENT PROTEIN KINASE TYPE II DELTA CII						

211	836789	GP36b glycoprotein [Homo sapiens] >pir G01447 G01447 GP36b glycoprotein - human >sp Q12907 Q12907 GP36B GLYCOPROTEIN PRECURSOR. Length = 356	gij 505652	1	849	99	99	HKAAD74
212	838577	binding protein [Oryctolagus cuniculus] >gij 82628 FK506-binding protein (FKBP) [Homo sapiens] >gij 82633 FKBP-12 protein [Homo sapiens] >gij 82649 FK506-binding protein 12 [Homo sapiens] >gij 288196 FKBP [Homo sapiens] >gij 665650 FK-506 binding protein [H	gij 65023	2	433	100	100	HCRQD09
213	838717			676	900			HE8UJ03
214	839008			2	997			HFOXSS2
215	840063	(AF006751) ES/130 [Homo sapiens] >sp O75300 O75300 ES/130. Length = 977	gij 3299885	3	2729	84	85	HWLHIX68
216	840533			183	482			HWLLU74
217	840669			474	1115			HPMGM71
218	841140	(AF081281) lysophospholipase [Homo sapiens] >sp O75608 O75608 LYSOPHOSPHOLIPASE. Length = 230 polypeptide GalNAc transferase-T4 [Mus musculus] >sp O08832 O08832 POLYPEPTIDE GALNAc TRANSFERASE-T4. Length = 578	gij 3415123	1	789	100	100	HAJCC51
219	841386		gij 2121220	491	1258	66	81	HMCCA66
220	841480			3	212			HDQET68
221	841509			3	662			HTELO87
222	841616			340	660			HWLFT95

223	841900	peptidylarginine deiminase (EC 3.5.3.15) [Rattus norvegicus] >pir A34339 DIRTR1 protein-arginine deiminase (EC 3.5.3.15) I - rat >sp P20717 PARD_RAT PROTEIN- ARGININE DEIMINASE (EC 3.5.3.15) (PEPTIDYLARGININE DEIMINASE). Length = 665	gij205960	2	439	87	90	HWLFR87
224	842054	ubiquinone-binding protein (QP) [Homo sapiens] >gij I90816 ubiquinone-binding protein precursor [Homo sapiens] >gij 37580 ubiquinone-binding protein (AA 1 - 111) [Homo sapiens] >pir A32450 A32450 ubiquinone-binding protein QP-C - human >gij 553631 ubiquinone (AB012933) acyl-CoA synthetase 5 [Rattus norvegicus] >sp O88813 LCFE_RAT LONG- CHAIN-FATTY-ACID--COA LIGASE 5 (EC 6.2.1.3) (LONG-CHAIN ACYL- COA SYNTHETASE 5) (LACS 5). Length = 683	gij I90802	1	369	100	100	HWHPF06
225	843061	(AB012933) acyl-CoA synthetase 5 [Rattus norvegicus] >sp O88813 LCFE_RAT LONG- CHAIN-FATTY-ACID--COA LIGASE 5 (EC 6.2.1.3) (LONG-CHAIN ACYL- COA SYNTHETASE 5) (LACS 5). Length = 683	gn PIPI d1034547	23	2308	81	92	HDAAV92
226	843544	(AF045573) FLI-LRR associated protein- I [Mus musculus] >sp O70323 O70323 FLIGHTLESS-I ASSOCIATED PROTEIN 1 (LRR DOMAIN) (FLI-LRR ASSOCIATED PROTEIN-1). Length = 628	gij3025718	28	391 837	65	83	HFLNB80 HTEKO43

228	844270	nuclear antigen EBNA-3B [Human herpesvirus 4] >pir[S2792]S27921 nuclear antigen EBNA-3B - human herpesvirus 4 >sp Q69139 Q69139 NUCLEAR ANTIGEN EBNA-3B. Length = 946	gi 330409	2	373	47	52	HWLBL06
229	844604	(AF071186) WW domain binding protein 11 [Mus musculus] >sp O88539 O88539 WW DOMAIN BINDING PROTEIN 11. Length = 389	gi 3550082	170	2110	66	70	HNTAD40
230	844685	immunoglobulin lambda heavy chain [Homo sapiens] >gi 567132 This CDS feature is included to show the translation of the corresponding C_region. Presently translation qualifiers on C_region features are illegal [Homo sapiens] (SUB 148-177) Length = 477	gnl PID e1227585	539	1564	94	94	HASAC08
231	844855	titin [Oryctolagus cuniculus] >sp E135530 E1355301 TITIN (FRAGMENT). Length = 2000	gnl PID e1355301	3	1634	34	54	HAICQ70
232	845101	(AF089814) growth suppressor related [Homo sapiens] >sp O75956 O75956 GROWTH SUPPRESSOR RELATED. Length = 126	gi 3661529	46	627	94	94	HHESZ77
233	845141	(AB011105) KIAA0533 protein [Homo sapiens] >sp O15230 O15230 KIAA0533 PROTEIN (LAMININ ALPHA 5 CHAIN) (FRAGMENT).	gnl PID d1026389	31	966	100	100	HWMF067
234	845220	>gnl PID e317479 laminin alpha 5 chain [Homo sapiens] (SUB 693-1645) Length = 1645	2	1096				HKADF64

235	845434	glutathione peroxidase [Synecocystis sp.] >pir[S75885 S75885 glutathione peroxidase homolog - Synecocystis sp. (PCC 6803)] >sp P74250 P74250 GLUTATHIONE PEROXIDASE (EC 1.11.1.9). Length = 169	gnl PID d1019077	3	590	50	61	HWAF112
236	845510	dipeptidyl peptidase III [Rattus norvegicus] >sp O55096 O55096 DIPEPTIDYL PEPTIDASE (EC 3.4.14.4) (DIPEPTIDYL PEPTIDASE III) (DIPEPTIDYL AMINOPEPTIDASE III) (DIPEPTIDYL ARYLAMIDASE III) (RED CELL ANGIOTENSINASE) (ENKEPHALINASE B). Length = 827	gnl PID d1025528	3	683	96	98	HEONN92
237	845600	procathepsin B [Homo sapiens] >pir A26498 K11HUB cathepsin B (EC 3.4.22.1) precursor - human >sp P07858 CATB_HUMAN CATHEPSIN B PRECURSOR (EC 3.4.22.1) (CATHEPSIN B1) (APP SECRETASE). >gij 181178 lysosomal proteinase cathepsin B [Homo sapiens] {SUB 131-33 (AF055666) kincsin light chain 2 [Mus musculus] >sp O88448 O88448 KINESIN LIGHT CHAIN 2. Length = 599	gij 181192	223	1254	99	99	HOEMIE38
238	845882		gij 347848	4	1155	68	75	HL.HCE82

239	846007	alpha-1-acid glycoprotein 2 [Homo sapiens] >pir JT0326[OMHU2 alpha-1-acid glycoprotein 2 precursor - human >sp P19652 A1A11_HUMAN ALPHA-1-ACID GLYCOPROTEIN 2 PRECURSOR (AGP 2) (OROSOMUCOID 2) (OMD 2). >gi 388511 alpha 1-acid glycoprotein [Homo sapiens] (SU	gi 177840	1	390	98	100	HILDBS16
240	846280	epididymal apical protein 1-precursor [Macaca fascicularis]		31	105			HCNAK57
241	846286	>pir S28258 S28258 androgen-regulated epididymal protein precursor - crab-eating macaque >sp Q28475 Q28475 EPIDIDYMAL APICAL PROTEIN 1- PRECURSOR. Length = 776	gi 38063	203	901	36	54	HASDA19
242	846388			3	1721			HIL3AA32
243	HCRNG17R			154	288			HCRNG17
244	HWMFG64R			1	315			HWMFG64
245	HAGCZ94R			13	102			HAGCZ94
246	HBJEJ74R			72	287			HBJEJ74
247	HUJBE67R			355	525			HUJBE67
248	HUTHM43R			2	55			HUTHM43
249	HLTGU75R			2	274			HLTGU75
250	HWLKF77R			51	134			HWLKF77
251	HWLLK67R			1	180			HWLLK67
252	HDQIE85R			3	203			HDQIE85
253	HWLFA67R			1	213			HWLFA67
254	HWLGX29R			136	351			HWLGX29
255	HWMFZ29R			324	404			HWMFZ29

256	HNTRR03R	(AB018797) calmodulin B [Haloecynthia roretzi] >sp D1034943 D1034943	1	363	HNTRR03
257	H6EEP19R	CALMODULIN B. Length = 149	2	103	H6EEP19
258	HJMAM83R	(AC005031) neuronal apoptosis inhibitory protein [Homo sapiens]	2	352	HJMAM83
259	HAGHF58R	>sp O75857 O75857 NEURONAL APOPTOSIS INHIBITORY PROTEIN (FRAGMENT). Length = 1178	1	88	HAGHF58
260	HDPHG48R	(AC005154) similar to protein U28928 (PID:g861306) [Homo sapiens]	1	98	HDPHG48
261	HWLUL19R	>sp O75223 O75223 WUGSC:H_DJ0777O23.1 PROTEIN. Length = 188	2	59	HWLUL19
262	HWLLI56R	(AD000684) liver-specific bHLH-Zip transcription factor [Homo sapiens]	1	61	HWLLI56
263	HWMAA87R	>sp O00112 O00112 LIVER-SPECIFIC bHLH-ZIP TRANSCRIPTION FACTOR (FRAGMENT). Length = 429	3	86	HWMAA87
264	HGLAT96R	(AF001904) 3-hydroxyacyl-CoA dehydrogenase isoform 2 [Homo sapiens]	165	91	HGLAT96

265	HCDMC32R	(AF014118) membrane-associated kinase [Homo sapiens] >sp O14731 O14731 MEMBRANE-ASSOCIATED KINASE. Length = 499	gi 2460023	3	272	100	100	HCDMC32
266	HCROF25R	(AF034800) liprin-alpha3 [Homo sapiens] >sp G3309535 G3309535 LIPRIN-ALPHA3 (FRAGMENT). Length = 443	gi 3309535	70	381	60	65	HCROF25
267	HTEQO80R	(AF035840) NADH:ubiquinone oxidoreductase B17 subunit [Homo sapiens] >sp G3800740 G3800740 NADH:UBIQUINONE. OXIDOREDUCTASE B17 SUBUNIT. Length = 128	gi 3800740	1	327	100	100	HTEQO80
268	H2LAU18R	(AF035940) similar to mago nashi [Homo sapiens] >gi 2330011 (AF007862) mm-Mago [Mus musculus] >gi 2909828 (AF035939) similar to mago nashi [Mus musculus] >sp O35169 O35169 MM-MAGO. >sp G2909830 G2909830 MAGOH. >sp P50606 MGN_HUMAN MAGO NASHI PROTEIN HOMOL	gi 2909830	2	592	100	100	H2LAU18
269	HTXPO87R	(AF038129) polyubiquitin [Ovis aries] >sp O46543 O46543 POLYUBIQUITTIN. >gnl PID e1263307 unnamed protein product [unidentified] {SUB 77-305} >gi 163575 polyubiquitin [Bos taurus] {SUB 142-305} >gi 1762374 polyubiquitin [Gallus gallus] {SUB 1-71} >gnl PID	gi 2707837	1	330	97	97	HTXPO87

270	H2LAR08R (AF040642) contains similarity to RNA recognition motifs (RNP) [Caenorhabditis elegans] >sp O44795 O44795 C50D2.5 PROTEIN. Length = 200	gi 2746787	188	514	75	90	H2LAR08
271	HADAF94R (AF044957) NADH:ubiquinone oxidoreductase B15 subunit [Homo sapiens] Length = 129	gi 4164446	88	135	88	88	HADAF94
272	HEMDA91R (AF047473) testis mitotic checkpoint BUB3 [Homo sapiens] >sp O43685 O43685 TESTIS MITOTIC CHECKPOINT BUB3. Length = 326	gi 3378104	132	431	85	85	HEMDA91
273	HWMFN58R (AF051426) slow delayed rectifier channel subunit [Homo sapiens] >sp O60607 O60607 SLOW DELAYED RECTIFIER CHANNEL SUBUNIT. Length = 548	gi 2961249	3	344	100	100	HWMFN58
274	HCNDJ66R (AF054643) lambda 1 immunoglobulin light chain variable region [Homo sapiens] >gi 3023109 (AF054643) lambda 1 immunoglobulin light chain variable region [Homo sapiens] Length = 125	gi 3023109	1	276	72	73	HCNDJ66
275	HOHDH05R (AF061833) aldehyde dehydrogenase; retinal dehydrogenase; class I aldehyde dehydrogenase; ALDH1 [Xenopus laevis] >sp G3818533 G3818533 ALDEHYDE DEHYDROGENASE (EC 1.2.1.3). >pir S1188 S1188 aldehyde dehydrogenase (NAD+) (EC 1.2.1.3), cytosolic - clawed f	gi 3818533	59	331	53	80	HOHDH05

276	HUFBP63R (AF062137) immunoglobulin heavy chain variable region [Homo sapiens] Length = 143	gi 3170737	17	463	92	96	HUFBP63
277	HUFBN90R (AF062211) immunoglobulin heavy chain variable region [Homo sapiens] Length = 149	gi 3170885	26	463	94	96	HUFBN90
278	HEBEJ57R (AF062214) immunoglobulin heavy chain variable region [Homo sapiens] Length = 142	gi 3170895	1	165	81	90	HEBEJ57
279	HDTDK65R (AF069048) immunoglobulin light chain variable region [Homo sapiens] Length = 120	gi 3328006	3	434	76	78	HDTDK65
280	HAIAD82R (AF069711) urokinase [Oryctolagus cuniculus] >sp G3982741 G3982741 UROKINASE (FRAGMENT). Length = 128	gi 3982741	1	156	68	71	HAIAD82
281	HFKHD61R (AF073298) 4F5rel [Homo sapiens] >gi 3641536 (AF073297) 4F5rel [Mus musculus] >sp O75918 O75918 4F5REL. >sp O88891 O88891 4F5REL. Length = 59	gi 3641538	3	203	100	100	HFKHD61
282	H2LAX28R (AF078817) high mobility group protein [Nannospalax ehrenbergi] >sp O88611 O88611 HIGH MOBILITY GROUP PROTEIN. Length = 215	gi 3342571	206	568	97	97	H2LAX28
283	HWLMY93R (AF078839) Rho related protein Rnd3/Rho8 [Sus scrofa] >sp O77683 O77683 RHO RELATED PROTEIN RND3/RHO8. Length = 244	gi 3386532	3	173	91	91	HWLMY93
284	HTXNL13R		3	356			HTXNL13

285	HDPWR89R	(AJ005259) homologous to Bombyx mori multiprotein bridging factor (EMBL: AB001078) [Homo sapiens] >sp O60869 O60869 EDF-1 PROTEIN. Length = 148	gnl PID e 286414	1	312	79	83	HDPWR89
286	H2LAK62R			22	165			H2LAK62
287	HWLKT15R	UBIQUINONE/MENAQUINONE BIOSYNTHESIS METHYLTRANSFERASE UBIE (ubiE) [Rickettsia prowazekii]	gnl PID e 342961	2	301	50	76	HWLKT15
288	HATAR77R	(AL021546) Cytochrome C Oxidase Polypeptide VIa-liver precursor (EC 1.9.3.1) [Homo sapiens]	gnl PID e 248288	3	413	70	73	HATAR77
289	HWLWN07R	(AL022237) bK1191B2.2 (BCL2-interacting killer (apoptosis-inducing) (NBK, BP4, BIP1)) [Homo sapiens] >sp E1359316 E1359316 BK1191B2.2 (BCL2-INTERACTING KILLER (APOPTOSIS-INDUCING) (NBK, BP4, BIP1)) (FRAGMENT). >gi 929655 NBK [Homo sapiens] {SUB 14-173} Le	gnl PID e 359316	1	183	82	88	HWLWN07
290	HWLDI18R	(AL023554) ribosomal protein [Schizosaccharomyces pombe] >sp O60118 O60118 RIBOSOMAL PROTEIN. Length = 157	gnl PID e 292696	3	206	43	59	HWLDI18
291	HWMEC68R			3	419			HWMEC68

292	HTXFO53R	11 beta-hydroxysteroid dehydrogenase type II [Homo sapiens] >pir 38858 38858.1 beta-hydroxysteroid dehydrogenase (EC 1.1.1.146) type 2 - human >sp P80365 DHI2_HUMAN CORTICOSTEROID 11-BETA-DEHYDROGENASE, ISOZYME 2 (EC 1.1.1.146) (11-DH2) (11-BETA-HYDROX	gij565082	3	236	88	94	HTXFO53
293	HWMEH18R	3',5'-cyclic-GMP phosphodiesterase (EC 3.1.4.35) alpha chain - human >gij3513491 (AF022380) rod photoreceptor cGMP phosphodiesterase alpha subunit [Homo sapiens] {SUB 1-122} Length = 859	pir B3461 B3461.1	3	203	92	92	HWMEH18
294	HCWFF03R	5' half of the product is homologues to Bacillus subtilis SAICAR synthetase, 3' half corresponds to the catalytic subunit of AIR carboxylase [Homo sapiens] >pir S14147 S14147 multifunctional purine biosynthesis protein - human Length = 425	gij28384	3	296	83	90	HCWFF03
295	HCNDP66R	A33 antigen precursor [Homo sapiens] >sp Q99795 A33_HUMAN CELL SURFACE A33 ANTIGEN PRECURSOR. Length = 319	gij1814277	3	503	73	75	HCNDP66
296	HCRMK82R	adenosine A2b receptor [Homo sapiens] >gij757911 A2b adenosine receptor [Homo sapiens] >pir JC1229 JC1229 adenosine receptor A2b - human >sp P29275 A2B_HUMAN	gij178150	2	427	100	100	HCRMK82

ADENOSINE A2B RECEPTOR. Length
= 332

297	HCDANI6R	alpha-1 collagen (I) [Gallus gallus] Length = 143	gij555432	2	133	77	88	HCDANI6
298	HCEOE88R	amplaxin [Homo sapiens] >pir A48063 A48063 mammary tumor/squamous cell carcinoma- associated protein EMS1 - human Length = 550	gij182087	1	291	93	94	HCEOE88
299	HALSK30R	angiogenin [Homo sapiens] >pir A90498 NRHUAG angiogenin precursor - human >sp P03950 ANGI_HUMAN ANGIOGENIN PRECURSOR (EC 3.1.27.-). Length = 147	gij178250	189	416	74	76	HALSK30
300	HDRME43R	anonymous [Homo sapiens] >pir 39463 39463 gene anonymous protein - human >sp Q13769 Q13769 ANONYMOUS. Length = 683	gij388012	2	346	94	95	HDRME43
301	HHEFA24R	APP-binding protein 1 [Rattus norvegicus] >sp G4099878 G4099878 APP-BINDING PROTEIN 1. Length = 534	gij4099878	10	177	63	65	HHEFA24
302	HSSGC52R	argininosuccinate synthetase [Bos taurus] >sp P14568 ASSY_BOVIN ARGININOSUCCINATE SYNTHASE (EC 6.3.4.5) (CITRULLINE-- ASPARTATE LIGASE). Length = 412	gij162697	1	438	94	95	HSSGC52

303	HCYBN49R	ATP synthase beta subunit precursor [Homo sapiens] >pir A33370 A33370 H+-transporting ATP synthase (EC 3.6.1.34) beta chain precursor, mitochondrial - human >sp P06576 ATPB_HUMAN ATP SYNTHASE BETA CHAIN. MITOCHONDRIAL PRECURSOR (EC 3.6.1.34). >gil28931 be	gil179281	56	445	97	97	HCYBN49
304	HWMGB90R	ATP synthase subunit e [Homo sapiens] >sp P56385 ATPJ_HUMAN ATP SYNTHASE E CHAIN, MITOCHONDRIAL (EC 3.6.1.34). {SUB 2-69} Length = 69	gil2605592	1	165	58	61	HWMGB90
305	HTEAW21R	ATPase coupling factor 6 subunit [Homo sapiens] >pir JT0563 JT0563 coupling factor 6 precursor, mitochondrial - human >sp P18859 ATPR_HUMAN ATP SYNTHASE COUPLING FACTOR 6, MITOCHONDRIAL PRECURSOR (EC 3.6.1.34) (F6). Length = 108	gil179275	47	259	93	93	HTEAW21
306	HCQCV96R	ATPase subunit 6 [Homo sapiens] >sp Q34772 Q34772 ATP SYNTHASE A CHAIN (EC 3.6.1.34). Length = 226	gnl PID d1007873	147	368	58	61	HCQCV96
307	HLTDN74R	autotaxin-1 [Homo sapiens] >sp Q13822 Q13822 AUTOTAXIN-T. >gnl PID d1008938 phosphodiesterase I alpha [Homo sapiens] {SUB 1-45} Length = 863	gil1160616	2	118	85	85	HLTDN74
308	HDABV61R	B-creatine kinase [Gallus gallus] Length = 65	gil211524	3	230	93	100	HDABV61

309	H2LAQ68R	beta prime cop [Bos taurus] >pir S35312 S35312 coatmer complex beta' chain - bovine >sp P35605 COPP_BOVIN COATOMER BETA' SUBUNIT (BETA'-COAT PROTEIN) (BETA'- COP) (PI02). {SUB 2-906} Length = 906	gil 312732	127	558	100	100	H2LAQ68
310	HDTLN42R	beta-2-microglobulin [Pan troglodytes] >gil 177065 beta-2-microglobulin [Gorilla gorilla] >gnl PID d1036168 (AB021288) beta 2-microglobulin [Homo sapiens] >pir A90976 MGHUB2 beta-2- microglobulin precursor - human >pir 136963 136963 beta-2-microglobulin pre	gil 176827	2	361	86	86	HDTLN42
311	HULFN47R	beta-2-microglobulin [Pan troglodytes] >gil 177065 beta-2-microglobulin [Gorilla gorilla] >gnl PID d1036168 (AB021288) beta 2-microglobulin [Homo sapiens] >pir A90976 MGHUB2 beta-2- microglobulin precursor - human >pir 136963 136963 beta-2-microglobulin pre	gil 176827	3	449	88	89	HULFN47
312	HCRM141R			1	528			HCRM141
313	HWLIP53R			2	499			HWLIP53
314	HBAAD60R			2	463			HBAAD60
315	HCROA35R			3	500			HCROA35
316	HCROM64R			201	512			HCROM64
317	HEOPS84R			2	388			HEOPS84
318	HKBAG82R			32	265			HKBAG82
319	HUTSB76R			188	418			HUTSB76

320	HWLJS67R	384	662	HWLJS67
321	HWLLZ82R	2	133	HWLLZ82
322	HCROM20R	351	557	HCROM20
323	HDQMC24R	1	144	HDQMC24
324	HOCTD89R	212	352	HOCTD89
325	HTGAZ53R	198	341	HTGAZ53
326	HWLKZ47R	429	626	HWLKZ47
327	HWLLLS1R	204	416	HWLLLS1
328	HRLAJ54R	207	548	HRLAJ54
329	HBAAD69R	1	456	HBAAD69
330	HWLJZ72R	25	453	HWLJZ72
331	HWMFG06R	43	321	HWMFG06
332	HPRTO65R	2	166	HPRTO65
	biliary glycoprotein a [Homo sapiens] >gnlPID[d1015047 biliary glycoprotein, BGPg [Homo sapiens] >gi 3172151 (AC004785) BGPg_HUMAN [Homo sapiens] >pir H0394 H0394 biliary glycoprotein g precursor - human Length = 417	gil179438	93	97
333	HUFDC01R	108	326	87
	biliary glycoprotein l precursor [Homo sapiens] >gi 37198 TM1-CEA preprotein [Homo sapiens] >gi 3172148 (AC004785) BGPI_HUMAN [Homo sapiens] >pir A32164 A32164 biliary glycoprotein l precursor - human >sp P13688 BGPI_HUMAN BILIARY GLYCOPROTEIN l PRECURSOR	gil179440	87	87
334	HWLHY44R	3	413	75
	bone-derived growth factor [Homo sapiens] >sp Q13876 Q13876 BONE- DERIVED GROWTH FACTOR (FRAGMENT). Length = 793	gil1203965	79	79
				HWLHY44

335	HWLGR92R	brain glycogen phosphorylase [Homo sapiens] >pir A29949 A29949 glycogen phosphorylase (EC 2.4.1.1), brain (astrocytoma cell line) - human Length = 863	gil307200	122	238	100	100	100	HWLGR92
336	HCNCQ71R	CAG-isl 7 [Homo sapiens] Length = 213	gil3126984	1	93	66	77	77	HCNCQ71
337	HBMCI28R	carbonic anhydrase I (EC 4.2.1.1) [Homo sapiens] >gil29600 carbonic anhydrase I (AA 1-261) [Homo sapiens] >pir JQ0786 CRHU1 carbonate dehydratase (EC 4.2.1.1) I - human >sp P00915 CAH1_HUMAN CARBONIC ANHYDRASE I (EC 4.2.1.1) (CARBONATE DEHYDRATASE I). ;SU	gil179793	81	293	84	84	84	HBMCI28
338	HWLENI1R	carbonic anhydrase I (EC 4.2.1.1) [Homo sapiens] >gil29600 carbonic anhydrase I (AA 1-261) [Homo sapiens] >pir JQ0786 CRHU1 carbonate dehydratase (EC 4.2.1.1) I - human >sp P00915 CAH1_HUMAN CARBONIC ANHYDRASE I (EC 4.2.1.1) (CARBONATE DEHYDRATASE I). ;SU	gil179793	84	347	80	80	80	HWLENI1
339	HMSDU92R	carbonic anhydrase II [Homo sapiens] >gil179780 carbonic anhydrase II [Homo sapiens] >gil179795 carbonic anhydrase II [Homo sapiens] >gil29587 carbonic anhydrase II (AA 1-260) [Homo sapiens] >pir A27175 CRHU2 carbonate dehydratase (EC 4.2.1.1) II - human	gil179772	1	360	76	83	83	HMSDU92

340	HCDBF89R	carbonic anhydrase IV [Homo sapiens] >gil409726 carbonic anhydrase IV [Homo sapiens] [SUB 73-294] Length = 294	gil409725	11	160	87	90	HCDBF89
341	HCNDP16R	carboxylesterase hCE-2 [Homo sapiens] >sp Q16859 Q16859 CARBOXYLESTERASE (EC 3.1.1.1) (AL1-ESTERASE) (B-ESTERASE) (MONOBUTYRASE) (COCAINE ESTERASE) (PROCAINE ESTERASE) (METHYLBUTYRASE). Length = 550	gil407780	1	252	70	71	HCNDP16
342	HWLGX53R	carcinoembryonic antigen [Homo sapiens] >gil178677 carcinoembryonic antigen precursor [Homo sapiens] >pir A36319 A36319 carcinoembryonic antigen precursor - human >sp P06731 CCEM_HUMAN CARCINOEMBRYONIC ANTIGEN PRECURSOR (CEA) (MECONIUM ANTIGEN 100) (CD66E)	gil180223	19	138	73	73	HWLGX53
343	HWLEH56R	carcinoembryonic antigen [Homo sapiens] >gnl PID e24945 carcinoembryonic antigen [Homo sapiens] >gil3702266 (AC005797) carcinoembryonic antigen CGM2 precursor - human [Homo sapiens] >pir A55811 A55811 carcinoembryonic antigen CGM2 precursor - human >sp Q	gil471077	1	453	86	87	HWLEH56

344	H2LAD26R	CaRG box-binding factor [Mus musculus] >gnl PID d1014884 CaRG-binding factor-A [Mus musculus] >pir Q0448 Q0448 CaRG-binding factor-A - mouse >sp Q99020 CABA_MOUSE CaRG-BINDING FACTOR-A (CBF-A). Length = 285	gi 840648	43	387	98	98	H2LAD26
345	HADAF48R	CD99 typell [Homo sapiens] >sp O00518 O00518 CD99 TYPEII. Length = 160	gi 2149135	2	151	59	59	HADAF48
346	HCRNV62R	Cdc6-related protein [Homo sapiens] >gi 2465437 (AF022109) HsCdc18p [Homo sapiens] >sp Q99741 Q99741 CDC6-RELATED PROTEIN. Length = 560	gi 1684903	2	442	90	91	HCRNV62
347	HCDC117R	chaperonin-like protein [Homo sapiens] >pir S48087 S48087 I-complex-type molecular chaperone CCT6 - human >gi 184462 chaperonin-like protein [Homo sapiens] {SUB 143-531; Length = 531	gi 517065	3	137	97	100	HCDC117
348	HJUAA02R	Cks1 protein homologue [Homo sapiens] >pir A36670 A36670 protein kinase cdc2 complex subunit CKS1 - human >sp P33551 CKS1_HUMAN CYCLIN-DEPENDENT KINASES REGULATORY SUBUNIT 1 (CKS-1). Length = 79	gi 29977	186	386	96	96	HJUAA02

349	HKAKO78R	Cks1 protein homologue [Homo sapiens] >pir B36670 B36670 protein kinase cdc2 complex subunit CKS2 - human >sp P33552 CKS2_HUMAN CYCLIN- DEPENDENT KINASES REGULATORY SUBUNIT 2 (CKS-2). Length = 79	gij29979	2	193	77	77	HKAKO78
350	H2CBD02R			58	522			H2CBD02
351	HWLCR90R	contains similarity to ATP/GTP-binding site motif (PS:PS00017) [Caenorhabditis elegans] >sp Q94180 Q94180 SIMILARITY TO ATP/GTP-BINDING SITE MOTIF. Length = 398	gij1519671	1	351	34	60	HWLCR90
352	H2LAK66R	core protein II precursor [Homo sapiens] >pir A32629 A32629 ubiquinol-- cytochrome-c reductase (EC 1.10.2.2) core protein II - human Length = 453	gij180928	126	632	79	79	H2LAK66
353	HSDKC65R	CoxII/D-loop DNA fusion protein [Homo sapiens] >sp Q34777 Q34777 COXII/D- LOOP DNA FUSION PROTEIN (FRAGMENT). Length = 125	gij1374867	179	346	95	97	HSDKC65
354	H2LAK52R	CUL-2 [Homo sapiens] >sp Q13617 CUL2_HUMAN CULLIN HOMOLOG 2 (CUL-2). Length = 745	gij1923243	24	608	100	100	H2LAK52
355	HKAEI2R	cyclin B1 - human >sp P14635 CGB1_HUMAN G2/MITOTIC-SPECIFIC CYCLIN B1. Length = 433	pir A32992 A32992 2	3	392	98	98	HKAEI2
356	HKADP43R	cyclin F [Homo sapiens] >sp P41002 CG2F_HUMAN G2/MITOTIC-SPECIFIC CYCLIN F. Length = 786	gij576781	1	375	71	71	HKADP43

357	HLXND10R	cystatin B [Homo sapiens] >gi 235678 cystatin B [Homo sapiens] >sp P04080 CYTB_HUMAN CYSTATIN B (LIVER THIOL PROTEINASE INHIBITOR) (CPI-B) (STEFIN B). Length = 98	gi 291927	2	355	100	100	100	HLXND10
358	HUSJE17R	cytochrome c oxidase subunit II [Pan troglodytes] >sp P26457 COX2_PANPA CYTOCHROME C OXIDASE POLYPEPTIDE II (EC 1.9.3.1). Length = 227	gi 336514	17	208	97	98	98	HUSJE17
359	HLJGH82R	cytochrome c oxidase subunit Va preprotein [Mus musculus] >pir S05495 S05495 cytochrome-c oxidase (EC 1.9.3.1) chain Va precursor - mouse >sp P12787 COXA_MOUSE CYTOCHROME C OXIDASE POLYPEPTIDE VA PRECURSOR (EC 1.9.3.1). Length = 145	gi 50527	2	106	94	94	94	HLJGH82
360	HHBEF06R	cytochrome oxidase III [Homo sapiens] >pir A00482 OTHU3 cytochrome-c oxidase (EC 1.9.3.1) chain III - human mitochondrion (SGC1) >sp P00414 COX3_HUMAN CYTOCHROME C OXIDASE POLYPEPTIDE III (EC 1.9.3.1). >gi 2245564 (AF004341) cytochrome c oxidase subunit I	gi 13010	167	373	75	80	80	HHBEF06

361	HISCW28R	cytochrome oxidase subunit II [Homo sapiens] >gi 530071 cytochrome oxidase subunit II [Homo sapiens] >gi 530073 cytochrome oxidase subunit II [Homo sapiens] >gi 530077 cytochrome oxidase subunit II [Homo sapiens] >gi 337187 cytochrome oxidase subunit II [gil530069	121	312	83	86	HISCW28
362	HODEN42R	cytochrome oxidase subunit II [Homo sapiens] >gi 530071 cytochrome oxidase subunit II [Homo sapiens] >gi 530073 cytochrome oxidase subunit II [Homo sapiens] >gi 530077 cytochrome oxidase subunit II [Homo sapiens] >gi 337187 cytochrome oxidase subunit II [gil530069	302	469	68	71	HODEN42
363	HOEMM43R	cytochrome oxidase subunit II [Homo sapiens] >gi 530071 cytochrome oxidase subunit II [Homo sapiens] >gi 530073 cytochrome oxidase subunit II [Homo sapiens] >gi 530077 cytochrome oxidase subunit II [Homo sapiens] >gi 337187 cytochrome oxidase subunit II [gil530069	1	180	64	67	HOEMM43
364	HPIAK29R	cytochrome oxidase subunit II [Homo sapiens] >gi 530071 cytochrome oxidase subunit II [Homo sapiens] >gi 530073 cytochrome oxidase subunit II [Homo sapiens] >gi 530077 cytochrome oxidase subunit II [Homo sapiens] >gi 337187 cytochrome oxidase subunit II [gil530069	295	441	63	70	HPIAK29

365	HUFAR71R	cytochrome oxidase subunit II [Homo sapiens] >gi 530071 cytochrome oxidase subunit II [Homo sapiens] >gi 530073 cytochrome oxidase subunit II [Homo sapiens] >gi 530077 cytochrome oxidase subunit II [Homo sapiens] >gi 337187 cytochrome oxidase subunit II [Homo sapiens]	gi 530069	128	367	82	85	HUFAR71
366	HHEUL74R	cytochrome oxidase subunit II [Homo sapiens] >sp Q37526 Q37526 CYTOCHROME C OXIDASE POLYPEPTIDE II (EC 1.9.3.1). Length = 227	gi 530075	3	227	70	74	HHEUL74
367	H2LAY36R	cytosolic malate dehydrogenase [Homo sapiens] >gi 3133269 malate dehydrogenase [Homo sapiens] >sp P40925 MDHC_HUMAN MALATE DEHYDROGENASE, CYTOPLASMIC (EC 1.1.1.37). {SUB 2-334} Length = 334	gnl PID d1010156	10	609	84	88	H2LAY36
368	HOECI21R	decay-accelerating factor precursor [Homo sapiens] >gnl PID d1023771 (AB003312) decay accelerating factor [Homo sapiens] {SUB 286-340} Length = 376	gi 181463	3	548	73	75	HOECI21
369	HKAFY51R	desmoglein 2 [Homo sapiens] >pir S38673 S38673 desmoglein 2 - human >sp Q14126 DSG2_HUMAN DESMOGLEIN 2 PRECURSOR (HDGC). Length = 1117	gi 416178	1	429	100	100	HKAFY51
370	HMCAR63R	diazepam binding inhibitor [Homo sapiens] Length = 104	gi 181478	3	335	100	100	HMCAR63

371	HWMAN06R	dopamine- and cAMP-regulated neuronal phosphoprotein [Sus scrofa] >sp Q29277 IPPD_PIG DOPAMINE-AND CAMP-REGULATED NEURONAL PHOSPHOPROTEIN (DARPP-32) (FRAGMENT). Length = 137	gi 972053	1	222	83	83	HWMAN06
372	HDPLD04R	early growth response 2 protein (EGR2) - human >gi 181987 early growth response 2 protein [Homo sapiens] {SUB 51-456} Length = 456	pir A40492 A40492	1	459	69	70	HDPLD04
373	HCEGK04R	elongation factor 2 [Gallus gallus] >sp Q90705 EF2_CHICK ELONGATION FACTOR 2 (EF-2). {SUB 2-858} Length = 858	gi 1184958	87	182	95	95	HCEGK04
374	HWLMB57R	epidermal growth factor receptor kinase substrate [Homo sapiens] >pir 38728 38728 epidermal growth factor receptor kinase substrate - human >sp Q12929 EPS8_HUMAN EPIDERMAL GROWTH FACTOR RECEPTOR KINASE SUBSTRATE EPS8. Length = 822	gi 530823	1	186	93	93	HWLMB57
375	HHFHF93R	epidermal growth factor receptor precursor [Homo sapiens] >sp P21860 ERB3_HUMAN ERBB-3 RECEPTOR PROTEIN-TYROSINE KINASE PRECURSOR (EC 2.7.1.112). >gi P1D1e304809 unnamed protein product [Homo sapiens] {SUB 1-27} Length = 1342	gi 181980	1	180	89	89	HHFHF93

376	HCDEM69R	epiligrin alpha 3 subunit [Homo sapiens] >pir A55347 A55347 adhesive ligand epiligrin, alpha-3 chain form A precursor - human >sp Q16787 LMA3_HUMAN LAMININ ALPHA-3 CHAIN PRECURSOR (EPIILIGRIN 170 KD SUBUNIT) (E170). Length = 1713	gij 551597	136	282	95	95	HCDEM69
377	HCHNP50R	epithelial cell marker protein 1 [Homo sapiens] >pir S38956 S38956 epithelial cell marker protein 1 - human Length = 248	gij 187302	54	218	94	94	HCHNP50
378	HAJAW27R	ERF-1 gene product [Homo sapiens] >pir S34854 S34854 epidermal growth factor-response factor 1 - human >gij 972116 ERF-1 protein [Sus scrofa] {SUB 299-337} Length = 338	gij 825653	3	488	100	100	HAJAW27
379	HAICY55R	G-rich sequence factor-1 [Homo sapiens] >gij 517196 G-rich sequence factor-1 [Homo sapiens] >sp Q12849 GRF1_HUMAN G-RICH SEQUENCE FACTOR-1 (GRSF-1). >pir S48081 S48081 GRSF-1 protein - human (fragment) {SUB 94-424} Length = 424	gij 517196	3	374	50	50	HAICY55
380	HWLIA38R	gap junction protein (aa 1-283) [Homo sapiens] >pir B29005 B29005 gap junction protein Cx32 - human >sp P08034 CX32_HUMAN GAP JUNCTION BETA-1 PROTEIN (CONNEXIN 32) (CX32) (GAP JUNCTION 28 KD LIVER PROTEIN). Length = 283	gij 31647	3	455	82	85	HWLIA38

381	HBXCL69R	glutamine--phenylpyruvate aminotransferase [Homo sapiens] >pir S6900 S52790 glutamine-- phenylpyruvate transaminase (EC 2.6.1.64) - human >sp Q16773 Q16773 GLUTAMINE--PIENYLPYRUVATE AMINOTRANSFERASE (EC 2.6.1.64) (GLUTAMINE TRANSAMINASE K). Length = 422	gi 758591	81	419	61	67	HBXCL69
382	H2LAP90R	glutathione peroxidase [Homo sapiens] Length = 202	gi 488476	234	545	97	97	H2LAP90
383	HCQCR94R	glutathione peroxidase-G1 [Homo sapiens] Length = 190	gi 579930	1	114	95	95	HCQCR94
384	HTELE03R	glutathione peroxidase-G1 [Homo sapiens] Length = 190	gi 579930	14	202	100	100	HTELE03
385	HJMBN86R	glutathione-insulin transhydrogenase (216 AA) [Homo sapiens] Length = 216	gi 31746	2	202	97	100	HJMBN86
386	HSKJC32R	GTP:AMP phosphotransferase (EC 2.7.4.10) [Bos taurus] >gnl PI D1001680 mitochondrial adenylate kinase isozyme 3 [Bos taurus] >pir A34442 A34442 nucleoside- triphosphate--adenylate kinase (EC 2.7.4.10) 3, mitochondrial - bovine >sp P08760 KAD3_BOVIN GTP:AM	gi 63528	1	642	89	94	HSKJC32
387	HOEAZ62R	GTP_binding protein [Sus scrofa] >sp Q29222 Q29222 GTP_BINDING PROTEIN (FRAGMENT). Length = 92	gi 971836	2	100	89	92	HOEAZ62

388	HAOAG76R	guanine nucleotide-binding protein G-s-alpha-4 [Homo sapiens] >gi31913 alpha-S1 (AA 1-380) [Homo sapiens] >pir[C31927/RGHUA1 GTP-binding regulatory protein Gs alpha chain (adenylate cyclase-stimulating). splice form 4 - human Length = 380 guanylin [Homo sapiens] >gi306824 guanylin [Homo sapiens] >pir[A46279/A46279 guanylin precursor - human >sp[Q02747/GUAN_HUMAN GUANYLIN PRECURSOR (GUANYLATE CYCLASE ACTIVATOR 2A). Length = 115 H+-ATP synthase subunit b [Homo sapiens] >pir[Q1144/Q1144 H+-transporting ATP synthase (EC 3.6.1.34) chain b precursor, mitochondrial - human >sp[P24539/ATPF_HUMAN ATP SYNTHASE B CHAIN, MITOCHONDRIAL PRECURSOR (EC 3.6.1.34). Length = 256	gi386746	1	369	86	86	86	HAOAG76
389	HCIAD45R	form 4 - human Length = 380 guanylin [Homo sapiens] >gi306824 guanylin [Homo sapiens] >pir[A46279/A46279 guanylin precursor - human >sp[Q02747/GUAN_HUMAN GUANYLIN PRECURSOR (GUANYLATE CYCLASE ACTIVATOR 2A). Length = 115 H+-ATP synthase subunit b [Homo sapiens] >pir[Q1144/Q1144 H+-transporting ATP synthase (EC 3.6.1.34) chain b precursor, mitochondrial - human >sp[P24539/ATPF_HUMAN ATP SYNTHASE B CHAIN, MITOCHONDRIAL PRECURSOR (EC 3.6.1.34). Length = 256	gi183415	2	262	75	81	81	HCIAD45
390	H2MAC82R	form 4 - human Length = 380 guanylin [Homo sapiens] >gi306824 guanylin [Homo sapiens] >pir[A46279/A46279 guanylin precursor - human >sp[Q02747/GUAN_HUMAN GUANYLIN PRECURSOR (GUANYLATE CYCLASE ACTIVATOR 2A). Length = 115 H+-ATP synthase subunit b [Homo sapiens] >pir[Q1144/Q1144 H+-transporting ATP synthase (EC 3.6.1.34) chain b precursor, mitochondrial - human >sp[P24539/ATPF_HUMAN ATP SYNTHASE B CHAIN, MITOCHONDRIAL PRECURSOR (EC 3.6.1.34). Length = 256	gi509291	214	513	95	96	96	H2MAC82
391	H2LAJ41R	heat shock protein [Homo sapiens] >pir[A32319/HHIU86 heat shock protein 90-alpha - human >gi184419 heat shock protein 86 [Homo sapiens] {SUB 1-312} >gnl[PID]d1014121 heat shock protein 90 [Homo sapiens] {SUB 582-732} Length = 732	gi703087	75	632	98	98	98	H2LAJ41
392	HWLGH40R	HKL1 [Homo sapiens] >sp[O60765/O60765 HKL1. Length = 605	gnl[PID]d1026110	1	597	92	93	93	HWLGH40

393	HBJFH33R	HLA DP4 beta-chain [Homo sapiens] >gi296648 pot. hla-dp-beta 1 [Homo sapiens] >pir A02229 HL.HUPB MHC class II histocompatibility antigen HLA-DP beta 1 chain (allele DPB4.1) precursor - human >sp P04440 HB2P_HUMAN HLA CLASS II HISTOCOMPATIBILITY ANTIGEN.	gij306858	97	369	88	92	HBJFH33
394	HISDV92R	homeobox c1 protein [Homo sapiens] >sp Q64081 Q64081 HOX-B HOX-2 {CLONE 17A1}. {SUB 137-196; Length = 217	gij306878	51	404	72	72	HISDV92
395	HMQCG89R	Hox5.4 gene product (AA 1-95) [Homo sapiens] >pir B32830 B32830 homeotic protein Hox D8 - human (fragment) >sp P13378 HDXD8_HUMAN HOMEBOX PROTEIN HOX-D8 (HOX-4E) (HOX-5.4) (FRAGMENT). Length = 95	gij32400	158 1	388 345	100	100	HMQCG89 HE9QB35
396	HE9QB35R							
397	HDABQ50R	hsOrc2p [Homo sapiens] >sp Q13416 ORC2_HUMAN ORIGIN RECOGNITION COMPLEX PROTEIN, SUBUNIT 2. Length = 577	gij113107	204	368	91	91	HDABQ50

398	HNTEG83R	hydroxymethylglutaryl-CoA lyase [Homo sapiens] >pir A45470 A45470 hydroxymethylglutaryl-CoA lyase (EC 4.1.3.4) - human >sp P35914 HMGL_HUMAN HYDROXYMETHYLGLUTARYL- COA LYASE PRECURSOR (EC 4.1.3.4) (HMG-COA LYASE) (HL) (3- HYDROXY-3-METHYLGLUTARATE- COA LYASE	gil184503	2	391	83	83	HNTEG83
399	HFVHM90R	hydroxymethylglutaryl-CoA synthase [Homo sapiens] >gil2463646 3-hydroxy- 3-methylglutaryl CoA synthase [Homo sapiens] >pir S71623 S71623 hydroxymethylglutaryl-CoA synthase (EC 4.1.3.5) precursor, mitochondrial - human >sp P54868 HMCM_HUMAN HYDROXYMETHYLGLU	gil619877	2	319	92	94	HFVHM90
400	HOSNF90R	hypothetical 18K protein (rRNA) - goldfish mitochondrion (SGC1) Length = 166	pir JC1348 JC1348	257	340	59	62	HOSNF90
401	HSDJE56R	hypothetical 18K protein (rRNA) - goldfish mitochondrion (SGC1) Length = 166	pir JC1348 JC1348	2	70	67	73	HSDJE56
402	HWLGC87R	hypothetical protein 2 (rRNA external transcribed spacer) - mouse Length = 153	pir S12206 S12206	1	135	96	96	HWLGC87

403	HTPAC28R	I-plastin [Homo sapiens] >pir A56536 A56536 plastin, intestine-specific - human >sp Q14651 PLSI_HUMAN I-PLASTIN (INTESTINE-SPECIFIC PLASTIN). Length = 629	gi 405230	68	325	92	93	HTPAC28
404	IIMCGN07R	ICK=INTRON-CONTAINING KALLIKREIN {ALTERNATIVELY SPLICED, INTRON 2}. Length = 216	sp G998972 G998972	1	498	98	99	IIMCGN07
405	HFIBV16R	Id1 gene product [Homo sapiens] >pir S47524 S47524 gene Id1 protein - human Length = 154	gi 457785	2	238	89	89	HFIBV16
406	HBMTT01R	Ig alpha-2 chain C region (allotype A2m(1)) - human >sp P01877 ALC2_HUMAN IG ALPHA-2 CHAIN C REGION. >gi 184761 Ig alpha-2 H-chain constant region (aa at 166) [Homo sapiens] {SUB 2-340} Length = 340	pir B22360 B22360	2	154	80	80	HBMTT01
407	HBMVM66R	Ig gamma chain C region - chimpanzee >gnl PID e40518 CH2 domain of IgG [Pan troglodytes] {SUB 25-134} >gnl PID e40517 CH3 domain of IgG [Pan troglodytes] {SUB 135-234} Length = 234	pir PT0207 PT0207	148	435	70	77	HBMVM66
408	HABGC21R	Ig heavy chain (DOT) - human (fragment) >gnl PID e4381 reading frame CH1 [Homo sapiens] {SUB 121-218} Length = 241	pir S69131 S69131	1	228	50	56	HABGC21
409	HWLGE72R	Ig kappa light chain (VJ) [Homo sapiens] >pir S40343 S40343 Ig kappa chain V-J region - human Length = 128	gi 441375	11	421	75	79	HWLGE72

410	HLJBX69R	IgM B-cell receptor associated protein (BAP) 37 [Mus musculus] >pir S46996 S46996 B-cell receptor-associated protein [BAP]37 - mouse >sp Q61336 Q61336 BCR-ASSOCIATED PROTEIN 37 (IGM B-CELL RECEPTOR ASSOCIATED PROTEIN 37) (BAP). Length = 298	gij541734	1	279	100	100	HLJBX69
411	HWAFW14R	immunoglobulin from VH4 family [Homo sapiens] >pir S13519 S13519 Ig heavy chain V region precursor - human >gij553385 immunoglobulin heavy chain [Homo sapiens] {SUB 24-125} Length = 147	gij37725	2	139	94	100	HWAFW14
412	HWAFK04R	immunoglobulin heavy chain [Homo sapiens] >pir E36005 E36005 Ig heavy chain V region (M72) - human {SUB 36-157} Length = 157	gij567126	48	473	78	86	HWAFK04
413	HEPNA09R	immunoglobulin heavy chain [Homo sapiens] >pir G36005 G36005 Ig heavy chain V region (M74) - human {SUB 38-158} Length = 158	gij567127	3	206	81	87	HEPNA09
414	HCRQD03R	immunoglobulin heavy chain [Homo sapiens] Length = 152	gij567128	1	573	76	82	HCRQD03
415	HAPSK08R	immunoglobulin heavy chain variable region [Homo sapiens] >gij903667 Ig heavy chain variable region VH [Homo sapiens] {SUB 1-97} >gij976311 This CDS feature is included to show the translation of the corresponding V_segment. Presently translation qualifie	gij1791017	1	363	79	81	HAPSK08

416	HBM1S11R	immunoglobulin IgH heavy chain Fd fragment [Homo sapiens] Length = 221	gi 468237	1	375	68	70	HBM1S11
417	HCNDR62R	immunoglobulin kappa light chain [Homo sapiens] >pir A37927 A37927 Ig kappa chain C region (allotype Inv(1.2)) - human (fragment) {SUB 138-236} Length = 236	gn PI e224083	245	337	100	100	HICNDR62
418	HNJBF13R	immunoglobulin lambda light chain gene product [Homo sapiens] >pir S25738 S25738 Ig lambda chain - human Length = 231	gi 33702	3	308	90	93	HNJBF13
419	HL YCD69R	immunoglobulin lambda light chain gene product [Homo sapiens] >pir S25743 S25743 Ig lambda chain - human (fragment) Length = 145	gi 33712	2	481	86	89	HL YCD69
420	HWAFK89R	immunoglobulin lambda light chain gene product [Homo sapiens] >pir S25750 S25750 Ig lambda chain - human Length = 235	gi 33730	2	460	87	92	HWAFK89
421	HWCAA53R	immunoglobulin light chain variable region [Homo sapiens] >gi 3142470 (AF063703) immunoglobulin lambda light chain variable region [Homo sapiens] {SUB 20-127} >gi 575243 immunoglobulin lambda chain precursor [Homo sapiens] {SUB 26-127} >gn PI d1020826 V	gi 465170	1	342	74	88	HWCAA53
422	HYAAY47R	immunoglobulin light chain variable region [Homo sapiens] Length = 154	gi 465168	2	292	70	74	HYAAY47
423	HMCJF14R			21	596			HMCJF14
424	HE8QU88R			13	141			HE8QU88

425	HFVGP11R	L-FABP [Homo sapiens] >pir A22289 FZHU.L fatty acid-binding protein, hepatic - human >sp P07148 FABL_HUMAN FATTY ACID-BINDING PROTEIN, LIVER (L- FABP). Length = 127	gij 182358	29	322	98	98	HFVGP11
426	HWLQH07R			3	554			HWLQH07
427	HSIGN24R	Irp gene product [Homo sapiens] >pir SS7723 SS7723 Irp protein - human >sp Q14764 MVP_HUMAN MAJOR VAULT PROTEIN (MVP) (LUNG RESISTANCE-RELATED PROTEIN). Length = 896	gij 895840	2	250	89	93	HSIGN24
428	HWLKH07R	lysophosphatidic acid acyltransferase- beta [Homo sapiens] Length = 278	gij 2155240	74	298	96	97	HWLKH07
429	HAPQC14R	macrophage capping protein [Homo sapiens] >pir A43358 A43358 macrophage capping protein - human >sp P40121 CAPG_HUMAN MACROPHAGE CAPPING PROTEIN (ACTIN-REGULATORY PROTEIN CAP-G). >gij 515505 Cap-G [Homo sapiens] {SUB 1-172} Length = 348	gij 187456	2	538	96	98	HAPQC14
430	HSODB48R	malonyl-CoA decarboxylase (EC 4.1.1.9) - goose >gij 305323 malonyl CoA decarboxylase [Anser anser] {SUB 33- 462} Length = 462	pir A33313 A33313 3	32	466	77	81	HSODB48
431	HBEAC75R	membrane glycoprotein [Homo sapiens] Length = 385	gij 307132	2	217	73	79	HBEAC75
432	HBGMJ24R	mitochondrial RNA polymerase [Homo sapiens] Length = 1230	gij 2114396	3	479	100	100	HBGMJ24

433	HBJEN94R	mitotic kinase-like protein-1 [Homo sapiens] >pir S28262 S28262 kinesin-related protein MKLP-1 - human >sp Q02241 MKLP_HUMAN MITOTIC KINESIN-LIKE PROTEIN-1. Length = 960	gij34672	1	327	89	89	HBJEN94
434	HICIAE73R	motor protein [Homo sapiens] Length = 721	gn PIPI d1005183	73	324	100	100	HICIAE73
435	HCNDN88R	mucin 2 precursor, intestinal - human (fragments) >gij186396 mucin [Homo sapiens] {SUB 626-1895} >gij186398 MUC2 [Homo sapiens] {SUB 2037-3020} >gij188874 intestinal mucin [Homo sapiens] {SUB 1916-2193} >gij188615 mucin-like protein [Homo sapiens] {SUB 23}	pir A49963 A4393 2	1	171	95	97	HCNDN88
436	HSIDX70R	N-benzoyl-L-tyrosyl-p-amino-benzoic acid hydrolase alpha subunit [Homo sapiens] >pir S60193 H_YHUMA meprin A (EC 3.4.24.18) alpha chain precursor - human >sp Q16819 MEPA_HUMAN MEPRIN A ALPHA-SUBUNIT PRECURSOR (EC 3.4.24.18) (ENDOPEPTIDASE-2) (N-BENZOYL-L-	gij535475	2	253	94	94	HSIDX70

437	HILWBC39R	Na ⁺ /H ⁺ exchanger NHE-1 isoform [human, heart, Peptide, 815 aa] [Homo sapiens] >pir 57487 57487 Na ⁺ /H ⁺ - exchanging protein NHE-1 - human >sp P19634 NAH1_HUMAN SODIUM/HYDROGEN EXCHANGER 1 (NA ⁺ /H ⁺) EXCHANGER 1 (NHE- 1) (NA ⁺ /H ⁺ ANTIporter, AMILORIDE-SENSI	bbs 43522	2	388	77	77	HILWBC39
438	HWLAA06R	NADH dehydrogenase (ubiquinone) (EC 1.6.5.3) chain 4 - chimpanzee mitochondrion (S611) (fragment) >sp P03906 NU4M_PANTR NADH- UBIQUINONE OXIDOREDUCTASE CHAIN 4 (EC 1.6.5.3) (FRAGMENT). Length = 152	pir A00435 A0043 5	66	194	86	97	HWLAA06
439	HASCH25R	NADH-UBIQUINONE OXIDOREDUCTASE 39 KD SUBUNIT PRECURSOR (EC 1.6.5.3) (EC 1.6.99.3) (COMPLEX I-39KD) (CI-39KD). >gi 189049 NADH dehydrogenase (ubiquinone) [Homo sapiens] {SUB 3- 377} Length = 377	sp Q16795 NUEM _HUMAN	57	143	78	82	HASCH25
440	HILQGB87R	NADPH--ferrihemoprotein reductase (EC 1.6.2.4) - human >sp P16435 NCPR_HUMAN NADPH- CYTOCHROME P450 REDUCTASE (EC 1.6.2.4) (CPR). {SUB 2-677} Length = 677	pir A3342 A6055 7	1	411	92	93	HILQGB87

441	IIIDMD17R	neutrophil gelatinase associated lipocalin [Homo sapiens] >sp P80188 NGAL_HUMAN NEUTROPHIL GELATINASE- ASSOCIATED LIPOCALIN PRECURSOR (NGAL) (P25) (25 KD ALPHA-2-MICROGLOBULIN- RELATED SUBUNIT OF MMP-9) (LIPOCALIN-2) (ONCOGENE 24P3). Length = 198	gi 929657	1	621	74	78	IIIDMD17
442	IIAOAC69R	nuclear autoantigen [Homo sapiens] >pir A37244 A37244 nuclear autoantigen Sp-100 - human Length = 480	gi 178689	3	209	88	88	IIAOAC69
443	IIWLEQ08R	Nuclear localization signal at AA 569- 573, 576-580, 579-583; acidic trans- activ. domain 620-640.; homeobox motif 653-676 [Homo sapiens] >pir A47456 A47456 down-regulated in adenoma (DRA) - human >sp P40879 DRA_HUMAN DRA PROTEIN (DOWN-REGULATED IN ADENO	gi 291964	191	364	75	84	IIWLEQ08
444	HKAAV70R	nucleic acid binding protein [Homo sapiens] >pir I38191 I38191 nucleic acid binding protein - human (fragment) >sp Q15410 Q15410 NUCLEIC ACID BINDING PROTEIN (FRAGMENT). Length = 163	gi 431953	1	432	73	73	HKAAV70

445	HOCTB64R	ORIGINAL PIGR [unidentified] >gi456346 Polymetric immunoglobulin receptor [Homo sapiens] >bbs62408 transmembrane secretory component, poly-Ig receptor, SC [human, colonic adenocarcinoma cell line, Peptide, 764 aa] [Homo sapiens] >bbs113253 transmembrane	gnl P D e307278	3	212	85	90	HOCTB64
446	HOFNB62R	ornithine decarboxylase [Bos taurus] >gi163449 ornithine decarboxylase [Bos taurus] >sp P27117 DCOR_BOVIN ORNITHINE DECARBOXYLASE (EC 4.1.1.17) (ODC). >gi604513 ornithine decarboxylase [Bos taurus] (SUB 1-34) Length = 461	gi 1036793	1	312	85	90	HOFNB62
447	HAUAU04R	p22 phagocyte b-cytochrome [Homo sapiens] >pir A28201 A28201 cytochrome b-245 alpha chain - human >sp P13498 C24A_HUMAN CYTOCHROME B-245 LIGIT CIIAIN (P22 PHAGOCYTE B-CYTOCHROME) (NEUTROPHIL CYTOCHROME B, 22 KD POLYPEPTIDE) (P22-PHOX) (CYTOCHROME B(558) AL	gi 189106	1	267	87	88	HAUAU04
448	HNFJE41R	p47-phox [Homo sapiens] >sp O43842 O43842 P47-PHOX. Length = 390	gi 2754713	1	423	94	97	HNFJE41
449	IICFOH92R	phosphoprotein phosphatase (EC 3.1.3.16) catalytic beta chain - pig (fragment) Length = 293	pir B27430 B2743 0	2	88	93	93	IICFOH92
450	HOUIDS3R	phosphorylation regulatory protein HP-10 - human Length = 492	pir A61382 A6138 2	85	213	45	49	HOUIDS3

451	HCRMW41R	polypeptide BM28 [Homo sapiens] Length = 892	gi 468704	1	282	100	100	IICRMW41
452	HOVAX78R	porin [Homo sapiens] >pir A45972 A45972 mitochondrial porin, long form - human >sp P45880 POR2_HUMAN VOLTAGE-DEPENDENT ANION- SELECTIVE CHANNEL PROTEIN 2 (VDAC2) (OUTER MITOCHONDRIAL MEMBRANE PROTEIN PORIN). >gi 190201 porin [Homo sapiens] [SUB 27-347] Len	gi 190200	2	214	94	98	HOVAX78
453	HWAEH57R	precursor [Homo sapiens] >sp P06314 KV4C_HUMAN IG KAPPA CHAIN PRECURSOR V-IV REGION (B17). Length = 134	gi 37910	1	462	91	93	HWAEH57
454	HHBJU76R	presenilin 1-463 [Homo sapiens] >pir S63683 S63683 presenilin 1-463 - human Length = 463	gi 1244638	1	303	98	98	HHBJU76
455	HBJFA18R	prosome P27K protein [Homo sapiens] >gnl P1D1d1002062 prosome subunit R-IOTA [Rattus sp.] >pir S30274 S30274 multicatalytic endopeptidase complex (EC 3.4.99.46) iota chain - human >pir JX0230 JX0230 multicatalytic endopeptidase complex (EC 3.4.99.46)	gi 35682	178	402	79	83	HBJFA18
456	HCRNF16R	protein kinase [Homo sapiens] >sp P51956 NEK3_HUMAN SERINE/THREONINE-PROTEIN KINASE NEK3 (EC 2.7.1.-) (NIMA- RELATED PROTEIN KINASE 3) (HSPK 36) (FRAGMENT). Length = 459	gi 479173	336	473	73	79	HCRNF16

457	HAHEK76R	putative surface glycoprotein [Homo sapiens] >sp P53801 C211_HUMAN PUTATIVE SURFACE GLYCOPROTEIN C21ORF1	gn PID e188111	33	440	83	86	HAHEK76
458	HEOPT38R	PRECURSOR (C21ORF3), Length = 180 renin-binding protein [Homo sapiens] >gij1302662 renin-binding protein [Homo sapiens] >pir JX0188 JX0188 renin-binding protein - human Length = 417	gn PID d1001551	2	316	100	100	HEOPT38
459	HOSCG81R	ribonucleoprotein La [Homo sapiens] >sp Q15367 Q15367 RIBONUCLEOPROTEIN (LA) (FRAGMENT). >gij1338496 SS-B/La protein [Homo sapiens] {SUB 121-171} Length = 355	gij1337457	1	297	96	96	HOSCG81
460	HTFMD43R	ribosomal protein L39 [Homo sapiens] >gn PID d1012131 ribosomal protein L39 [Homo sapiens] >gij1575382 ribosomal protein L39 [Rattus norvegicus] >pir JC4229 R6RT39 ribosomal protein L39 - rat >pir G02654 G02654 ribosomal protein L39 - human Length = 51	gij1373419	3	242	100	100	HTFMD43
461	HDTGQ68R	ribosomal protein L7a large subunit [Homo sapiens] >gij134203 L7a protein [Homo sapiens] >gij135512 PLA-X polypeptide [Homo sapiens] >gij136647 ribosomal protein L7a [Homo sapiens] >gij156956 ribosomal protein L7a (AA 1- 266) [Rattus rattus] >pir S19717 R5HU7A	gij1337495	43	291	100	100	HDTGQ68

462	H2LAR73R	ribosomal protein S15a [Rattus norvegicus] >pir JC2234 JC2234 ribosomal protein S15a - rat Length = 130	gij495273	23	505	100	100	100	H2LAR73
463	HAMFM26R	ribosomal protein S6 kinase 1 [Homo sapiens] >pir S1901 S1901 ribosomal protein S6 kinase 2 - human >sp Q15418 KS61_HUMAN RIBOSOMAL PROTEIN S6 KINASE II ALPHA 1 (EC 2.7.1.-) (S6KII-ALPHA 1) (P90-RSK 1) (RIBOSOMAL S6 KINASE 1) (RSK1) (PP90RSK1). Length =	gij292457	3	458	97	97	97	HAMFM26
464	HBMTM61R	Rieske Fe-S protein [Homo sapiens] Length = 274	gij488299	1	219	53	53	55	HBMTM61
465	I1W1PK71R	RIP [Homo sapiens] >pir 38992 38992 receptor interacting protein RIP - human (fragment) Length = 372	gij829617	198	320	56	56	64	I1W1PK71
466	HWBBJ39R	Sec23 protein [Homo sapiens] Length = 767	gnl PID c236014	2	127	81	81	84	HWBBJ39
467	HSLJJ36R	selenium donor protein [Homo sapiens] Length = 383	gij1000284	2	319	96	96	98	HSLJJ36
468	HSODD94R	selenoprotein P [Homo sapiens] Length = 381	gnl PID c1192260	2	232	61	61	70	HSODD94
469	HMIAG25R	serine kinase [Homo sapiens] >pir S45337 S45337 serine protein kinase SRPK1 - human >sp Q12890 Q12890 SERINE KINASE. Length = 655	gij507213	1	330	82	82	82	HMIAG25
470	HWLEM94R	serine protease [Homo sapiens] Length = 492	gij2507613	2	304	78	78	82	HWLEM94

471	HCNDW17R	Sm protein G [Homo sapiens] >pir S55054 S55054 Sm protein G - human >sp Q15357 Q15357 SM PROTEIN G. Length = 76	gi 806366	1	240	100	100	HCNDW17
472	HWLEY08R	SNAP23A protein [Homo sapiens] >gnl PID e1331767 (AJ011915) synaptosome associated protein of 23 kilodaltons, isoform A [Homo sapiens] >pir JC5296 JC5296 vesicle-membrane fusion protein SNAP-23A - human >sp O00161 O00161 VESICLE-MEMBRANE FUSION PROTEIN SN	gnl PID e290695	222	608	97	97	HWLEY08
473	HULFN68R	sorcin CP-22 [Homo sapiens] >gi 459836 sorcin [Homo sapiens] >pir S52094 S52094 sorcin - human >gi 2772536 (AC003991) calcium binding protein amplified in multidrug-resistant cells [Homo sapiens] {SUB 1-68} Length = 198	gi 338482	2	409	88	91	HULFN68
474	HMEJD77R	SRp30c [Homo sapiens] >gnl PID e1248292 (AL021546) pre-mRNA splicing factor SRp30c [Homo sapiens] >gi 4099429 splicing factor SRp30c [Homo sapiens] >pir S59075 S59075 splicing factor SRp30c - human >sp G4099429 G4099429 SPLICING FACTOR SRP30C. Length = 22	gi 1049078	3	263	46	48	HMEJD77
475	HS2AD15R	stimulator of TAR RNA binding [Homo sapiens] Length = 539	gi 1200184	1	336	87	88	HS2AD15

476	HTEJJ32R	STM-7 [Homo sapiens] >sp Q92749 Q92749 TYPE I PHOSPHATIDYLINOSITOL-4- PHOSPHATE 5-KINASE BETA (EC 2.7.1.68) (STM-7 PROTEIN). >gi 1743883 type I phosphatidylinositol- 4-phosphate 5-kinase beta [Homo sapiens] {SUB 112-502} >gi 1743879 type I phosphatidylinosi	gnl PID e206448	3	341	100	100	100	HTEJJ32
477	HETIF46R	sulfate transporter [Homo sapiens] >sp P50443 DTD_HUMAN SULFATE TRANSPORTER (DIASTROPHIC DYSPLASIA PROTEIN). Length = 739 thrombospondin 2 [Homo sapiens] >pir A47379 TSITUP2 thrombospondin 2 precursor - human Length = 1172 thymosin beta-4 precursor [Rattus norvegicus] >pir 52084 52084 thymosin beta-4 precursor - rat (fragment) >gi 339689 thymosin beta-4 [Homo sapiens] {SUB 13-56} >pir A0152 TNBOB4 thymosin beta-4 - bovine {SUB 14-56} >gi 825683 open reading frame [Homo s	gi 549988	1	228	71	71	71	HETIF46
478	H2CBS58R		gi 307506	3	455	96	96	97	H2CBS58
479	H2LAB77R		gi 207318	98	265	100	100	100	H2LAB77
480	IIODAJ23R	tissue-specific secretory protein [unidentified] >gi 32051 HE4 protein [Homo sapiens] >pir S25454 S25454 HE4 protein - human >sp Q14508 EP4_HUMAN MAJOR EPIDIDYMIS-SPECIFIC PROTEIN E4 PRECURSOR (HE4) (EPIDIDYMAL SECRETORY PROTEIN E4). Length =	gi 583141	2	223	62	62	62	IIODAJ23

125

76

481	HWAFP88R	TRANSCRIPTION FACTOR BTF3 (RNA POLYMERASE B TRANSCRIPTION FACTOR 3). Length = 204	sp Q64152 BTF3_1 MOUSE	85	471	92	93	HWAFP88
482	HDTTH51R	transcription factor-like protein 4 - human Length = 298	pir JC5333 JC5333	2	565	82	86	IIDTII51
483	HWMEB67R	tryptase-III [Homo sapiens] >sp Q15664 Q15664 TRYPTASE-III (FRAGMENT). Length = 267	gi 339985	21	218	92	92	HWMEB67
484	HTXOU93R	tumor susceptibility protein [Homo sapiens] >sp Q99816 Q99816 TUMOR SUSCEPTIBILITY PROTEIN. Length = 390	gi 3184258	2	439	100	100	HTXOU93
485	HANKB37R	ubiquitin [Plasmodium falciparum] >sp Q26029 Q26029 UBIQUITIN. Length = 77	gi 552237	11	115	70	73	HANKB37
486	HWLHN38R	ubiquitin-conjugating enzyme [Mus musculus] >sp Q88738 Q88738 UBIQUITIN-CONJUGATING ENZYME. Length = 4845	gn P1D1e1311091	129	347	77	83	HWLHN38

487	HOSDZ35R	UDP-GalNAc:polypeptide N-acetyl-galactosaminyltransferase [Homo sapiens] >sp Q14435 Q14435 POLYPEPTIDE N-ACETYL-GALACTOSAMINYLTRANSFERASE (EC 2.4.1.41) (PROTEIN-UDP ACETYL-GALACTOSAMINYLTRANSFERASE) (UDP-GALNAc:POLYPEPTIDE, N-ACETYL-GALACTOSAMINYLTRANSFERASE)	gnl PID c20971	2	286	85	85	HOSDZ35
488	HKMAA52R	UDP-glucuronosyltransferase [Homo sapiens] >pir A31340 A31340 glucuronosyltransferase (EC 2.4.1.17) UGT1A1 precursor - human >sp G245274 G245274 PHENOL TRANSFERASE=UGT1F: PRODUCT: {SUB 1-286} >gij 264549 (AF014112) phenol UDP-glucuronosyltransferase [Homo	gij 624725	3	284	98	98	HKMAA52
489	H2LAB37R			93	290			H2LAB37
490	H2LAP46R			206	568			H2LAP46
491	H6BSE61R			67	369			H6BSE61
492	H6EEE76R			149	277			H6EEE76
493	H6EEV26R			2	88			H6EEV26
494	HABAF88R			40	216			HABAF88
495	IIABGD41R			1	147			IIABGD41
496	HACBS75R			5	187			HACBS75
497	HACCA48R			5	91			IIACCA48
498	HACCS19R			3	341			HACCS19
499	IIADAB25R			1	261			IIADAB25
500	HAGGL96R			3	347			HAGGL96

501	HAGGT37R	3	113	IIAGGT37
502	HAHDR66R	27	347	HAHDR66
503	HAJCC53R	164	418	HAIJCC53
504	HAIJCL80R	3	122	HAIJCL80
505	HANKF43R	372	566	HANKF43
506	HAPCM11R	69	152	HAPCM11
507	HAPNT66R	1	66	HAPNT66
508	HAQAG47R	2	148	HAQAG47
509	HAQBW58R	3	260	HAQBW58
510	HAQMH45R	91	363	HAQMH45
511	HAQMI94R	1	183	HAQMI94
512	HARNC74R	84	272	HARNC74
513	HATBA87R	98	202	HATBA87
514	HATBG77R	174	392	HATBG77
515	HBAGQ79R	1	231	HBAGQ79
516	HBCAN64R	2	82	HBCAN64
517	HBGCA44R	1	123	HBGCA44
518	HBGFX27R	3	281	HBGFX27
519	HBGMU38R	40	429	HBGMU38
520	HBIBO10R	1	93	HBIBO10
521	HBJCC53R	2	106	HBJCC53
522	HBJED55R	1	252	HBJED55
523	HBJGR39R	2	106	HBJGR39
524	HBJLU30R	39	344	HBJLU30
525	HBKEC78R	93	245	HBKEC78
526	HBMST81R	1	192	HBMST81
527	HBMTJ51R	150	323	HBMTJ51
528	HBMWLF72R	1	111	HBMWLF72
529	HBWBD78R	2	226	HBWBD78
530	HBXCU02R	2	79	HBXCU02
531	HCDAK65R	1	138	HCDAK65

532	HCDBM08R	130	339	HCDBM08
533	HCDCP10R	72	206	HCDCP10
534	HCDDQ63R	3	116	HCDDQ63
535	HCEEH05R	204	380	HCEEH05
536	HCEIQ92R	1	90	HCEIQ92
537	HCPCD01R	28	228	HCPCD01
538	HCFCR43R	64	360	HCFCR43
539	HCFLT83R	3	104	HCFLT83
540	HCHAO92R	193	342	HCHAO92
541	HCHOH49R	183	344	HCHOH49
542	HCHPG05R	365	616	HCHPG05
543	HCIAD24R	98	301	HCIAD24
544	HCNCA90R	380	532	HCNCA90
545	HCNCN80R	120	353	HCNCN80
546	HCNCY51R	184	267	HCNCY51
547	HCNCY63R	1	81	HCNCY63
548	HCNDO71R	1	213	HCNDO71
549	HCNDV83R	64	303	HCNDV83
550	HCNUB26R	119	289	HCNUB26
551	HCQBN22R	2	94	HCQBN22
552	HCQCL27R	116	235	HCQCL27
553	HCQCL48R	57	251	HCQCL48
554	HCQCL96R	287	430	HCQCL96
555	HCQDC74R	145	360	HCQDC74
556	HCQDH94R	20	76	HCQDH94
557	HCQDJ42R	149	388	HCQDJ42
558	HCNMD77R	3	185	HCNMD77
559	HCNME02R	3	293	HCNME02
560	HCNMX88R	3	284	HCNMX88
561	HCNNA70R	40	204	HCNNA70
562	HCNRP66R	3	431	HCNRP66

563	HCRNX32R	2	196	HCRNX32
564	HCROH25R	3	128	HCROH25
565	HCROJ05R	66	170	HCROJ05
566	HCROJ68R	3	239	HCROJ68
567	HCROK68R	2	208	HCROK68
568	HCROK94R	1	210	HCROK94
569	HCROM30R	3	365	HCROM30
570	HCROQ34R	29	136	HCROQ34
571	HCROQ54R	3	98	HCROQ54
572	HCROZ66R	239	427	HCROZ66
573	HCRPC61R	3	194	HCRPC61
574	HCRPG28R	95	229	HCRPG28
575	HCRPL80R	59	235	HCRPL80
576	HCRPN52R	3	191	HCRPN52
577	HCRPS40R	208	321	HCRPS40
578	HCRPV74R	179	409	HCRPV74
579	HCRQC89R	2	85	HCRQC89
580	HCWDS78R	322	558	HCWDS78
581	HDCAA21R	1	120	HDCAA21
582	HDDAA85R	139	258	HDDAA85
583	HDPGO03R	110	352	HDPGO03
584	HDPLB08R	142	360	HDPLB08
585	HDQDB15R	220	417	HDQDB15
586	HDQEX80R	274	492	HDQEX80
587	HDRMI91R	3	116	HDRMI91
588	HDTJO85R	36	197	HDTJO85
589	HDTMJ22R	192	608	HDTMJ22
590	HE6CS28R	40	213	HE6CS28
591	HE6DJ45R	2	64	HE6DJ45
592	HE7TJ40R	62	268	HE7TJ40
593	HE9FH12R	182	307	HE9FH12

594	HE9HJ57R	3	74	HE9HJ57
595	HE9QH08R	360	596	HE9QH08
596	HE9TC50R	198	425	HE9TC50
597	HEAAL59R	1	150	HEAAL59
598	HEGAR32R	448	675	HEGAR32
599	HEGAR85R	361	534	HEGAR85
600	HELFE05R	32	187	HELFE05
601	HEMF188R	26	343	HEMF188
602	HEMFR18R	83	397	HEMFR18
603	HEONL43R	2	76	HEONL43
604	HESAC53R	3	116	HESAC53
605	HETJB05R	1	138	HETJB05
606	HETJC36R	1	102	HETJC36
607	HFADM62R	1	78	HFADM62
608	HFATE31R	2	361	HFATE31
609	HFATZ30R	3	152	HFATZ30
610	HFCEL77R	3	278	HFCEL77
611	HFEBN43R	174	491	HFEBN43
612	HFCAF10R	272	469	HFCAF10
613	HFEC01R	1	144	HFEC01
614	HFIR75R	317	427	HFIR75
615	HFUB90R	2	124	HFUB90
616	HFUM71R	37	159	HFUM71
617	HFOXLS3R	1	117	HFOXLS3
618	HFPBO66R	196	408	HFPBO66
619	HFTBI57R	47	220	HFTBI57
620	HFTCC22R	1	126	HFTCC22
621	HFXGX46R	1	114	HFXGX46
622	HGAME72R	2	199	HGAME72
623	HGBCS53R	142	279	HGBCS53
624	HGBHP81R	87	221	HGBHP81

625	IGCOX03R	323	511	HGCOX03
626	HHBES92R	349	483	HHBES92
627	HHBEW72R	13	219	HHBEW72
628	HHERT59R	2	88	HHERT59
629	HHMMD64R	31	252	HHMMD64
630	HHSGT13R	428	619	HHSGT13
631	HISED82R	1	126	HISED82
632	IJMAH76R	2	253	IJMAH76
633	IJMAN56R	1	180	IJMAN56
634	IJMAO54R	1	291	IJMAO54
635	HKDAD56R	2	109	HKDAD56
636	HKLSD93R	89	298	HKLSD93
637	HLMFH16R	1	447	HLMFH16
638	HLQBD52R	1	195	HLQBD52
639	HLQCQ73R	3	350	HLQCQ73
640	HLQEF47R	348	503	HLQEF47
641	HLQFM50R	136	291	HLQFM50
642	HLQFY61R	411	575	HLQFY61
643	HLQGA76R	210	404	HLQGA76
644	HLQGE53R	1	66	HLQGE53
645	HLTEV09R	210	371	HLTEV09
646	HLXNE63R	142	258	HLXNE63
647	HLXTF64R	2	136	HLXTF64
648	HMACF85R	23	430	HMACF85
649	HMAIA15R	108	452	HMAIA15
650	HMCHZ07R	247	402	HMCHZ07
651	HMCIS54R	84	242	HMCIS54
652	HMSFW88R	1	69	HMSFW88
653	HMSMW71R	290	514	HMSMW71
654	HNHMR05R	77	598	HNHMR05
655	HNJBB78R	91	282	HNJBB78

656	HINTMA96R	3	362	HINTMA96
657	HINTRL32R	130	291	HINTRL32
658	HINTST76R	2	397	HINTST76
659	HOCNC55R	67	156	HOCNC55
660	HOCND06R	147	275	HOCND06
661	HOCND49R	133	273	HOCND49
662	HODEH30R	2	154	HODEH30
663	HODFA26R	263	550	HODFA26
664	HODHL89R	106	279	HODHL89
665	HOEJM67R	2	364	HOEJM67
666	HOGBN48R	147	380	HOGBN48
667	HOHCX95R	2	364	HOHCX95
668	HORBP43R	3	365	HORBP43
669	HOUHN53R	235	345	HOUHN53
670	HOUJE10R	72	254	HOUJE10
671	HPBEE63R	107	211	HPBEE63
672	HIPEBO20R	1	237	HIPEBO20
673	HPJBE91R	1	312	HPJBE91
674	HPTRW82R	32	133	HPTRW82
675	HPWDC51R	33	272	HPWDC51
676	HPWDK52R	1	330	HPWDK52
677	HRDBJ82R	2	334	HRDBJ82
678	HIRODH93R	2	121	HIRODH93
679	HS2AD53R	1	120	HS2AD53
680	HSATR92R	3	203	HSATR92
681	HSIDZG83R	5	136	HSIDZG83
682	HSICQ60R	2	118	HSICQ60
683	HSIFA64R	3	449	HSIFA64
684	HSKNN36R	108	527	HSKNN36
685	HSKYE52R	2	124	HSKYE52
686	HSLJA55R	2	169	HSLJA55

687	HSODA95R	2	169	HSODA95
688	HSPBS19R	1	372	HSPBS19
689	HSSGK43R	3	155	HSSGK43
690	HSXFJ91R	3	242	HSXFJ91
691	HTFEMB57R	168	410	HTFEMB57
692	HTGBR05R	37	138	HTGBR05
693	HTLGA72R	3	455	HTLGA72
694	HTLIX61R	1	102	HTLIX61
695	HTNTE25R	307	426	HTNTE25
696	HTWCP79R	91	180	HTWCP79
697	HTXFA64R	3	263	HTXFA64
698	HUSJF91R	218	412	HUSJF91
699	HUSJN48R	259	462	HUSJN48
700	HUSJX68R	98	493	HUSJX68
701	HUSZN23R	36	131	HUSZN23
702	IUTSD20R	104	256	IUTSD20
703	HWACH10R	66	275	HWACH10
704	HWAF163R	3	272	HWAF163
705	HWAGZ89R	176	385	HWAGZ89
706	HWBAQ20R	1	177	HWBAQ20
707	HWHHM83R	2	298	HWHHM83
708	HWLAC24R	11	133	HWLAC24
709	HWLAC81R	64	360	HWLAC81
710	HWLBF27R	3	149	HWLBF27
711	HWLBS90R	195	347	HWLBS90
712	HWLCU10R	55	120	HWLCU10
713	HWLEH13R	2	379	HWLEH13
714	HWLEJ67R	375	527	HWLEJ67
715	HWLEM49R	244	354	HWLEM49
716	HWLFP27R	2	79	HWLFP27
717	HWLGG20R	92	208	HWLGG20

718	HWLGK22R	209	373	HWLGK22
719	HWLGM21R	244	354	HWLGM21
720	HWLGP37R	8	181	HWLGP37
721	HWLGS46R	40	324	HWLGS46
722	HWLGU40R	2	202	HWLGU40
723	HWLGX65R	3	230	HWLGX65
724	HWLJD09R	2	310	HWLJD09
725	HWLHD50R	3	98	HWLHD50
726	HWLHM40R	2	208	HWLHM40
727	HWLHW89R	56	382	HWLHW89
728	HWLJD17R	64	276	HWLJD17
729	HWLIM20R	3	158	HWLIM20
730	HWLJA26R	34	135	HWLJA26
731	HWLJA28R	1	108	HWLJA28
732	HWLJG57R	240	404	HWLJG57
733	HWLJL19R	119	292	HWLJL19
734	HWLJP50R	1	147	HWLJP50
735	HWLKG82R	1	360	HWLKG82
736	HWLKG95R	1	300	HWLKG95
737	HWLKI53R	1	144	HWLKI53
738	HWLKM09R	2	100	HWLKM09
739	HWLKM86R	44	226	HWLKM86
740	HWLKM95R	2	184	HWLKM95
741	HWLKU25R	3	137	HWLKU25
742	HWLQS83R	1	117	HWLQS83
743	HWLQU65R	361	558	HWLQU65
744	HWLRL59R	1	225	HWLRL59
745	HWLRP86R	2	253	HWLRP86
746	HWLRQ49R	3	158	HWLRQ49
747	HWLUF60R	84	218	HWLUF60
748	HWLUI37R	51	263	HWLUI37

Accession	Protein Name	Length	Score	Value	Protein Name	Length	Score	Value
749	HWLUR4IR	33	155		HWLUR4I	33	155	
750	HWLVD60R	1	174		HWLVD60	1	174	
751	HWLVV50R	1	72		HWLVV50	1	72	
752	HWMAN61R	3	107		HWMAN61	3	107	
753	HWMEB47R	87	185		HWMEB47	87	185	
754	HWMEH13R	2	256		HWMEH13	2	256	
755	HWMEH26R	168	341		HWMEH26	168	341	
756	HWMEI50R	131	400		HWMEI50	131	400	
757	HWMEFB3IR	100	285		HWMEFB3I	100	285	
758	IWMFL66R	61	153		IWMFL66	61	153	
759	HWMFO93R	2	79		HWMFO93	2	79	
760	HWMP01R	120	284		HWMP01	120	284	
761	HZAAD81R	1	144		HZAAD81	1	144	
762	HWLHN70R	2	160		HWLHN70	2	160	
763	HFIXK57R	2	211		HFIXK57	2	211	
	URF 3 (NADH dehydrogenase subunit) [Homo sapiens] >gi 506832 protein 3 [Homo sapiens] >pir A00422 DNHUN3 NADH dehydrogenase (ubiquinone) (EC 1.6.5.3) chain 3 - human mitochondrion (SGCI) >sp P03897 NU3M_HUMAN NADH-UBIQUINONE OXIDOREDUCTASE CHAIN 3 (EC 1.6 URF 3 (NADH dehydrogenase subunit)	gi 13011	47	205	90	100		
764	HMAFE48R	47	205	90	HMAFE48	47	205	100

765	HRDJ88R	URF 3 (NADH dehydrogenase subunit) [Homo sapiens] >gi 506832 protein 3 [Homo sapiens] >pir A00422 DNHUN3 NADH dehydrogenase (ubiquinone) (EC 1.6.5.3) chain 3 - human mitochondrion (SGC1) >sp P03897 NU3M_HUMAN NADH-UBIQUINONE	gi 13011	55	213	83	94	HRDJ88
766	HWLAR3IR	OXIDOREDUCTASE CHAIN 3 (EC 1.6 URF 3 (NADH dehydrogenase subunit) [Homo sapiens] >gi 506832 protein 3 [Homo sapiens] >pir A00422 DNHUN3 NADH dehydrogenase (ubiquinone) (EC 1.6.5.3) chain 3 - human mitochondrion (SGC1) >sp P03897 NU3M_HUMAN NADH-UBIQUINONE	gi 13011	56	214	91	100	HWLAR3I
767	HNHLH26R	OXIDOREDUCTASE CHAIN 3 (EC 1.6 v-SNARE [Cricetulus griseus] >sp O08522 O08522 v-SNARE. Length = 250	gi 1912453	73	243	64	76	HNHLH26
768	H2LAU24R	weakly similar to gastrula zinc finger protein [Caenorhabditis elegans] >sp Q09998 Q09998 PUTATIVE 55.5 KD ZINC FINGER PROTEIN R144.3 IN CHROMOSOME III. Length = 492	gi 746495	78	488	45	60	H2LAU24
769	HATDR94R	X box binding protein-1 [Homo sapiens] >pir A36299 A36299 transcription factor hXBP-1 - human Length = 260	gi 306893	2	367	95	100	HATDR94
770	HWLLI85R	X-linked deafness dystonia protein [Homo sapiens] >sp O60220 O60220 X- LINKED DEAFNESS DYSTONIA PROTEIN. Length = 97	gi 3123843	410	580	60	80	HWLLI85

771	HBHMF67R	XP-C repair complementing protein (p58/11HR23B) [Homo sapiens] >pir S44346 S44346 RAI23 protein homolog - human Length = 409	gnl PID d1005181	3	191	96	96	HBHMF67
772	HSYCH41R	yeast methionyl-tRNA synthetase homolog [Homo sapiens] >pir JC5224 JC5224 methionine--tRNA ligase (EC 6.1.1.10) - human >gi 804996 mitoxantrone-resistance associated gene [Homo sapiens] {SUB 423-900} Length = 900	gnl PID e218477	2	373	90	90	HSYCH41
773	HWLJR53R	zinc finger protein PZF [Mus musculus] >pir 48724 48724 zinc finger protein PZF - mouse >sp Q62511 Q62511 ZINC FINGER PROTEIN PZF: Length = 455	gi 453376	1	552	81	83	HWLJR53

The first column of Table 1 shows the "SEQ ID NO:" for each of the 773 colon cancer antigen polynucleotide sequences of the invention.

The second column in Table 1, provides a unique "Sequence/Contig ID" identification for each colon and/or colon cancer associated sequence. The third column in Table 1, "Gene Name," provides a putative identification of the gene based on the sequence similarity of its translation product to an amino acid sequence found in a publicly accessible gene database, such as GenBank (NCBI). The great majority of the cDNA sequences reported in Table 1 are unrelated to any sequences previously described in the literature. The fourth column, in Table 1, "Overlap," provides the database accession no. for the database sequence having similarity. The fifth and sixth columns in Table 1 provide the location (nucleotide position nos. within the contig), "Start" and "End", in the polynucleotide sequence "SEQ ID NO:X" that delineate the preferred ORF shown in the sequence listing as SEQ ID NO:Y. In one embodiment, the invention provides a protein comprising, or alternatively consisting of, a polypeptide encoded by the portion of SEQ ID NO:X delineated by the nucleotide position nos. "Start" and "End". Also provided are polynucleotides encoding such proteins and the complementary strand thereto. The seventh and eighth columns provide the "% Identity" (percent identity) and "% Similarity" (percent similarity) observed between the aligned sequence segments of the translation product of SEQ ID NO:X and the database sequence.

The ninth column of Table 1 provides a unique "Clone ID" for a clone related to each contig sequence. This clone ID references the cDNA clone which contains at least the 5' most sequence of the assembled contig and at least a portion of SEQ ID NO:X was determined by directly sequencing the referenced clone. The reference clone may have more sequence than described in the sequence listing or the clone may have less. In the vast majority of cases, however, the clone is believed to encode a full-length polypeptide. In the case where a clone is not full-length, a full-length cDNA can be obtained by methods described elsewhere herein.

Table 3 indicates public ESTs, of which at least one, two, three, four, five, ten, or more of any one or more of these public ESTs are optionally excluded from the invention.

SEQ ID NO:X (where X may be any of the polynucleotide sequences disclosed in the sequence listing as SEQ ID NO:1 through SEQ ID NO:773) and the translated SEQ ID NO:Y (where Y may be any of the polypeptide sequences disclosed in the sequence listing as SEQ ID NO:774 through SEQ ID NO:1546) are sufficiently accurate and otherwise suitable for a

variety of uses well known in the art and described further below. For instance, SEQ ID NO:X has uses including, but not limited to, in designing nucleic acid hybridization probes that will detect nucleic acid sequences contained in SEQ ID NO:X or the related cDNA clone contained in a library deposited with the ATCC. These probes will also hybridize to nucleic acid molecules in biological samples, thereby enabling immediate applications in chromosome mapping, linkage analysis, tissue identification and/or typing, and a variety of forensic and diagnostic methods of the invention. Similarly, polypeptides identified from SEQ ID NO:Y have uses that include, but are not limited to, generating antibodies which bind specifically to the colon cancer antigen polypeptides, or fragments thereof, and/or to the colon cancer antigen polypeptides encoded by the cDNA clones identified in Table 1.

Nevertheless, DNA sequences generated by sequencing reactions can contain sequencing errors. The errors exist as misidentified nucleotides, or as insertions or deletions of nucleotides in the generated DNA sequence. The erroneously inserted or deleted nucleotides cause frame shifts in the reading frames of the predicted amino acid sequence. In these cases, the predicted amino acid sequence diverges from the actual amino acid sequence, even though the generated DNA sequence may be greater than 99.9% identical to the actual DNA sequence (for example, one base insertion or deletion in an open reading frame of over 1000 bases).

Accordingly, for those applications requiring precision in the nucleotide sequence or the amino acid sequence, the present invention provides not only the generated nucleotide sequence identified as SEQ ID NO:X, the predicted translated amino acid sequence identified as SEQ ID NO:Y, but also a sample of plasmid DNA containing the related cDNA clone (deposited with the ATCC, as set forth in Table 1). The nucleotide sequence of each deposited clone can readily be determined by sequencing the deposited clone in accordance with known methods. Further, techniques known in the art can be used to verify the nucleotide sequences of SEQ ID NO:X.

The predicted amino acid sequence can then be verified from such deposits. Moreover, the amino acid sequence of the protein encoded by a particular clone can also be directly determined by peptide sequencing or by expressing the protein in a suitable host cell containing the deposited human cDNA, collecting the protein, and determining its sequence.

The present invention also relates to vectors or plasmids which include such DNA sequences, as well as the use of the DNA sequences. The material deposited with the ATCC on:

5 **Table 2**

ATCC Deposits	Deposit Date	ATCC Designation Number
LP01, LP02, LP03, LP04, LP05, LP06, LP07, LP08, LP09, LP10, LP11,	May-20-97	209059, 209060, 209061, 209062, 209063, 209064, 209065, 209066, 209067, 209068, 209069
LP12	Jan-12-98	209579
LP13	Jan-12-98	209578
LP14	Jul-16-98	203067
LP15	Jul-16-98	203068
LP16	Feb-1-99	203609
LP17	Feb-1-99	203610
LP20	Nov-17-98	203485
LP21	Jun-18-99	PTA-252
LP22	Jun-18-99	PTA-253
LP23	Dec-22-99	PTA-1081

each is a mixture of cDNA clones derived from a variety of human tissue and cloned in either a plasmid vector or a phage vector, as shown in Table 5. These deposits are referred to as "the deposits" herein. The tissues from which the clones were derived are listed in Table 5, and the vector in which the cDNA is contained is also indicated in Table 5. The deposited material includes the cDNA clones which were partially sequenced and are related to the SEQ ID NO:X described in Table 1 (column 9). Thus, a clone which is isolatable from the ATCC Deposits by use of a sequence listed as SEQ ID NO:X may include the entire coding region of a human gene or in other cases such clone may include a substantial portion of the coding region of a human gene. Although the sequence listing lists only a portion of the DNA sequence in a clone included in the ATCC Deposits, it is well within the ability of one skilled in the art to complete the sequence of the DNA included in a clone isolatable from the

ATCC Deposits by use of a sequence (or portion thereof) listed in Table 1 by procedures hereinafter further described, and others apparent to those skilled in the art.

Also provided in Table 5 is the name of the vector which contains the cDNA clone. Each vector is routinely used in the art. The following additional information is provided for convenience.

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128, 256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., *Nucleic Acids Res.* 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., *Nucleic Acids Res.* 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., *Strategies* 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Phagemid pBS may be excised from the Lambda Zap and Uni-Zap XR vectors, and phagemid pBK may be excised from the Zap Express vector. Both phagemids may be transformed into *E. coli* strain XL-1 Blue, also available from Stratagene.

Vectors pSport1, pCMVSPORT 1.0, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, also available from Life Technologies. See, for instance, Gruber, C. E., et al., *Focus* 15:59 (1993). Vector lafmid BA (Bento Soares, Columbia University, New York, NY) contains an ampicillin resistance gene and can be transformed into *E. coli* strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an ampicillin resistance gene and may be transformed into *E. coli* strain DH10B, available from Life Technologies. See, for instance, Clark, J. M., *Nuc. Acids Res.* 16:9677-9686 (1988) and Mead, D. et al., *Bio/Technology* 9: (1991).

The present invention also relates to the genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, and/or the cDNA contained in a deposited cDNA clone. The corresponding gene can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include, but are not limited to, preparing probes or primers from the disclosed sequence and identifying or amplifying the corresponding gene from appropriate sources of genomic material.

Also provided in the present invention are allelic variants, orthologs, and/or species homologs. Procedures known in the art can be used to obtain full-length genes, allelic variants, splice variants, full-length coding portions, orthologs, and/or species homologs of genes corresponding to SEQ ID NO:X, SEQ ID NO:Y, and/or the cDNA contained in the related cDNA clone in the deposit, using information from the sequences disclosed herein or the clones deposited with the ATCC. For example, allelic variants and/or species homologs may be isolated and identified by making suitable probes or primers from the sequences provided herein and screening a suitable nucleic acid source for allelic variants and/or the desired homologue.

The present invention provides a polynucleotide comprising, or alternatively consisting of, the nucleic acid sequence of SEQ ID NO:X, and/or the related cDNA clone (See, e.g., columns 1 and 9 of Table 1). The present invention also provides a polypeptide comprising, or alternatively, consisting of, the polypeptide sequence of SEQ ID NO:Y, a polypeptide encoded by SEQ ID NO:X, and/or a polypeptide encoded by the cDNA in the related cDNA clone contained in a deposited library. Polynucleotides encoding a polypeptide comprising, or alternatively consisting of, the polypeptide sequence of SEQ ID NO:Y, a polypeptide encoded by SEQ ID NO:X, and/or a polypeptide encoded by the cDNA in the related cDNA clone contained in a deposited library, are also encompassed by the invention. The present invention further encompasses a polynucleotide comprising, or alternatively consisting of, the complement of the nucleic acid sequence of SEQ ID NO:X, and/or the complement of the coding strand of the related cDNA clone contained in a deposited library.

Many polynucleotide sequences, such as EST sequences, are publicly available and accessible through sequence databases and may have been publicly available prior to conception of the present invention. Preferably, such related polynucleotides are specifically excluded from the scope of the present invention. To list every related sequence would unduly burden the disclosure of this application. Accordingly, for each "Contig Id" listed in the first column of Table 3, preferably excluded are one or more polynucleotides comprising a nucleotide sequence described in the second column of Table 3 by the general formula of a-b, each of which are uniquely defined for the SEQ ID NO:X corresponding to that Contig Id in Table 1. Additionally, specific embodiments are directed to polynucleotide sequences excluding at least one, two, three, four, five, ten, or more of the specific polynucleotide sequences referenced by the Genbank Accession No. for each Contig Id which may be

included in column 3 of Table 3. In no way is this listing meant to encompass all of the sequences which may be excluded by the general formula, it is just a representative example.

Table 3.

Sequence/ Contig ID	General formula	Genbank Accession No.
500802	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 619 of SEQ ID NO:1, b is an integer of 15 to 633, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:1, and where b is greater than or equal to a + 14.	
531091	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 281 of SEQ ID NO:2, b is an integer of 15 to 295, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:2, and where b is greater than or equal to a + 14.	
553147	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 428 of SEQ ID NO:3, b is an integer of 15 to 442, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:3, and where b is greater than or equal to a + 14.	
558860	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 740 of SEQ ID NO:4, b is an integer of 15 to 754, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:4, and where b is greater than or equal to a + 14.	
561730	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 379 of SEQ ID NO:5, b is an integer of 15 to 393, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:5, and where b is greater than or equal to a + 14.	
585938	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 525 of SEQ ID NO:6, b is an integer of 15 to 539, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:6, and where b is greater than or equal to a + 14.	
587785	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 790 of SEQ ID	

	NO:7, b is an integer of 15 to 804, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:7, and where b is greater than or equal to a + 14.	
588916	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 706 of SEQ ID NO:8, b is an integer of 15 to 720, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:8, and where b is greater than or equal to a + 14.	
613825	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 526 of SEQ ID NO:9, b is an integer of 15 to 540, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:9, and where b is greater than or equal to a + 14.	
639090	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 547 of SEQ ID NO:10, b is an integer of 15 to 561, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:10, and where b is greater than or equal to a + 14.	
651644	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 379 of SEQ ID NO:11, b is an integer of 15 to 393, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:11, and where b is greater than or equal to a + 14.	
659544	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 308 of SEQ ID NO:12, b is an integer of 15 to 322, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:12, and where b is greater than or equal to a + 14.	
659739	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1893 of SEQ ID NO:13, b is an integer of 15 to 1907, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:13, and where b is greater than or equal to a + 14.	
661057	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1126 of SEQ ID NO:14, b is an integer of 15 to 1140, where both a	

	and b correspond to the positions of nucleotide residues shown in SEQ ID NO:14, and where b is greater than or equal to a + 14.	
661313	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1994 of SEQ ID NO:15, b is an integer of 15 to 2008, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:15, and where b is greater than or equal to a + 14.	
666316	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 357 of SEQ ID NO:16, b is an integer of 15 to 371, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:16, and where b is greater than or equal to a + 14.	
669229	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 749 of SEQ ID NO:17, b is an integer of 15 to 763, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:17, and where b is greater than or equal to a + 14.	
670471	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1912 of SEQ ID NO:18, b is an integer of 15 to 1926, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:18, and where b is greater than or equal to a + 14.	
676611	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2287 of SEQ ID NO:19, b is an integer of 15 to 2301, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:19, and where b is greater than or equal to a + 14.	
691240	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 524 of SEQ ID NO:20, b is an integer of 15 to 538, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:20, and where b is greater than or equal to a + 14.	
702977	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1389 of SEQ ID NO:21, b is an integer of 15 to 1403, where both a and b correspond to the positions of nucleotide	

	residues shown in SEQ ID NO:21. and where b is greater than or equal to $a + 14$.	
709517	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 464 of SEQ ID NO:22, b is an integer of 15 to 478, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:22, and where b is greater than or equal to $a + 14$.	
714730	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1238 of SEQ ID NO:23, b is an integer of 15 to 1252, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:23, and where b is greater than or equal to $a + 14$.	
714834	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1060 of SEQ ID NO:24, b is an integer of 15 to 1074, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:24, and where b is greater than or equal to $a + 14$.	
715016	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1172 of SEQ ID NO:25, b is an integer of 15 to 1186, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:25, and where b is greater than or equal to $a + 14$.	
719584	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 874 of SEQ ID NO:26, b is an integer of 15 to 888, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:26, and where b is greater than or equal to $a + 14$.	
724637	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 775 of SEQ ID NO:27, b is an integer of 15 to 789, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:27, and where b is greater than or equal to $a + 14$.	
728392	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 833 of SEQ ID NO:28, b is an integer of 15 to 847, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:28, and where b is greater than	

	or equal to $a + 14$.	
738716	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 652 of SEQ ID NO:29, b is an integer of 15 to 666, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:29, and where b is greater than or equal to $a + 14$.	
739056	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 503 of SEQ ID NO:30, b is an integer of 15 to 517, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:30, and where b is greater than or equal to $a + 14$.	
739143	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2661 of SEQ ID NO:31, b is an integer of 15 to 2675, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:31, and where b is greater than or equal to $a + 14$.	
742329	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 263 of SEQ ID NO:32, b is an integer of 15 to 277, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:32, and where b is greater than or equal to $a + 14$.	
742557	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 907 of SEQ ID NO:33, b is an integer of 15 to 921, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:33, and where b is greater than or equal to $a + 14$.	
745481	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1453 of SEQ ID NO:34, b is an integer of 15 to 1467, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:34, and where b is greater than or equal to $a + 14$.	
746035	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2063 of SEQ ID NO:35, b is an integer of 15 to 2077, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:35, and where b is greater than or equal to $a + 14$.	

753731	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 370 of SEQ ID NO:36, b is an integer of 15 to 384, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:36, and where b is greater than or equal to a + 14.	
754383	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 454 of SEQ ID NO:37, b is an integer of 15 to 468, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:37, and where b is greater than or equal to a + 14.	
756749	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1081 of SEQ ID NO:38, b is an integer of 15 to 1095, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:38, and where b is greater than or equal to a + 14.	
757980	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1743 of SEQ ID NO:39, b is an integer of 15 to 1757, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:39, and where b is greater than or equal to a + 14.	R38216, R63249, R78721, H01441, H02557, H02640, H86258, H86321, N21599, W16868, W31882, W56228, N90610, AA047227, AA056107, AA058568, AA100609, AA115890
764818	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1931 of SEQ ID NO:40, b is an integer of 15 to 1945, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:40, and where b is greater than or equal to a + 14.	
765140	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 574 of SEQ ID NO:41, b is an integer of 15 to 588, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:41, and where b is greater than or equal to a + 14.	
766893	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1554 of SEQ ID NO:42, b is an integer of 15 to 1568, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:42, and where b is greater than or equal to a + 14.	R69702, R76994, R77002, H01357
771338	Preferably excluded from the present invention are	

	one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1046 of SEQ ID NO:43, b is an integer of 15 to 1060, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:43, and where b is greater than or equal to a + 14.	
771412	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1330 of SEQ ID NO:44, b is an integer of 15 to 1344, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:44, and where b is greater than or equal to a + 14.	
772226	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 878 of SEQ ID NO:45, b is an integer of 15 to 892, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:45, and where b is greater than or equal to a + 14.	
773057	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 482 of SEQ ID NO:46, b is an integer of 15 to 496, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:46, and where b is greater than or equal to a + 14.	N41725
773173	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1215 of SEQ ID NO:47, b is an integer of 15 to 1229, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:47, and where b is greater than or equal to a + 14.	
780154	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1397 of SEQ ID NO:48, b is an integer of 15 to 1411, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:48, and where b is greater than or equal to a + 14.	
780768	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1671 of SEQ ID NO:49, b is an integer of 15 to 1685, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:49, and where b is greater than or equal to a + 14.	
780779	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide	

	sequence described by the general formula of a-b, where a is any integer between 1 to 646 of SEQ ID NO:50, b is an integer of 15 to 660, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:50, and where b is greater than or equal to a + 14.	
782394	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1558 of SEQ ID NO:51, b is an integer of 15 to 1572, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:51, and where b is greater than or equal to a + 14.	R24689, R25853, R34457, R66839, R68536, H22874, H45555, N50184, AA015963, AA028939, AA028938
783160	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 621 of SEQ ID NO:52, b is an integer of 15 to 635, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:52, and where b is greater than or equal to a + 14.	
783506	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1353 of SEQ ID NO:53, b is an integer of 15 to 1367, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:53, and where b is greater than or equal to a + 14.	
784446	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 364 of SEQ ID NO:54, b is an integer of 15 to 378, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:54, and where b is greater than or equal to a + 14.	
784832	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1044 of SEQ ID NO:55, b is an integer of 15 to 1058, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:55, and where b is greater than or equal to a + 14.	
786813	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 668 of SEQ ID NO:56, b is an integer of 15 to 682, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:56, and where b is greater than or equal to a + 14.	W44740, AA235981
792139	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b.	

	where a is any integer between 1 to 630 of SEQ ID NO:57, b is an integer of 15 to 644, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:57, and where b is greater than or equal to a + 14.	
793987	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 752 of SEQ ID NO:58, b is an integer of 15 to 766, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:58, and where b is greater than or equal to a + 14.	
805715	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2347 of SEQ ID NO:59, b is an integer of 15 to 2361, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:59, and where b is greater than or equal to a + 14.	
811111	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1458 of SEQ ID NO:60, b is an integer of 15 to 1472, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:60, and where b is greater than or equal to a + 14.	R11325, R11326, R43655, R43655, R72437, R78096, H23850, N20947, N22686, N25829, N27270, N31401, N40002, N46020, W92748, W92871, AA461202, AA461382
811113	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1658 of SEQ ID NO:61, b is an integer of 15 to 1672, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:61, and where b is greater than or equal to a + 14.	
823902	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1526 of SEQ ID NO:62, b is an integer of 15 to 1540, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:62, and where b is greater than or equal to a + 14.	
826518	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1030 of SEQ ID NO:63, b is an integer of 15 to 1044, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:63, and where b is greater than or equal to a + 14.	T60163, T60223, T61894, R12251, T81471, T81679, T95899, R98321, R98322, H52605, H59085, N27268, N31506, N53499, N54486, N58236, N92460, AA027189, AA045077, AA127016, AA418935, AA426582
826704	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 837 of SEQ ID	

	NO:64. b is an integer of 15 to 851, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:64. and where b is greater than or equal to a + 14.	
827720	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2779 of SEQ ID NO:65. b is an integer of 15 to 2793, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:65. and where b is greater than or equal to a + 14.	
828102	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 289 of SEQ ID NO:66. b is an integer of 15 to 303. where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:66. and where b is greater than or equal to a + 14.	
828180	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1396 of SEQ ID NO:67. b is an integer of 15 to 1410. where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:67. and where b is greater than or equal to a + 14.	
828386	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1010 of SEQ ID NO:68. b is an integer of 15 to 1024. where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:68. and where b is greater than or equal to a + 14.	
828658	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1834 of SEQ ID NO:69. b is an integer of 15 to 1848. where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:69. and where b is greater than or equal to a + 14.	
828919	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2668 of SEQ ID NO:70. b is an integer of 15 to 2682. where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:70. and where b is greater than or equal to a + 14.	T66771, T66772, T71638, R08935, R09044, R09373, T80114, T85695, R00758, R00759, R12645, R19577, R20545, R22041, R22097, R20545, R59701, R59811, R60034, R60096, R60694, R76255, R81371, R81370, H04390, H04415, H05912, H47622, H47647, R83679, H71735, H72298, N25487, N35542, N49731, N52660, N67681, N75596, W03490, AA044638, AA044702, AA165090, AA164628, AA215698, AA215699, AA233182, AA233196, AA236759, AA256822, AA429489.

		AA428534
829572	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 398 of SEQ ID NO:71, b is an integer of 15 to 412, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:71, and where b is greater than or equal to a + 14.	T63032
830138	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1347 of SEQ ID NO:72, b is an integer of 15 to 1361, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:72, and where b is greater than or equal to a + 14.	
830208	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 914 of SEQ ID NO:73, b is an integer of 15 to 928, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:73, and where b is greater than or equal to a + 14.	R01611, N76461, W74577, W79757, AA045350, AA056064, AA190524
830248	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1172 of SEQ ID NO:74, b is an integer of 15 to 1186, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:74, and where b is greater than or equal to a + 14.	
830275	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 919 of SEQ ID NO:75, b is an integer of 15 to 933, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:75, and where b is greater than or equal to a + 14.	
830286	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1950 of SEQ ID NO:76, b is an integer of 15 to 1964, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:76, and where b is greater than or equal to a + 14.	T90376, R46154, R46154, AA224239, AA467906, AA483293, AA502593, AA513313, AA594445, AA594570, AA594876, AA579404, AA720893, AA767344, AA857646, AA877489, AA954868, AA991634, A1014751, C02074, AA093141
830347	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1788 of SEQ ID NO:77, b is an integer of 15 to 1802, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:77, and where b is greater than or equal to a + 14.	

830348	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 981 of SEQ ID NO:78, b is an integer of 15 to 995, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:78, and where b is greater than or equal to a + 14.	AA983601
830364	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1201 of SEQ ID NO:79, b is an integer of 15 to 1215, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:79, and where b is greater than or equal to a + 14.	
830394	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2646 of SEQ ID NO:80, b is an integer of 15 to 2660, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:80, and where b is greater than or equal to a + 14.	
830398	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1776 of SEQ ID NO:81, b is an integer of 15 to 1790, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:81, and where b is greater than or equal to a + 14.	
830412	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1336 of SEQ ID NO:82, b is an integer of 15 to 1350, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:82, and where b is greater than or equal to a + 14.	
830436	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1732 of SEQ ID NO:83, b is an integer of 15 to 1746, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:83, and where b is greater than or equal to a + 14.	T89041, R38418, R51559, R62385, R63785, H21426, N55384, AA009460, AA039527, AA039526, AA490811, AA588539, AA574253, AA827525, AA975094, D79482, D79908, N55964, C14631, C14891, C14892
830464	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1477 of SEQ ID NO:84, b is an integer of 15 to 1491, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:84, and where b is greater than or equal to a + 14.	H06247, H19227, W52470
830471	Preferably excluded from the present invention are	R28064, R28282, AA143044, AA151127.

	one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 954 of SEQ ID NO:85, b is an integer of 15 to 968, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:85, and where b is greater than or equal to a + 14.	AA165093, AA164631, AA256943, AA765384, D80554
830477	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3054 of SEQ ID NO:86, b is an integer of 15 to 3068, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:86, and where b is greater than or equal to a + 14.	T71686, R81413, R81414, H52583, H84987, H87923, H88319, H88319, W74073, W79680, AA021098, AA179389, AA182649, AA188175, AA191449, AA228943, AA228942, AA594459, AA737972, C02737
830500	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2216 of SEQ ID NO:87, b is an integer of 15 to 2230, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:87, and where b is greater than or equal to a + 14.	
830509	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1149 of SEQ ID NO:88, b is an integer of 15 to 1163, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:88, and where b is greater than or equal to a + 14.	
830528	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1925 of SEQ ID NO:89, b is an integer of 15 to 1939, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:89, and where b is greater than or equal to a + 14.	
830542	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2018 of SEQ ID NO:90, b is an integer of 15 to 2032, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:90, and where b is greater than or equal to a + 14.	T60268, T61648, T68371, T88743, R00503, R13392, R40908, R40908, H02114, H07926, H29767, H29768, H38826, H93354, W42415, W42513, W61060, W72566, W76560, AA011078, AA011079, AA031697, AA031863, AA058529, AA100913, AA100912, AA129619, AA129593, AA129330, AA128581, AA160087, AA160675, AA173629, AA173985, AA186698, AA188326, AA480672, AA587251, AA576938, AA743161, AA834774, AA872783, AA877207, AA878505, AA923685, AA934427, AA962214, AA995455, AA995857, N88876
830564	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b.	

	where a is any integer between 1 to 1774 of SEQ ID NO:91, b is an integer of 15 to 1788, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:91, and where b is greater than or equal to a + 14.	
830611	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 481 of SEQ ID NO:92, b is an integer of 15 to 495, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:92, and where b is greater than or equal to a + 14.	
830618	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1363 of SEQ ID NO:93, b is an integer of 15 to 1377, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:93, and where b is greater than or equal to a + 14.	R43709, R43709, H09113, H43746, N92632, AA022453, AA120876, AA120889, AA493651, AA493785, AA494347, AA565392, AA743179, AA769161
830620	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2805 of SEQ ID NO:94, b is an integer of 15 to 2819, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:94, and where b is greater than or equal to a + 14.	
830630	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 691 of SEQ ID NO:95, b is an integer of 15 to 705, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:95, and where b is greater than or equal to a + 14.	
830654	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3458 of SEQ ID NO:96, b is an integer of 15 to 3472, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:96, and where b is greater than or equal to a + 14.	
830660	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1202 of SEQ ID NO:97, b is an integer of 15 to 1216, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:97, and where b is greater than or equal to a + 14.	
830661	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1172 of SEQ ID	

	NO:98, b is an integer of 15 to 1186, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:98, and where b is greater than or equal to a + 14.	
830704	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1106 of SEQ ID NO:99, b is an integer of 15 to 1120, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:99, and where b is greater than or equal to a + 14.	
830765	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1211 of SEQ ID NO:100, b is an integer of 15 to 1225, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:100, and where b is greater than or equal to a + 14.	
830778	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1199 of SEQ ID NO:101, b is an integer of 15 to 1213, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:101, and where b is greater than or equal to a + 14.	
830784	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1550 of SEQ ID NO:102, b is an integer of 15 to 1564, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:102, and where b is greater than or equal to a + 14.	R63323, R66534, AA491630
830800	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1443 of SEQ ID NO:103, b is an integer of 15 to 1457, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:103, and where b is greater than or equal to a + 14.	
830821	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 771 of SEQ ID NO:104, b is an integer of 15 to 785, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:104, and where b is greater than or equal to a + 14.	
830849	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 907 of SEQ ID NO:105, b is an integer of 15 to 921, where both a	AA258128, AA259034, AA262104, AA742612, AA804402

	and b correspond to the positions of nucleotide residues shown in SEQ ID NO:105, and where b is greater than or equal to a + 14.	
830903	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 578 of SEQ ID NO:106, b is an integer of 15 to 592, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:106, and where b is greater than or equal to a + 14.	
830913	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2234 of SEQ ID NO:107, b is an integer of 15 to 2248, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:107, and where b is greater than or equal to a + 14.	R06463, R06517, R48006, R51455, R61502, R72398, R72399, R74489, R74599, H07933, H08039, H61149, H62056, H90758, H90809, N32837, N42283, W40284, W45325, AA079353, AA079592, AA100814, AA102342, AA111844, AA122150, AA134127, AA134128, AA148738, AA148709, AA164240, AA164899, AA164275, AA171881, AA179310, AA179453, AA180811, AA180955, AA187432, AA190377, AA190791, AA190383, AA458475, AA427428, AA468548, AA554518, AA595768, AA595893, AA640601, AA574035, AA658143, AA863401, AA906604, AA995159, C03746, C04875, C05396, AA033510
830920	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 771 of SEQ ID NO:108, b is an integer of 15 to 785, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:108, and where b is greater than or equal to a + 14.	
830938	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 597 of SEQ ID NO:109, b is an integer of 15 to 611, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:109, and where b is greater than or equal to a + 14.	AA053612
830980	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 650 of SEQ ID NO:110, b is an integer of 15 to 664, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:110, and where b is greater than or equal to a + 14.	
831014	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 4051 of SEQ ID NO:111, b is an integer of 15 to 4065, where both a	

	and b correspond to the positions of nucleotide residues shown in SEQ ID NO:111, and where b is greater than or equal to a + 14.	
831026	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1478 of SEQ ID NO:112, b is an integer of 15 to 1492, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:112, and where b is greater than or equal to a + 14.	
831031	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1468 of SEQ ID NO:113, b is an integer of 15 to 1482, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:113, and where b is greater than or equal to a + 14.	R46004, R46004, H06850, N27532, N30567, N30842, N34647, N40349, N41369, N49777, N52708, N62958, W68355, W68490, AA054602, AA193410, AA193648, AA503204, AA688236, AA730103, AA736540, AA747555, AA811522, AA863169, N79861
831055	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3717 of SEQ ID NO:114, b is an integer of 15 to 3731, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:114, and where b is greater than or equal to a + 14.	
831057	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1301 of SEQ ID NO:115, b is an integer of 15 to 1315, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:115, and where b is greater than or equal to a + 14.	R69415, R69546, H14127, H62767, N62927, N63320, W00649, W01189, AA053293, AA058396, AA149075, AA458528, AA418699, AA418770, AA505598, AA576507, AA730033, AA805864, AA988279, AA991217, D82661, C21298
831062	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1306 of SEQ ID NO:116, b is an integer of 15 to 1320, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:116, and where b is greater than or equal to a + 14.	
831117	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2011 of SEQ ID NO:117, b is an integer of 15 to 2025, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:117, and where b is greater than or equal to a + 14.	R80585, R80586, N49020, AA173625, AA173981, AA557142, AA627866, AA847195, A1015673
831122	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1281 of SEQ ID NO:118, b is an integer of 15 to 1295, where both a and b correspond to the positions of nucleotide	R72079, R72128, AA715820, AA804163, AA809123, AA641490

	residues shown in SEQ ID NO:118, and where b is greater than or equal to a + 14.	
831125	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1243 of SEQ ID NO:119, b is an integer of 15 to 1257, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:119, and where b is greater than or equal to a + 14.	N80647, AA114140, AA143553, AA156386, N68188, AA070867
831132	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 383 of SEQ ID NO:120, b is an integer of 15 to 397, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:120, and where b is greater than or equal to a + 14.	
831152	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 862 of SEQ ID NO:121, b is an integer of 15 to 876, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:121, and where b is greater than or equal to a + 14.	AA765155
831157	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1264 of SEQ ID NO:122, b is an integer of 15 to 1278, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:122, and where b is greater than or equal to a + 14.	T57943, R34275, R35472, R77406, R77405, N23203, N59015, AA160841, AA610280, AA857624, A1089936, A1094724, A1094954
831160	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3101 of SEQ ID NO:123, b is an integer of 15 to 3115, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:123, and where b is greater than or equal to a + 14.	
831193	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 365 of SEQ ID NO:124, b is an integer of 15 to 379, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:124, and where b is greater than or equal to a + 14.	
831197	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1253 of SEQ ID NO:125, b is an integer of 15 to 1267, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:125, and where b is	AA134613

	greater than or equal to $a + 14$.	
831217	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 827 of SEQ ID NO:126, b is an integer of 15 to 841, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:126, and where b is greater than or equal to $a + 14$.	
831239	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1158 of SEQ ID NO:127, b is an integer of 15 to 1172, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:127, and where b is greater than or equal to $a + 14$.	T68487, T88923, T88994, R09550, R09663, R26714, R26937, H27046, H28228, H30272, H30335, N27966, N36884, N46156, N93575, W21407, W44513, W44514, W47626, W47627, W56215, W60528, W80465, W80574, W92729, AA002237, AA002076, AA099290, AA099291, AA127753, AA127706, AA128275, AA128572, AA148737, AA149497, AA419078, AA423819, AA506117, AA534694, AA552105, AA552219, AA583468, AA622094, AA633205, AA878663, AA911544, AA916173, AA974873, AA988860, AI056396, AI074163, W92753
831248	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 877 of SEQ ID NO:128, b is an integer of 15 to 891, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:128, and where b is greater than or equal to $a + 14$.	
831313	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2447 of SEQ ID NO:129, b is an integer of 15 to 2461, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:129, and where b is greater than or equal to $a + 14$.	T61093, T97774, R13148, R31511, R32943, R33906, R33921, R37053, R44148, R44148, R74449, R79209, R79476, H12271, H27631, H30122, R84834, H63166, H71003, H71015, H83387, N23726, N23730, N23773, N52416, N66497, N67917, N68137, N73801, N99428, W95944, AA018712, AA020879, AA429721, AA470397, AA493243, AA507952, AA515358, AA583463, AA617991, AA618186, AA631437, AA566089, AA746085, AA837997, AA878863, AA922678, AA985597, AA947992, AI074096, C03207, C17030, C18106
831369	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2183 of SEQ ID NO:130, b is an integer of 15 to 2197, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:130, and where b is greater than or equal to $a + 14$.	
831371	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide	

	sequence described by the general formula of a-b, where a is any integer between 1 to 450 of SEQ ID NO:131, b is an integer of 15 to 464, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:131, and where b is greater than or equal to a + 14.	
831373	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1936 of SEQ ID NO:132, b is an integer of 15 to 1950, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:132, and where b is greater than or equal to a + 14.	T50786, T50949, T53797, T53916, T64650, T71681, T71836, T71876, T71877, T74596, T74656, H30426, H46449, H46671, H46670, H46990, H50500, AA419051, AA423809, AA928986
831387	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2079 of SEQ ID NO:133, b is an integer of 15 to 2093, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:133, and where b is greater than or equal to a + 14.	
831410	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 715 of SEQ ID NO:134, b is an integer of 15 to 729, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:134, and where b is greater than or equal to a + 14.	
831448	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1175 of SEQ ID NO:135, b is an integer of 15 to 1189, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:135, and where b is greater than or equal to a + 14.	
831450	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1452 of SEQ ID NO:136, b is an integer of 15 to 1466, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:136, and where b is greater than or equal to a + 14.	
831472	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 126 of SEQ ID NO:137, b is an integer of 15 to 140, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:137, and where b is greater than or equal to a + 14.	
831473	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b,	

	where a is any integer between 1 to 4128 of SEQ ID NO:138, b is an integer of 15 to 4142, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:138, and where b is greater than or equal to a + 14.	
831474	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1733 of SEQ ID NO:139, b is an integer of 15 to 1747, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:139, and where b is greater than or equal to a + 14.	T66054, T89542, R10967, T78297, T83524, T97793, R13138, H08701, H10662, R82956, R96295, R98912, H66237, H79525, N31425, N36736, W76142, W81053, AA010227, AA011652, AA057613, AA057653, AA069088, AA083946, AA084193, AA126186, H70618, H79526, W72916, W80802, AA011433, AA057699, AA057752, AA069023
831494	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1226 of SEQ ID NO:140, b is an integer of 15 to 1240, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:140, and where b is greater than or equal to a + 14.	H14081, H14102, N34979, N42213, N43740, N68241, W69584, W69583, AA507828, AA877181, AA975100, A1000204
831506	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 657 of SEQ ID NO:141, b is an integer of 15 to 671, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:141, and where b is greater than or equal to a + 14.	AA035596, AA577792, AA903617, AA972775, AA996054, C00084
831533	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3251 of SEQ ID NO:142, b is an integer of 15 to 3265, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:142, and where b is greater than or equal to a + 14.	
831539	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 751 of SEQ ID NO:143, b is an integer of 15 to 765, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:143, and where b is greater than or equal to a + 14.	
831556	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1680 of SEQ ID NO:144, b is an integer of 15 to 1694, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:144, and where b is greater than or equal to a + 14.	H01879, H01880, H43546, H43547, H43548, N58813, N75148, AA428902, AA429101, AA278337, AA662009, AA928907, AA988624
831594	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide	

	sequence described by the general formula of a-b, where a is any integer between 1 to 809 of SEQ ID NO:145, b is an integer of 15 to 823, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:145, and where b is greater than or equal to a + 14.	
831598	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1120 of SEQ ID NO:146, b is an integer of 15 to 1134, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:146, and where b is greater than or equal to a + 14.	
831608	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1472 of SEQ ID NO:147, b is an integer of 15 to 1486, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:147, and where b is greater than or equal to a + 14.	
831613	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 139 of SEQ ID NO:148, b is an integer of 15 to 153, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:148, and where b is greater than or equal to a + 14.	
831622	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 868 of SEQ ID NO:149, b is an integer of 15 to 882, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:149, and where b is greater than or equal to a + 14.	T40013, T40117, T55842, T55892, T58738, T58764, T58805, T58835, T58963, T60293, T60386, T61270, T61322, T61371, T61395, T61404, T61721, T61734, T61735, T61841, T61856, T61857, T61884, T62049, T62065, T62070, T62087, T62113, T62126, T62146, T41021, T62664, T62668, T62669, T62676, T62816, T62819, T62820, T62827, T64118, T64230, T64368, T64422, T64678, T64698, T64747, T67429, T67590, T67709, T67724, T67754, T67785, T67831, T67863, T67888, T67996, T68022, T68038, T68104, T68142, T68217, T68418, T68465, T68484, T68531, T68548, T68557, T68575, T68623, T68633, T68648, T68653, T68760, T68826, T68895, T68969, T68981, T69056, T69126, T69184, T69428, T69605, T69622, T69678, T69699, T70483, T70907, T70960, T71019, T71080, T71224, T71297, T71437, T71660, T71885, T71903, T71985, T72050, T72115, T72129, T72147, T72158, T72263, T72310, T72415, T72769, T72775, T72802, T72897, T72903, T72922, T72924, T73035, T73068, T73167, T73224, T73305, T73392, T73458, T73473, T73482, T73525, T73540, T73541, T73551, T73560, T73599, T73606, T73619, T73637, T73644, T73655, T73659, T73660, T73800.

		T73887, T73913, T73945, T73950, T74048, T74200, T74201, T74423, T74477, T74559, T74706, T74827, T99112, R05781, R05867, H47944, R95831, H60131, H65347, H65551, H68454, H68777, H73380, H73381, H79275, H79386, H82213, H82307, H93202, H93992, H93991, H94491, H94804, H95257, H95307, H95341, N28274, N58244, N68733, N77623, N80767, N91623, W07555, W80697, AA004677, AA004255, AA033869, AA034057, AA234464, AA491842, C20927
831631	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1494 of SEQ ID NO:150, b is an integer of 15 to 1508, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:150, and where b is greater than or equal to a + 14.	
831632	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1218 of SEQ ID NO:151, b is an integer of 15 to 1232, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:151, and where b is greater than or equal to a + 14.	T60158, T60218, T62213, T62652, T62877, T62966, T63329, T63951, T64542, T64634, T65965, T90119, T91565, T91610, T92138, T94160, T94999, T90219, T83025, T84028, T84029, T84511, R22325, R22619, R22620, R25250, R25595, R26992, R27328, R32850, R32954, R33282, R44282, R47779, R48151, R48152, R48322, R48428, R48538, R50415, R52277, R52278, R54608, R44282, R55376, R70352, R72103, R72155, R72280, R72317, R72367, R72368, R72371, R72372, R72716, R73784, R74375, R77393, R77394, R77892, R77987, R81485, R81725, H05676, H15941, H22149, H22193, H24533, H25059, H26810, H27743, H27803, H28012, H28066, H28290, H28291, H30654, H39748, H39761, H41932, H41979, H42063, H42642, H42766, H42767, H44628, H45776, H45777, H46386, H46404, R93135, R93942, R94660, R94661, H50708, H50709, H50720, H50812, H50811, H50826, H61352, H62379, H63665, H63944, H66336, H66385, H70746, H73887, H74080, H74176, H82646, H82647, H86555, H87065, H87719, H91147, H91197, H93078, H93211, H98788, N24993, N25111, N30229, N32159, N34033, N36553, N41829, N42292, N46951, N49340, N52921, N55462, N57121, N69863, N76837, N80667, N92844, N93333, N93683, N94449, N95075, W16427, W15325, W23470, W23480, W25070, W25186, W30795.

		W38675, W39219, W39393, W69270, W69557, AA019864, AA022662, AA022669, AA022768, AA025335, AA024417, AA031282, AA031281, AA032192, AA039752, AA040328, AA040307, AA041359, AA041442, AA057720, AA074855, AA086192, AA099717, AA099716, AA100416, AA142927, AA143150, AA149895, AA150239, AA150313, AA176193, AA459294, AA464165, AA425845, AA425899, AA428397, AA430393, AA427364, AA469113, AA505259, AA515918, AA516032, AA527677, AA533908, AA541266, AA554671, AA555247, AA557794, AA565267, AA582247, AA584415, AA588477, AA593255, AA595311, AA595376, AA604354, AA622137, AA573444, AA574244, AA732469, AA740323, AA741360, AA742872, AA749432, AA807903, AA808285, AA872498, AA873181, AA878139, AA878294, AA909748, AA937058, AA987672, AA994225, A1076066, W07696
831653	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 985 of SEQ ID NO:152, b is an integer of 15 to 999, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:152, and where b is greater than or equal to a + 14.	
831655	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1198 of SEQ ID NO:153, b is an integer of 15 to 1212, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:153, and where b is greater than or equal to a + 14.	N95539, W24228, W37689, AA019086, AA430215
831708	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2347 of SEQ ID NO:154, b is an integer of 15 to 2361, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:154, and where b is greater than or equal to a + 14.	
831738	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1817 of SEQ ID NO:155, b is an integer of 15 to 1831, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:155, and where b is greater than or equal to a + 14.	

831741	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1172 of SEQ ID NO:156, b is an integer of 15 to 1186, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:156, and where b is greater than or equal to a + 14.	T47689, T80213, H11356, H13411, R86865, R87546, N35663, AA081442, AA161001, C17978, C18946
831754	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1434 of SEQ ID NO:157, b is an integer of 15 to 1448, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:157, and where b is greater than or equal to a + 14.	
831760	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 990 of SEQ ID NO:158, b is an integer of 15 to 1004, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:158, and where b is greater than or equal to a + 14.	R73907, R74000, N64405, AA196765, AA232516, AA806432, AA837776, A1017699
831780	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1495 of SEQ ID NO:159, b is an integer of 15 to 1509, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:159, and where b is greater than or equal to a + 14.	AA100654, AA112750, AA594472, AA731487
831796	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2146 of SEQ ID NO:160, b is an integer of 15 to 2160, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:160, and where b is greater than or equal to a + 14.	H14891, W74005, AA623010, D80585, A1096496, W38434
831800	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3595 of SEQ ID NO:161, b is an integer of 15 to 3609, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:161, and where b is greater than or equal to a + 14.	
831807	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1589 of SEQ ID NO:162, b is an integer of 15 to 1603, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:162, and where b is greater than or equal to a + 14.	
831812	Preferably excluded from the present invention are	

	one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 839 of SEQ ID NO:163, b is an integer of 15 to 853, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:163, and where b is greater than or equal to a + 14.	
831813	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1903 of SEQ ID NO:164, b is an integer of 15 to 1917, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:164, and where b is greater than or equal to a + 14.	H14269, AA069213, AA808661
831830	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2406 of SEQ ID NO:165, b is an integer of 15 to 2420, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:165, and where b is greater than or equal to a + 14.	H04695, AA112742, AA251641, AA506539
831860	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2047 of SEQ ID NO:166, b is an integer of 15 to 2061, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:166, and where b is greater than or equal to a + 14.	
831872	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2553 of SEQ ID NO:167, b is an integer of 15 to 2567, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:167, and where b is greater than or equal to a + 14.	R15368, R36227, R36228, R36669, R39751, H12331, H12382, H47986, R84945, R97224, R97223, W78107, AA149874, AA193466, AA193348, AA287444, AA535607, AA687414, AA689396, AA748665, AA809715
831896	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2310 of SEQ ID NO:168, b is an integer of 15 to 2324, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:168, and where b is greater than or equal to a + 14.	R59635, N28389, AA158646, AA158659, AA188594, AA190705, AA459426, AA465652
831928	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1770 of SEQ ID NO:169, b is an integer of 15 to 1784, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:169, and where b is greater than or equal to a + 14.	
831949	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide	

	sequence described by the general formula of a-b, where a is any integer between 1 to 1282 of SEQ ID NO:170, b is an integer of 15 to 1296, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:170, and where b is greater than or equal to a + 14.	
831950	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1883 of SEQ ID NO:171, b is an integer of 15 to 1897, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:171, and where b is greater than or equal to a + 14.	
831953	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1709 of SEQ ID NO:172, b is an integer of 15 to 1723, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:172, and where b is greater than or equal to a + 14.	
831975	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1402 of SEQ ID NO:173, b is an integer of 15 to 1416, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:173, and where b is greater than or equal to a + 14.	
832036	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1942 of SEQ ID NO:174, b is an integer of 15 to 1956, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:174, and where b is greater than or equal to a + 14.	R60820, R78776, R79082, H01912, H04427, N34789, N44513, W20183, W35150, AA159701, AA159628, AA470753, AA659808
832047	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1675 of SEQ ID NO:175, b is an integer of 15 to 1689, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:175, and where b is greater than or equal to a + 14.	R21952, R21968, R26963, R78028, H75703, H75632, H84015, H88136, H88135, H94007, H95012, N24834, N30818, N31761, N41592, N79533, W16686, W24639, W38979, W87777, W87875, AA121146, AA122426, AA131874, AA131978, AA147083, AA147140, AA282507, AA282605, AA558945, H84016, AA587558, AA830662, AA866026, AA917653, AI017813, C06340
832078	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1002 of SEQ ID NO:176, b is an integer of 15 to 1016, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:176, and where b is greater than or equal to a + 14.	

832100	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1350 of SEQ ID NO:177, b is an integer of 15 to 1364, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:177, and where b is greater than or equal to a + 14.	
832104	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 726 of SEQ ID NO:178, b is an integer of 15 to 740, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:178, and where b is greater than or equal to a + 14.	
832268	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1396 of SEQ ID NO:179, b is an integer of 15 to 1410, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:179, and where b is greater than or equal to a + 14.	
832270	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1479 of SEQ ID NO:180, b is an integer of 15 to 1493, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:180, and where b is greater than or equal to a + 14.	
832279	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2026 of SEQ ID NO:181, b is an integer of 15 to 2040, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:181, and where b is greater than or equal to a + 14.	
832317	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 955 of SEQ ID NO:182, b is an integer of 15 to 969, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:182, and where b is greater than or equal to a + 14.	R81508, H12476, H86945, AA053747, AA115783, AA133749, AA134163, AA134164, AA224985, AA228334, AA228423, AA229297, AA640471, AA657793, AA687568, AA904162, AA983632
832354	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1438 of SEQ ID NO:183, b is an integer of 15 to 1452, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:183, and where b is greater than or equal to a + 14.	
832364	Preferably excluded from the present invention are	

	one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2105 of SEQ ID NO:184, b is an integer of 15 to 2119, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:184, and where b is greater than or equal to a + 14.	
832378	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1311 of SEQ ID NO:185, b is an integer of 15 to 1325, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:185, and where b is greater than or equal to a + 14.	
832385	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 419 of SEQ ID NO:186, b is an integer of 15 to 433, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:186, and where b is greater than or equal to a + 14.	
832428	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 845 of SEQ ID NO:187, b is an integer of 15 to 859, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:187, and where b is greater than or equal to a + 14.	AA031420
832485	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 819 of SEQ ID NO:188, b is an integer of 15 to 833, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:188, and where b is greater than or equal to a + 14.	R63025, R66741, H53264, H53265, H53769, H53822, H54405, H54489, H81182, H91282, AA526672, H81181
832494	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2197 of SEQ ID NO:189, b is an integer of 15 to 2211, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:189, and where b is greater than or equal to a + 14.	T61040, T61591, T90055, T90157, T92840, T93714, T96177, T77726, H04686, H05450, H06997, H20176, H20366, R92666, H65144, H92413, N64053, N64060, N66714, N71338, N71388, N79742, N95497, N99884, W07259, W24989, W37394, W37657, W40208, W40260, W40532, W45430, W56165, W60427, W60986, W61080, W63739, W72328, W73757, W74394, AA025512, AA026057, AA065019, AA069295, AA069798, AA069845, AA070441, AA075793, AA083393, AA083394, AA084576, AA086181, AA099019, AA099097, AA099493, AA102003, AA100395, AA100554, AA100555, AA100638, AA101578, AA113226, AA113811, AA115645, AA115646.

		AA115888, AA115889, AA122231, AA121108, AA121596, AA121671, AA121743, AA126075, AA126102, AA126181, AA126295, AA126404, AA129470, AA129665, AA133945, AA133946, AA146752, AA155947, AA157140, AA157228, AA159947, AA160900, AA164889, AA164890, AA164840, AA164839, AA172107, AA182040, AA171714, AA187244, AA187376, AA186418, AA188846, AA189131, AA196155, AA196257, AA196611, AA196789, AA196961, AA223155, AA223415, AA226816, AA226856, AA227026, AA227109, AA227208, AA243161, AA243205, AA428759, AA429347, AA514858, AA535250, AA555125, AA565075, AA565168, AA581531, AA587192, AA576761, AA580523, AA659699, AA688240, AA689484, AA689543, AA689313, AA729979, AA740203, AA747258, AA747399, AA747993, AA837961, AA865930, AA906561, AA910350, AA919085, AA931143, AA999884, AI051141, F19298, W22294, W22759, W22970, W25820, W73709, C02713, C02766, C03390, C03613, C04202, C05262, C05272, R28954, R29028, R29032, AA062628, AA090039, C18989
832512	<p>Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1645 of SEQ ID NO:190, b is an integer of 15 to 1659, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:190, and where b is greater than or equal to a + 14.</p>	
832515	<p>Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3880 of SEQ ID NO:191, b is an integer of 15 to 3894, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:191, and where b is greater than or equal to a + 14.</p>	
832526	<p>Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 681 of SEQ ID NO:192, b is an integer of 15 to 695, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:192, and where b is greater than or equal to a + 14.</p>	
832575	<p>Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide</p>	R28543, R28684, R55782, R55862, R62797, R62843, R67670, R71154.

	sequence described by the general formula of a-b, where a is any integer between 1 to 3117 of SEQ ID NO:193, b is an integer of 15 to 3131, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:193, and where b is greater than or equal to a + 14.	R71651, N20642, N24838, N25562, N29014, N31768, N34161, N57560, N72111, W00338, W00374, W30889, W52729, W59982, W68047, W68189, AA019459, AA043870, AA044336, AA045040, AA045041, AA115599, AA115134, AA131177, AA165259, AA165260, AA165191, AA165192, AA164549, AA164550, AA261988, AA424972, AA279863, AA458832, AA459024, AA505193, AA507542, AA514388, AA622542, AA689232, AA689233, AA804910, AA807169, AA832321, AA878091, AA904023, AA936069, AA936071, AA946621, C00143, N86645, AA010988, AA641236, AA641464, C18301
832576	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2044 of SEQ ID NO:194, b is an integer of 15 to 2058, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:194, and where b is greater than or equal to a + 14.	
832588	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 817 of SEQ ID NO:195, b is an integer of 15 to 831, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:195, and where b is greater than or equal to a + 14.	
832634	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 947 of SEQ ID NO:196, b is an integer of 15 to 961, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:196, and where b is greater than or equal to a + 14.	
832728	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 592 of SEQ ID NO:197, b is an integer of 15 to 606, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:197, and where b is greater than or equal to a + 14.	
833094	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 379 of SEQ ID NO:198, b is an integer of 15 to 393, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:198, and where b is greater than or equal to a + 14.	

833395	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1047 of SEQ ID NO:199, b is an integer of 15 to 1061, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:199, and where b is greater than or equal to a + 14.	
834326	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1345 of SEQ ID NO:200, b is an integer of 15 to 1359, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:200, and where b is greater than or equal to a + 14.	
834583	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 712 of SEQ ID NO:201, b is an integer of 15 to 726, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:201, and where b is greater than or equal to a + 14.	
834944	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2700 of SEQ ID NO:202, b is an integer of 15 to 2714, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:202, and where b is greater than or equal to a + 14.	
835012	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 408 of SEQ ID NO:203, b is an integer of 15 to 422, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:203, and where b is greater than or equal to a + 14.	
835104	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2325 of SEQ ID NO:204, b is an integer of 15 to 2339, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:204, and where b is greater than or equal to a + 14.	
835332	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1641 of SEQ ID NO:205, b is an integer of 15 to 1655, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:205, and where b is greater than or equal to a + 14.	
835487	Preferably excluded from the present invention are	

	one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 5131 of SEQ ID NO:206, b is an integer of 15 to 5145, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:206, and where b is greater than or equal to a + 14.	
836182	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 473 of SEQ ID NO:207, b is an integer of 15 to 487, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:207, and where b is greater than or equal to a + 14.	
836522	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2282 of SEQ ID NO:208, b is an integer of 15 to 2296, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:208, and where b is greater than or equal to a + 14.	
836655	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 611 of SEQ ID NO:209, b is an integer of 15 to 625, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:209, and where b is greater than or equal to a + 14.	
836787	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1537 of SEQ ID NO:210, b is an integer of 15 to 1551, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:210, and where b is greater than or equal to a + 14.	W56241, W56321, AA009901, AA521313, AA732599, AA730271, AA766911, AA767313, W27009
836789	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 997 of SEQ ID NO:211, b is an integer of 15 to 1011, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:211, and where b is greater than or equal to a + 14.	T68817, R22374, R27362, H38950, R89148, R91088, H68416, H93594, N33889, N47045, N56761, W19886, W44630, W61370, W86385, AA036993, AA065062, AA101017, AA121107, AA130485, AA147474, AA160596, AA282977
838577	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1625 of SEQ ID NO:212, b is an integer of 15 to 1639, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:212, and where b is greater than or equal to a + 14.	T53501, T40735, T63398, T63985, T64053, T64155, T64284, T93511, T94941, T94995, T96340, R00890, R01553, R12738, R12739, R39790, R54423, R66373, R66595, R67104, R67219, R79151, R79152, R82180, R82224, R82470, R82471, H01963, H02048, H02758, H02759, H05982, H19484, H19567, H19882, H19900, H44901, H44938, H44978, H46289, H46871, H49538.

		H49781, H53114, H53220, H54300, H56079, H56279, H79695, H79696, N23140, N25755, N25850, N26983, N29784, N32719, N36477, N40104, N42924, N44580, N50724, N55052, N67751, N93444, N98425, N98537, W02803, W21105, W23673, W30688, W30899, W35106, W45448, W45449, W45661, W44441, W46823, W46872, W47373, W47374, W52205, W58331, W58652, W96332, AA007386, AA007676, AA011363, AA016311, AA017511, AA018464, AA019899, AA025040, AA025039, AA029796, AA029797, AA031472, AA035395, AA035396, AA037272, AA040791, AA041228, AA042893, AA043029, AA055565, AA056185, AA056186, AA056621, AA056726, AA069193, AA079705, AA082517, AA084044, AA084043, AA115273, AA115056, AA132031, AA132153, AA149267, AA149284, AA149378, AA158093, AA158103, AA158364, AA158904, AA158905, AA165106, AA220957, AA235312, AA251169, AA421302, AA421425, AA428706, AA429291, AA513790, AA531603, AA551736, AA554236, AA605236, AA604674, AA604939, AA612935, AA617731, AA627300, AA687527, AA732095, AA740760, AA765135, AA765136, AA765296, AA765891, AA888144, AA908665, AA928038, AA936934, AA961143, AA987647, AA975856, W03595, C03206, C18055, AA164690, AA218956, AA291352, AA292329, AA293276, AA393988, AA398076, AA410772, D12417, AA442678, AA442969, AA454814, AA454888, AA482370, AA486098, AA486161, AA625879, AA678365, AA679281, AA703505, AA722872, AA732793, AA989559, A1003448, A1014938, A1022070, A1084792, A1092360
838717	<p>Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2113 of SEQ ID NO:213, b is an integer of 15 to 2127, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:213, and where b is greater than or equal to a + 14.</p>	
839008	<p>Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1152 of SEQ ID</p>	

	NO:214, b is an integer of 15 to 1166, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:214, and where b is greater than or equal to a + 14.	
840063	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3309 of SEQ ID NO:215, b is an integer of 15 to 3323, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:215, and where b is greater than or equal to a + 14.	
840533	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1394 of SEQ ID NO:216, b is an integer of 15 to 1408, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:216, and where b is greater than or equal to a + 14.	
840669	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2097 of SEQ ID NO:217, b is an integer of 15 to 2111, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:217, and where b is greater than or equal to a + 14.	T71029, T79145, T79226, T99989, R59589, R61735, R61734, R66190, R67070, H16201, H16200, H22960, H84137, H85574, H98850, N23572, N26340, N56614, W72249, W76334, W86530, W87654, W87653, AA057869, AA122103, AA129545, AA136524, AA137122, AA429808, AA525242, AA558970, H99223, AA584317, AA595168, AA825180, AA931521, AA938437, A1017369, N29659, N68604, W86674, AA007246
841140	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2479 of SEQ ID NO:218, b is an integer of 15 to 2493, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:218, and where b is greater than or equal to a + 14.	
841386	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1245 of SEQ ID NO:219, b is an integer of 15 to 1259, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:219, and where b is greater than or equal to a + 14.	AA429393, AA429394, AA493187, AA807096, AA836046
841480	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1835 of SEQ ID NO:220, b is an integer of 15 to 1849, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:220, and where b is greater than or equal to a + 14.	
841509	Preferably excluded from the present invention are	

	one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1253 of SEQ ID NO:221, b is an integer of 15 to 1267, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:221, and where b is greater than or equal to a + 14.	
841616	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 740 of SEQ ID NO:222, b is an integer of 15 to 754, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:222, and where b is greater than or equal to a + 14.	
841900	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1244 of SEQ ID NO:223, b is an integer of 15 to 1258, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:223, and where b is greater than or equal to a + 14.	R87848, AA806230, Z28656
842054	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 570 of SEQ ID NO:224, b is an integer of 15 to 584, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:224, and where b is greater than or equal to a + 14.	
843061	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 3435 of SEQ ID NO:225, b is an integer of 15 to 3449, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:225, and where b is greater than or equal to a + 14.	
843544	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1852 of SEQ ID NO:226, b is an integer of 15 to 1866, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:226, and where b is greater than or equal to a + 14.	
844092	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1050 of SEQ ID NO:227, b is an integer of 15 to 1064, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:227, and where b is greater than or equal to a + 14.	
844270	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide	

	sequence described by the general formula of a-b, where a is any integer between 1 to 359 of SEQ ID NO:228, b is an integer of 15 to 373, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:228, and where b is greater than or equal to a + 14.	
844604	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2830 of SEQ ID NO:229, b is an integer of 15 to 2844, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:229, and where b is greater than or equal to a + 14.	
844685	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1784 of SEQ ID NO:230, b is an integer of 15 to 1798, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:230, and where b is greater than or equal to a + 14.	
844855	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1809 of SEQ ID NO:231, b is an integer of 15 to 1823, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:231, and where b is greater than or equal to a + 14.	
845101	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 956 of SEQ ID NO:232, b is an integer of 15 to 970, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:232, and where b is greater than or equal to a + 14.	
845141	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 953 of SEQ ID NO:233, b is an integer of 15 to 967, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:233, and where b is greater than or equal to a + 14.	
845220	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2149 of SEQ ID NO:234, b is an integer of 15 to 2163, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:234, and where b is greater than or equal to a + 14.	R70310, H02204, H28992, H29096, W67797, W67855, W72320, AA459289, AA459519, AA430385, AA746169
845434	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b,	

	where a is any integer between 1 to 1307 of SEQ ID NO:235, b is an integer of 15 to 1321, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:235, and where b is greater than or equal to $a + 14$.	
845510	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 669 of SEQ ID NO:236, b is an integer of 15 to 683, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:236, and where b is greater than or equal to $a + 14$.	
845600	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 2101 of SEQ ID NO:237, b is an integer of 15 to 2115, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:237, and where b is greater than or equal to $a + 14$.	
845882	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1628 of SEQ ID NO:238, b is an integer of 15 to 1642, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:238, and where b is greater than or equal to $a + 14$.	
846007	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 454 of SEQ ID NO:239, b is an integer of 15 to 468, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:239, and where b is greater than or equal to $a + 14$.	H81424
846280	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1315 of SEQ ID NO:240, b is an integer of 15 to 1329, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:240, and where b is greater than or equal to $a + 14$.	
846286	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1638 of SEQ ID NO:241, b is an integer of 15 to 1652, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:241, and where b is greater than or equal to $a + 14$.	
846388	Preferably excluded from the present invention are one or more polynucleotides comprising a nucleotide sequence described by the general formula of a-b, where a is any integer between 1 to 1932 of SEQ ID	

	NO:242. b is an integer of 15 to 1946, where both a and b correspond to the positions of nucleotide residues shown in SEQ ID NO:242, and where b is greater than or equal to a + 14.	
--	--	--

Polynucleotide and Polypeptide Variants

The present invention is directed to variants of the polynucleotide sequence disclosed in SEQ ID NO:X or the complementary strand thereto, and/or the cDNA sequence contained in a cDNA clone contained in the deposit.

- 5 The present invention also encompasses variants of a colon and/or colon cancer polypeptide sequence disclosed in SEQ ID NO:Y, a polypeptide sequence encoded by the polynucleotide sequence in SEQ ID NO:X, and/or a polypeptide sequence encoded by the cDNA in the related cDNA clone contained in the deposit.

- 10 "Variant" refers to a polynucleotide or polypeptide differing from the polynucleotide or polypeptide of the present invention, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the polynucleotide or polypeptide of the present invention.

- 15 The present invention is also directed to nucleic acid molecules which comprise, or alternatively consist of, a nucleotide sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100%, identical to, for example, the nucleotide coding sequence in SEQ ID NO:X or the complementary strand thereto, the nucleotide coding sequence of the related cDNA contained in a deposited library or the complementary strand thereto, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding a polypeptide sequence encoded by the nucleotide sequence in SEQ ID NO:X, a nucleotide sequence encoding the polypeptide encoded by the cDNA in the related cDNA contained in a deposited library, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein). Polypeptides encoded by these nucleic acid molecules are also encompassed by the invention. In another embodiment, the invention encompasses nucleic acid molecules which comprise or alternatively consist of, a polynucleotide which hybridizes under stringent hybridization conditions, or alternatively, under low stringency conditions, to the nucleotide coding sequence in SEQ ID NO:X, the nucleotide coding sequence of the related cDNA clone contained in a deposited library, a nucleotide sequence encoding the polypeptide of SEQ ID NO:Y, a nucleotide sequence encoding a polypeptide sequence encoded by the nucleotide sequence in SEQ ID NO:X, a nucleotide sequence encoding the polypeptide encoded by the cDNA in the related cDNA clone contained in a deposited library, and/or polynucleotide fragments of any of these nucleic acid molecules (e.g., those fragments described herein). Polynucleotides which
- 20
- 25
- 30

hybridize to the complement of these nucleic acid molecules under stringent hybridization conditions or alternatively, under lower stringency conditions. are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

The present invention is also directed to polypeptides which comprise, or alternatively
5 consist of, an amino acid sequence which is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to, for example, the polypeptide sequence shown in SEQ ID NO:Y, a polypeptide sequence encoded by the nucleotide sequence in SEQ ID NO:X, a polypeptide sequence encoded by the cDNA in the related cDNA clone contained in a deposited library, and/or polypeptide fragments of any of these polypeptides (e.g., those fragments described
10 herein). Polynucleotides which hybridize to the complement of the nucleic acid molecules encoding these polypeptides under stringent hybridization conditions, or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides.

By a nucleic acid having a nucleotide sequence at least, for example, 95% "identical"
15 to a reference nucleotide sequence of the present invention, it is intended that the nucleotide sequence of the nucleic acid is identical to the reference sequence except that the nucleotide sequence may include up to five point mutations per each 100 nucleotides of the reference nucleotide sequence encoding the polypeptide. In other words, to obtain a nucleic acid having a nucleotide sequence at least 95% identical to a reference nucleotide sequence, up to
20 5% of the nucleotides in the reference sequence may be deleted or substituted with another nucleotide, or a number of nucleotides up to 5% of the total nucleotides in the reference sequence may be inserted into the reference sequence. The query sequence may be, for example, an entire sequence referred to in Table 1, an ORF (open reading frame), or any fragment specified as described herein.

25 As a practical matter, whether any particular nucleic acid molecule or polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a nucleotide sequence of the present invention can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present invention) and a subject sequence, also referred to as a global
30 sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245 (1990)). In a sequence alignment the query and subject sequences are both DNA sequences. An RNA sequence can be

compared by converting U's to T's. The result of said global sequence alignment is in percent identity. Preferred parameters used in a FASTDB alignment of DNA sequences to calculate percent identity are: Matrix=Unitary, k-tuple=4, Mismatch Penalty=1, Joining Penalty=30. Randomization Group Length=0, Cutoff Score=1, Gap Penalty=5, Gap Size
5 Penalty 0.05, Window Size=500 or the length of the subject nucleotide sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence because of 5' or 3' deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for 5' and 3' truncations of the
10 subject sequence when calculating percent identity. For subject sequences truncated at the 5' or 3' ends, relative to the query sequence, the percent identity is corrected by calculating the number of bases of the query sequence that are 5' and 3' of the subject sequence, which are not matched/aligned, as a percent of the total bases of the query sequence. Whether a nucleotide is matched/aligned is determined by results of the FASTDB sequence alignment.
15 This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This corrected score is what is used for the purposes of the present invention. Only bases outside the 5' and 3' bases of the subject sequence, as displayed by the FASTDB alignment, which are not matched/aligned with the query sequence, are calculated for the purposes of
20 manually adjusting the percent identity score.

For example, a 90 base subject sequence is aligned to a 100 base query sequence to determine percent identity. The deletions occur at the 5' end of the subject sequence and therefore, the FASTDB alignment does not show a matched/alignment of the first 10 bases at 5' end. The 10 unpaired bases represent 10% of the sequence (number of bases at the 5' and
25 3' ends not matched/total number of bases in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 bases were perfectly matched the final percent identity would be 90%. In another example, a 90 base subject sequence is compared with a 100 base query sequence. This time the deletions are internal deletions so that there are no bases on the 5' or 3' of the subject sequence which
30 are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only bases 5' and 3' of the subject sequence which are not matched/aligned with the query sequence are manually corrected for. No other

manual corrections are to made for the purposes of the present invention.

By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that
5 the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, (indels) or substituted with another amino acid. These alterations of the reference sequence may occur
10 at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence in SEQ ID
15 NO:Y or a fragment thereof, the amino acid sequence encoded by the nucleotide sequence in SEQ ID NO:X or a fragment thereof, or the amino acid sequence encoded by the cDNA in the related cDNA clone contained in a deposited library, or a fragment thereof, can be determined conventionally using known computer programs. A preferred method for determining the best overall match between a query sequence (a sequence of the present
20 invention) and a subject sequence, also referred to as a global sequence alignment, can be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci.6:237- 245(1990)). In a sequence alignment the query and subject sequences are either both nucleotide sequences or both amino acid sequences. The result of said global sequence alignment is in percent identity. Preferred parameters used in a
25 FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter.

If the subject sequence is shorter than the query sequence due to N- or C-terminal
30 deletions, not because of internal deletions, a manual correction must be made to the results. This is because the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences

truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. Whether a residue is
5 matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of the present invention. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the
10 query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C- terminal residues of the subject sequence.

For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the
15 subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C- termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent
20 identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject
25 sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for. No other manual corrections are to made for the purposes of the present invention.

The variants may contain alterations in the coding regions, non-coding regions, or both. Especially preferred are polynucleotide variants containing alterations which produce
30 silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Nucleotide variants produced by silent substitutions due to the degeneracy of the genetic code are preferred. Moreover, variants in which less than 50, less

than 40, less than 30, less than 20, less than 10, or 5-50, 5-25, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination are also preferred. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize codon expression for a particular host (change codons in the human mRNA to those preferred by a bacterial host such as *E. coli*).

Naturally occurring variants are called "allelic variants," and refer to one of several alternate forms of a gene occupying a given locus on a chromosome of an organism. (Genes II. Lewin, B., ed., John Wiley & Sons, New York (1985).) These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the polypeptides of the present invention. For instance, as discussed herein, one or more amino acids can be deleted from the N-terminus or C-terminus of the polypeptide of the present invention without substantial loss of biological function. The authors of Ron et al., *J. Biol. Chem.* 268: 2984-2988 (1993), reported variant KGF proteins having heparin binding activity even after deleting 3, 8, or 27 amino-terminal amino acid residues. Similarly, Interferon gamma exhibited up to ten times higher activity after deleting 8-10 amino acid residues from the carboxy terminus of this protein. (Dobeli et al., *J. Biotechnology* 7:199-216 (1988).)

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein. For example, Gayle and coworkers (*J. Biol. Chem.* 268:22105-22111 (1993)) conducted extensive mutational analysis of human cytokine IL-1a. They used random mutagenesis to generate over 3,500 individual IL-1a mutants that averaged 2.5 amino acid changes per variant over the entire length of the molecule. Multiple mutations were examined at every possible amino acid position. The investigators found that "[m]ost of the molecule could be altered with little effect on either [binding or biological activity]." (See, Abstract.) In fact, only 23 unique amino acid sequences, out of more than 3,500 nucleotide sequences examined, produced a protein that significantly differed in activity from wild-type.

Furthermore, as discussed herein, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more

biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which show a functional activity (e.g., biological activity) of the polypeptide of the invention of which they are a variant. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as have little effect on activity.

The present application is directed to nucleic acid molecules at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the nucleic acid sequences disclosed herein or fragments thereof, (e.g., including but not limited to fragments encoding a polypeptide having the amino acid sequence of an N and/or C terminal deletion), irrespective of whether they encode a polypeptide having functional activity. This is because even where a particular nucleic acid molecule does not encode a polypeptide having functional activity, one of skill in the art would still know how to use the nucleic acid molecule, for instance, as a hybridization probe or a polymerase chain reaction (PCR) primer. Uses of the nucleic acid molecules of the present invention that do not encode a polypeptide having functional activity include, inter alia, (1) isolating a gene or allelic or splice variants thereof in a cDNA library; (2) in situ hybridization (e.g., "FISH") to metaphase chromosomal spreads to provide precise chromosomal location of the gene, as described in Verma et al., Human Chromosomes: A Manual of Basic Techniques, Pergamon Press, New York (1988); and (3) Northern Blot analysis for detecting mRNA expression in specific tissues.

Preferred, however, are nucleic acid molecules having sequences at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identical to the nucleic acid sequences disclosed herein, which do, in fact, encode a polypeptide having a functional activity of a polypeptide of the invention.

Of course, due to the degeneracy of the genetic code, one of ordinary skill in the art will immediately recognize that a large number of the nucleic acid molecules having a sequence at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% identical to, for example, the nucleic acid sequence of the cDNA in the related cDNA clone contained in a

deposited library, the nucleic acid sequence referred to in Table 1 (SEQ ID NO:X), or fragments thereof, will encode polypeptides "having functional activity." In fact, since degenerate variants of any of these nucleotide sequences all encode the same polypeptide, in many instances, this will be clear to the skilled artisan even without performing the above described comparison assay. It will be further recognized in the art that, for such nucleic acid molecules that are not degenerate variants, a reasonable number will also encode a polypeptide having functional activity. This is because the skilled artisan is fully aware of amino acid substitutions that are either less likely or not likely to significantly effect protein function (e.g., replacing one aliphatic amino acid with a second aliphatic amino acid), as further described below.

For example, guidance concerning how to make phenotypically silent amino acid substitutions is provided in Bowie et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

The first strategy exploits the tolerance of amino acid substitutions by natural selection during the process of evolution. By comparing amino acid sequences in different species, conserved amino acids can be identified. These conserved amino acids are likely important for protein function. In contrast, the amino acid positions where substitutions have been tolerated by natural selection indicates that these positions are not critical for protein function. Thus, positions tolerating amino acid substitution could be modified while still maintaining biological activity of the protein.

The second strategy uses genetic engineering to introduce amino acid changes at specific positions of a cloned gene to identify regions critical for protein function. For example, site directed mutagenesis or alanine-scanning mutagenesis (introduction of single alanine mutations at every residue in the molecule) can be used. (Cunningham and Wells, *Science* 244:1081-1085 (1989).) The resulting mutant molecules can then be tested for biological activity.

As the authors state, these two strategies have revealed that proteins are surprisingly tolerant of amino acid substitutions. The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side

chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly. Besides conservative amino acid substitution, variants of the present invention include (i) substitutions with one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) substitution with one or more of amino acid residues having a substituent group, or (iii) fusion of the mature polypeptide with another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), or (iv) fusion of the polypeptide with additional amino acids, such as, for example, an IgG Fc fusion region peptide, or leader or secretory sequence, or a sequence facilitating purification. Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. (Pinckard et al., Clin. Exp. Immunol. 2:331-340 (1967); Robbins et al., Diabetes 36: 838-845 (1987); Cleland et al., Crit. Rev. Therapeutic Drug Carrier Systems 10:307-377 (1993).)

A further embodiment of the invention relates to a polypeptide which comprises the amino acid sequence of a polypeptide having an amino acid sequence which contains at least one amino acid substitution, but not more than 50 amino acid substitutions, even more preferably, not more than 40 amino acid substitutions, still more preferably, not more than 30 amino acid substitutions, and still even more preferably, not more than 20 amino acid substitutions. Of course it is highly preferable for a polypeptide to have an amino acid sequence which comprises the amino acid sequence of a polypeptide of SEQ ID NO:Y, an amino acid sequence encoded by SEQ ID NO:X, and/or the amino acid sequence encoded by the cDNA in the related cDNA clone contained in a deposited library which contains, in order of ever-increasing preference, at least one, but not more than 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1

amino acid substitutions. In specific embodiments, the number of additions, substitutions, and/or deletions in the amino acid sequence of SEQ ID NO:Y or fragments thereof (e.g., the mature form and/or other fragments described herein), an amino acid sequence encoded by SEQ ID NO:X or fragments thereof, and/or the amino acid sequence encoded by the cDNA in the related cDNA clone contained in a deposited library or fragments thereof, is 1-5, 5-10, 5-25, 5-50, 10-50 or 50-150, conservative amino acid substitutions are preferable.

Polynucleotide and Polypeptide Fragments

The present invention is also directed to polynucleotide fragments of the colon and/or colon cancer polynucleotides (nucleic acids) of the invention. In the present invention, a "polynucleotide fragment" refers, for example, to a polynucleotide having a nucleic acid sequence which: is a portion of the cDNA contained in a deposited cDNA clone; or is a portion of a polynucleotide sequence encoding the polypeptide encoded by the cDNA contained in a deposited cDNA clone; or is a portion of the polynucleotide sequence in SEQ ID NO:X or the complementary strand thereto; or is a polynucleotide sequence encoding a portion of the polypeptide of SEQ ID NO:Y; or is a polynucleotide sequence encoding a portion of a polypeptide encoded by SEQ ID NO:X or the complementary strand thereto. The nucleotide fragments of the invention are preferably at least about 15 nt, and more preferably at least about 20 nt, still more preferably at least about 30 nt, and even more preferably, at least about 40 nt, at least about 50 nt, at least about 75 nt, at least about 100 nt, at least about 125 nt or at least about 150 nt in length. A fragment "at least 20 nt in length," for example, is intended to include 20 or more contiguous bases from, for example, the sequence contained in the cDNA in a related cDNA clone contained in a deposited library, the nucleotide sequence shown in SEQ ID NO:X or the complementary stand thereto. In this context "about" includes the particularly recited value or a value larger or smaller by several (5, 4, 3, 2, or 1) nucleotides. These nucleotide fragments have uses that include, but are not limited to, as diagnostic probes and primers as discussed herein. Of course, larger fragments (e.g., at least 150, 175, 200, 250, 500, 600, 1000, or 2000 nucleotides in length) are also encompassed by the invention.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-

400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, 2001-2050, 2051-2100, 2101-2150, 2151-2200, 2201-2250, 2251-2300, 2301-2350, 2351-2400, 2401-2450, 2451-2500, 2501-2550, 2551-2600, 2601-2650, 2651-2700, 2701-2750, 2751-2800, 2801-2850, 2851-2900, 2901-2950, 2951-3000, 3001-3050, 3051-3100, 3101-3150, 3151-3200, 3201-3250, 3251-3300, 3301-3350, 3351-3400, 3401-3450, 3451-3500, 3501-3550, 3551-3600, 3601-3650, 3651-3700, 3701-3750, 3751-3800, 3801-3850, 3851-3900, 3901-3950, 3951-4000, 4001-4050, 4051-4100, and 4101 to the end of SEQ ID NO:X, or the complementary strand thereto. In this context "about" includes the particularly recited range or a range larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has a functional activity (e.g., biological activity) of the polypeptide encoded by the polynucleotide of which the sequence is a portion. More preferably, these fragments can be used as probes or primers as discussed herein. Polynucleotides which hybridize to one or more of these nucleic acid molecules under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides or fragments.

Moreover, representative examples of polynucleotide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, a sequence from about nucleotide number 1-50, 51-100, 101-150, 151-200, 201-250, 251-300, 301-350, 351-400, 401-450, 451-500, 501-550, 551-600, 651-700, 701-750, 751-800, 800-850, 851-900, 901-950, 951-1000, 1001-1050, 1051-1100, 1101-1150, 1151-1200, 1201-1250, 1251-1300, 1301-1350, 1351-1400, 1401-1450, 1451-1500, 1501-1550, 1551-1600, 1601-1650, 1651-1700, 1701-1750, 1751-1800, 1801-1850, 1851-1900, 1901-1950, 1951-2000, 2001-2050, 2051-2100, 2101-2150, 2151-2200, 2201-2250, 2251-2300, 2301-2350, 2351-2400, 2401-2450, 2451-2500, 2501-2550, 2551-2600, 2601-2650, 2651-2700, 2701-2750, 2751-2800, 2801-2850, 2851-2900, 2901-2950, 2951-3000, 3001-3050, 3051-3100, 3101-3150, 3151-3200, 3201-3250, 3251-3300, 3301-3350, 3351-3400, 3401-3450, 3451-3500, 3501-3550, 3551-3600, 3601-3650, 3651-3700, 3701-3750, 3751-3800, 3801-3850, 3851-3900, 3901-3950, 3951-4000, 4001-4050, 4051-4100, and 4101 to the end of the cDNA nucleotide

sequence contained in the deposited cDNA clone, or the complementary strand thereto. In this context "about" includes the particularly recited range, or a range larger or smaller by several (5, 4, 3, 2, or 1) nucleotides, at either terminus or at both termini. Preferably, these fragments encode a polypeptide which has a functional activity (e.g., biological activity) of the polypeptide encoded by the cDNA nucleotide sequence contained in the deposited cDNA clone. More preferably, these fragments can be used as probes or primers as discussed herein. Polynucleotides which hybridize to one or more of these fragments under stringent hybridization conditions or alternatively, under lower stringency conditions, are also encompassed by the invention, as are polypeptides encoded by these polynucleotides or fragments.

In the present invention, a "polypeptide fragment" refers to an amino acid sequence which is a portion of that contained in SEQ ID NO:Y, a portion of an amino acid sequence encoded by the polynucleotide sequence of SEQ ID NO:X, and/or encoded by the cDNA contained in the related cDNA clone contained in a deposited library. Protein (polypeptide) fragments may be "free-standing," or comprised within a larger polypeptide of which the fragment forms a part or region, most preferably as a single continuous region. Representative examples of polypeptide fragments of the invention, include, for example, fragments comprising, or alternatively consisting of, an amino acid sequence from about amino acid number 1-20, 21-40, 41-60, 61-80, 81-100, 102-120, 121-140, 141-160, 161-180, 181-200, 201-220, 221-240, 241-260, 261-280, 281-300, 301-320, 321-340, 341-360, 361-380, 381-400, 401-420, 421-440, 441-460, 461-480, 481-500, 501-520, 521-540, 541-560, 561-580, 581-600, 601-620, 621-640, 641-660, 661-680, 681-700, 701-720, 721-740, 741-760, 761-780, 781-800, 801-820, 821-840, 841-860, 861-880, 881-900, 901-920, 921-940, 941-960, 961-980, 981-1000, 1001-1020, 1021-1040, 1041-1060, 1061-1080, 1081-1100, 1101-1120, 1121-1140, 1141-1160, 1161-1180, 1181-1200, 1201-1220, 1221-1240, 1241-1260, 1261-1280, 1281-1300, 1301-1320, 1321-1340, 1341-1360, and 1361 to the end of SEQ ID NO:Y. Moreover, polypeptide fragments of the invention may be at least about 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 100, 110, 120, 130, 140, or 150 amino acids in length. In this context "about" includes the particularly recited ranges or values, or ranges or values larger or smaller by several (5, 4, 3, 2, or 1) amino acids, at either terminus or at both termini. Polynucleotides encoding these polypeptide fragments are also encompassed by the invention.

Even if deletion of one or more amino acids from the N-terminus of a protein results in modification or loss of one or more biological functions of the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example, the ability of shortened muteins to induce and/or bind to antibodies
5 which recognize the complete or mature forms of the polypeptides generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a
10 mutein with a large number of deleted N-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

Accordingly, polypeptide fragments of the invention include the secreted protein as well as the mature form. Further preferred polypeptide fragments include the secreted protein
15 or the mature form having a continuous series of deleted residues from the amino or the carboxy terminus, or both. For example, any number of amino acids, ranging from 1-60, can be deleted from the amino terminus of either the secreted polypeptide or the mature form. Similarly, any number of amino acids, ranging from 1-30, can be deleted from the carboxy terminus of the secreted protein or mature form. Furthermore, any combination of the above
20 amino and carboxy terminus deletions are preferred. Similarly, polynucleotides encoding these polypeptide fragments are also preferred.

The present invention further provides polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of a polypeptide disclosed herein (e.g., a polypeptide of SEQ ID NO:Y, a polypeptide encoded by the polynucleotide
25 sequence contained in SEQ ID NO:X, and/or a polypeptide encoded by the cDNA contained in the related cDNA clone contained in a deposited library). In particular, N-terminal deletions may be described by the general formula $m-q$, where q is a whole integer representing the total number of amino acid residues in a polypeptide of the invention (e.g., the polypeptide disclosed in SEQ ID NO:Y), and m is defined as any integer ranging from 2
30 to $q-6$. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Also as mentioned above, even if deletion of one or more amino acids from the C-terminus of a protein results in modification or loss of one or more biological functions of

the protein, other functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example the ability of the shortened mutein to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptide generally will be retained when less than the majority of the residues of the complete or mature polypeptide are removed from the C-terminus. Whether a particular polypeptide lacking C-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a mutein with a large number of deleted C-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

Accordingly, the present invention further provides polypeptides having one or more residues from the carboxy terminus of the amino acid sequence of a polypeptide disclosed herein (e.g., a polypeptide of SEQ ID NO:Y, a polypeptide encoded by the polynucleotide sequence contained in SEQ ID NO:X, and/or a polypeptide encoded by the cDNA contained in the related cDNA referenced in Table 1). In particular, C-terminal deletions may be described by the general formula 1-n, where n is any whole integer ranging from 6 to q-1, and where n corresponds to the position of an amino acid residue in a polypeptide of the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

In addition, any of the above described N- or C-terminal deletions can be combined to produce a N- and C-terminal deleted polypeptide. The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini, which may be described generally as having residues m-n of a polypeptide encoded by SEQ ID NO:X (e.g., including, but not limited to, the preferred polypeptide disclosed as SEQ ID NO:Y), and/or the cDNA in the related cDNA clone contained in a deposited library, where n and m are integers as described above. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Any polypeptide sequence contained in the polypeptide of SEQ ID NO:Y, encoded by the polynucleotide sequences set forth as SEQ ID NO:X, or encoded by the cDNA in the related cDNA clone contained in a deposited library may be analyzed to determine certain preferred regions of the polypeptide. For example, the amino acid sequence of a polypeptide encoded by a polynucleotide sequence of SEQ ID NO:X, or the cDNA in a deposited cDNA

clone may be analyzed using the default parameters of the DNASTAR computer algorithm (DNASTAR, Inc., 1228 S. Park St., Madison, WI 53715 USA; <http://www.dnastar.com/>).

Polypeptide regions that may be routinely obtained using the DNASTAR computer algorithm include, but are not limited to, Garnier-Robson alpha-regions, beta-regions, turn-regions, and coil-regions, Chou-Fasman alpha-regions, beta-regions, and turn-regions, Kyte-Doolittle hydrophilic regions and hydrophobic regions, Eisenberg alpha- and beta-amphipathic regions, Karplus-Schulz flexible regions, Emini surface-forming regions and Jameson-Wolf regions of high antigenic index. Among highly preferred polynucleotides of the invention in this regard are those that encode polypeptides comprising regions that combine several structural features, such as several (e.g., 1, 2, 3 or 4) of the features set out above.

Additionally, Kyte-Doolittle hydrophilic regions and hydrophobic regions, Emini surface-forming regions, and Jameson-Wolf regions of high antigenic index (i.e., containing four or more contiguous amino acids having an antigenic index of greater than or equal to 1.5, as identified using the default parameters of the Jameson-Wolf program) can routinely be used to determine polypeptide regions that exhibit a high degree of potential for antigenicity. Regions of high antigenicity are determined from data by DNASTAR analysis by choosing values which represent regions of the polypeptide which are likely to be exposed on the surface of the polypeptide in an environment in which antigen recognition may occur in the process of initiation of an immune response.

Preferred polypeptide fragments of the invention are fragments comprising, or alternatively consisting of, an amino acid sequence that displays a functional activity of the polypeptide sequence of which the amino acid sequence is a fragment.

By a polypeptide demonstrating a "functional activity" is meant, a polypeptide capable of displaying one or more known functional activities associated with a full-length (complete) protein of the invention. Such functional activities include, but are not limited to, biological activity, antigenicity [ability to bind (or compete with a polypeptide for binding) to an anti-polypeptide antibody], immunogenicity (ability to generate antibody which binds to a specific polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide.

Other preferred polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an

activity of the polypeptide of the present invention. The biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

In preferred embodiments, polypeptides of the invention comprise, or alternatively consist of, one, two, three, four, five or more of the antigenic fragments of the polypeptide of
5 SEQ ID NO:Y, or portions thereof. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Table 4.

Sequence/ Contig ID	Predicted Epitopes
500802	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 774 as residues: Gln-1 to Ser-17, Ser-19 to Ile-25, Leu-29 to Arg-41, Ser-46 to Glu-57.
553147	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 776 as residues: Phe-1 to Ile-20.
558860	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 777 as residues: Ser-6 to Arg-11.
561730	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 778 as residues: Asn-1 to Arg-7, Leu-28 to Pro-45.
585938	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 779 as residues: Arg-10 to Ser-23, Gln-69 to His-74.
587785	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 780 as residues: Ile-1 to Ser-11, Leu-20 to Thr-30, Cys-74 to Cys-82, Leu-94 to Glu-110.
588916	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 781 as residues: Val-43 to Pro-55, Glu-92 to Ser-99.
613825	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 782 as residues: Asn-1 to Trp-11, Ser-15 to Gln-22, Ser-43 to Ala-51, Lys-58 to Gly-66.
639090	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 783 as residues: Ser-29 to Ser-35, Pro-43 to Gly-48, Gln-60 to Ser-65.
659544	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 785 as residues: Leu-10 to Glu-15, His-19 to Glu-26.
659739	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 786 as residues: Lys-70 to His-78, Lys-149 to Asn-154, Gly-209 to Leu-217, Lys-248 to Val-255, Ile-259 to Arg-264, Arg-280 to Ala-287.
661057	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 787 as residues: Cys-59 to Arg-64, Gly-110 to Asp-115, Pro-127 to Trp-132.
661313	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 788 as residues: Glu-1 to Phe-7, Lys-42 to Leu-48.
666316	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 789 as residues: Lys-27 to Asn-52.
669229	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 790 as residues: Asp-1 to Phe-12, Val-92 to Ser-103.
670471	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 791 as residues: Lys-75 to Asp-81, Glu-145 to Gln-156, Glu-163 to Arg-170, Lys-225 to Leu-231.
676611	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 792 as residues: Tyr-4 to Lys-12, Thr-23 to Asn-31, Val-52 to Thr-63, Arg-90 to Met-95.
691240	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 793 as residues: Pro-74 to Glu-79, Ser-116 to Lys-121.
702977	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 794 as residues: Pro-8 to Tyr-20.
709517	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 795 as residues: Leu-7 to Gly-12, Cys-20 to His-27.
714730	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 796 as residues: Pro-14 to Arg-23, Ala-171 to Ser-178.
714834	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 797 as residues: Ala-6 to Gly-12, Gln-18 to Arg-32.
719584	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 799 as residues: Pro-22 to Ile-31.
724637	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 800 as residues: Val-11 to Arg-34, Asn-54 to Cys-59.
728392	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 801 as

	residues: Arg-31 to Glu-45, Glv-76 to Pro-88, Asn-143 to Asp-148.
738716	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 802 as residues: Pro-40 to Pro-46.
739056	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 803 as residues: Ser-28 to Ala-33, Pro-44 to Phe-49, Arg-113 to Gly-118, Pro-131 to Arg-142, Asp-155 to Leu-166.
739143	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 804 as residues: Ala-1 to Gly-14, Glu-21 to Gly-27, Asp-54 to Lys-59, Lys-64 to Glu-71, Gln-92 to Leu-97, Asn-114 to His-120, Leu-135 to Asp-142, Glu-149 to Ser-154, Ser-256 to Thr-261, Asp-290 to Lys-301, Glu-315 to Gln-323, Lys-331 to Asn-342, Arg-346 to Met-361.
742329	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 805 as residues: Arg-7 to Ala-13, Gln-21 to Ser-27, Gln-68 to Glv-73, Pro-75 to Val-88.
745481	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 807 as residues: Asn-1 to Lys-14, Arg-32 to His-39, Asn-46 to Gly-51.
753731	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 809 as residues: Arg-22 to Ser-39, Val-42 to Thr-54, Gln-61 to His-69.
754383	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 810 as residues: Ala-2 to Glv-12.
756749	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 811 as residues: His-1 to Thr-11, Thr-13 to Ser-18, Gly-25 to Gly-30, Pro-63 to Pro-69, Glu-84 to Tyr-101, Asn-110 to Ala-140.
757980	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 812 as residues: Phe-9 to His-21.
764818	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 813 as residues: Pro-12 to Trp-17, Asn-22 to Ala-37, Arg-45 to Gly-54, Asp-72 to Thr-95, Pro-97 to Glu-116, Gly-137 to Lys-151, Glu-164 to Asp-171, Ser-175 to Gly-185, Glu-187 to Gly-213, Lys-270 to Glu-276, Leu-281 to Lys-286, Asp-314 to Gly-321, Glu-324 to Glu-331, Val-333 to Arg-340.
765140	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 814 as residues: Thr-15 to Asp-27.
766893	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 815 as residues: Arg-6 to Leu-11, Arg-21 to Tyr-27, Phe-37 to Lys-46, Gly-59 to Glv-64.
771412	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 817 as residues: Pro-1 to His-6, Pro-37 to Arg-47.
772226	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 818 as residues: Phe-16 to Arg-30, Glu-35 to Trp-58, Lys-60 to Gln-68, Pro-80 to Tyr-85.
773057	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 819 as residues: Gly-37 to Arg-43.
773173	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 820 as residues: Pro-19 to Asn-26.
780154	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 821 as residues: Arg-20 to Ile-31, Pro-34 to Ala-59, Glu-66 to Pro-125, Leu-132 to Lys-137, Lys-155 to Arg-259.
780768	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 822 as residues: Phe-12 to Lys-17.
780779	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 823 as residues: Ser-1 to Ser-11, Gln-64 to Gln-69, Arg-117 to Arg-127.
782394	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 824 as residues: Phe-18 to Gly-24.
783160	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 825 as residues: Lys-35 to Lys-41, Thr-50 to His-56, Thr-110 to Gly-119.
783506	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 826 as residues: Thr-3 to Thr-9.
792139	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 830 as residues: Arg-1 to Thr-13, Arg-21 to Pro-30, Ser-70 to Arg-79, Asp-89 to Arg-101.

805715	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 832 as residues: Met-7 to Ala-17, Arg-26 to Leu-32, Lys-47 to Lys-52, Asn-67 to Asn-72, Val-77 to Tyr-82, Pro-101 to Arg-107, Arg-137 to Arg-146, Ser-168 to Thr-173, Asp-189 to Lys-199.
811111	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 833 as residues: His-24 to Asn-31.
811113	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 834 as residues: Gln-1 to Ala-9, Cys-56 to Gly-61, Trp-105 to Thr-110, Arg-150 to Thr-155, Leu-189 to Lys-195.
823902	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 835 as residues: Thr-18 to Glu-23.
826518	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 836 as residues: Ile-20 to Lys-26, Cys-39 to Arg-46.
826704	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 837 as residues: His-14 to Phe-20, Glu-70 to Leu-83.
828180	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 840 as residues: Glu-38 to Arg-52, Ser-56 to Val-62.
828658	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 842 as residues: Asp-1 to Pro-12, Gly-59 to Lys-64, Asp-70 to Leu-76, Pro-160 to Pro-166, Thr-174 to Asn-179.
828919	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 843 as residues: Thr-49 to Val-54, Leu-83 to Lys-91, Gly-121 to Thr-130, Asp-165 to Glu-172, Thr-180 to Gly-188.
830208	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 846 as residues: Lys-49 to Asn-56, Glu-61 to Ala-67.
830248	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 847 as residues: Pro-17 to Asp-36, Pro-102 to Glu-108, Pro-122 to Lys-128, His-150 to Gly-155, Asn-162 to Tyr-168, Pro-186 to Gln-193, Ser-205 to Pro-211, Gln-305 to Gly-317.
830275	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 848 as residues: Ser-16 to Glu-22, Asn-45 to Ser-50, Thr-121 to Gly-136, Lys-150 to Arg-157, Ser-175 to Cys-181, Glu-198 to Ser-203.
830286	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 849 as residues: His-11 to Pro-18, Thr-241 to Thr-258, Ala-352 to Ala-365.
830347	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 850 as residues: Asp-33 to Ala-39.
830348	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 851 as residues: Gln-5 to Arg-15, Ile-96 to Asn-101, Asp-122 to Gly-128.
830364	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 852 as residues: Val-76 to Asn-82, Lys-87 to Tyr-94, Glu-118 to Gln-125, Pro-140 to Ile-145, Gly-149 to Pro-173, Ala-215 to Lys-222, Lys-230 to Gly-235, Pro-250 to Asn-256, Ser-302 to Arg-307, Ser-321 to Glu-332.
830394	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 853 as residues: Thr-37 to Thr-44, Leu-57 to Ser-63, Ser-74 to Lys-86, Gln-107 to Leu-112, Lys-140 to Ala-145, Asp-154 to Ser-163.
830412	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 855 as residues: His-65 to Gly-74, Asp-85 to Ser-97, Leu-133 to Glu-138, Glu-144 to Asp-153, Arg-170 to Ser-175, Gly-184 to Arg-189, Gln-202 to Tyr-208.
830464	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 857 as residues: Val-3 to Val-11, Gln-16 to Gln-27, Glu-41 to Asp-51.
830471	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 858 as residues: Glu-10 to His-22, Ser-37 to Lys-45.
830477	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 859 as residues: Lys-1 to Cys-13, Thr-32 to Cys-37, Ser-44 to Glu-50, Glu-57 to Asn-64, Glu-85 to Glu-93, Ala-129 to Ser-139, Gln-157 to Thr-185, Gln-199 to Gly-215, Ile-241 to Leu-247, Asp-254 to Leu-263, Gln-265 to Gln-270, Glu-298 to Gln-309, Glu-316 to Ala-321, Leu-325 to Glu-334, Glu-340 to Ser-345, Leu-348 to His-367, Lys-384 to Arg-391.

	Leu-409 to Asn-417, Arg-431 to Arg-437, Phe-441 to Leu-448, Ala-456 to Glu-484, Lys-509 to Val-519, Glu-521 to Asp-528, Asp-546 to Phe-553, Glu-558 to Phe-567, Pro-573 to Thr-588.
830500	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 860 as residues: Gln-27 to Gly-34.
830509	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 861 as residues: Pro-2 to Asp-7, Gln-13 to Gln-29, Pro-35 to Trp-41.
830528	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 862 as residues: Gln-1 to Arg-12, Asp-22 to Pro-44, Lys-52 to Asp-62, Pro-68 to Lys-93, Pro-99 to Pro-129, Ala-138 to Ser-150, Lys-156 to Val-194, Ile-197 to Glu-210, Ala-213 to Ala-287, Leu-289 to Lys-327, Lys-330 to Gly-340, Asp-344 to Gln-360, Ile-396 to Thr-401, Lys-409 to Asp-418, Met-450 to Ala-460, Glu-468 to Gly-475.
830542	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 863 as residues: Val-1 to Gly-10, Arg-24 to Asp-36, Leu-225 to Trp-231, Val-249 to Met-258, Glu-262 to Thr-269, Val-279 to Gly-284, Asp-307 to Asn-313, Arg-411 to Lys-416.
830564	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 864 as residues: Trp-103 to Glu-113, Lys-118 to Tyr-125.
830611	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 865 as residues: Glu-51 to Ser-57, Arg-128 to Ala-133.
830620	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 867 as residues: Lys-54 to Arg-59, Arg-66 to Arg-71.
830630	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 868 as residues: Pro-12 to Gly-17.
830654	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 869 as residues: Leu-1 to Asp-6.
830660	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 870 as residues: Lys-111 to Trp-116, Glu-139 to Gly-148, Arg-182 to Ser-189.
830704	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 872 as residues: Asn-1 to Glu-8, Ala-38 to Gly-46, Gln-58 to Asp-71, Ala-75 to Cys-103, Met-106 to Ala-140, Gln-153 to Ile-159.
830765	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 873 as residues: Ser-19 to Thr-26, Pro-47 to Thr-59.
830778	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 874 as residues: Asp-35 to Gly-40, Glu-104 to Glu-109, Ser-226 to Tyr-231.
830784	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 875 as residues: Pro-34 to Leu-41.
830800	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 876 as residues: Ser-16 to Lys-24, Gly-91 to Thr-96.
830821	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 877 as residues: Leu-2 to Thr-8, Asp-15 to Gly-26, Phe-64 to Ser-70, Pro-77 to Trp-82, Pro-85 to Lys-90.
830849	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 878 as residues: Leu-2 to Ser-18, Gly-31 to Ser-40, Asn-56 to Thr-86, Asp-114 to Arg-120.
830903	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 879 as residues: Thr-21 to Thr-33.
830913	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 880 as residues: Gly-48 to Pro-53, Gln-66 to Pro-74, Thr-151 to Gly-156, Asn-292 to Asn-297.
830920	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 881 as residues: Asp-15 to Ser-25, Ser-33 to Val-38, Lys-181 to Phe-187.
830938	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 882 as residues: Thr-65 to Asp-70, Leu-89 to Ala-95.
831014	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 884 as residues: Ala-2 to Gln-11, Glu-71 to Leu-78, Leu-89 to Trp-98, Ser-163 to Ala-170, Glu-261 to Asp-269, Phe-286 to Val-292.
831026	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 885 as residues: Lys-41 to Gly-46, Tyr-64 to Phe-75.

831055	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 887 as residues: Trp-37 to His-50, Lys-108 to Phe-114, Lys-131 to Thr-137, Arg-351 to Ser-356, Pro-363 to Cys-369, Glu-390 to Asp-397.
831057	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 888 as residues: Arg-1 to Gly-14, Thr-19 to Gly-25, Ala-31 to Ala-41, Glu-53 to Ile-62, Val-66 to Glu-75, Ser-103 to Asp-113, Ala-135 to Asp-140.
831062	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 889 as residues: Ser-24 to Ala-31.
831117	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 890 as residues: Lys-50 to Tyr-55.
831122	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 891 as residues: Phe-8 to Gly-14, Arg-58 to Gly-68, Lys-107 to Ser-131, Gln-151 to Val-160, Lys-180 to Lys-186, Lys-211 to Thr-223.
831132	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 893 as residues: Gly-1 to Ser-16.
831152	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 894 as residues: Ser-8 to Arg-13, Lys-59 to Ala-65, Glu-71 to Glu-86, Leu-98 to His-108, Arg-118 to Ile-126, His-138 to Ala-145, Pro-148 to Tyr-156, Pro-170 to Ala-175, Val-187 to Lys-194, Glu-206 to Val-217, Gly-221 to Ser-226, Asp-250 to Lys-255.
831157	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 895 as residues: Val-1 to Asn-11, Glu-13 to Gly-25, Ser-31 to Ala-49, Arg-61 to Gly-66, Ala-84 to Ala-90.
831160	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 896 as residues: His-1 to Ala-7, Asp-43 to Lys-52, Tyr-98 to Gly-103, Glu-118 to Leu-125, Phe-183 to Tyr-195, Gln-209 to Arg-220, Ile-257 to Gly-262, Glu-278 to Thr-284, Ile-309 to Pro-314, Leu-339 to Asp-347, Ala-358 to Gln-388, Gln-401 to Leu-414, Glu-425 to Ala-440, Ala-448 to Glu-453, Ile-460 to Gln-465, Glu-482 to Glu-492, Ala-498 to Glu-511, Pro-520 to Val-526, Gly-556 to Gln-577, Leu-587 to His-598, Glu-605 to Asp-630.
831197	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 898 as residues: Ser-28 to Leu-39, Phe-48 to Phe-55, Pro-60 to Gln-66, Arg-73 to Thr-78.
831217	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 899 as residues: Asp-52 to Val-63, Asn-75 to Glu-83.
831248	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 901 as residues: Pro-24 to Gly-34, Lys-108 to Arg-118.
831369	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 903 as residues: Ala-1 to Gly-8.
831371	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 904 as residues: Arg-39 to Ser-44, Arg-66 to Arg-76.
831373	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 905 as residues: Gly-7 to Ser-13, Gln-40 to Trp-45, Lys-109 to Gly-116, Gly-134 to Arg-141, Arg-149 to Arg-164, Arg-174 to Phe-181, Lys-202 to Lys-210, Glu-263 to Leu-272, Pro-274 to Leu-280, Glu-289 to Glu-296, Pro-334 to His-341, Tyr-413 to Pro-426, Glu-432 to Lys-449.
831387	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 906 as residues: Tyr-21 to Leu-28, Cys-51 to Phe-72, Ser-107 to Leu-113, Leu-125 to Leu-134, Ser-142 to Ala-152, His-159 to Tyr-164, Arg-276 to Val-290.
831410	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 907 as residues: Arg-7 to Lys-13, Pro-28 to Cys-34, Gly-100 to Asn-109, Cys-155 to Arg-162.
831448	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 908 as residues: Ala-10 to Cys-20, Tyr-36 to Lys-41, Asp-68 to Ala-75, Ala-84 to Arg-89, Glu-112 to Ser-119.
831450	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 909 as residues: Pro-23 to Gly-28, Thr-52 to Pro-63.
831472	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 910 as residues: Ser-16 to Ala-26.

831473	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 911 as residues: Arg-37 to Gln-42, Asn-59 to Asn-65, Asn-109 to Val-121, Arg-191 to Glu-199, Lys-205 to Ile-214.
831474	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 912 as residues: Glu-1 to Leu-8, Ser-50 to Arg-56, Thr-61 to Arg-66, Val-69 to Arg-82.
831494	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 913 as residues: Arg-21 to Ser-27, Arg-77 to Asp-82, Glu-116 to Ile-134, Ser-139 to Ser-162, Leu-167 to Gly-190, Cys-192 to Gly-205.
831506	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 914 as residues: Val-6 to Tyr-12, Lys-77 to Ala-82, Ser-102 to Arg-108, Ser-145 to Ser-151.
831533	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 915 as residues: Thr-9 to Cys-16, Arg-52 to Tyr-57, Ser-61 to Ser-69.
831539	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 916 as residues: Thr-32 to Arg-39, Cys-44 to Arg-60, Lys-65 to Gln-70, Gly-78 to Ile-86, Lys-126 to Thr-134, Leu-140 to Glu-148.
831556	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 917 as residues: Gly-45 to Asp-52.
831598	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 919 as residues: Asn-1 to Val-6, Phe-76 to Tyr-83, Gly-129 to Gln-135, Thr-145 to Asp-153, Pro-213 to Gln-220, Thr-230 to Asn-236, Lys-242 to Ala-248.
831608	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 920 as residues: Thr-23 to Pro-34, Glu-39 to Asp-83, Asn-89 to Lys-99, Asp-118 to Asp-128, Asn-135 to Glu-150, Glu-153 to Gly-168, Gly-181 to Thr-187, Arg-200 to Asp-205, Arg-273 to Ile-279, Thr-295 to Asp-300, Thr-316 to Cys-321.
831613	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 921 as residues: Pro-1 to Glu-7, Arg-9 to Phe-15, Thr-27 to Gly-34.
831655	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 926 as residues: Tyr-31 to Gln-38.
831708	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 927 as residues: Glu-22 to Ile-27, Gly-43 to Gly-49, His-83 to Arg-105.
831741	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 929 as residues: Asp-22 to Asp-27, Pro-64 to Gln-74, Ser-126 to Gly-131, Lys-134 to Arg-143, Arg-150 to Gly-162, Gln-180 to Tyr-196, Asp-209 to Leu-224, Gly-233 to Gly-241, Pro-246 to Arg-251.
831754	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 930 as residues: Arg-40 to Glu-50, Gly-57 to Gly-68, Phe-72 to Tyr-79.
831760	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 931 as residues: His-24 to Asp-39.
831780	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 932 as residues: Arg-92 to Thr-101.
831796	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 933 as residues: Pro-1 to Ser-8.
831800	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 934 as residues: Asp-1 to Ser-6, Glu-16 to Ser-26, Lys-66 to Pro-76, Leu-93 to Arg-99, Val-153 to Lys-164, Glu-177 to Asp-183, Ser-188 to Leu-193, Arg-210 to Ser-220, Thr-229 to Ser-244, Pro-283 to Phe-297.
831813	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 937 as residues: Pro-20 to Ala-30.
831830	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 938 as residues: Arg-12 to Lys-17, Gln-51 to Phe-60, Asp-97 to Trp-102, Glu-132 to Cys-137, Asp-160 to Leu-168, Glu-210 to Gln-219, Lys-302 to Pro-308, Phe-416 to Asp-421, Leu-444 to Leu-449, Val-457 to Asn-464, Leu-466 to Trp-472, Ile-474 to Trp-480, Ser-527 to Ser-533, Pro-558 to Phe-565, Ile-578 to Trp-584, Asp-614 to Asp-627, Asn-698 to Asp-710, Pro-738 to Ser-744.
831860	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 939 as residues: Pro-19 to Tyr-25.

831896	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 941 as residues: Ser-18 to Phe-30, Leu-34 to Asn-41, Ala-48 to Tyr-56, Leu-103 to Ala-110, Asp-124 to Val-130, Ile-141 to Leu-150, Leu-188 to Ser-196, Glu-229 to Asn-238, Thr-248 to Cys-259.
831928	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 942 as residues: Asn-55 to Asp-60.
831949	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 943 as residues: Arg-1 to Glu-9, Glu-19 to Arg-32, Ala-77 to Thr-90, Thr-95 to Thr-104, Lys-106 to Ser-119, Leu-136 to Arg-141, Tyr-165 to Asn-174.
831950	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 944 as residues: Ser-18 to Glu-26, Phe-93 to Arg-102, Leu-137 to Gln-143, Pro-148 to Glv-157.
831975	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 946 as residues: His-41 to Thr-48.
832047	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 948 as residues: Arg-57 to Glu-62, Pro-73 to Gly-80.
832078	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 949 as residues: Pro-14 to Leu-21, Cys-34 to Gly-39.
832100	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 950 as residues: Tyr-37 to Val-45.
832104	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 951 as residues: Thr-1 to Ser-6, Arg-14 to Cys-20.
832279	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 954 as residues: Ser-28 to Pro-34, Pro-134 to Ser-139, Gln-178 to Gly-183, Thr-193 to Gly-198, His-244 to Gly-257, Asp-263 to Tyr-273, Lys-337 to Arg-347, Pro-366 to Lys-372, Ala-382 to Asp-387.
832317	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 955 as residues: Thr-32 to Gln-39, Asn-58 to Trp-71, Glu-96 to Trp-108, Cys-126 to Glv-133.
832364	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 957 as residues: Glu-2 to Met-9, Asp-17 to Asn-22, Leu-27 to Val-35.
832428	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 960 as residues: Arg-35 to Glv-41.
832485	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 961 as residues: Ser-121 to Cys-127.
832494	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 962 as residues: Ser-10 to Leu-28, Ser-31 to Asp-40, Ser-55 to Thr-62, Thr-94 to Asn-102, Asp-124 to Phe-135, Asn-175 to Lys-193, Glu-238 to Leu-243, Val-250 to Ala-259, Lys-291 to Asn-308, Ser-318 to Gly-327, Lys-335 to Asp-346, Tyr-404 to Ile-410, Gln-420 to Gln-430, Thr-476 to Phe-482, Pro-536 to Val-561, Tyr-563 to Leu-568.
832512	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 963 as residues: Arg-1 to Ala-7, Leu-9 to Ser-24, Glu-32 to Asp-43, Glu-71 to Glu-86, Val-92 to Ile-104, Asp-143 to Ser-154, Lys-190 to Glu-202, Glu-218 to Lys-241.
832515	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 964 as residues: Glu-3 to Gly-12, Arg-20 to Gln-30, Leu-34 to Gln-39, Asp-51 to Arg-58, Gln-69 to Val-77, Gly-105 to Lys-117, Cys-123 to Phe-132.
832526	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 965 as residues: Pro-15 to Asn-25, Glu-48 to Phe-59.
832575	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 966 as residues: Thr-24 to Arg-29, Ala-55 to Tyr-60, Tyr-77 to Asp-89, Leu-108 to Gly-115, Thr-142 to Gly-149.
832576	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 967 as residues: Arg-1 to Leu-11, Pro-21 to Gly-28, Pro-37 to His-47, Lys-79 to Gln-88, Pro-108 to Glv-116, Pro-179 to Thr-188, Arg-207 to Asn-213.
832634	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 969 as residues: Leu-2 to Ser-12, Pro-125 to Asp-133.
832728	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 970 as residues: Gln-16 to Glv-32, Leu-100 to Gly-106, Glv-118 to Lys-132, Pro-156 to Leu-

	162.
833395	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 972 as residues: Ser-3 to Gly-9.
834326	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 973 as residues: Ser-1 to Trp-19, Asn-148 to Leu-153, Tyr-235 to Trp-244.
834944	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 975 as residues: Glu-42 to Gln-51, Pro-115 to Asp-120, Arg-127 to Gly-133, Gln-199 to Gln-211.
835104	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 977 as residues: Thr-1 to Arg-14, Val-18 to Pro-23, Thr-37 to Met-44, Gln-51 to Leu-57.
835332	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 978 as residues: Thr-1 to Glu-13, Arg-135 to Asp-142, Thr-150 to Gln-155, Cys-173 to Cys-183, Cys-203 to Asp-214.
835487	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 979 as residues: Ala-13 to Arg-22, Pro-43 to Glu-57, Ala-73 to Pro-90, Arg-102 to Ser-109, Pro-114 to Gly-122, Arg-127 to Arg-138, Glu-153 to Gly-158, Pro-165 to Pro-171, Gly-185 to Arg-190, Pro-211 to Pro-216, Glu-231 to Asn-261, Ala-280 to Pro-291, Pro-303 to Gly-311, Arg-313 to Gly-326, Ala-358 to Ala-364, Pro-369 to Gly-377, Pro-390 to Gly-407, Tyr-420 to Tyr-441, Glu-461 to Thr-470, Pro-479 to Trp-487, Asp-489 to Cys-494, Gln-515 to Lys-532, Ala-572 to Asn-582, Asp-588 to Leu-594, Cys-625 to Trp-632, Tyr-639 to Arg-646.
836182	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 980 as residues: Ala-7 to Thr-17, Arg-31 to Thr-36.
836522	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 981 as residues: Gly-59 to Cys-65.
836789	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 984 as residues: Gly-18 to Gly-25, Glu-59 to Glu-64.
838577	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 985 as residues: Pro-15 to Trp-20, Pro-46 to Gln-57, Glu-68 to Phe-83.
839008	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 987 as residues: Arg-1 to Arg-13, Gln-125 to Glu-131, Asn-137 to Val-142, Gly-183 to Tyr-188, Asn-245 to Ser-251, Gln-302 to Asn-311.
840063	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 988 as residues: Gly-1 to Gly-31.
840533	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 989 as residues: Thr-16 to Pro-23, Pro-39 to Trp-48, Arg-50 to Lys-55, Gly-73 to Gly-79.
840669	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 990 as residues: Met-27 to Gln-33, Gln-49 to Gly-56, Thr-63 to Leu-70, Thr-115 to Arg-127, Pro-174 to Asn-184.
841140	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 991 as residues: Arg-17 to Phe-24, Pro-113 to Gly-121, Thr-235 to Met-240.
841386	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 992 as residues: Val-58 to Met-66, Pro-134 to Lys-143, Tyr-163 to Ala-170, Val-178 to Lys-187, Pro-207 to Gly-212.
841900	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 996 as residues: Ile-2 to Phe-12.
842054	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 997 as residues: Asp-27 to Trp-32, Pro-89 to Glu-99, Arg-112 to Lys-123.
843061	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 998 as residues: Leu-3 to Gly-18, His-36 to His-57, Lys-136 to Leu-145, Gly-174 to Trp-184, Lys-188 to Tyr-196, Lys-204 to Asp-211, Pro-293 to Ser-305, Glu-321 to Asp-333, Gly-342 to Lys-348, Ala-371 to Asp-377, Asp-439 to Leu-449, Ala-521 to Gly-529, Tyr-583 to Trp-599, Asn-639 to Ser-644, Leu-738 to Leu-745.
843544	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 999 as residues: Tyr-11 to Phe-18, Ser-34 to Lys-43.
844092	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1000 as

	residues: Gln-1 to Lys-6, Glu-30 to Glu-37, Glu-40 to Thr-53.
844270	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1001 as residues: Thr-10 to Glu-20, Pro-44 to Thr-50.
844604	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1002 as residues: Gly-8 to Phe-20, Pro-23 to Arg-43, Asp-62 to Asp-67, Pro-73 to Asn-80, Val-83 to Phe-95, Glu-103 to Ile-109, Tyr-120 to Ala-125, Thr-176 to Thr-183, Pro-200 to Pro-214, Pro-232 to Met-240, Gln-248 to Asp-292, Arg-297 to Ser-310, Pro-320 to Glu-332, Glu-347 to Ser-390, Ala-392 to Pro-404, Pro-425 to Gly-435, Pro-438 to Gly-443, Gly-467 to Pro-480, Pro-486 to Pro-499, Pro-506 to Met-512, Pro-572 to Glu-580, Arg-592 to Glu-597, Ala-601 to Ser-610, Ala-618 to Pro-623.
844685	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1003 as residues: Ser-14 to Ser-19, Pro-25 to Gly-32, Asn-98 to Lys-108.
844855	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1004 as residues: Ala-9 to Ser-15, Pro-21 to Arg-26.
845101	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1005 as residues: Ala-2 to Glu-13, Pro-31 to Pro-42, Gln-89 to Tyr-95, Gln-169 to Leu-189.
845141	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1006 as residues: Gly-13 to Met-26, Arg-34 to Gly-39, Ile-60 to Ser-80, Ala-85 to Thr-98.
845220	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1007 as residues: Pro-14 to Gly-24, Glu-33 to Ala-39, Asp-145 to Pro-168, Ala-238 to Arg-250, Pro-258 to Phe-269, Arg-285 to Pro-290, Ala-340 to Cys-364.
845434	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1008 as residues: Ala-1 to Glu-7, Gln-29 to Phe-34, Gly-67 to Ala-75, Gln-78 to Leu-83, Asn-96 to Ile-109, Thr-144 to Trp-151.
845510	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1009 as residues: Arg-79 to Leu-86, Met-114 to Asp-122, Leu-129 to Leu-134, Gln-145 to Arg-152.
845600	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1010 as residues: Ala-22 to Phe-28.
845882	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1011 as residues: Ala-1 to Gly-7, Arg-29 to Lys-35, Lys-72 to Ala-79, Leu-94 to Val-101, Gly-137 to Asn-142, Arg-145 to Leu-150, Gly-180 to Lys-187, Glu-194 to Gly-208, Arg-257 to Ser-267, Ser-278 to Asp-290, Gly-312 to Ser-319, Leu-338 to Lys-351, Tyr-358 to Ser-363.
846007	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1012 as residues: Tyr-16 to Ala-24, Arg-59 to Ser-66, Thr-78 to Glu-83, Glu-90 to Ser-103, Gln-108 to Thr-113, Ser-115 to Cys-124.
HCRNG17R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1016 as residues: Pro-16 to Asp-21.
HWMFG64R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1017 as residues: Ser-70 to Asp-76, Lys-87 to Leu-95.
HAGCZ94R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1018 as residues: Val-3 to Lys-9.
HBJEJ74R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1019 as residues: Pro-1 to Asp-8.
HUTHM43R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1021 as residues: Pro-7 to Arg-15.
HLTGU75R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1022 as residues: Ser-1 to Gly-11.
HWLKF77R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1023 as residues: Leu-10 to Asn-28.
HWLGX29R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1027 as residues: Val-3 to Ile-10, Pro-34 to Gln-40.
HWMFZ29R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1028 as residues: Leu-7 to Leu-13.
H6EEP19R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1030 as

	residues: Ala-1 to Trp-8, Lys-10 to Asp-27.
HJMAM83R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1031 as residues: Ser-1 to Val-11, Glu-19 to Ala-29, Asp-52 to Ala-68, Gly-78 to Lys-94.
HAGHF58R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1032 as residues: Lys-1 to Val-7.
HDPHG48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1033 as residues: Gly-24 to Lys-34.
HCDMC32R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1038 as residues: Pro-2 to Arg-17, Lys-36 to Pro-47, Phe-61 to Trp-68, Gln-72 to Ala-86.
HTEQO80R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1040 as residues: Gly-1 to Val-15, Pro-17 to Pro-23, Leu-32 to Met-41, Lys-102 to His-109.
H2LAR08R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1043 as residues: Asn-58 to Gly-64.
HWMFN58R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1046 as residues: Glu-6 to Asn-14, Arg-22 to Asp-31, Gly-49 to Thr-56.
HUFBP63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1049 as residues: Pro-1 to Gln-8, Thr-57 to Gly-64, Arg-69 to Arg-74, Gly-80 to Asp-91, Asp-105 to Gln-110, Arg-130 to Tyr-148.
HUFBN90R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1050 as residues: Glu-34 to Ala-40, Arg-111 to Ala-116.
HFKHD61R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1054 as residues: Arg-11 to Gly-38, Arg-44 to Glu-50, Gln-53 to Lys-67.
HTXNL13R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1057 as residues: Ser-48 to Arg-57, Glu-89 to Pro-95, Ser-102 to Asn-107.
H2LAK62R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1059 as residues: Pro-20 to Ser-25.
HATAR77R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1061 as residues: Gly-2 to Arg-16.
HWMEH18R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1066 as residues: Gln-61 to Ser-67.
HCNDP66R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1068 as residues: Leu-8 to Arg-15, Gln-46 to Pro-54.
HCRMK82R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1069 as residues: Ser-32 to Arg-38, Ala-72 to Lys-79, Arg-103 to Phe-111.
HSSGC52R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1075 as residues: Gly-1 to Pro-6, Arg-25 to Ile-30.
HCYBN49R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1076 as residues: Gly-16 to Gly-21, Ile-99 to Gln-109.
HWMGB90R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1077 as residues: Gly-1 to Ala-7, Asp-17 to Arg-27, Glu-32 to Leu-40.
HTEAW21R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1078 as residues: Glu-1 to Gly-6, Gln-19 to Leu-37.
H2LAQ68R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1082 as residues: Val-2 to Trp-10, Leu-25 to Lys-33.
HBAAD60R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1087 as residues: Pro-1 to Lys-32.
HCROA35R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1088 as residues: Gly-6 to Lys-12.
HCROM64R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1089 as residues: Asn-1 to Arg-7.
HKBAG82R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1091 as residues: Pro-9 to Gly-28.
HUTSB76R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1092 as residues: Lys-1 to Ser-17.
HWLJS67R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1093 as residues: Gln-3 to Lys-18, Gln-44 to Glu-49.

HTGAZ53R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1098 as residues: Ser-1 to Ala-16, Gln-36 to Thr-48.
HWLL51R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1100 as residues: Gln-6 to Glu-18.
HWLJZ72R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1103 as residues: Ile-1 to Ser-19.
HWMFG06R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1104 as residues: Arg-1 to Lys-14, Gln-40 to Glu-45, Arg-65 to Arg-80.
HPRT065R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1105 as residues: Thr-12 to Thr-17, Cys-35 to Ser-40.
HUFDC01R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1106 as residues: Pro-11 to Glu-26.
HWLHY44R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1107 as residues: Pro-14 to Gln-24, Cys-34 to Leu-39, Thr-72 to Val-77, Glu-94 to Thr-99, Asp-101 to Met-107, Lys-109 to Pro-116.
HWLGR92R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1108 as residues: Pro-17 to Gly-22.
HCNCQ71R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1109 as residues: Glu-22 to Leu-30.
HWLEN11R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1111 as residues: Pro-6 to Lys-21, Ala-26 to Val-34, Lys-37 to Ser-46.
HWLEH56R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1116 as residues: Thr-23 to Ala-28, Asn-88 to Trp-98, Cys-114 to Asp-131.
H2LAD26R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1117 as residues: Pro-20 to Gly-31.
H2LAK66R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1125 as residues: Pro-33 to Leu-39, Glu-54 to Val-59, Gly-69 to Ser-76.
HSDKC65R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1126 as residues: Asn-32 to Pro-39, Pro-41 to Pro-49.
H2LAK52R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1127 as residues: Pro-20 to Ala-28.
HKAEG12R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1128 as residues: Asp-47 to Lys-52.
HKADP43R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1129 as residues: Pro-7 to Pro-15, Arg-35 to Val-44.
HUSJE17R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1131 as residues: Pro-26 to Gln-32.
HHBEF06R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1133 as residues: Pro-1 to Gly-6.
HISCW28R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1134 as residues: Pro-26 to Gln-32.
HPIAK29R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1137 as residues: Thr-1 to Tyr-7.
HUFAR71R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1138 as residues: Pro-26 to Gln-32.
HOEC121R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1141 as residues: Asn-11 to Pro-20, Pro-22 to Thr-30, Glu-49 to Glu-70, Ser-84 to Thr-96, Thr-108 to Thr-113.
HMCAR63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1143 as residues: Ala-1 to Gly-9, Lys-41 to Glu-47, Asn-65 to Gly-70, Glu-85 to Asp-93, Glu-103 to Tyr-109.
HAICY55R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1152 as residues: Glu-2 to His-9.
HWLIA38R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1153 as residues: Arg-60 to Gly-74, Ser-80 to Ile-88, Leu-92 to Ser-98.
HBXCL69R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1154 as

	residues: Ser-2 to Cys-8, Pro-10 to Leu-17.
H2LAP90R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1155 as residues: Thr-3 to Gln-9, Asn-11 to Pro-19, Gln-35 to Glu-42.
HTELE03R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1157 as residues: Asp-1 to Gln-9, Asn-11 to Arg-16, Cys-28 to Ser-44, Gln-50 to Gln-56.
HJMBN86R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1158 as residues: Ser-31 to Glu-47.
HSKJC32R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1159 as residues: Gln-151 to Glu-158, Glu-168 to Pro-173, Ser-188 to Ile-195.
HAOAG76R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1161 as residues: Gly-1 to Ala-14.
HC1AD45R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1162 as residues: Pro-1 to Lys-23, Pro-43 to Leu-49.
H2MAC82R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1163 as residues: Lys-54 to Lys-59.
H2LAJ41R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1164 as residues: Met-20 to Val-36, Ser-82 to Lys-93, Pro-101 to Arg-106.
HBJFH33R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1166 as residues: Gly-10 to Tyr-26, Asn-29 to Leu-37, Thr-52 to His-59.
HISDV92R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1167 as residues: Pro-3 to Ser-8, Asn-48 to Tyr-54.
HE9QB35R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1169 as residues: Gly-1 to Asp-6, Pro-20 to Gln-33, Tyr-46 to Arg-52, Asn-72 to Lys-85, Gln-91 to Ala-110.
HDABQ50R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1170 as residues: Ser-9 to Lys-17, Lys-41 to Arg-46.
HTPAC28R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1176 as residues: Lys-10 to Thr-15, Thr-17 to Leu-23.
HMCGN07R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1177 as residues: Asn-88 to Ser-98, Pro-123 to Val-129.
HBMVM66R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1180 as residues: Ser-2 to Gly-7, Arg-10 to Phe-24, Ala-36 to Arg-41.
HEPNA09R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1186 as residues: Ser-1 to Pro-6.
HCNDR62R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1190 as residues: Pro-14 to Ser-21.
HNJBF13R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1191 as residues: Asp-18 to Asp-28.
HLYCD69R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1192 as residues: Gly-90 to Thr-109.
HWCAAS3R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1194 as residues: Ser-22 to Gly-28, Glu-37 to Ile-45, Val-67 to Arg-85, Asn-91 to Trp-99.
HFVGP11R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1198 as residues: Ala-4 to Asn-13.
HWLQH07R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1199 as residues: Lys-1 to Lys-25.
HWLKH07R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1201 as residues: Pro-49 to Asp-58.
HAPQC14R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1202 as residues: Lys-1 to Met-8.
HSODB48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1203 as residues: Ser-24 to Gly-31, Ala-37 to Ser-44, Pro-57 to Ser-64, Pro-97 to Gly-104.
HBEAC75R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1204 as residues: Pro-1 to Arg-9.
HBGMJ24R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1205 as residues: Tyr-11 to Val-17, Thr-30 to Phe-48, Gln-150 to Thr-155.

HBJEN94R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1206 as residues: Gln-1 to Asn-6.
HLQGB87R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1213 as residues: Lys-2 to Ser-7.
HAOAC69R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1215 as residues: Ser-2 to Arg-10.
HWLEQ08R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1216 as residues: Glu-21 to His-31.
HKAAY70R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1217 as residues: Gly-6 to Thr-93, Glu-95 to Glu-104, Asp-117 to Asp-125.
HNFJE41R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1221 as residues: Arg-15 to His-21, Pro-48 to Ala-58, Asn-61 to Leu-66, Val-92 to Thr-110, Pro-114 to Thr-120.
HCRMW41R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1224 as residues: Phe-14 to Asn-19.
HOVAX78R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1225 as residues: Gly-1 to Thr-8.
HWAEH57R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1226 as residues: Ser-54 to Tyr-60, Gln-65 to Pro-72, Thr-81 to Gly-92.
HAHEK76R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1230 as residues: Cys-20 to Cys-28.
HOSCG81R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1232 as residues: Thr-8 to Asn-13.
HTFMD43R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1233 as residues: Lys-44 to Ile-52, Arg-57 to Lys-77.
H2LAR73R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1235 as residues: Pro-20 to Arg-27, Asn-47 to Lys-53, Asp-116 to Asn-123, Glu-145 to Gly-154.
HWHPK71R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1238 as residues: Asp-15 to His-24, Pro-27 to Leu-39.
HWBBJ39R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1239 as residues: His-1 to Lys-6.
HSODD94R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1241 as residues: Gly-7 to Glu-15, Gly-29 to Lys-41, Pro-43 to Ser-52, Pro-68 to His-73.
HMIAG25R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1242 as residues: Arg-19 to Ser-41, Pro-43 to Glu-54, Ser-59 to Gly-74.
HCNDW17R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1244 as residues: Lys-7 to Lys-15, Thr-54 to Asn-59.
HWLEY08R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1245 as residues: Glu-9 to Arg-14, Thr-19 to Arg-27, Asp-48 to Ile-57, Gln-63 to Leu-75, Cys-89 to Thr-104, Gly-106 to Pro-113.
HULFN68R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1246 as residues: Ser-1 to Cys-16, Lys-18 to Gly-23, Pro-31 to Tyr-37, Gly-53 to Pro-58.
HTEJJ32R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1249 as residues: Ser-17 to Cys-23, Gln-42 to Leu-51, Ser-68 to Asp-73.
H2CBS58R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1251 as residues: Ser-82 to Phe-88, Lys-110 to Gly-118.
H2LAB77R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1252 as residues: Met-13 to Asp-18, Glu-23 to Ser-43, Glu-45 to Gly-54.
HWAFP88R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1254 as residues: Arg-8 to Lys-13, Gly-35 to Lys-42, Ala-48 to Lys-54.
HWMEB67R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1256 as residues: Arg-9 to Arg-16.
HKMAA52R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1261 as residues: Gly-2 to Lys-10, Asp-36 to Asn-42.
H2LAB37R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1262 as residues: Glu-52 to Thr-59.

H2LAP46R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1263 as residues: Pro-40 to Asn-46, Tyr-71 to Arg-79.
H6BSE61R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1264 as residues: Ile-36 to Asp-41, Ala-54 to Pro-63.
HACBS75R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1269 as residues: Arg-20 to Ser-27, Arg-45 to Trp-59.
HACCA48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1270 as residues: Lys-12 to Lys-26.
HACCS19R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1271 as residues: Gly-1 to Gly-10.
HAGGL96R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1273 as residues: Ser-74 to Phe-88.
HAGGT37R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1274 as residues: Phe-17 to Pro-22.
HAHDR66R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1275 as residues: Gly-11 to Ala-18.
HAJCL80R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1277 as residues: Asn-22 to Phe-32.
HAQMH45R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1283 as residues: Pro-2 to Tyr-13, Leu-21 to Gly-47, Val-49 to Gly-55, Pro-63 to Glu-78.
HBGCA44R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1290 as residues: Thr-20 to Trp-25, Lys-32 to Leu-40.
HBGFX27R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1291 as residues: Ser-1 to Pro-6.
HBGMU38R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1292 as residues: Gln-1 to Phe-8, Thr-34 to Trp-53, Arg-56 to Gly-63, Arg-86 to Cys-102.
HBJED55R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1295 as residues: Arg-6 to Pro-14.
HBMTJ51R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1300 as residues: Cys-8 to Asp-13.
HBWBD78R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1302 as residues: Pro-51 to Ala-58.
HCDDQ63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1307 as residues: Gln-1 to Lys-10.
HCFC01R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1310 as residues: Ser-1 to Thr-6.
HCFCR43R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1311 as residues: Arg-10 to Thr-20.
HCHAO92R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1313 as residues: Asn-19 to Arg-25.
HCHOH49R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1314 as residues: Asn-19 to Asp-30.
HCHPG05R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1315 as residues: Pro-6 to Ser-11.
HCIAD24R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1316 as residues: Lys-1 to Gly-7.
HCNCY51R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1319 as residues: Lys-10 to Arg-16.
HCNCY63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1320 as residues: Gly-1 to Lys-9.
HCNDO71R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1321 as residues: Lys-33 to Ile-42, Arg-51 to Phe-64.
HCQBN22R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1324 as residues: Lys-1 to Asn-11.
HCQCL27R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1325 as residues: Gly-7 to His-27.

HCQCL48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1326 as residues: Ala-1 to Thr-13.
HCQDJ42R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1330 as residues: Glu-8 to Asn-13, Arg-16 to Glu-24.
HCRMD77R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1331 as residues: Asn-4 to Asn-10.
HCROJ68R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1339 as residues: Ile-2 to His-8.
HCROM30R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1342 as residues: Glu-1 to Glu-7, Pro-26 to Leu-32, Gly-37 to Gln-44, Thr-84 to Thr-92.
HCROQ34R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1343 as residues: Asn-1 to Asp-11.
HCROZ66R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1345 as residues: Arg-7 to Lys-13.
HCRPC61R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1346 as residues: Ala-3 to Gly-8.
HCRPG28R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1347 as residues: Pro-26 to Ser-32.
HCRPN52R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1349 as residues: Ser-24 to Lys-30, Lys-54 to Ser-61.
HDCAA21R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1354 as residues: Phe-6 to Val-12, Ile-15 to Phe-20.
HDDAA85R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1355 as residues: Lys-18 to Lys-24.
HDPGO03R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1356 as residues: Ala-4 to Gln-17.
HDPLB08R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1357 as residues: Pro-2 to Tyr-13, Leu-21 to Ala-36.
HDQEX80R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1359 as residues: Arg-1 to Arg-6, Phe-27 to Arg-32, Pro-37 to Lys-42, Arg-47 to Trp-53, Arg-55 to Ser-61.
HDRMI91R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1360 as residues: Thr-1 to Lys-8.
HE6DJ45R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1364 as residues: Pro-1 to Asn-8.
HE9FH12R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1366 as residues: Asn-12 to Ser-20.
HEAAL59R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1370 as residues: Gln-20 to Asn-25.
HEGAR32R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1371 as residues: Lys-9 to Ser-19.
HEGAR85R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1372 as residues: Ser-16 to His-46, Arg-49 to Thr-58.
HELFE05R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1373 as residues: Tyr-8 to Leu-16.
HEMFI88R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1374 as residues: Pro-6 to Ala-13.
HEMFR18R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1375 as residues: Ala-1 to Ala-10, Pro-12 to Gly-17, Ala-22 to Cys-27, Glu-30 to Arg-35, Pro-43 to Ser-50.
HEONL43R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1376 as residues: Arg-1 to Val-10.
HFADM62R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1380 as residues: Lys-6 to Lys-14.
HFATE31R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1381 as residues: Asp-1 to Arg-9, Arg-20 to Arg-26, Glu-33 to Gly-40.

HFCEL77R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1383 as residues: Glu-33 to Ser-48, Ile-54 to Ile-63, Leu-79 to Asp-84.
HFTBI57R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1392 as residues: Pro-18 to Ser-23.
HFXGX46R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1394 as residues: Pro-11 to Gln-28.
HHBEW72R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1400 as residues: Pro-20 to Thr-27.
HHERT59R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1401 as residues: Arg-1 to Trp-9.
HJMAH76R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1405 as residues: Cys-10 to Ala-15.
HJMAN56R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1406 as residues: Ala-45 to Asp-60.
HJMAO54R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1407 as residues: Pro-28 to Gln-39, Pro-65 to Cys-80.
HKLSD93R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1409 as residues: Gly-11 to Gly-17.
HLMFH16R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1410 as residues: Gly-1 to Asp-8.
HLQCQ73R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1412 as residues: Glu-1 to Gly-6, Arg-8 to Phe-13.
HLQEF47R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1413 as residues: Leu-8 to Leu-13.
HLQFM50R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1414 as residues: Gly-29 to Asp-34.
HLQGA76R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1416 as residues: Ser-16 to Ser-33.
HLTEV09R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1418 as residues: Arg-9 to Asn-17.
HMACF85R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1421 as residues: Glu-29 to Lys-34, Leu-113 to Gln-120.
HMAIA15R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1422 as residues: Lys-15 to Gln-21, Ile-51 to Gly-57, Lys-72 to Gly-83.
HMCIS54R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1424 as residues: Lys-3 to His-24.
HNHMR05R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1427 as residues: Pro-9 to Gly-20, Thr-26 to Arg-42, Ala-48 to Ser-54.
HNJBB78R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1428 as residues: Thr-6 to Lys-13, Leu-48 to Asn-54.
HOCND06R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1433 as residues: Pro-2 to Tyr-13, Leu-21 to Ala-35.
HOCND49R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1434 as residues: Asn-2 to Gly-12, Ile-14 to Ala-30.
HODFA26R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1436 as residues: Glu-1 to His-6, Gly-19 to Asp-29, Leu-44 to Leu-49.
HODHL89R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1437 as residues: Ser-16 to His-46, Arg-49 to Thr-58.
HOEJM67R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1438 as residues: Ser-19 to Lys-25, Asp-29 to Glu-55, Ser-102 to Thr-107.
HGOBN48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1439 as residues: Lys-14 to Arg-19, Asp-25 to Phe-32.
HOUHN53R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1442 as residues: Glu-1 to His-6, Gly-19 to Trp-31.
HPBEE63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1444 as residues: Pro-14 to Gly-20, His-28 to Arg-35.

HPJBE91R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1446 as residues: Ser-15 to Asn-20, Ala-22 to Ile-49, Lys-52 to Val-57, Tyr-71 to Cys-83, Thr-90 to Tyr-95.
HSDZG83R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1454 as residues: Val-17 to Lys-22.
HSICQ60R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1455 as residues: Val-12 to Gly-17.
HSIFA64R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1456 as residues: His-17 to Ile-22, Leu-33 to Pro-40.
HSKYE52R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1458 as residues: Pro-2 to Ser-7.
HSODA95R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1460 as residues: Ser-14 to His-44, Arg-47 to Thr-56.
HSSGK43R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1462 as residues: Ser-24 to Leu-35, Pro-38 to Ser-45.
HTXFA64R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1470 as residues: Thr-1 to Glu-8.
HUSJF91R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1471 as residues: Glv-1 to Gly-6.
HUSJN48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1472 as residues: Ser-16 to Tyr-24.
HUSZN23R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1474 as residues: Ser-16 to Lys-24.
HUTSD20R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1475 as residues: Arg-10 to Asn-20.
HWAFI63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1477 as residues: Pro-15 to Gly-24, Pro-26 to Arg-45.
HWAGZ89R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1478 as residues: Ser-47 to Lys-52.
HWHHM83R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1480 as residues: Leu-1 to Gly-6.
HWLBS90R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1484 as residues: Lys-37 to Asn-44.
HWLEH13R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1486 as residues: Gln-22 to Glu-29.
HWLEJ67R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1487 as residues: Asn-5 to Trp-13.
HWLEM49R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1488 as residues: Glu-1 to His-6, Gly-19 to Trp-31.
HWLGM21R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1492 as residues: Glu-1 to His-6, Gly-19 to Trp-31.
HWLGS46R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1494 as residues: Glu-17 to Asn-23, Glu-38 to Gly-49.
HWLGU40R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1495 as residues: His-10 to Pro-15.
HWLGX63R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1496 as residues: Glu-1 to Asn-7.
HWLHD09R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1497 as residues: Pro-6 to Ala-37, Arg-40 to Ser-49.
HWLHW89R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1500 as residues: Asn-1 to Lys-16, Glu-32 to Ser-41, Leu-57 to Gly-71.
HWLJL19R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1506 as residues: Arg-46 to Phe-58.
HWLKG82R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1508 as residues: Pro-5 to Gly-25, Ser-29 to Leu-36, Arg-49 to Phe-55.
HWLKM86R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1512 as

	residues: Arg-10 to Lys-23.
HWLQS83R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1515 as residues: Ala-1 to Arg-6.
HWLRP86R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1518 as residues: Tyr-3 to Gly-10.
HWLRQ49R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1519 as residues: Pro-19 to Ser-26. Gln-44 to Lys-52.
HWLUF60R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1520 as residues: Gln-7 to Lys-31.
HWLUR41R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1522 as residues: Ser-24 to Trp-30.
HWLVD60R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1523 as residues: Cys-15 to Lys-51.
HWMAN61R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1525 as residues: Ser-21 to Asp-26.
HWMEH26R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1528 as residues: Ser-16 to His-46. Arg-49 to Thr-58.
HWMEL50R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1529 as residues: Pro-24 to Thr-40. Phe-63 to Arg-69.
HWMFB31R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1530 as residues: Asn-2 to Lys-10. Cys-16 to Pro-28. Ser-36 to Glu-41.
HWMFO93R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1532 as residues: Ser-8 to Gln-14.
HMAFE48R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1537 as residues: Glu-9 to Gly-17.
HRODJ88R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1538 as residues: Gly-6 to Tyr-14.
HWLAR31R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1539 as residues: Glu-9 to Gly-17.
H2LAU24R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1541 as residues: Glu-11 to Gly-19.
HATDR94R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1542 as residues: Glu-14 to Lys-19. Asn-21 to Gly-27.
HWLLI85R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1543 as residues: Val-19 to Asn-32.
HSYCH41R	Preferred epitopes include those comprising a sequence shown in SEQ ID NO. 1545 as residues: Thr-71 to Ile-79.

The present invention encompasses polypeptides comprising, or alternatively consisting of, an epitope of the polypeptide sequence shown in SEQ ID NO:Y, or an epitope of the polypeptide sequence encoded by the cDNA in the related cDNA clone contained in a deposited library or encoded by a polynucleotide that hybridizes to the complement of an epitope encoding sequence of SEQ ID NO:X, or an epitope encoding sequence contained in the deposited cDNA clone under stringent hybridization conditions, or alternatively, under lower stringency hybridization conditions, as defined supra. The present invention further encompasses polynucleotide sequences encoding an epitope of a polypeptide sequence of the invention (such as, for example, the sequence disclosed in SEQ ID NO:X), polynucleotide sequences of the complementary strand of a polynucleotide sequence encoding an epitope of the invention, and polynucleotide sequences which hybridize to this complementary strand under stringent hybridization conditions or alternatively, under lower stringency hybridization conditions, as defined supra.

The term "epitopes," as used herein, refers to portions of a polypeptide having antigenic or immunogenic activity in an animal, preferably a mammal, and most preferably in a human. In a preferred embodiment, the present invention encompasses a polypeptide comprising an epitope, as well as the polynucleotide encoding this polypeptide. An "immunogenic epitope," as used herein, is defined as a portion of a protein that elicits an antibody response in an animal, as determined by any method known in the art, for example, by the methods for generating antibodies described infra. (See, for example, Geysen et al., Proc. Natl. Acad. Sci. USA 81:3998- 4002 (1983)). The term "antigenic epitope," as used herein, is defined as a portion of a protein to which an antibody can immunospecifically bind its antigen as determined by any method well known in the art, for example, by the immunoassays described herein. Immunospecific binding excludes non-specific binding but does not necessarily exclude cross- reactivity with other antigens. Antigenic epitopes need not necessarily be immunogenic.

Fragments which function as epitopes may be produced by any conventional means. (See, e.g., Houghten, R. A., Proc. Natl. Acad. Sci. USA 82:5131-5135 (1985) further described in U.S. Patent No. 4,631,211.)

In the present invention, antigenic epitopes preferably contain a sequence of at least 4, at least 5, at least 6, at least 7, more preferably at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 40, at

least 50, and, most preferably, between about 15 to about 30 amino acids. Preferred polypeptides comprising immunogenic or antigenic epitopes are at least 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 amino acid residues in length. Additional non-exclusive preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as portions thereof. Antigenic epitopes are useful, for example, to raise antibodies, including monoclonal antibodies, that specifically bind the epitope. Preferred antigenic epitopes include the antigenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these antigenic epitopes. Antigenic epitopes can be used as the target molecules in immunoassays. (See, for instance, Wilson et al., Cell 5 37:767-778 (1984); Sutcliffe et al., Science 219:660-666 (1983)).

Similarly, immunogenic epitopes can be used, for example, to induce antibodies according to methods well known in the art. (See, for instance, Sutcliffe et al., supra; Wilson et al., supra; Chow et al., Proc. Natl. Acad. Sci. USA 82:910-914; and Bittle et al., J. Gen. Virol. 66:2347-2354 (1985). Preferred immunogenic epitopes include the immunogenic epitopes disclosed herein, as well as any combination of two, three, four, five or more of these immunogenic epitopes. The polypeptides comprising one or more immunogenic epitopes may be presented for eliciting an antibody response together with a carrier protein, such as an albumin, to an animal system (such as rabbit or mouse), or, if the polypeptide is of sufficient length (at least about 25 amino acids), the polypeptide may be presented without a carrier. However, immunogenic epitopes comprising as few as 8 to 10 amino acids have 15 been shown to be sufficient to raise antibodies capable of binding to, at the very least, linear epitopes in a denatured polypeptide (e.g., in Western blotting).

Epitope-bearing polypeptides of the present invention may be used to induce antibodies according to methods well known in the art including, but not limited to, in vivo immunization, in vitro immunization, and phage display methods. See, e.g., Sutcliffe et al., 25 supra; Wilson et al., supra, and Bittle et al., J. Gen. Virol., 66:2347-2354 (1985). If in vivo immunization is used, animals may be immunized with free peptide; however, anti-peptide antibody titer may be boosted by coupling the peptide to a macromolecular carrier, such as keyhole limpet hemacyanin (KLH) or tetanus toxoid. For instance, peptides containing cysteine residues may be coupled to a carrier using a linker such as maleimidobenzoyl- N-hydroxysuccinimide ester (MBS), while other peptides may be coupled to carriers using a more general linking agent such as glutaraldehyde. Animals such as rabbits, rats and mice 30

are immunized with either free or carrier- coupled peptides, for instance, by intraperitoneal and/or intradermal injection of emulsions containing about 100 µg of peptide or carrier protein and Freund's adjuvant or any other adjuvant known for stimulating an immune response. Several booster injections may be needed, for instance, at intervals of about two weeks, to provide a useful titer of anti-peptide antibody which can be detected, for example, by ELISA assay using free peptide adsorbed to a solid surface. The titer of anti-peptide antibodies in serum from an immunized animal may be increased by selection of anti-peptide antibodies, for instance, by adsorption to the peptide on a solid support and elution of the selected antibodies according to methods well known in the art.

As one of skill in the art will appreciate, and as discussed above, the polypeptides of the present invention, and immunogenic and/or antigenic epitope fragments thereof can be fused to other polypeptide sequences. For example, the polypeptides of the present invention may be fused with the constant domain of immunoglobulins (IgA, IgE, IgG, IgM), or portions thereof (CH1, CH2, CH3, or any combination thereof and portions thereof) resulting in chimeric polypeptides. Such fusion proteins may facilitate purification and may increase half-life in vivo. This has been shown for chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. See, e.g., EP 394,827; Traunecker et al., *Nature*, 331:84-86 (1988). Enhanced delivery of an antigen across the epithelial barrier to the immune system has been demonstrated for antigens (e.g., insulin) conjugated to an FcRn binding partner such as IgG or Fc fragments (see, e.g., PCT Publications WO 96/22024 and WO 99/04813). IgG Fusion proteins that have a disulfide-linked dimeric structure due to the IgG portion disulfide bonds have also been found to be more efficient in binding and neutralizing other molecules than monomeric polypeptides or fragments thereof alone. See, e.g., Fountoulakis et al., *J. Biochem.*, 270:3958-3964 (1995).

Similarly, EP-A-O 464 533 (Canadian counterpart 2045869) discloses fusion proteins comprising various portions of constant region of immunoglobulin molecules together with another human protein or part thereof. In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP-A 0232 262.) Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, may be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for

immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, D. Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); K. Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995).)

5 Moreover, the polypeptides of the present invention can be fused to marker sequences, such as a peptide which facilitates purification of the fused polypeptide. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for
10 convenient purification of the fusion protein. Another peptide tag useful for purification, the "HA" tag, corresponds to an epitope derived from the influenza hemagglutinin protein. (Wilson et al., *Cell* 37:767 (1984).)

15 Thus, any of these above fusions can be engineered using the polynucleotides or the polypeptides of the present invention.

20 Nucleic acids encoding the above epitopes can also be recombined with a gene of interest as an epitope tag (e.g., the hemagglutinin ("HA") tag or flag tag) to aid in detection and purification of the expressed polypeptide. For example, a system described by Janknecht et al. allows for the ready purification of non-denatured fusion proteins expressed in human cell lines (Janknecht et al., *Proc. Natl. Acad. Sci. USA* 88:8972- 897 (1991)). In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the open reading frame of the gene is translationally fused to an amino-terminal tag consisting of six histidine residues. The tag serves as a matrix binding domain for the fusion protein. Extracts from cells infected with the recombinant vaccinia virus are loaded onto
25 Ni²⁺ nitriloacetic acid-agarose column and histidine-tagged proteins can be selectively eluted with imidazole-containing buffers.

30 Additional fusion proteins of the invention may be generated through the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or codon-shuffling (collectively referred to as "DNA shuffling"). DNA shuffling may be employed to modulate the activities of polypeptides of the invention, such methods can be used to generate polypeptides with altered activity, as well as agonists and antagonists of the polypeptides. See, generally, U.S. Patent Nos. 5,605,793; 5,811,238; 5,830,721; 5,834,252; and 5,837,458, and Patten et al.,

Curr. Opin. Biotechnol. 8:724-33 (1997); Harayama, Trends Biotechnol. 16(2):76-82 (1998); Hansson, et al., J. Mol. Biol. 287:265-76 (1999); and Lorenzo and Blasco, Biotechniques 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference in its entirety). In one embodiment, alteration of polynucleotides
5 corresponding to SEQ ID NO:X and the polypeptides encoded by these polynucleotides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments by homologous or site-specific recombination to generate variation in the polynucleotide sequence. In another embodiment, polynucleotides of the invention, or the encoded polypeptides, may be altered by being subjected to random mutagenesis by error-prone
10 PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of a polynucleotide encoding a polypeptide of the invention may be recombined with one or more components, motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules.

15 As discussed herein, any polypeptide of the present invention can be used to generate fusion proteins. For example, the polypeptide of the present invention, when fused to a second protein, can be used as an antigenic tag. Antibodies raised against the polypeptide of the present invention can be used to indirectly detect the second protein by binding to the polypeptide. Moreover, because secreted proteins target cellular locations based on
20 trafficking signals, polypeptides of the present invention which are shown to be secreted can be used as targeting molecules once fused to other proteins.

Examples of domains that can be fused to polypeptides of the present invention include not only heterologous signal sequences, but also other heterologous functional regions. The fusion does not necessarily need to be direct, but may occur through linker
25 sequences.

In certain preferred embodiments, proteins of the invention comprise fusion proteins wherein the polypeptides are N and/or C-terminal deletion mutants. In preferred embodiments, the application is directed to nucleic acid molecules at least 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to the nucleic acid sequences encoding polypeptides
30 having the amino acid sequence of the specific N- and C-terminal deletions mutants. Polynucleotides encoding these polypeptides are also encompassed by the invention.

Moreover, fusion proteins may also be engineered to improve characteristics of the polypeptide of the present invention. For instance, a region of additional amino acids, particularly charged amino acids, may be added to the N-terminus of the polypeptide to improve stability and persistence during purification from the host cell or subsequent handling and storage. Also, peptide moieties may be added to the polypeptide to facilitate purification. Such regions may be removed prior to final preparation of the polypeptide. The addition of peptide moieties to facilitate handling of polypeptides are familiar and routine techniques in the art.

10 **Vectors, Host Cells, and Protein Production**

The present invention also relates to vectors containing the polynucleotide of the present invention, host cells, and the production of polypeptides by recombinant techniques. The vector may be, for example, a phage, plasmid, viral, or retroviral vector. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host cells.

The polynucleotides of the invention may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector is introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and then transduced into host cells.

The polynucleotide insert should be operatively linked to an appropriate promoter, such as the phage lambda PL promoter, the E. coli lac, trp, phoA and tac promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs will preferably include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors will preferably include at least one selectable marker. Such markers include dihydrofolate reductase, G418 or neomycin resistance for eukaryotic cell culture and tetracycline, kanamycin or ampicillin resistance genes for culturing in E. coli and other bacteria. Representative examples of appropriate hosts include,

but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS, 293, and Bowes melanoma cells; and plant cells.

5 Appropriate culture mediums and conditions for the above-described host cells are known in the art.

Among vectors preferred for use in bacteria include pQE70, pQE60 and pQE-9, available from QIAGEN, Inc.; pBluescript vectors, Phagescript vectors, pNH8A, pNH16a, pNH18A, pNH46A, available from Stratagene Cloning Systems, Inc.; and ptrc99a, pKK223-
10 3, pKK233-3, pDR540, pRIT5 available from Pharmacia Biotech, Inc. Among preferred eukaryotic vectors are pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; and pSVK3, pBPV, pMSG and pSVL available from Pharmacia. Preferred expression vectors for use in yeast systems include, but are not limited to pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalph, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1,
15 pPIC3.5K, pPIC9K, and PAO815 (all available from Invitrogen, Carlsbad, CA). Other suitable vectors will be readily apparent to the skilled artisan.

Introduction of the construct into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in
20 many standard laboratory manuals, such as Davis et al., Basic Methods In Molecular Biology (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

A polypeptide of this invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid
25 extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography ("HPLC") is employed for purification.

Polypeptides of the present invention can also be recovered from: products purified
30 from natural sources, including bodily fluids, tissues and cells, whether directly isolated or cultured; products of chemical synthetic procedures; and products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast,

higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, polypeptides of the invention may also include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

In one embodiment, the yeast *Pichia pastoris* is used to express polypeptides of the invention in a eukaryotic system. *Pichia pastoris* is a methylotrophic yeast which can metabolize methanol as its sole carbon source. A main step in the methanol metabolism pathway is the oxidation of methanol to formaldehyde using O₂. This reaction is catalyzed by the enzyme alcohol oxidase. In order to metabolize methanol as its sole carbon source, *Pichia pastoris* must generate high levels of alcohol oxidase due, in part, to the relatively low affinity of alcohol oxidase for O₂. Consequently, in a growth medium depending on methanol as a main carbon source, the promoter region of one of the two alcohol oxidase genes (*AOX1*) is highly active. In the presence of methanol, alcohol oxidase produced from the *AOX1* gene comprises up to approximately 30% of the total soluble protein in *Pichia pastoris*. See, Ellis, S.B., et al., *Mol. Cell. Biol.* 5:1111-21 (1985); Koutz, P.J., et al., *Yeast* 5:167-77 (1989); Tschopp, J.F., et al., *Nucl. Acids Res.* 15:3859-76 (1987). Thus, a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, under the transcriptional regulation of all or part of the *AOX1* regulatory sequence is expressed at exceptionally high levels in *Pichia* yeast grown in the presence of methanol.

In one example, the plasmid vector pPIC9K is used to express DNA encoding a polypeptide of the invention, as set forth herein, in a *Pichea* yeast system essentially as described in "*Pichia* Protocols: Methods in Molecular Biology," D.R. Higgins and J. Cregg, eds. The Humana Press. Totowa, NJ, 1998. This expression vector allows expression and secretion of a polypeptide of the invention by virtue of the strong *AOX1* promoter linked to

the *Pichia pastoris* alkaline phosphatase (PHO) secretory signal peptide (i.e., leader) located upstream of a multiple cloning site.

Many other yeast vectors could be used in place of pPIC9K, such as, pYES2, pYD1, pTEF1/Zeo, pYES2/GS, pPICZ, pGAPZ, pGAPZalpha, pPIC9, pPIC3.5, pHIL-D2, pHIL-S1, pPIC3.5K, and PAO815. as one skilled in the art would readily appreciate, as long as the proposed expression construct provides appropriately located signals for transcription, translation, secretion (if desired), and the like, including an in-frame AUG as required.

In another embodiment, high-level expression of a heterologous coding sequence, such as, for example, a polynucleotide of the present invention, may be achieved by cloning the heterologous polynucleotide of the invention into an expression vector such as, for example, pGAPZ or pGAPZalpha, and growing the yeast culture in the absence of methanol.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., coding sequence), and/or to include genetic material (e.g., heterologous polynucleotide sequences) that is operably associated with polynucleotides of the invention, and which activates, alters, and/or amplifies endogenous polynucleotides. For example, techniques known in the art may be used to operably associate heterologous control regions (e.g., promoter and/or enhancer) and endogenous polynucleotide sequences via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996; International Publication No. WO 94/12650, published August 4, 1994; Koller et al., Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); and Zijlstra et al., Nature 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

In addition, polypeptides of the invention can be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, Proteins: Structures and Molecular Principles, W.H. Freeman & Co., N.Y., and Hunkapiller et al., Nature, 310:105-111 (1984)). For example, a polypeptide corresponding to a fragment of a polypeptide can be synthesized by use of a peptide synthesizer. Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs can be introduced as a substitution or addition into the

polypeptide sequence. Non-classical amino acids include, but are not limited to, to the D-isomers of the common amino acids, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, Abu, 2-amino butyric acid, γ -Abu, e-Ahx, 6-amino hexanoic acid, Aib, 2-amino isobutyric acid, 3-amino propionic acid, ornithine, norleucine, norvaline, 5 hydroxyproline, sarcosine, citrulline, homocitrulline, cysteic acid, t-butylglycine, t-butylalanine, phenylglycine, cyclohexylalanine, b-alanine, fluoro-amino acids, designer amino acids such as b-methyl amino acids, Ca-methyl amino acids, Na-methyl amino acids, and amino acid analogs in general. Furthermore, the amino acid can be D (dextrorotary) or L (levorotary).

10 Non-naturally occurring variants may be produced using art-known mutagenesis techniques, which include, but are not limited to oligonucleotide mediated mutagenesis, alanine scanning, PCR mutagenesis, site directed mutagenesis (*see, e.g., Carter et al., Nucl. Acids Res. 13:4331 (1986); and Zoller et al., Nucl. Acids Res. 10:6487 (1982)*), cassette mutagenesis (*see, e.g., Wells et al., Gene 34:315 (1985)*), restriction selection mutagenesis 15 (*see, e.g., Wells et al., Philos. Trans. R. Soc. London SerA 317:415 (1986)*).

The invention additionally, encompasses polypeptides of the present invention which are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous 20 chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH_4 ; acetylation, formylation, oxidation, reduction; metabolic synthesis in the presence of tunicamycin; etc.

Additional post-translational modifications encompassed by the invention include, for 25 example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of procaryotic host cell expression. The polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, 30 isotopic or affinity label to allow for detection and isolation of the protein.

Also provided by the invention are chemically modified derivatives of the polypeptides of the invention which may provide additional advantages such as increased

solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent No. 4,179,337). The chemical moieties for derivitization may be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol and the like. The polypeptides may be modified at random positions within the molecule, or at predetermined positions within the molecule and may include one, two, three or more attached chemical moieties.

The polymer may be of any molecular weight, and may be branched or unbranched. For polyethylene glycol, the preferred molecular weight is between about 1 kDa and about 100 kDa (the term "about" indicating that in preparations of polyethylene glycol, some molecules will weigh more, some less, than the stated molecular weight) for ease in handling and manufacturing. Other sizes may be used, depending on the desired therapeutic profile (e.g., the duration of sustained release desired, the effects, if any on biological activity, the ease in handling, the degree or lack of antigenicity and other known effects of the polyethylene glycol to a therapeutic protein or analog). For example, the polyethylene glycol may have an average molecular weight of about 200; 500; 1000; 1500; 2000; 2500; 3000; 3500; 4000; 4500; 5000; 5500; 6000; 6500; 7000; 7500; 8000; 8500; 9000; 9500; 10,000; 10,500; 11,000; 11,500; 12,000; 12,500; 13,000; 13,500; 14,000; 14,500; 15,000; 15,500; 16,000; 16,500; 17,000; 17,500; 18,000; 18,500; 19,000; 19,500; 20,000; 25,000; 30,000; 35,000; 40,000; 50,000; 55,000; 60,000; 65,000; 70,000; 75,000; 80,000; 85,000; 90,000; 95,000; or 100,000 kDa.

As noted above, the polyethylene glycol may have a branched structure. Branched polyethylene glycols are described, for example, in U.S. Patent No. 5,643,575; Morpurgo *et al.*, *Appl. Biochem. Biotechnol.* 56:59-72 (1996); Vorobjev *et al.*, *Nucleosides Nucleotides* 18:2745-2750 (1999); and Caliceti *et al.*, *Bioconjug. Chem.* 10:638-646 (1999), the disclosures of each of which are incorporated herein by reference.

The polyethylene glycol molecules (or other chemical moieties) should be attached to the protein with consideration of effects on functional or antigenic domains of the protein. There are a number of attachment methods available to those skilled in the art, e.g., EP 0 401 384, herein incorporated by reference (coupling PEG to G-CSF), see also Malik *et al.*, *Exp. Hematol.* 20:1028-1035 (1992) (reporting pegylation of GM-CSF using tresyl chloride). For example, polyethylene glycol may be covalently bound through amino acid residues via a

reactive group, such as, a free amino or carboxyl group. Reactive groups are those to which an activated polyethylene glycol molecule may be bound. The amino acid residues having a free amino group may include lysine residues and the N-terminal amino acid residues; those having a free carboxyl group may include aspartic acid residues glutamic acid residues and the C-terminal amino acid residue. Sulfhydryl groups may also be used as a reactive group for attaching the polyethylene glycol molecules. Preferred for therapeutic purposes is attachment at an amino group, such as attachment at the N-terminus or lysine group.

As suggested above, polyethylene glycol may be attached to proteins via linkage to any of a number of amino acid residues. For example, polyethylene glycol can be linked to a proteins via covalent bonds to lysine, histidine, aspartic acid, glutamic acid, or cysteine residues. One or more reaction chemistries may be employed to attach polyethylene glycol to specific amino acid residues (e.g., lysine, histidine, aspartic acid, glutamic acid, or cysteine) of the protein or to more than one type of amino acid residue (e.g., lysine, histidine, aspartic acid, glutamic acid, cysteine and combinations thereof) of the protein.

One may specifically desire proteins chemically modified at the N-terminus. Using polyethylene glycol as an illustration of the present composition, one may select from a variety of polyethylene glycol molecules (by molecular weight, branching, etc.), the proportion of polyethylene glycol molecules to protein (polypeptide) molecules in the reaction mix, the type of pegylation reaction to be performed, and the method of obtaining the selected N-terminally pegylated protein. The method of obtaining the N-terminally pegylated preparation (i.e., separating this moiety from other monopegylated moieties if necessary) may be by purification of the N-terminally pegylated material from a population of pegylated protein molecules. Selective proteins chemically modified at the N-terminus modification may be accomplished by reductive alkylation which exploits differential reactivity of different types of primary amino groups (lysine versus the N-terminal) available for derivatization in a particular protein. Under the appropriate reaction conditions, substantially selective derivatization of the protein at the N-terminus with a carbonyl group containing polymer is achieved.

As indicated above, pegylation of the proteins of the invention may be accomplished by any number of means. For example, polyethylene glycol may be attached to the protein either directly or by an intervening linker. Linkerless systems for attaching polyethylene glycol to proteins are described in Delgado *et al.*, *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-

304 (1992); Francis *et al.*, *Intern. J. of Hematol.* 68:1-18 (1998); U.S. Patent No. 4,002,531; U.S. Patent No. 5,349,052; WO 95/06058; and WO 98/32466, the disclosures of each of which are incorporated herein by reference.

One system for attaching polyethylene glycol directly to amino acid residues of proteins without an intervening linker employs tresylated MPEG, which is produced by the modification of monmethoxy polyethylene glycol (MPEG) using tresylchloride ($\text{ClSO}_2\text{CH}_2\text{CF}_3$). Upon reaction of protein with tresylated MPEG, polyethylene glycol is directly attached to amine groups of the protein. Thus, the invention includes protein-polyethylene glycol conjugates produced by reacting proteins of the invention with a polyethylene glycol molecule having a 2,2,2-trifluoroethane sulphonyl group.

Polyethylene glycol can also be attached to proteins using a number of different intervening linkers. For example, U.S. Patent No. 5,612,460, the entire disclosure of which is incorporated herein by reference, discloses urethane linkers for connecting polyethylene glycol to proteins. Protein-polyethylene glycol conjugates wherein the polyethylene glycol is attached to the protein by a linker can also be produced by reaction of proteins with compounds such as MPEG-succinimidylsuccinate, MPEG activated with 1,1'-carbonyldiimidazole, MPEG-2,4,5-trichloropenylcarbonate, MPEG-p-nitrophenolcarbonate, and various MPEG-succinate derivatives. A number additional polyethylene glycol derivatives and reaction chemistries for attaching polyethylene glycol to proteins are described in WO 98/32466, the entire disclosure of which is incorporated herein by reference. Pegylated protein products produced using the reaction chemistries set out herein are included within the scope of the invention.

The number of polyethylene glycol moieties attached to each protein of the invention (*i.e.*, the degree of substitution) may also vary. For example, the pegylated proteins of the invention may be linked, on average, to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12, 15, 17, 20, or more polyethylene glycol molecules. Similarly, the average degree of substitution within ranges such as 1-3, 2-4, 3-5, 4-6, 5-7, 6-8, 7-9, 8-10, 9-11, 10-12, 11-13, 12-14, 13-15, 14-16, 15-17, 16-18, 17-19, or 18-20 polyethylene glycol moieties per protein molecule. Methods for determining the degree of substitution are discussed, for example, in Delgado *et al.*, *Crit. Rev. Thera. Drug Carrier Sys.* 9:249-304 (1992).

The colon cancer antigen polypeptides of the invention may be in monomers or multimers (*i.e.*, dimers, trimers, tetramers and higher multimers). Accordingly, the present

invention relates to monomers and multimers of the polypeptides of the invention, their preparation, and compositions (preferably, Therapeutics) containing them. In specific embodiments, the polypeptides of the invention are monomers, dimers, trimers or tetramers. In additional embodiments, the multimers of the invention are at least dimers, at least trimers, or at least tetramers.

Multimers encompassed by the invention may be homomers or heteromers. As used herein, the term homomer, refers to a multimer containing only polypeptides corresponding to the amino acid sequence of SEQ ID NO:Y or an amino acid sequence encoded by SEQ ID NO:X, and/or an amino acid sequence encoded by the cDNA in a related cDNA clone contained in a deposited library (including fragments, variants, splice variants, and fusion proteins, corresponding to any one of these as described herein). These homomers may contain polypeptides having identical or different amino acid sequences. In a specific embodiment, a homomer of the invention is a multimer containing only polypeptides having an identical amino acid sequence. In another specific embodiment, a homomer of the invention is a multimer containing polypeptides having different amino acid sequences. In specific embodiments, the multimer of the invention is a homodimer (e.g., containing polypeptides having identical or different amino acid sequences) or a homotrimer (e.g., containing polypeptides having identical and/or different amino acid sequences). In additional embodiments, the homomeric multimer of the invention is at least a homodimer, at least a homotrimer, or at least a homotetramer.

As used herein, the term heteromer refers to a multimer containing one or more heterologous polypeptides (i.e., polypeptides of different proteins) in addition to the polypeptides of the invention. In a specific embodiment, the multimer of the invention is a heterodimer, a heterotrimer, or a heterotetramer. In additional embodiments, the heteromeric multimer of the invention is at least a heterodimer, at least a heterotrimer, or at least a heterotetramer.

Multimers of the invention may be the result of hydrophobic, hydrophilic, ionic and/or covalent associations and/or may be indirectly linked, by for example, liposome formation. Thus, in one embodiment, multimers of the invention, such as, for example, homodimers or homotrimers, are formed when polypeptides of the invention contact one another in solution. In another embodiment, heteromultimers of the invention, such as, for example, heterotrimers or heterotetramers, are formed when polypeptides of the invention

contact antibodies to the polypeptides of the invention (including antibodies to the heterologous polypeptide sequence in a fusion protein of the invention) in solution. In other embodiments, multimers of the invention are formed by covalent associations with and/or between the polypeptides of the invention. Such covalent associations may involve one or more amino acid residues contained in the polypeptide sequence (e.g., that recited in SEQ ID NO:Y, or contained in a polypeptide encoded by SEQ ID NO:X, and/or by the cDNA in the related cDNA clone contained in a deposited library). In one instance, the covalent associations are cross-linking between cysteine residues located within the polypeptide sequences which interact in the native (i.e., naturally occurring) polypeptide. In another instance, the covalent associations are the consequence of chemical or recombinant manipulation. Alternatively, such covalent associations may involve one or more amino acid residues contained in the heterologous polypeptide sequence in a fusion protein. In one example, covalent associations are between the heterologous sequence contained in a fusion protein of the invention (see, e.g., US Patent Number 5,478,925). In a specific example, the covalent associations are between the heterologous sequence contained in a Fc fusion protein of the invention (as described herein). In another specific example, covalent associations of fusion proteins of the invention are between heterologous polypeptide sequence from another protein that is capable of forming covalently associated multimers, such as for example, oseteoprotegerin (see, e.g., International Publication NO: WO 98/49305, the contents of which are herein incorporated by reference in its entirety). In another embodiment, two or more polypeptides of the invention are joined through peptide linkers. Examples include those peptide linkers described in U.S. Pat. No. 5,073,627 (hereby incorporated by reference). Proteins comprising multiple polypeptides of the invention separated by peptide linkers may be produced using conventional recombinant DNA technology.

Another method for preparing multimer polypeptides of the invention involves use of polypeptides of the invention fused to a leucine zipper or isoleucine zipper polypeptide sequence. Leucine zipper and isoleucine zipper domains are polypeptides that promote multimerization of the proteins in which they are found. Leucine zippers were originally identified in several DNA-binding proteins (Landschulz et al., Science 240:1759, (1988)), and have since been found in a variety of different proteins. Among the known leucine zippers are naturally occurring peptides and derivatives thereof that dimerize or trimerize. Examples of leucine zipper domains suitable for producing soluble multimeric proteins of the

invention are those described in PCT application WO 94/10308, hereby incorporated by reference. Recombinant fusion proteins comprising a polypeptide of the invention fused to a polypeptide sequence that dimerizes or trimerizes in solution are expressed in suitable host cells, and the resulting soluble multimeric fusion protein is recovered from the culture supernatant using techniques known in the art.

Trimeric polypeptides of the invention may offer the advantage of enhanced biological activity. Preferred leucine zipper moieties and isoleucine moieties are those that preferentially form trimers. One example is a leucine zipper derived from lung surfactant protein D (SPD), as described in Hoppe et al. (FEBS Letters 344:191, (1994)) and in U.S. patent application Ser. No. 08/446,922, hereby incorporated by reference. Other peptides derived from naturally occurring trimeric proteins may be employed in preparing trimeric polypeptides of the invention.

In another example, proteins of the invention are associated by interactions between Flag® polypeptide sequence contained in fusion proteins of the invention containing Flag® polypeptide sequence. In a further embodiment, associations proteins of the invention are associated by interactions between heterologous polypeptide sequence contained in Flag® fusion proteins of the invention and anti-Flag® antibody.

The multimers of the invention may be generated using chemical techniques known in the art. For example, polypeptides desired to be contained in the multimers of the invention may be chemically cross-linked using linker molecules and linker molecule length optimization techniques known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, multimers of the invention may be generated using techniques known in the art to form one or more inter-molecule cross-links between the cysteine residues located within the sequence of the polypeptides desired to be contained in the multimer (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Further, polypeptides of the invention may be routinely modified by the addition of cysteine or biotin to the C-terminus or N-terminus of the polypeptide and techniques known in the art may be applied to generate multimers containing one or more of these modified polypeptides (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). Additionally, techniques known in the art may be applied to generate liposomes containing the polypeptide

components desired to be contained in the multimer of the invention (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Alternatively, multimers of the invention may be generated using genetic engineering techniques known in the art. In one embodiment, polypeptides contained in multimers of the invention are produced recombinantly using fusion protein technology described herein or otherwise known in the art (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In a specific embodiment, polynucleotides coding for a homodimer of the invention are generated by ligating a polynucleotide sequence encoding a polypeptide of the invention to a sequence encoding a linker polypeptide and then further to a synthetic polynucleotide encoding the translated product of the polypeptide in the reverse orientation from the original C-terminus to the N-terminus (lacking the leader sequence) (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety). In another embodiment, recombinant techniques described herein or otherwise known in the art are applied to generate recombinant polypeptides of the invention which contain a transmembrane domain (or hydrophobic or signal peptide) and which can be incorporated by membrane reconstitution techniques into liposomes (see, e.g., US Patent Number 5,478,925, which is herein incorporated by reference in its entirety).

Antibodies

Further polypeptides of the invention relate to antibodies and T-cell antigen receptors (TCR) which immunospecifically bind a polypeptide, polypeptide fragment, or variant of SEQ ID NO:Y, and/or an epitope, of the present invention (as determined by immunoassays well known in the art for assaying specific antibody-antigen binding). Antibodies of the invention include, but are not limited to, polyclonal, monoclonal, multispecific, human, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab') fragments, fragments produced by a Fab expression library, anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id antibodies to antibodies of the invention), and epitope-binding fragments of any of the above. The term "antibody," as used herein, refers to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antigen binding site that immunospecifically binds an antigen. The immunoglobulin molecules of the invention can be of any type (e.g., IgG,

IgE, IgM, IgD, IgA and IgY), class (e.g., IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2) or subclass of immunoglobulin molecule.

Most preferably the antibodies are human antigen-binding antibody fragments of the present invention and include, but are not limited to, Fab, Fab' and F(ab')₂, Fd, single-chain Fvs (scFv), single-chain antibodies, disulfide-linked Fvs (sdFv) and fragments comprising either a VL or VH domain. Antigen-binding antibody fragments, including single-chain antibodies, may comprise the variable region(s) alone or in combination with the entirety or a portion of the following: hinge region, CH1, CH2, and CH3 domains. Also included in the invention are antigen-binding fragments also comprising any combination of variable region(s) with a hinge region, CH1, CH2, and CH3 domains. The antibodies of the invention may be from any animal origin including birds and mammals. Preferably, the antibodies are human, murine (e.g., mouse and rat), donkey, ship rabbit, goat, guinea pig, camel, horse, or chicken. As used herein, "human" antibodies include antibodies having the amino acid sequence of a human immunoglobulin and include antibodies isolated from human immunoglobulin libraries or from animals transgenic for one or more human immunoglobulin and that do not express endogenous immunoglobulins, as described infra and, for example in, U.S. Patent No. 5,939,598 by Kucherlapati et al.

The antibodies of the present invention may be monospecific, bispecific, trispecific or of greater multispecificity. Multispecific antibodies may be specific for different epitopes of a polypeptide of the present invention or may be specific for both a polypeptide of the present invention as well as for a heterologous epitope, such as a heterologous polypeptide or solid support material. See, e.g., PCT publications WO 93/17715; WO 92/08802; WO 91/00360; WO 92/05793; Tutt, et al., J. Immunol. 147:60-69 (1991); U.S. Patent Nos. 4,474,893; 4,714,681; 4,925,648; 5,573,920; 5,601,819; Kostelny et al., J. Immunol. 148:1547-1553 (1992).

Antibodies of the present invention may be described or specified in terms of the epitope(s) or portion(s) of a polypeptide of the present invention which they recognize or specifically bind. The epitope(s) or polypeptide portion(s) may be specified as described herein, e.g., by N-terminal and C-terminal positions, or by size in contiguous amino acid residues. Antibodies which specifically bind any epitope or polypeptide of the present invention may also be excluded. Therefore, the present invention includes antibodies that

specifically bind polypeptides of the present invention, and allows for the exclusion of the same.

Antibodies of the present invention may also be described or specified in terms of their cross-reactivity. Antibodies that do not bind any other analog, ortholog, or homolog of a polypeptide of the present invention are included. Antibodies that bind polypeptides with at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 65%, at least 60%, at least 55%, and at least 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In specific embodiments, antibodies of the present invention cross-react with murine, rat and/or rabbit homologs of human proteins and the corresponding epitopes thereof. Antibodies that do not bind polypeptides with less than 95%, less than 90%, less than 85%, less than 80%, less than 75%, less than 70%, less than 65%, less than 60%, less than 55%, and less than 50% identity (as calculated using methods known in the art and described herein) to a polypeptide of the present invention are also included in the present invention. In a specific embodiment, the above-described cross-reactivity is with respect to any single specific antigenic or immunogenic polypeptide, or combination(s) of 2, 3, 4, 5, or more of the specific antigenic and/or immunogenic polypeptides disclosed herein. Further included in the present invention are antibodies which bind polypeptides encoded by polynucleotides which hybridize to a polynucleotide of the present invention under stringent hybridization conditions (as described herein). Antibodies of the present invention may also be described or specified in terms of their binding affinity to a polypeptide of the invention. Preferred binding affinities include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, or 10^{-15} M.

The invention also provides antibodies that competitively inhibit binding of an antibody to an epitope of the invention as determined by any method known in the art for determining competitive binding, for example, the immunoassays described herein. In preferred embodiments, the antibody competitively inhibits binding to the epitope by at least 95%, at least 90%, at least 85 %, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50%.

Antibodies of the present invention may act as agonists or antagonists of the polypeptides of the present invention. For example, the present invention includes antibodies which disrupt the receptor/ligand interactions with the polypeptides of the invention either partially or fully. Preferably, antibodies of the present invention bind an antigenic epitope disclosed herein, or a portion thereof. The invention features both receptor-specific antibodies and ligand-specific antibodies. The invention also features receptor-specific antibodies which do not prevent ligand binding but prevent receptor activation. Receptor activation (i.e., signaling) may be determined by techniques described herein or otherwise known in the art. For example, receptor activation can be determined by detecting the phosphorylation (e.g., tyrosine or serine/threonine) of the receptor or its substrate by immunoprecipitation followed by western blot analysis (for example, as described supra). In specific embodiments, antibodies are provided that inhibit ligand activity or receptor activity by at least 95%, at least 90%, at least 85%, at least 80%, at least 75%, at least 70%, at least 60%, or at least 50% of the activity in absence of the antibody.

The invention also features receptor-specific antibodies which both prevent ligand binding and receptor activation as well as antibodies that recognize the receptor-ligand complex, and, preferably, do not specifically recognize the unbound receptor or the unbound ligand. Likewise, included in the invention are neutralizing antibodies which bind the ligand and prevent binding of the ligand to the receptor, as well as antibodies which bind the ligand, thereby preventing receptor activation, but do not prevent the ligand from binding the receptor. Further included in the invention are antibodies which activate the receptor. These antibodies may act as receptor agonists, i.e., potentiate or activate either all or a subset of the biological activities of the ligand-mediated receptor activation, for example, by inducing dimerization of the receptor. The antibodies may be specified as agonists, antagonists or inverse agonists for biological activities comprising the specific biological activities of the peptides of the invention disclosed herein. The above antibody agonists can be made using methods known in the art. See, e.g., PCT publication WO 96/40281; U.S. Patent No. 5,811,097; Deng et al., *Blood* 92(6):1981-1988 (1998); Chen et al., *Cancer Res.* 58(16):3668-3678 (1998); Harrop et al., *J. Immunol.* 161(4):1786-1794 (1998); Zhu et al., *Cancer Res.* 58(15):3209-3214 (1998); Yoon et al., *J. Immunol.* 160(7):3170-3179 (1998); Prat et al., *J. Cell. Sci.* 111(Pt2):237-247 (1998); Pitard et al., *J. Immunol. Methods* 205(2):177-190 (1997); Liautard et al., *Cytokine* 9(4):233-241 (1997); Carlson et al., *J. Biol.*

Chem. 272(17):11295-11301 (1997); Taryman et al., Neuron 14(4):755-762 (1995); Muller et al., Structure 6(9):1153-1167 (1998); Bartunek et al., Cytokine 8(1):14-20 (1996) (which are all incorporated by reference herein in their entireties).

Antibodies of the present invention may be used, for example, but not limited to, to
5 purify, detect, and target the polypeptides of the present invention, including both in vitro and
in vivo diagnostic and therapeutic methods. For example, the antibodies have use in
immunoassays for qualitatively and quantitatively measuring levels of the polypeptides of the
present invention in biological samples. See, e.g., Harlow et al., Antibodies: A Laboratory
Manual. (Cold Spring Harbor Laboratory Press, 2nd ed. 1988) (incorporated by reference
10 herein in its entirety).

As discussed in more detail below, the antibodies of the present invention may be
used either alone or in combination with other compositions. The antibodies may further be
recombinantly fused to a heterologous polypeptide at the N- or C-terminus or chemically
conjugated (including covalently and non-covalently conjugations) to polypeptides or other
15 compositions. For example, antibodies of the present invention may be recombinantly fused
or conjugated to molecules useful as labels in detection assays and effector molecules such as
heterologous polypeptides, drugs, radionuclides, or toxins. See, e.g., PCT publications WO
92/08495; WO 91/14438; WO 89/12624; U.S. Patent No. 5,314,995; and EP 396,387.

The antibodies of the invention include derivatives that are modified, i.e., by the
20 covalent attachment of any type of molecule to the antibody such that covalent attachment
does not prevent the antibody from generating an anti-idiotypic response. For example, but
not by way of limitation, the antibody derivatives include antibodies that have been modified,
e.g., by glycosylation, acetylation, pegylation, phosphorylation, amidation, derivatization by
known protecting/blocking groups, proteolytic cleavage, linkage to a cellular ligand or other
25 protein, etc. Any of numerous chemical modifications may be carried out by known
techniques, including, but not limited to specific chemical cleavage, acetylation, formylation,
metabolic synthesis of tunicamycin, etc. Additionally, the derivative may contain one or
more non-classical amino acids.

The antibodies of the present invention may be generated by any suitable method
30 known in the art. Polyclonal antibodies to an antigen-of-interest can be produced by various
procedures well known in the art. For example, a polypeptide of the invention can be
administered to various host animals including, but not limited to, rabbits, mice, rats, etc. to

induce the production of sera containing polyclonal antibodies specific for the antigen. Various adjuvants may be used to increase the immunological response, depending on the host species, and include but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants are also well known in the art.

Monoclonal antibodies can be prepared using a wide variety of techniques known in the art including the use of hybridoma, recombinant, and phage display technologies, or a combination thereof. For example, monoclonal antibodies can be produced using hybridoma techniques including those known in the art and taught, for example, in Harlow et al., *Antibodies: A Laboratory Manual*, (Cold Spring Harbor Laboratory Press, 2nd ed. 1988); Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) (said references incorporated by reference in their entireties). The term "monoclonal antibody" as used herein is not limited to antibodies produced through hybridoma technology. The term "monoclonal antibody" refers to an antibody that is derived from a single clone, including any eukaryotic, prokaryotic, or phage clone, and not the method by which it is produced.

Methods for producing and screening for specific antibodies using hybridoma technology are routine and well known in the art and are discussed in detail in the Examples. In a non-limiting example, mice can be immunized with a polypeptide of the invention or a cell expressing such peptide. Once an immune response is detected, e.g., antibodies specific for the antigen are detected in the mouse serum, the mouse spleen is harvested and splenocytes isolated. The splenocytes are then fused by well known techniques to any suitable myeloma cells, for example cells from cell line SP20 available from the ATCC. Hybridomas are selected and cloned by limited dilution. The hybridoma clones are then assayed by methods known in the art for cells that secrete antibodies capable of binding a polypeptide of the invention. Ascites fluid, which generally contains high levels of antibodies, can be generated by immunizing mice with positive hybridoma clones.

Accordingly, the present invention provides methods of generating monoclonal antibodies as well as antibodies produced by the method comprising culturing a hybridoma cell secreting an antibody of the invention wherein, preferably, the hybridoma is generated by

fusing splenocytes isolated from a mouse immunized with an antigen of the invention with myeloma cells and then screening the hybridomas resulting from the fusion for hybridoma clones that secrete an antibody able to bind a polypeptide of the invention.

Antibody fragments which recognize specific epitopes may be generated by known techniques. For example, Fab and F(ab')₂ fragments of the invention may be produced by proteolytic cleavage of immunoglobulin molecules, using enzymes such as papain (to produce Fab fragments) or pepsin (to produce F(ab')₂ fragments). F(ab')₂ fragments contain the variable region, the light chain constant region and the CH1 domain of the heavy chain.

For example, the antibodies of the present invention can also be generated using various phage display methods known in the art. In phage display methods, functional antibody domains are displayed on the surface of phage particles which carry the polynucleotide sequences encoding them. In a particular embodiment, such phage can be utilized to display antigen binding domains expressed from a repertoire or combinatorial antibody library (e.g., human or murine). Phage expressing an antigen binding domain that binds the antigen of interest can be selected or identified with antigen, e.g., using labeled antigen or antigen bound or captured to a solid surface or bead. Phage used in these methods are typically filamentous phage including fd and M13 binding domains expressed from phage with Fab, Fv or disulfide stabilized Fv antibody domains recombinantly fused to either the phage gene III or gene VIII protein. Examples of phage display methods that can be used to make the antibodies of the present invention include those disclosed in Brinkman et al., J. Immunol. Methods 182:41-50 (1995); Ames et al., J. Immunol. Methods 184:177-186 (1995); Kettleborough et al., Eur. J. Immunol. 24:952-958 (1994); Persic et al., Gene 187 9-18 (1997); Burton et al., Advances in Immunology 57:191-280 (1994); PCT application No. PCT/GB91/01134; PCT publications WO 90/02809; WO 91/10737; WO 92/01047; WO 92/18619; WO 93/11236; WO 95/15982; WO 95/20401; and U.S. Patent Nos. 5,698,426; 5,223,409; 5,403,484; 5,580,717; 5,427,908; 5,750,753; 5,821,047; 5,571,698; 5,427,908; 5,516,637; 5,780,225; 5,658,727; 5,733,743 and 5,969,108; each of which is incorporated herein by reference in its entirety.

As described in the above references, after phage selection, the antibody coding regions from the phage can be isolated and used to generate whole antibodies, including human antibodies, or any other desired antigen binding fragment, and expressed in any desired host, including mammalian cells, insect cells, plant cells, yeast, and bacteria, e.g., as

described in detail below. For example, techniques to recombinantly produce Fab, Fab' and F(ab')₂ fragments can also be employed using methods known in the art such as those disclosed in PCT publication WO 92/22324; Mullinax et al., *BioTechniques* 12(6):864-869 (1992); and Sawai et al., *AJRI* 34:26-34 (1995); and Better et al., *Science* 240:1041-1043 (1988) (said references incorporated by reference in their entireties).

Examples of techniques which can be used to produce single-chain Fvs and antibodies include those described in U.S. Patents 4,946,778 and 5,258,498; Huston et al., *Methods in Enzymology* 203:46-88 (1991); Shu et al., *PNAS* 90:7995-7999 (1993); and Skerra et al., *Science* 240:1038-1040 (1988). For some uses, including in vivo use of antibodies in humans and in vitro detection assays, it may be preferable to use chimeric, humanized, or human antibodies. A chimeric antibody is a molecule in which different portions of the antibody are derived from different animal species, such as antibodies having a variable region derived from a murine monoclonal antibody and a human immunoglobulin constant region. Methods for producing chimeric antibodies are known in the art. See e.g., Morrison, *Science* 229:1202 (1985); Oi et al., *BioTechniques* 4:214 (1986); Gillies et al., (1989) *J. Immunol. Methods* 125:191-202; U.S. Patent Nos. 5,807,715; 4,816,567; and 4,816,397, which are incorporated herein by reference in their entirety. Humanized antibodies are antibody molecules from non-human species antibody that binds the desired antigen having one or more complementarity determining regions (CDRs) from the non-human species and a framework regions from a human immunoglobulin molecule. Often, framework residues in the human framework regions will be substituted with the corresponding residue from the CDR donor antibody to alter, preferably improve, antigen binding. These framework substitutions are identified by methods well known in the art, e.g., by modeling of the interactions of the CDR and framework residues to identify framework residues important for antigen binding and sequence comparison to identify unusual framework residues at particular positions. (See, e.g., Queen et al., U.S. Patent No. 5,585,089; Riechmann et al., *Nature* 332:323 (1988), which are incorporated herein by reference in their entireties.) Antibodies can be humanized using a variety of techniques known in the art including, for example, CDR-grafting (EP 239,400; PCT publication WO 91/09967; U.S. Patent Nos. 5,225,539; 5,530,101; and 5,585,089), veneering or resurfacing (EP 592,106; EP 519,596; Padlan, *Molecular Immunology* 28(4/5):489-498 (1991); Studnicka et al., *Protein*

Engineering 7(6):805-814 (1994); Roguska. et al., PNAS 91:969-973 (1994)), and chain shuffling (U.S. Patent No. 5,565,332).

5 Completely human antibodies are particularly desirable for therapeutic treatment of human patients. Human antibodies can be made by a variety of methods known in the art including phage display methods described above using antibody libraries derived from human immunoglobulin sequences. See also, U.S. Patent Nos. 4,444,887 and 4,716,111; and PCT publications WO 98/46645, WO 98/50433, WO 98/24893, WO 98/16654, WO 96/34096, WO 96/33735, and WO 91/10741; each of which is incorporated herein by reference in its entirety.

10 Human antibodies can also be produced using transgenic mice which are incapable of expressing functional endogenous immunoglobulins, but which can express human immunoglobulin genes. For example, the human heavy and light chain immunoglobulin gene complexes may be introduced randomly or by homologous recombination into mouse embryonic stem cells. Alternatively, the human variable region, constant region, and
15 diversity region may be introduced into mouse embryonic stem cells in addition to the human heavy and light chain genes. The mouse heavy and light chain immunoglobulin genes may be rendered non-functional separately or simultaneously with the introduction of human immunoglobulin loci by homologous recombination. In particular, homozygous deletion of the JH region prevents endogenous antibody production. The modified embryonic stem cells
20 are expanded and microinjected into blastocysts to produce chimeric mice. The chimeric mice are then bred to produce homozygous offspring which express human antibodies. The transgenic mice are immunized in the normal fashion with a selected antigen, e.g., all or a portion of a polypeptide of the invention. Monoclonal antibodies directed against the antigen can be obtained from the immunized, transgenic mice using conventional hybridoma
25 technology. The human immunoglobulin transgenes harbored by the transgenic mice rearrange during B cell differentiation, and subsequently undergo class switching and somatic mutation. Thus, using such a technique, it is possible to produce therapeutically useful IgG, IgA, IgM and IgE antibodies. For an overview of this technology for producing human antibodies, see Lonberg and Huszar, Int. Rev. Immunol. 13:65-93 (1995). For a
30 detailed discussion of this technology for producing human antibodies and human monoclonal antibodies and protocols for producing such antibodies, see, e.g., PCT publications WO 98/24893; WO 92/01047; WO 96/34096; WO 96/33735; European Patent

No. 0 598 877; U.S. Patent Nos. 5,413,923; 5,625,126; 5,633,425; 5,569,825; 5,661,016; 5,545,806; 5,814,318; 5,885,793; 5,916,771; and 5,939,598, which are incorporated by reference herein in their entirety. In addition, companies such as Abgenix, Inc. (Freemont, CA) and Genpharm (San Jose, CA) can be engaged to provide human antibodies directed
5 against a selected antigen using technology similar to that described above.

Completely human antibodies which recognize a selected epitope can be generated using a technique referred to as "guided selection." In this approach a selected non-human monoclonal antibody, e.g., a mouse antibody, is used to guide the selection of a completely human antibody recognizing the same epitope. (Jespers et al., Bio/technology 12:899-903
10 (1988)).

Further, antibodies to the polypeptides of the invention can, in turn, be utilized to generate anti-idiotypic antibodies that "mimic" polypeptides of the invention using techniques well known to those skilled in the art. (See, e.g., Greenspan & Bona, FASEB J. 7(5):437-444; (1989) and Nissinoff, J. Immunol. 147(8):2429-2438 (1991)). For example, antibodies
15 which bind to and competitively inhibit polypeptide multimerization and/or binding of a polypeptide of the invention to a ligand can be used to generate anti-idiotypes that "mimic" the polypeptide multimerization and/or binding domain and, as a consequence, bind to and neutralize polypeptide and/or its ligand. Such neutralizing anti-idiotypes or Fab fragments of such anti-idiotypes can be used in therapeutic regimens to neutralize polypeptide ligand. For
20 example, such anti-idiotypic antibodies can be used to bind a polypeptide of the invention and/or to bind its ligands/receptors, and thereby block its biological activity.

Polynucleotides Encoding Antibodies

The invention further provides polynucleotides comprising a nucleotide sequence
25 encoding an antibody of the invention and fragments thereof. The invention also encompasses polynucleotides that hybridize under stringent or alternatively, under lower stringency hybridization conditions, e.g., as defined supra, to polynucleotides that encode an antibody, preferably, that specifically binds to a polypeptide of the invention, preferably, an antibody that binds to a polypeptide having the amino acid sequence of SEQ ID NO:Y.

30 The polynucleotides may be obtained, and the nucleotide sequence of the polynucleotides determined, by any method known in the art. For example, if the nucleotide sequence of the antibody is known, a polynucleotide encoding the antibody may be

assembled from chemically synthesized oligonucleotides (e.g., as described in Kutmeier et al., *BioTechniques* 17:242 (1994)), which, briefly, involves the synthesis of overlapping oligonucleotides containing portions of the sequence encoding the antibody, annealing and ligating of those oligonucleotides, and then amplification of the ligated oligonucleotides by PCR.

Alternatively, a polynucleotide encoding an antibody may be generated from nucleic acid from a suitable source. If a clone containing a nucleic acid encoding a particular antibody is not available, but the sequence of the antibody molecule is known, a nucleic acid encoding the immunoglobulin may be chemically synthesized or obtained from a suitable source (e.g., an antibody cDNA library, or a cDNA library generated from, or nucleic acid, preferably poly A+ RNA, isolated from, any tissue or cells expressing the antibody, such as hybridoma cells selected to express an antibody of the invention) by PCR amplification using synthetic primers hybridizable to the 3' and 5' ends of the sequence or by cloning using an oligonucleotide probe specific for the particular gene sequence to identify, e.g., a cDNA clone from a cDNA library that encodes the antibody. Amplified nucleic acids generated by PCR may then be cloned into replicable cloning vectors using any method well known in the art.

Once the nucleotide sequence and corresponding amino acid sequence of the antibody is determined, the nucleotide sequence of the antibody may be manipulated using methods well known in the art for the manipulation of nucleotide sequences, e.g., recombinant DNA techniques, site directed mutagenesis, PCR, etc. (see, for example, the techniques described in Sambrook et al., 1990, *Molecular Cloning, A Laboratory Manual*, 2d Ed., Cold Spring Harbor Laboratory, Cold Spring Harbor, NY and Ausubel et al., eds., 1998, *Current Protocols in Molecular Biology*, John Wiley & Sons, NY, which are both incorporated by reference herein in their entireties), to generate antibodies having a different amino acid sequence, for example to create amino acid substitutions, deletions, and/or insertions.

In a specific embodiment, the amino acid sequence of the heavy and/or light chain variable domains may be inspected to identify the sequences of the complementarity determining regions (CDRs) by methods that are well known in the art. e.g., by comparison to known amino acid sequences of other heavy and light chain variable regions to determine the regions of sequence hypervariability. Using routine recombinant DNA techniques, one or more of the CDRs may be inserted within framework regions, e.g., into human framework

regions to humanize a non-human antibody, as described supra. The framework regions may be naturally occurring or consensus framework regions, and preferably human framework regions (see, e.g., Chothia et al., J. Mol. Biol. 278: 457-479 (1998) for a listing of human framework regions). Preferably, the polynucleotide generated by the combination of the framework regions and CDRs encodes an antibody that specifically binds a polypeptide of the invention. Preferably, as discussed supra, one or more amino acid substitutions may be made within the framework regions, and, preferably, the amino acid substitutions improve binding of the antibody to its antigen. Additionally, such methods may be used to make amino acid substitutions or deletions of one or more variable region cysteine residues participating in an intrachain disulfide bond to generate antibody molecules lacking one or more intrachain disulfide bonds. Other alterations to the polynucleotide are encompassed by the present invention and within the skill of the art.

In addition, techniques developed for the production of "chimeric antibodies" (Morrison et al., Proc. Natl. Acad. Sci. 81:851-855 (1984); Neuberger et al., Nature 312:604-608 (1984); Takeda et al., Nature 314:452-454 (1985)) by splicing genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity can be used. As described supra, a chimeric antibody is a molecule in which different portions are derived from different animal species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region, e.g., humanized antibodies.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patent No. 4,946,778; Bird, Science 242:423-42 (1988); Huston et al., Proc. Natl. Acad. Sci. USA 85:5879-5883 (1988); and Ward et al., Nature 334:544-54 (1989)) can be adapted to produce single chain antibodies. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single chain polypeptide. Techniques for the assembly of functional Fv fragments in E. coli may also be used (Skerra et al., Science 242:1038-1041 (1988)).

Methods of Producing Antibodies

The antibodies of the invention can be produced by any method known in the art for the synthesis of antibodies, in particular, by chemical synthesis or preferably, by recombinant expression techniques.

Recombinant expression of an antibody of the invention, or fragment, derivative or analog thereof, (e.g., a heavy or light chain of an antibody of the invention or a single chain antibody of the invention), requires construction of an expression vector containing a polynucleotide that encodes the antibody. Once a polynucleotide encoding an antibody molecule or a heavy or light chain of an antibody, or portion thereof (preferably containing the heavy or light chain variable domain), of the invention has been obtained, the vector for the production of the antibody molecule may be produced by recombinant DNA technology using techniques well known in the art. Thus, methods for preparing a protein by expressing a polynucleotide containing an antibody encoding nucleotide sequence are described herein.

Methods which are well known to those skilled in the art can be used to construct expression vectors containing antibody coding sequences and appropriate transcriptional and translational control signals. These methods include, for example, in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. The invention, thus, provides replicable vectors comprising a nucleotide sequence encoding an antibody molecule of the invention, or a heavy or light chain thereof, or a heavy or light chain variable domain, operably linked to a promoter. Such vectors may include the nucleotide sequence encoding the constant region of the antibody molecule (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S. Patent No. 5,122,464) and the variable domain of the antibody may be cloned into such a vector for expression of the entire heavy or light chain.

The expression vector is transferred to a host cell by conventional techniques and the transfected cells are then cultured by conventional techniques to produce an antibody of the invention. Thus, the invention includes host cells containing a polynucleotide encoding an antibody of the invention, or a heavy or light chain thereof, or a single chain antibody of the invention, operably linked to a heterologous promoter. In preferred embodiments for the expression of double-chained antibodies, vectors encoding both the heavy and light chains may be co-expressed in the host cell for expression of the entire immunoglobulin molecule, as detailed below.

A variety of host-expression vector systems may be utilized to express the antibody molecules of the invention. Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express an antibody molecule of the invention in situ. These include but are not

limited to microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing antibody coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transformed with recombinant yeast expression vectors containing antibody coding sequences; insect cell
5 systems infected with recombinant virus expression vectors (e.g., baculovirus) containing antibody coding sequences; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing antibody coding sequences; or mammalian cell systems (e.g., COS, CHO, BHK, 293, 3T3 cells) harboring
10 recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Preferably, bacterial cells such as *Escherichia coli*, and more preferably, eukaryotic cells, especially for the expression of whole recombinant antibody molecule, are used for the expression of a recombinant antibody
15 molecule. For example, mammalian cells such as Chinese hamster ovary cells (CHO), in conjunction with a vector such as the major intermediate early gene promoter element from human cytomegalovirus is an effective expression system for antibodies (Foecking et al., *Gene* 45:101 (1986); Cockett et al., *Bio/Technology* 8:2 (1990)).

In bacterial systems, a number of expression vectors may be advantageously selected
20 depending upon the use intended for the antibody molecule being expressed. For example, when a large quantity of such a protein is to be produced, for the generation of pharmaceutical compositions of an antibody molecule, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Such vectors include, but are not limited, to the *E. coli* expression vector pUR278 (Ruther et al., *EMBO J.*
25 2:1791 (1983)), in which the antibody coding sequence may be ligated individually into the vector in frame with the lac Z coding region so that a fusion protein is produced; pIN vectors (Inouye & Inouye, *Nucleic Acids Res.* 13:3101-3109 (1985); Van Heeke & Schuster, *J. Biol. Chem.* 24:5503-5509 (1989)); and the like. pGEX vectors may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general,
30 such fusion proteins are soluble and can easily be purified from lysed cells by adsorption and binding to matrix glutathione-agarose beads followed by elution in the presence of free

glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an insect system, Autographa californica nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells.

- 5 The antibody coding sequence may be cloned individually into non-essential regions (for example the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter).

- In mammalian host cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, the antibody coding sequence
10 of interest may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the antibody molecule in infected hosts. (e.g., see Logan & Shenk,
15 Proc. Natl. Acad. Sci. USA 81:355-359 (1984)). Specific initiation signals may also be required for efficient translation of inserted antibody coding sequences. These signals include the ATG initiation codon and adjacent sequences. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can
20 be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, transcription terminators, etc. (see Bittner et al., Methods in Enzymol. 153:51-544 (1987)).

- In addition, a host cell strain may be chosen which modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion
25 desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the protein. Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end,
30 eukaryotic host cells which possess the cellular machinery for proper processing of the primary transcript, glycosylation, and phosphorylation of the gene product may be used. Such mammalian host cells include but are not limited to CHO, VERY, BHK, Hela, COS,

MDCK, 293, 3T3, WI38, and in particular, breast cancer cell lines such as, for example, BT483, Hs578T, HTB2, BT20 and T47D, and normal mammary gland cell line such as, for example, CRL7030 and Hs578Bst.

For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines which stably express the antibody molecule may be engineered. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of the foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a selective media. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. This method may advantageously be used to engineer cell lines which express the antibody molecule. Such engineered cell lines may be particularly useful in screening and evaluation of compounds that interact directly or indirectly with the antibody molecule.

A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler et al., Cell 11:223 (1977)), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, Proc. Natl. Acad. Sci. USA 48:202 (1992)), and adenine phosphoribosyltransferase (Lowy et al., Cell 22:817 (1980)) genes can be employed in tk-, hgp^rt- or ap^rt- cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for the following genes: dhfr, which confers resistance to methotrexate (Wigler et al., Natl. Acad. Sci. USA 77:357 (1980); O'Hare et al., Proc. Natl. Acad. Sci. USA 78:1527 (1981)); gpt, which confers resistance to mycophenolic acid (Mulligan & Berg, Proc. Natl. Acad. Sci. USA 78:2072 (1981)); neo, which confers resistance to the aminoglycoside G-418 Clinical Pharmacy 12:488-505; Wu and Wu, Biotherapy 3:87-95 (1991); Tolstoshev, Ann. Rev. Pharmacol. Toxicol. 32:573-596 (1993); Mulligan, Science 260:926-932 (1993); and Morgan and Anderson, Ann. Rev. Biochem. 62:191-217 (1993); May, 1993, TIB TECH 11(5):155-215; and hyg^r, which confers resistance to hygromycin (Santerre et al., Gene 30:147 (1984)). Methods commonly known in the art of recombinant DNA technology may be routinely applied to select the desired recombinant clone, and such methods are described, for example, in Ausubel et al. (eds.),

Current Protocols in Molecular Biology, John Wiley & Sons, NY (1993); Kriegler, Gene Transfer and Expression, A Laboratory Manual, Stockton Press, NY (1990); and in Chapters 12 and 13, Dracopoli et al. (eds), Current Protocols in Human Genetics, John Wiley & Sons, NY (1994); Colberre-Garapin et al., J. Mol. Biol. 150:1 (1981), which are incorporated by
5 reference herein in their entirety.

The expression levels of an antibody molecule can be increased by vector amplification (for a review, see Bebbington and Hentschel, The use of vectors based on gene amplification for the expression of cloned genes in mammalian cells in DNA cloning, Vol.3. (Academic Press, New York, 1987)). When a marker in the vector system expressing
10 antibody is amplifiable, increase in the level of inhibitor present in culture of host cell will increase the number of copies of the marker gene. Since the amplified region is associated with the antibody gene, production of the antibody will also increase (Crouse et al., Mol. Cell. Biol. 3:257 (1983)).

The host cell may be co-transfected with two expression vectors of the invention, the
15 first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide. The two vectors may contain identical selectable markers which enable equal expression of heavy and light chain polypeptides. Alternatively, a single vector may be used which encodes, and is capable of expressing, both heavy and light chain polypeptides. In such situations, the light chain should be placed before the heavy chain to
20 avoid an excess of toxic free heavy chain (Proudfoot, Nature 322:52 (1986); Kohler, Proc. Natl. Acad. Sci. USA 77:2197 (1980)). The coding sequences for the heavy and light chains may comprise cDNA or genomic DNA.

Once an antibody molecule of the invention has been produced by an animal, chemically synthesized, or recombinantly expressed, it may be purified by any method
25 known in the art for purification of an immunoglobulin molecule, for example, by chromatography (e.g., ion exchange, affinity, particularly by affinity for the specific antigen after Protein A, and sizing column chromatography), centrifugation, differential solubility, or by any other standard technique for the purification of proteins. In addition, the antibodies of the present invention or fragments thereof can be fused to heterologous polypeptide
30 sequences described herein or otherwise known in the art. to facilitate purification.

The present invention encompasses antibodies recombinantly fused or chemically conjugated (including both covalently and non-covalently conjugations) to a polypeptide (or

portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention to generate fusion proteins. The fusion does not necessarily need to be direct, but may occur through linker sequences. The antibodies may be specific for antigens other than polypeptides (or portion thereof, preferably at least 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 amino acids of the polypeptide) of the present invention. For example, antibodies may be used to target the polypeptides of the present invention to particular cell types, either in vitro or in vivo, by fusing or conjugating the polypeptides of the present invention to antibodies specific for particular cell surface receptors. Antibodies fused or conjugated to the polypeptides of the present invention may also be used in in vitro immunoassays and purification methods using methods known in the art. See e.g., Harbor et al., supra, and PCT publication WO 93/21232; EP 439,095; Naramura et al., Immunol. Lett. 39:91-99 (1994); U.S. Patent 5,474,981; Gillies et al., PNAS 89:1428-1432 (1992); Fell et al., J. Immunol. 146:2446-2452(1991), which are incorporated by reference in their entireties.

The present invention further includes compositions comprising the polypeptides of the present invention fused or conjugated to antibody domains other than the variable regions. For example, the polypeptides of the present invention may be fused or conjugated to an antibody Fc region, or portion thereof. The antibody portion fused to a polypeptide of the present invention may comprise the constant region, hinge region, CH1 domain, CH2 domain, and CH3 domain or any combination of whole domains or portions thereof. The polypeptides may also be fused or conjugated to the above antibody portions to form multimers. For example, Fc portions fused to the polypeptides of the present invention can form dimers through disulfide bonding between the Fc portions. Higher multimeric forms can be made by fusing the polypeptides to portions of IgA and IgM. Methods for fusing or conjugating the polypeptides of the present invention to antibody portions are known in the art. See, e.g., U.S. Patent Nos. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851; 5,112,946; EP 307,434; EP 367,166; PCT publications WO 96/04388; WO 91/06570; Ashkenazi et al., Proc. Natl. Acad. Sci. USA 88:10535-10539 (1991); Zheng et al., J. Immunol. 154:5590-5600 (1995); and Vil et al., Proc. Natl. Acad. Sci. USA 89:11337-11341(1992) (said references incorporated by reference in their entireties).

As discussed, supra, the polypeptides corresponding to a polypeptide, polypeptide fragment, or a variant of SEQ ID NO:Y may be fused or conjugated to the above antibody portions to increase the in vivo half life of the polypeptides or for use in immunoassays using

methods known in the art. Further, the polypeptides corresponding to SEQ ID NO:Y may be fused or conjugated to the above antibody portions to facilitate purification. One reported example describes chimeric proteins consisting of the first two domains of the human CD4-polypeptide and various domains of the constant regions of the heavy or light chains of mammalian immunoglobulins. (EP 394,827; Traunecker et al., *Nature* 331:84-86 (1988)). The polypeptides of the present invention fused or conjugated to an antibody having disulfide-linked dimeric structures (due to the IgG) may also be more efficient in binding and neutralizing other molecules, than the monomeric secreted protein or protein fragment alone. (Fountoulakis et al., *J. Biochem.* 270:3958-3964 (1995)). In many cases, the Fc part in a fusion protein is beneficial in therapy and diagnosis, and thus can result in, for example, improved pharmacokinetic properties. (EP A 232,262). Alternatively, deleting the Fc part after the fusion protein has been expressed, detected, and purified, would be desired. For example, the Fc portion may hinder therapy and diagnosis if the fusion protein is used as an antigen for immunizations. In drug discovery, for example, human proteins, such as hIL-5, have been fused with Fc portions for the purpose of high-throughput screening assays to identify antagonists of hIL-5. (See, Bennett et al., *J. Molecular Recognition* 8:52-58 (1995); Johanson et al., *J. Biol. Chem.* 270:9459-9471 (1995)).

Moreover, the antibodies or fragments thereof of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. In preferred embodiments, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz et al., *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., *Cell* 37:767 (1984)) and the "flag" tag.

The present invention further encompasses antibodies or fragments thereof conjugated to a diagnostic or therapeutic agent. The antibodies can be used diagnostically to, for example, monitor the development or progression of a tumor as part of a clinical testing procedure to, e.g., determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent

materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the antibody (or fragment thereof) or indirectly, through an intermediate (such as, for example, a linker known in the art) using techniques known in the art. See, for example, U.S. Patent No. 4,741,900 for metal ions which can be conjugated to antibodies for use as diagnostics according to the present invention. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin; and examples of suitable radioactive material include ¹²⁵I, ¹³¹I, ¹¹¹In or ⁹⁹Tc.

Further, an antibody or fragment thereof may be conjugated to a therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytotoxic agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ²¹³Bi. A cytotoxin or cytotoxic agent includes any agent that is detrimental to cells. Examples include paclitaxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine).

The conjugates of the invention can be used for modifying a given biological response, the therapeutic agent or drug moiety is not to be construed as limited to classical

chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, pseudomonas exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, 5 tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (See, International Publication No. WO 97/33899), AIM II (See, International Publication No. WO 97/34911), Fas Ligand (Takahashi *et al.*, *Int. Immunol.*, 6:1567-1574 (1994)), VEGF (See, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, biological response modifiers such as, for example, 10 lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other growth factors.

Antibodies may also be attached to solid supports, which are particularly useful for immunoassays or purification of the target antigen. Such solid supports include, but are not 15 limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Amon *et al.*, "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld *et al.* (eds.), pp. 243-56 20 (Alan R. Liss, Inc. 1985); Hellstrom *et al.*, "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson *et al.* (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera *et al.* (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of 25 Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin *et al.* (eds.), pp. 303-16 (Academic Press 1985), and Thorpe *et al.*, "The Preparation And Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.* 62:119-58 (1982).

Alternatively, an antibody can be conjugated to a second antibody to form an antibody 30 heteroconjugate as described by Segal in U.S. Patent No. 4,676,980, which is incorporated herein by reference in its entirety.

An antibody, with or without a therapeutic moiety conjugated to it, administered alone or in combination with cytotoxic factor(s) and/or cytokine(s) can be used as a therapeutic.

Immunophenotyping

5 The antibodies of the invention may be utilized for immunophenotyping of cell lines and biological samples. The translation product of the gene of the present invention may be useful as a cell specific marker, or more specifically as a cellular marker that is differentially expressed at various stages of differentiation and/or maturation of particular cell types. Monoclonal antibodies directed against a specific epitope, or combination of epitopes, will
10 allow for the screening of cellular populations expressing the marker. Various techniques can be utilized using monoclonal antibodies to screen for cellular populations expressing the marker(s), and include magnetic separation using antibody-coated magnetic beads, "panning" with antibody attached to a solid matrix (i.e., plate), and flow cytometry (See, e.g., U.S. Patent 5,985,660; and Morrison *et al.*, *Cell*, 96:737-49 (1999)).

15 These techniques allow for the screening of particular populations of cells, such as might be found with hematological malignancies (i.e. minimal residual disease (MRD) in acute leukemic patients) and "non-self" cells in transplantations to prevent Graft-versus-Host Disease (GVHD). Alternatively, these techniques allow for the screening of hematopoietic stem and progenitor cells capable of undergoing proliferation and/or differentiation, as might
20 be found in human umbilical cord blood.

Assays For Antibody Binding

 The antibodies of the invention may be assayed for immunospecific binding by any method known in the art. The immunoassays which can be used include but are not limited
25 to competitive and non-competitive assay systems using techniques such as western blots, radioimmunoassays, ELISA (enzyme linked immunosorbent assay), "sandwich" immunoassays, immunoprecipitation assays, precipitin reactions, gel diffusion precipitin reactions, immunodiffusion assays, agglutination assays, complement-fixation assays, immunoradiometric assays, fluorescent immunoassays, protein A immunoassays, to name
30 but a few. Such assays are routine and well known in the art (see, e.g., Ausubel et al, eds. 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York,

which is incorporated by reference herein in its entirety). Exemplary immunoassays are described briefly below (but are not intended by way of limitation).

Immunoprecipitation protocols generally comprise lysing a population of cells in a lysis buffer such as RIPA buffer (1% NP-40 or Triton X-100, 1% sodium deoxycholate, 0.1% SDS, 0.15 M NaCl, 0.01 M sodium phosphate at pH 7.2, 1% Trasyllol) supplemented with protein phosphatase and/or protease inhibitors (e.g., EDTA, PMSF, aprotinin, sodium vanadate), adding the antibody of interest to the cell lysate, incubating for a period of time (e.g., 1-4 hours) at 4° C, adding protein A and/or protein G sepharose beads to the cell lysate, incubating for about an hour or more at 4° C, washing the beads in lysis buffer and resuspending the beads in SDS/sample buffer. The ability of the antibody of interest to immunoprecipitate a particular antigen can be assessed by, e.g., western blot analysis. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the binding of the antibody to an antigen and decrease the background (e.g., pre-clearing the cell lysate with sepharose beads). For further discussion regarding immunoprecipitation protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.16.1.

Western blot analysis generally comprises preparing protein samples, electrophoresis of the protein samples in a polyacrylamide gel (e.g., 8%-20% SDS-PAGE depending on the molecular weight of the antigen), transferring the protein sample from the polyacrylamide gel to a membrane such as nitrocellulose, PVDF or nylon, blocking the membrane in blocking solution (e.g., PBS with 3% BSA or non-fat milk), washing the membrane in washing buffer (e.g., PBS-Tween 20), blocking the membrane with primary antibody (the antibody of interest) diluted in blocking buffer, washing the membrane in washing buffer, blocking the membrane with a secondary antibody (which recognizes the primary antibody, e.g., an anti-human antibody) conjugated to an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) or radioactive molecule (e.g., ³²P or ¹²⁵I) diluted in blocking buffer, washing the membrane in wash buffer, and detecting the presence of the antigen. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected and to reduce the background noise. For further discussion regarding western blot protocols see, e.g., Ausubel et al, eds, 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 10.8.1.

ELISAs comprise preparing antigen, coating the well of a 96 well microtiter plate with the antigen, adding the antibody of interest conjugated to a detectable compound such as an enzymatic substrate (e.g., horseradish peroxidase or alkaline phosphatase) to the well and incubating for a period of time, and detecting the presence of the antigen. In ELISAs the antibody of interest does not have to be conjugated to a detectable compound: instead, a second antibody (which recognizes the antibody of interest) conjugated to a detectable compound may be added to the well. Further, instead of coating the well with the antigen, the antibody may be coated to the well. In this case, a second antibody conjugated to a detectable compound may be added following the addition of the antigen of interest to the coated well. One of skill in the art would be knowledgeable as to the parameters that can be modified to increase the signal detected as well as other variations of ELISAs known in the art. For further discussion regarding ELISAs see, e.g., Ausubel et al, eds. 1994, Current Protocols in Molecular Biology, Vol. 1, John Wiley & Sons, Inc., New York at 11.2.1.

The binding affinity of an antibody to an antigen and the off-rate of an antibody-antigen interaction can be determined by competitive binding assays. One example of a competitive binding assay is a radioimmunoassay comprising the incubation of labeled antigen (e.g., ^3H or ^{125}I) with the antibody of interest in the presence of increasing amounts of unlabeled antigen, and the detection of the antibody bound to the labeled antigen. The affinity of the antibody of interest for a particular antigen and the binding off-rates can be determined from the data by scatchard plot analysis. Competition with a second antibody can also be determined using radioimmunoassays. In this case, the antigen is incubated with antibody of interest conjugated to a labeled compound (e.g., ^3H or ^{125}I) in the presence of increasing amounts of an unlabeled second antibody.

25 *Therapeutic Uses*

The present invention is further directed to antibody-based therapies which involve administering antibodies of the invention to an animal, preferably a mammal, and most preferably a human, patient for treating one or more of the disclosed diseases, disorders, or conditions. Therapeutic compounds of the invention include, but are not limited to, antibodies of the invention (including fragments, analogs and derivatives thereof as described herein) and nucleic acids encoding antibodies of the invention (including fragments, analogs and derivatives thereof and anti-idiotypic antibodies as described herein). The antibodies of

the invention can be used to treat, inhibit or prevent diseases, disorders or conditions associated with aberrant expression and/or activity of a polypeptide of the invention, including, but not limited to, any one or more of the diseases, disorders, or conditions described herein. The treatment and/or prevention of diseases, disorders, or conditions associated with aberrant expression and/or activity of a polypeptide of the invention includes, but is not limited to, alleviating symptoms associated with those diseases, disorders or conditions. Antibodies of the invention may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth factors (such as, e.g., IL-2, IL-3 and IL-7), for example, which serve to increase the number or activity of effector cells which interact with the antibodies.

The antibodies of the invention may be administered alone or in combination with other types of treatments (e.g., radiation therapy, chemotherapy, hormonal therapy, immunotherapy and anti-tumor agents). Generally, administration of products of a species origin or species reactivity (in the case of antibodies) that is the same species as that of the patient is preferred. Thus, in a preferred embodiment, human antibodies, fragments derivatives, analogs, or nucleic acids, are administered to a human patient for therapy or prophylaxis.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention, fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides of the invention, including fragments thereof. Preferred binding affinities

include those with a dissociation constant or K_d less than 5×10^{-2} M, 10^{-2} M, 5×10^{-3} M, 10^{-3} M, 5×10^{-4} M, 10^{-4} M, 5×10^{-5} M, 10^{-5} M, 5×10^{-6} M, 10^{-6} M, 5×10^{-7} M, 10^{-7} M, 5×10^{-8} M, 10^{-8} M, 5×10^{-9} M, 10^{-9} M, 5×10^{-10} M, 10^{-10} M, 5×10^{-11} M, 10^{-11} M, 5×10^{-12} M, 10^{-12} M, 5×10^{-13} M, 10^{-13} M, 5×10^{-14} M, 10^{-14} M, 5×10^{-15} M, and 10^{-15} M.

5

Gene Therapy

In a specific embodiment, nucleic acids comprising sequences encoding antibodies or functional derivatives thereof, are administered to treat, inhibit or prevent a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention, by way of gene therapy. Gene therapy refers to therapy performed by the administration to a subject of an expressed or expressible nucleic acid. In this embodiment of the invention, the nucleic acids produce their encoded protein that mediates a therapeutic effect.

Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

For general reviews of the methods of gene therapy, see Goldspiel et al., *Clinical Pharmacy* 12:488-505 (1993); Wu and Wu, *Biotherapy* 3:87-95 (1991); Tolstoshev, *Ann. Rev. Pharmacol. Toxicol.* 32:573-596 (1993); Mulligan, *Science* 260:926-932 (1993); and Morgan and Anderson, *Ann. Rev. Biochem.* 62:191-217 (1993); May, *TIBTECH* 11(5):155-215 (1993). Methods commonly known in the art of recombinant DNA technology which can be used are described in Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, NY (1993); and Kriegler, *Gene Transfer and Expression, A Laboratory Manual*, Stockton Press, NY (1990).

In a preferred aspect, the compound comprises nucleic acid sequences encoding an antibody, said nucleic acid sequences being part of expression vectors that express the antibody or fragments or chimeric proteins or heavy or light chains thereof in a suitable host. In particular, such nucleic acid sequences have promoters operably linked to the antibody coding region, said promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, nucleic acid molecules are used in which the antibody coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the antibody encoding nucleic acids (Koller and Smithies, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); Zijlstra et al., *Nature* 342:435-438 (1989).

In specific embodiments, the expressed antibody molecule is a single chain antibody; alternatively, the nucleic acid sequences include sequences encoding both the heavy and light chains, or fragments thereof, of the antibody.

Delivery of the nucleic acids into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vectors, or indirect, in which case, cells are first transformed with the nucleic acids in vitro, then transplanted into the patient. These two approaches are known, respectively, as in vivo or ex vivo gene therapy.

In a specific embodiment, the nucleic acid sequences are directly administered in vivo, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing them as part of an appropriate nucleic acid expression vector and administering it so that they become intracellular, e.g., by infection using defective or attenuated retrovirals or other viral vectors (see U.S. Patent No. 4,980,286), or by direct injection of naked DNA, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes, microparticles, or microcapsules, or by administering them in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see, e.g., Wu and Wu, J. Biol. Chem. 262:4429-4432 (1987)) (which can be used to target cell types specifically expressing the receptors), etc. In another embodiment, nucleic acid-ligand complexes can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted in vivo for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188, WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (Koller and Smithies, Proc. Natl. Acad. Sci. USA 86:8932-8935 (1989); Zijlstra et al., Nature 342:435-438 (1989)).

In a specific embodiment, viral vectors that contains nucleic acid sequences encoding an antibody of the invention are used. For example, a retroviral vector can be used (see Miller et al., Meth. Enzymol. 217:581-599 (1993)). These retroviral vectors contain the components necessary for the correct packaging of the viral genome and integration into the

host cell DNA. The nucleic acid sequences encoding the antibody to be used in gene therapy are cloned into one or more vectors, which facilitates delivery of the gene into a patient. More detail about retroviral vectors can be found in Boesen et al., *Biotherapy* 6:291-302 (1994), which describes the use of a retroviral vector to deliver the *mdr1* gene to hematopoietic stem cells in order to make the stem cells more resistant to chemotherapy. Other references illustrating the use of retroviral vectors in gene therapy are: Clowes et al., *J. Clin. Invest.* 93:644-651 (1994); Kiem et al., *Blood* 83:1467-1473 (1994); Salmons and Gunzberg, *Human Gene Therapy* 4:129-141 (1993); and Grossman and Wilson, *Curr. Opin. in Genetics and Devel.* 3:110-114 (1993).

Adenoviruses are other viral vectors that can be used in gene therapy. Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting non-dividing cells. Kozarsky and Wilson, *Current Opinion in Genetics and Development* 3:499-503 (1993) present a review of adenovirus-based gene therapy. Bout et al., *Human Gene Therapy* 5:3-10 (1994) demonstrated the use of adenovirus vectors to transfer genes to the respiratory epithelia of rhesus monkeys. Other instances of the use of adenoviruses in gene therapy can be found in Rosenfeld et al., *Science* 252:431-434 (1991); Rosenfeld et al., *Cell* 68:143-155 (1992); Mastrangeli et al., *J. Clin. Invest.* 91:225-234 (1993); PCT Publication WO94/12649; and Wang, et al., *Gene Therapy* 2:775-783 (1995). In a preferred embodiment, adenovirus vectors are used.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy (Walsh et al., *Proc. Soc. Exp. Biol. Med.* 204:289-300 (1993); U.S. Patent No. 5,436,146).

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration in vivo of the resulting recombinant cell. Such introduction can be carried out by any method

known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleic acid sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc. Numerous techniques are known in the art for the introduction of foreign genes
5 into cells (see, e.g., Loeffler and Behr, *Meth. Enzymol.* 217:599-618 (1993); Cohen et al., *Meth. Enzymol.* 217:618-644 (1993); Cline, *Pharmac. Ther.* 29:69-92m (1985) and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is
10 expressible by the cell and preferably heritable and expressible by its cell progeny.

The resulting recombinant cells can be delivered to a patient by various methods known in the art. Recombinant blood cells (e.g., hematopoietic stem or progenitor cells) are preferably administered intravenously. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

15 Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as Tlymphocytes, Blymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic
20 stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc.

In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, nucleic acid sequences encoding an antibody are introduced into the cells such that they are expressible
25 by the cells or their progeny, and the recombinant cells are then administered in vivo for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem and/or progenitor cells which can be isolated and maintained in vitro can potentially be used in accordance with this embodiment of the present invention (see e.g. PCT Publication WO 94/08598; Stemple and Anderson, *Cell* 71:973-985 (1992); Rheinwald, *Meth. Cell Bio.*
30 21A:229 (1980); and Pittelkow and Scott, *Mayo Clinic Proc.* 61:771 (1986)).

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy comprises an inducible promoter operably linked to the coding region, such that

expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription. **Demonstration of Therapeutic or Prophylactic Activity**

The compounds or pharmaceutical compositions of the invention are preferably tested in vitro, and then in vivo for the desired therapeutic or prophylactic activity, prior to use in
5 humans. For example, in vitro assays to demonstrate the therapeutic or prophylactic utility of a compound or pharmaceutical composition include, the effect of a compound on a cell line or a patient tissue sample. The effect of the compound or composition on the cell line and/or tissue sample can be determined utilizing techniques known to those of skill in the art including, but not limited to, rosette formation assays and cell lysis assays. In accordance
10 with the invention, in vitro assays which can be used to determine whether administration of a specific compound is indicated, include in vitro cell culture assays in which a patient tissue sample is grown in culture, and exposed to or otherwise administered a compound, and the effect of such compound upon the tissue sample is observed.

15 ***Therapeutic/Prophylactic Administration and Composition***

The invention provides methods of treatment, inhibition and prophylaxis by administration to a subject of an effective amount of a compound or pharmaceutical composition of the invention, preferably a polypeptide or antibody of the invention. In a preferred aspect, the compound is substantially purified (e.g., substantially free from
20 substances that limit its effect or produce undesired side-effects). The subject is preferably an animal, including but not limited to animals such as cows, pigs, horses, chickens, cats, dogs, etc., and is preferably a mammal, and most preferably human.

Formulations and methods of administration that can be employed when the compound comprises a nucleic acid or an immunoglobulin are described above; additional
25 appropriate formulations and routes of administration can be selected from among those described herein below.

Various delivery systems are known and can be used to administer a compound of the invention, e.g., encapsulation in liposomes, microparticles, microcapsules, recombinant cells capable of expressing the compound, receptor-mediated endocytosis (see, e.g., Wu and Wu,
30 J. Biol. Chem. 262:4429-4432 (1987)). construction of a nucleic acid as part of a retroviral or other vector, etc. Methods of introduction include but are not limited to intradermal, intramuscular, intraperitoneal, intravenous, subcutaneous, intranasal, epidural, and oral

routes. The compounds or compositions may be administered by any convenient route, for example by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local. In addition, it may be desirable to introduce the pharmaceutical compounds or compositions of the invention into the central nervous system by any suitable route, including intraventricular and intrathecal injection; intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir, such as an Ommaya reservoir. Pulmonary administration can also be employed, e.g., by use of an inhaler or nebulizer, and formulation with an aerosolizing agent.

In a specific embodiment, it may be desirable to administer the pharmaceutical compounds or compositions of the invention locally to the area in need of treatment; this may be achieved by, for example, and not by way of limitation, local infusion during surgery, topical application, e.g., in conjunction with a wound dressing after surgery, by injection, by means of a catheter, by means of a suppository, or by means of an implant, said implant being of a porous, non-porous, or gelatinous material, including membranes, such as sialastic membranes, or fibers. Preferably, when administering a protein, including an antibody, of the invention, care must be taken to use materials to which the protein does not absorb.

In another embodiment, the compound or composition can be delivered in a vesicle, in particular a liposome (see Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353- 365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*)

In yet another embodiment, the compound or composition can be delivered in a controlled release system. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, *CRC Crit. Rev. Biomed. Eng.* 14:201 (1987); Buchwald et al., *Surgery* 88:507 (1980); Saudek et al., *N. Engl. J. Med.* 321:574 (1989)). In another embodiment, polymeric materials can be used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres., Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J., *Macromol. Sci. Rev. Macromol. Chem.* 23:61 (1983); see also Levy et al., *Science* 228:190 (1985); During et al., *Ann. Neurol.* 25:351 (1989); Howard et al.,

J.Neurosurg. 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, *supra*, vol. 2, pp. 115-138 (1984)).

5 Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

 In a specific embodiment where the compound of the invention is a nucleic acid encoding a protein, the nucleic acid can be administered in vivo to promote expression of its encoded protein, by constructing it as part of an appropriate nucleic acid expression vector
10 and administering it so that it becomes intracellular, e.g., by use of a retroviral vector (see U.S. Patent No. 4,980,286), or by direct injection, or by use of microparticle bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, or by administering it in linkage to a homeobox- like peptide which is known to enter the nucleus (see e.g., Joliot et al., Proc. Natl. Acad. Sci. USA 88:1864-1868
15 (1991)), etc. Alternatively, a nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination.

 The present invention also provides pharmaceutical compositions. Such compositions comprise a therapeutically effective amount of a compound, and a pharmaceutically acceptable carrier. In a specific embodiment, the term "pharmaceutically acceptable" means
20 approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic
25 origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol
30 monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene glycol, water, ethanol and the like. The composition, if desired, can also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions can take the form of

solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like. The composition can be formulated as a suppository, with traditional binders and carriers such as triglycerides. Oral formulation can include standard carriers such as pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharine, cellulose, magnesium carbonate, etc. Examples of suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin. Such compositions will contain a therapeutically effective amount of the compound, preferably in purified form, together with a suitable amount of carrier so as to provide the form for proper administration to the patient. The formulation should suit the mode of administration.

In a preferred embodiment, the composition is formulated in accordance with routine procedures as a pharmaceutical composition adapted for intravenous administration to human beings. Typically, compositions for intravenous administration are solutions in sterile isotonic aqueous buffer. Where necessary, the composition may also include a solubilizing agent and a local anesthetic such as lignocaine to ease pain at the site of the injection. Generally, the ingredients are supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline can be provided so that the ingredients may be mixed prior to administration.

The compounds of the invention can be formulated as neutral or salt forms. Pharmaceutically acceptable salts include those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

The amount of the compound of the invention which will be effective in the treatment, inhibition and prevention of a disease or disorder associated with aberrant expression and/or activity of a polypeptide of the invention can be determined by standard clinical techniques. In addition, in vitro assays may optionally be employed to help identify optimal dosage ranges. The precise dose to be employed in the formulation will also depend

on the route of administration, and the seriousness of the disease or disorder, and should be decided according to the judgment of the practitioner and each patient's circumstances. Effective doses may be extrapolated from dose-response curves derived from in vitro or animal model test systems.

5 For antibodies, the dosage administered to a patient is typically 0.1 mg/kg to 100 mg/kg of the patient's body weight. Preferably, the dosage administered to a patient is between 0.1 mg/kg and 20 mg/kg of the patient's body weight, more preferably 1 mg/kg to 10 mg/kg of the patient's body weight. Generally, human antibodies have a longer half-life
10 foreign polypeptides. Thus, lower dosages of human antibodies and less frequent administration is often possible. Further, the dosage and frequency of administration of antibodies of the invention may be reduced by enhancing uptake and tissue penetration (e.g., into the brain) of the antibodies by modifications such as, for example, lipidation.

The invention also provides a pharmaceutical pack or kit comprising one or more
15 containers filled with one or more of the ingredients of the pharmaceutical compositions of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration.

20

Diagnosis and Imaging

Labeled antibodies, and derivatives and analogs thereof, which specifically bind to a polypeptide of interest can be used for diagnostic purposes to detect, diagnose, or monitor
25 of a polypeptide of the invention. The invention provides for the detection of aberrant expression of a polypeptide of interest, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide
30 gene expression level compared to the standard expression level is indicative of aberrant expression.

The invention provides a diagnostic assay for diagnosing a disorder, comprising (a) assaying the expression of the polypeptide of interest in cells or body fluid of an individual using one or more antibodies specific to the polypeptide interest and (b) comparing the level of gene expression with a standard gene expression level, whereby an increase or decrease in the assayed polypeptide gene expression level compared to the standard expression level is indicative of a particular disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Antibodies of the invention can be used to assay protein levels in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{125}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium (^{112}In), and technetium (^{99}Tc); luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

One aspect of the invention is the detection and diagnosis of a disease or disorder associated with aberrant expression of a polypeptide of interest in an animal, preferably a mammal and most preferably a human. In one embodiment, diagnosis comprises: a) administering (for example, parenterally, subcutaneously, or intraperitoneally) to a subject an effective amount of a labeled molecule which specifically binds to the polypeptide of interest; b) waiting for a time interval following the administering for permitting the labeled molecule to preferentially concentrate at sites in the subject where the polypeptide is expressed (and for unbound labeled molecule to be cleared to background level); c) determining background level; and d) detecting the labeled molecule in the subject, such that detection of labeled molecule above the background level indicates that the subject has a particular disease or disorder associated with aberrant expression of the polypeptide of interest. Background level can be determined by various methods including, comparing the

amount of labeled molecule detected to a standard value previously determined for a particular system.

It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of ^{99m}Tc. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which contain the specific protein. In vivo tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments." (Chapter 13 in Tumor Imaging: The Radiochemical Detection of Cancer, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982).

Depending on several variables, including the type of label used and the mode of administration, the time interval following the administration for permitting the labeled molecule to preferentially concentrate at sites in the subject and for unbound labeled molecule to be cleared to background level is 6 to 48 hours or 6 to 24 hours or 6 to 12 hours. In another embodiment the time interval following administration is 5 to 20 days or 5 to 10 days.

In an embodiment, monitoring of the disease or disorder is carried out by repeating the method for diagnosing the disease or disorder, for example, one month after initial diagnosis, six months after initial diagnosis, one year after initial diagnosis, etc.

Presence of the labeled molecule can be detected in the patient using methods known in the art for in vivo scanning. These methods depend upon the type of label used. Skilled artisans will be able to determine the appropriate method for detecting a particular label. Methods and devices that may be used in the diagnostic methods of the invention include, but are not limited to, computed tomography (CT), whole body scan such as positron emission tomography (PET), magnetic resonance imaging (MRI), and sonography.

In a specific embodiment, the molecule is labeled with a radioisotope and is detected in the patient using a radiation responsive surgical instrument (Thurston et al., U.S. Patent No. 5,441,050). In another embodiment, the molecule is labeled with a fluorescent compound and is detected in the patient using a fluorescence responsive scanning instrument. In another embodiment, the molecule is labeled with a positron emitting metal and is detected in the patient using positron emission-tomography. In yet another embodiment, the molecule

is labeled with a paramagnetic label and is detected in a patient using magnetic resonance imaging (MRI).

Kits

5 The present invention provides kits that can be used in the above methods. In one embodiment, a kit comprises an antibody of the invention, preferably a purified antibody, in one or more containers. In a specific embodiment, the kits of the present invention contain a substantially isolated polypeptide comprising an epitope which is specifically immunoreactive with an antibody included in the kit. Preferably, the kits of the present
10 invention further comprise a control antibody which does not react with the polypeptide of interest. In another specific embodiment, the kits of the present invention contain a means for detecting the binding of an antibody to a polypeptide of interest (e.g., the antibody may be conjugated to a detectable substrate such as a fluorescent compound, an enzymatic substrate, a radioactive compound or a luminescent compound, or a second antibody which recognizes
15 the first antibody may be conjugated to a detectable substrate).

 In another specific embodiment of the present invention, the kit is a diagnostic kit for use in screening serum containing antibodies specific against proliferative and/or cancerous polynucleotides and polypeptides. Such a kit may include a control antibody that does not react with the polypeptide of interest. Such a kit may include a substantially isolated
20 polypeptide antigen comprising an epitope which is specifically immunoreactive with at least one anti-polypeptide antigen antibody. Further, such a kit includes means for detecting the binding of said antibody to the antigen (e.g., the antibody may be conjugated to a fluorescent compound such as fluorescein or rhodamine which can be detected by flow cytometry). In specific embodiments, the kit may include a recombinantly produced or chemically
25 synthesized polypeptide antigen. The polypeptide antigen of the kit may also be attached to a solid support.

 In a more specific embodiment the detecting means of the above-described kit includes a solid support to which said polypeptide antigen is attached. Such a kit may also include a non-attached reporter-labeled anti-human antibody. In this embodiment, binding of
30 the antibody to the polypeptide antigen can be detected by binding of the said reporter-labeled antibody.

In an additional embodiment, the invention includes a diagnostic kit for use in screening serum containing antigens of the polypeptide of the invention. The diagnostic kit includes a substantially isolated antibody specifically immunoreactive with polypeptide or polynucleotide antigens, and means for detecting the binding of the polynucleotide or polypeptide antigen to the antibody. In one embodiment, the antibody is attached to a solid support. In a specific embodiment, the antibody may be a monoclonal antibody. The detecting means of the kit may include a second, labeled monoclonal antibody. Alternatively, or in addition, the detecting means may include a labeled, competing antigen.

In one diagnostic configuration, test serum is reacted with a solid phase reagent having a surface-bound antigen obtained by the methods of the present invention. After binding with specific antigen antibody to the reagent and removing unbound serum components by washing, the reagent is reacted with reporter-labeled anti-human antibody to bind reporter to the reagent in proportion to the amount of bound anti-antigen antibody on the solid support. The reagent is again washed to remove unbound labeled antibody, and the amount of reporter associated with the reagent is determined. Typically, the reporter is an enzyme which is detected by incubating the solid phase in the presence of a suitable fluorometric, luminescent or colorimetric substrate (Sigma, St. Louis, MO).

The solid surface reagent in the above assay is prepared by known techniques for attaching protein material to solid support material, such as polymeric beads, dip sticks, 96-well plate or filter material. These attachment methods generally include non-specific adsorption of the protein to the support or covalent attachment of the protein, typically through a free amine group, to a chemically reactive group on the solid support, such as an activated carboxyl, hydroxyl, or aldehyde group. Alternatively, streptavidin coated plates can be used in conjunction with biotinylated antigen(s).

Thus, the invention provides an assay system or kit for carrying out this diagnostic method. The kit generally includes a support with surface-bound recombinant antigens, and a reporter-labeled anti-human antibody for detecting surface-bound anti-antigen antibody.

Uses of the Polynucleotides

Each of the polynucleotides identified herein can be used in numerous ways as reagents. The following description should be considered exemplary and utilizes known techniques.

The colon cancer antigen polynucleotides of the present invention are useful for chromosome identification. There exists an ongoing need to identify new chromosome markers, since few chromosome marking reagents, based on actual sequence data (repeat polymorphisms), are presently available. Each sequence is specifically targeted to and can
5 hybridize with a particular location on an individual human chromosome, thus each polynucleotide of the present invention can routinely be used as a chromosome marker using techniques known in the art.

Briefly, sequences can be mapped to chromosomes by preparing PCR primers (preferably at least 15 bp (e.g., 15-25 bp) from the sequences shown in SEQ ID NO:X, or the
10 complement thereto. Primers can optionally be selected using computer analysis so that primers do not span more than one predicted exon in the genomic DNA. These primers are then used for PCR screening of somatic cell hybrids containing individual human chromosomes. Only those hybrids containing the human gene corresponding to SEQ ID NO:X will yield an amplified fragment.

15 Similarly, somatic hybrids provide a rapid method of PCR mapping the polynucleotides to particular chromosomes. Three or more clones can be assigned per day using a single thermal cycler. Moreover, sublocalization of the polynucleotides can be achieved with panels of specific chromosome fragments. Other gene mapping strategies that can be used include in situ hybridization, prescreening with labeled flow-sorted
20 chromosomes, preselection by hybridization to construct chromosome specific-cDNA libraries, and computer mapping techniques (See, e.g., Shuler, Trends Biotechnol 16:456-459 (1998) which is hereby incorporated by reference in its entirety).

Precise chromosomal location of the polynucleotides can also be achieved using fluorescence in situ hybridization (FISH) of a metaphase chromosomal spread. This
25 technique uses polynucleotides as short as 500 or 600 bases; however, polynucleotides 2,000-4,000 bp are preferred. For a review of this technique, see Verma et al., "Human Chromosomes: a Manual of Basic Techniques," Pergamon Press, New York (1988).

For chromosome mapping, the polynucleotides can be used individually (to mark a single chromosome or a single site on that chromosome) or in panels (for marking multiple
30 sites and/or multiple chromosomes).

Thus, the present invention also provides a method for chromosomal localization which involves (a) preparing PCR primers from the polynucleotide sequences in Table 3 and SEQ ID NO:X and (b) screening somatic cell hybrids containing individual chromosomes.

The polynucleotides of the present invention would likewise be useful for radiation
5 hybrid mapping, HAPPY mapping, and long range restriction mapping. For a review of these techniques and others known in the art, see, e.g. Dear, "Genome Mapping: A Practical Approach," IRL Press at Oxford University Press, London (1997); Aydin. J. Mol. Med. 77:691-694 (1999); Hacia et al., Mol. Psychiatry 3:483-492 (1998); Herrick et al., Chromosome Res. 7:409-423 (1999); Hamilton et al., Methods Cell Biol. 62:265-280 (2000);
10 and/or Ott, J. Hered. 90:68-70 (1999) each of which is hereby incorporated by reference in its entirety.

Once a polynucleotide has been mapped to a precise chromosomal location, the physical position of the polynucleotide can be used in linkage analysis. Linkage analysis establishes coinheritance between a chromosomal location and presentation of a particular
15 disease. (Disease mapping data are found, for example, in V. McKusick, Mendelian Inheritance in Man (available on line through Johns Hopkins University Welch Medical Library).) Assuming 1 megabase mapping resolution and one gene per 20 kb, a cDNA precisely localized to a chromosomal region associated with the disease could be one of 50-500 potential causative genes.

20 Thus, once coinheritance is established, differences in a polynucleotide of the invention and the corresponding gene between affected and unaffected individuals can be examined. First, visible structural alterations in the chromosomes, such as deletions or translocations, are examined in chromosome spreads or by PCR. If no structural alterations exist, the presence of point mutations are ascertained. Mutations observed in some or all
25 affected individuals, but not in normal individuals, indicates that the mutation may cause the disease. However, complete sequencing of the polypeptide and the corresponding gene from several normal individuals is required to distinguish the mutation from a polymorphism. If a new polymorphism is identified, this polymorphic polypeptide can be used for further linkage analysis.

30 Furthermore, increased or decreased expression of the gene in affected individuals as compared to unaffected individuals can be assessed using the polynucleotides of the

invention. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

Thus, the invention provides a method of detecting increased or decreased expression levels of the colon cancer polynucleotides in affected individuals as compared to unaffected
5 individuals using polynucleotides of the present invention and techniques known in the art, including but not limited to the method described in Example 11. Any of these alterations (altered expression, chromosomal rearrangement, or mutation) can be used as a diagnostic or prognostic marker.

Thus, the invention also provides a diagnostic method useful during diagnosis of a
10 colon related disorder, including colon cancer, involving measuring the expression level of colon cancer polynucleotides in colon tissue or other cells or body fluid from an individual and comparing the measured gene expression level with a standard colon cancer polynucleotide expression level, whereby an increase or decrease in the gene expression level compared to the standard is indicative of a colon related disorder.

15 In still another embodiment, the invention includes a kit for analyzing samples for the presence of proliferative and/or cancerous polynucleotides derived from a test subject. In a general embodiment, the kit includes at least one polynucleotide probe containing a nucleotide sequence that will specifically hybridize with a polynucleotide of the invention and a suitable container. In a specific embodiment, the kit includes two polynucleotide probes
20 defining an internal region of the polynucleotide of the invention, where each probe has one strand containing a 31'-mer-end internal to the region. In a further embodiment, the probes may be useful as primers for polymerase chain reaction amplification.

Where a diagnosis of a colon related disorder, including, for example, diagnosis of a tumor, has already been made according to conventional methods, the present invention is
25 useful as a prognostic indicator, whereby patients exhibiting enhanced or depressed colon cancer polynucleotide expression will experience a worse clinical outcome relative to patients expressing the gene at a level nearer the standard level.

By "measuring the expression level of colon cancer polynucleotides" is intended qualitatively or quantitatively measuring or estimating the level of the colon cancer
30 polypeptide or the level of the mRNA encoding the colon cancer polypeptide in a first biological sample either directly (e.g., by determining or estimating absolute protein level or mRNA level) or relatively (e.g., by comparing to the colon cancer polypeptide level or

mRNA level in a second biological sample). Preferably, the colon cancer polypeptide level or mRNA level in the first biological sample is measured or estimated and compared to a standard colon cancer polypeptide level or mRNA level, the standard being taken from a second biological sample obtained from an individual not having the colon related disorder or
5 being determined by averaging levels from a population of individuals not having a colon related disorder. As will be appreciated in the art, once a standard colon cancer polypeptide level or mRNA level is known, it can be used repeatedly as a standard for comparison.

By "biological sample" is intended any biological sample obtained from an individual, body fluid, cell line, tissue culture, or other source which contains colon cancer polypeptide
10 or the corresponding mRNA. As indicated, biological samples include body fluids (such as lymph, sera, plasma, urine, bile, synovial fluid and spinal fluid) which contain the colon cancer polypeptide, colon tissue, and other tissue sources found to express the colon cancer polypeptide. Methods for obtaining tissue biopsies and body fluids from mammals are well known in the art. Where the biological sample is to include mRNA, a tissue biopsy is the
15 preferred source.

The method(s) provided above may preferably be applied in a diagnostic method and/or kits in which polynucleotides and/or polypeptides of the invention are attached to a solid support. In one exemplary method, the support may be a "gene chip" or a "biological chip" as described in US Patents 5,837,832, 5,874,219, and 5,856,174. Further, such a gene
20 chip with colon cancer polynucleotides attached may be used to identify polymorphisms between the colon cancer polynucleotide sequences, with polynucleotides isolated from a test subject. The knowledge of such polymorphisms (i.e. their location, as well as, their existence) would be beneficial in identifying disease loci for many disorders, such as for example, in neural disorders, immune system disorders, muscular disorders, reproductive
25 disorders, gastrointestinal disorders, pulmonary disorders, cardiovascular disorders, renal disorders, proliferative disorders, and/or cancerous diseases and conditions, though most preferably in colon related proliferative, and/or cancerous diseases and conditions. Such a method is described in US Patents 5,858,659 and 5,856,104. The US Patents referenced supra are hereby incorporated by reference in their entirety herein.

30 The present invention encompasses colon cancer polynucleotides that are chemically synthesized, or reproduced as peptide nucleic acids (PNA), or according to other methods known in the art. The use of PNAs would serve as the preferred form if the polynucleotides

of the invention are incorporated onto a solid support, or gene chip. For the purposes of the present invention, a peptide nucleic acid (PNA) is a polyamide type of DNA analog and the monomeric units for adenine, guanine, thymine and cytosine are available commercially (Perceptive Biosystems). Certain components of DNA, such as phosphorus, phosphorus
5 oxides, or deoxyribose derivatives, are not present in PNAs. As disclosed by P. E. Nielsen, M. Egholm, R. H. Berg and O. Buchardt, *Science* 254, 1497 (1991); and M. Egholm, O. Buchardt, L. Christensen, C. Behrens, S. M. Freier, D. A. Driver, R. H. Berg, S. K. Kim, B. Norden, and P. E. Nielsen, *Nature* 365, 666 (1993), PNAs bind specifically and tightly to complementary DNA strands and are not degraded by nucleases. In fact, PNA binds more
10 strongly to DNA than DNA itself does. This is probably because there is no electrostatic repulsion between the two strands, and also the polyamide backbone is more flexible. Because of this, PNA/DNA duplexes bind under a wider range of stringency conditions than DNA/DNA duplexes, making it easier to perform multiplex hybridization. Smaller probes can be used than with DNA due to the strong binding. In addition, it is more likely that single
15 base mismatches can be determined with PNA/DNA hybridization because a single mismatch in a PNA/DNA 15-mer lowers the melting point ($T_{sub.m}$) by 8°-20° C, vs. 4°-16° C for the DNA/DNA 15-mer duplex. Also, the absence of charge groups in PNA means that hybridization can be done at low ionic strengths and reduce possible interference by salt during the analysis.

20 The present invention have uses which include, but are not limited to, detecting cancer in mammals. In particular the invention is useful during diagnosis of pathological cell proliferative neoplasias which include, but are not limited to: acute myelogenous leukemias including acute monocytic leukemia, acute myeloblastic leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute erythroleukemia, acute megakaryocytic
25 leukemia, and acute undifferentiated leukemia, etc.; and chronic myelogenous leukemias including chronic myelomonocytic leukemia, chronic granulocytic leukemia, etc. Preferred mammals include monkeys, apes, cats, dogs, cows, pigs, horses, rabbits and humans. Particularly preferred are humans.

Pathological cell proliferative disorders are often associated with inappropriate
30 activation of proto-oncogenes. (Germann, E. P. et al., "The Etiology of Acute Leukemia: Molecular Genetics and Viral Oncology," in *Neoplastic Diseases of the Blood*, Vol 1., Wiernik, P. H. et al. eds., 161-182 (1985)). Neoplasias are now believed to result from the

qualitative alteration of a normal cellular gene product, or from the quantitative modification of gene expression by insertion into the chromosome of a viral sequence, by chromosomal translocation of a gene to a more actively transcribed region, or by some other mechanism. (Gelmann et al., supra) It is likely that mutated or altered expression of specific genes is involved in the pathogenesis of some leukemias, among other tissues and cell types. (Gelmann et al., supra) Indeed, the human counterparts of the oncogenes involved in some animal neoplasias have been amplified or translocated in some cases of human leukemia and carcinoma. (Gelmann et al., supra)

For example, c-myc expression is highly amplified in the non-lymphocytic leukemia cell line HL-60. When HL-60 cells are chemically induced to stop proliferation, the level of c-myc is found to be downregulated. (International Publication Number WO 91/15580). However, it has been shown that exposure of HL-60 cells to a DNA construct that is complementary to the 5' end of c-myc or c-myb blocks translation of the corresponding mRNAs which downregulates expression of the c-myc or c-myb proteins and causes arrest of cell proliferation and differentiation of the treated cells. (International Publication Number WO 91/15580; Wickstrom et al., Proc. Natl. Acad. Sci. 85:1028 (1988); Anfossi et al., Proc. Natl. Acad. Sci. 86:3379 (1989)). However, the skilled artisan would appreciate the present invention's usefulness is not limited to treatment of proliferative disorders of hematopoietic cells and tissues, in light of the numerous cells and cell types of varying origins which are known to exhibit proliferative phenotypes.

In addition to the foregoing, a colon cancer antigen polynucleotide can be used to control gene expression through triple helix formation or through antisense DNA or RNA. Antisense techniques are discussed, for example, in Okano, J. Neurochem. 56: 560 (1991); "Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance Lee et al., Nucleic Acids Research 6: 3073 (1979); Cooney et al., Science 241: 456 (1988); and Dervan et al., Science 251: 1360 (1991). Both methods rely on binding of the polynucleotide to a complementary DNA or RNA. For these techniques, preferred polynucleotides are usually oligonucleotides 20 to 40 bases in length and complementary to either the region of the gene involved in transcription (triple helix - see Lee et al., Nucl. Acids Res. 6:3073 (1979); Cooney et al., Science 241:456 (1988); and Dervan et al., Science 251:1360 (1991)) or to the mRNA itself (antisense - Okano, J. Neurochem. 56:560 (1991); Oligodeoxy-nucleotides as Antisense

Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988).) Triple helix formation optimally results in a shut-off of RNA transcription from DNA, while antisense RNA hybridization blocks translation of an mRNA molecule into polypeptide. The oligonucleotide described above can also be delivered to cells such that the antisense RNA or DNA may be
5 expressed in vivo to inhibit production of polypeptide of the present invention antigens. Both techniques are effective in model systems, and the information disclosed herein can be used to design antisense or triple helix polynucleotides in an effort to treat disease, and in particular, for the treatment of proliferative diseases and/or conditions.

Polynucleotides of the present invention are also useful in gene therapy. One goal of
10 gene therapy is to insert a normal gene into an organism having a defective gene, in an effort to correct the genetic defect. The polynucleotides disclosed in the present invention offer a means of targeting such genetic defects in a highly accurate manner. Another goal is to insert a new gene that was not present in the host genome, thereby producing a new trait in the host cell.

15 The polynucleotides are also useful for identifying individuals from minute biological samples. The United States military, for example, is considering the use of restriction fragment length polymorphism (RFLP) for identification of its personnel. In this technique, an individual's genomic DNA is digested with one or more restriction enzymes, and probed on a Southern blot to yield unique bands for identifying personnel. This method does not
20 suffer from the current limitations of "Dog Tags" which can be lost, switched, or stolen, making positive identification difficult. The polynucleotides of the present invention can be used as additional DNA markers for RFLP.

The polynucleotides of the present invention can also be used as an alternative to RFLP, by determining the actual base-by-base DNA sequence of selected portions of an
25 individual's genome. These sequences can be used to prepare PCR primers for amplifying and isolating such selected DNA, which can then be sequenced. Using this technique, individuals can be identified because each individual will have a unique set of DNA sequences. Once an unique ID database is established for an individual, positive identification of that individual, living or dead, can be made from extremely small tissue
30 samples.

Forensic biology also benefits from using DNA-based identification techniques as disclosed herein. DNA sequences taken from very small biological samples such as tissues,

e.g., hair or skin, or body fluids, e.g., blood, saliva, semen, synovial fluid, amniotic fluid, breast milk, lymph, pulmonary sputum or surfactant, urine, fecal matter, etc., can be amplified using PCR. In one prior art technique, gene sequences amplified from polymorphic loci, such as DQa class II HLA gene, are used in forensic biology to identify individuals. (Erlich, H., PCR Technology, Freeman and Co. (1992).) Once these specific polymorphic loci are amplified, they are digested with one or more restriction enzymes, yielding an identifying set of bands on a Southern blot probed with DNA corresponding to the DQa class II HLA gene. Similarly, polynucleotides of the present invention can be used as polymorphic markers for forensic purposes.

10 There is also a need for reagents capable of identifying the source of a particular tissue. Such need arises, for example, in forensics when presented with tissue of unknown origin. Appropriate reagents can comprise, for example, DNA probes or primers specific to colon or colon cancer polynucleotides prepared from the sequences of the present invention. Panels of such reagents can identify tissue by species and/or by organ type. In a similar
15 fashion, these reagents can be used to screen tissue cultures for contamination.

 The polynucleotides of the present invention are also useful as hybridization probes for differential identification of the tissue(s) or cell type(s) present in a biological sample. Similarly, polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of the tissue(s) (e.g.,
20 immunohistochemistry assays) or cell type(s) (e.g., immunocytochemistry assays). In addition, for a number of disorders of the above tissues or cells, significantly higher or lower levels of gene expression of the polynucleotides/polypeptides of the present invention may be detected in certain tissues (e.g., tissues expressing polypeptides and/or polynucleotides of the present invention, colon and colon cancer tissues and/or cancerous and/or wounded tissues)
25 or bodily fluids (e.g., serum, plasma, urine, synovial fluid or spinal fluid) taken from an individual having such a disorder, relative to a "standard" gene expression level, i.e., the expression level in healthy tissue from an individual not having the disorder.

 Thus, the invention provides a diagnostic method of a disorder, which involves: (a) assaying gene expression level in cells or body fluid of an individual; (b) comparing the gene
30 expression level with a standard gene expression level, whereby an increase or decrease in the assayed gene expression level compared to the standard expression level is indicative of a disorder.

In the very least, the polynucleotides of the present invention can be used as molecular weight markers on Southern gels, as diagnostic probes for the presence of a specific mRNA in a particular cell type, as a probe to "subtract-out" known sequences in the process of discovering novel polynucleotides, for selecting and making oligomers for attachment to a "gene chip" or other support, to raise anti-DNA antibodies using DNA immunization techniques, and as an antigen to elicit an immune response.

Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

Polypeptides and antibodies directed to polypeptides of the present invention are useful to provide immunological probes for differential identification of the tissue(s) (e.g., immunohistochemistry assays such as, for example, ABC immunoperoxidase (Hsu et al., J. Histochem. Cytochem. 29:577-580 (1981)) or cell type(s) (e.g., immunocytochemistry assays).

Antibodies can be used to assay levels of polypeptides encoded by polynucleotides of the invention in a biological sample using classical immunohistological methods known to those of skill in the art (e.g., see Jalkanen, et al., J. Cell. Biol. 101:976-985 (1985); Jalkanen, et al., J. Cell. Biol. 105:3087-3096 (1987)). Other antibody-based methods useful for detecting protein gene expression include immunoassays, such as the enzyme linked immunosorbent assay (ELISA) and the radioimmunoassay (RIA). Suitable antibody assay labels are known in the art and include enzyme labels, such as, glucose oxidase; radioisotopes, such as iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium ($^{115\text{m}}\text{In}$, $^{113\text{m}}\text{In}$, ^{112}In , ^{111}In), and technetium (^{99}Tc , $^{99\text{m}}\text{Tc}$), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F), ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru ; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

In addition to assaying levels of polypeptide of the present invention in a biological sample, proteins can also be detected in vivo by imaging. Antibody labels or markers for in vivo imaging of protein include those detectable by X-radiography, NMR or ESR. For X-radiography, suitable labels include radioisotopes such as barium or cesium, which emit

detectable radiation but are not overtly harmful to the subject. Suitable markers for NMR and ESR include those with a detectable characteristic spin, such as deuterium, which may be incorporated into the antibody by labeling of nutrients for the relevant hybridoma.

A protein-specific antibody or antibody fragment which has been labeled with an appropriate detectable imaging moiety, such as a radioisotope (for example, ^{131}I , ^{112}In , $^{99\text{m}}\text{Tc}$, ^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), sulfur (^{35}S), tritium (^3H), indium ($^{115\text{m}}\text{In}$, $^{113\text{m}}\text{In}$, ^{112}In , ^{111}In), and technetium (^{99}Tc , $^{99\text{m}}\text{Tc}$), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F , ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru), a radio-opaque substance, or a material detectable by nuclear magnetic resonance, is introduced (for example, parenterally, subcutaneously or intraperitoneally) into the mammal to be examined for immune system disorder. It will be understood in the art that the size of the subject and the imaging system used will determine the quantity of imaging moiety needed to produce diagnostic images. In the case of a radioisotope moiety, for a human subject, the quantity of radioactivity injected will normally range from about 5 to 20 millicuries of $^{99\text{m}}\text{Tc}$. The labeled antibody or antibody fragment will then preferentially accumulate at the location of cells which express the polypeptide encoded by a polynucleotide of the invention. *In vivo* tumor imaging is described in S.W. Burchiel et al., "Immunopharmacokinetics of Radiolabeled Antibodies and Their Fragments" (Chapter 13 in *Tumor Imaging: The Radiochemical Detection of Cancer*, S.W. Burchiel and B. A. Rhodes, eds., Masson Publishing Inc. (1982)).

In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (e.g., polypeptides encoded by polynucleotides of the invention and/or antibodies) that are associated with heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention in association with toxins or cytotoxic prodrugs.

By "toxin" is meant one or more compounds that bind and activate endogenous cytotoxic effector systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin, momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. "Toxin" also includes a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ^{213}Bi , or other radioisotopes such as, for example, ^{103}Pd , ^{133}Xe , ^{131}I , ^{68}Ge , ^{57}Co , ^{65}Zn , ^{85}Sr , ^{32}P , ^{35}S , ^{90}Y , ^{153}Sm , ^{153}Gd , ^{169}Yb , ^{51}Cr , ^{54}Mn , ^{75}Se , ^{113}Sn , $^{90}\text{Yttrium}$, ^{117}Tin , $^{186}\text{Rhenium}$, $^{166}\text{Holmium}$, and $^{188}\text{Rhenium}$; luminescent labels, such as luminol; and fluorescent labels, such as fluorescein and rhodamine, and biotin.

Techniques known in the art may be applied to label polypeptides of the invention (including antibodies). Such techniques include, but are not limited to, the use of bifunctional conjugating agents (see e.g., U.S. Patent Nos. 5,756,065; 5,714,631; 5,696,239; 5,652,361; 5,505,931; 5,489,425; 5,435,990; 5,428,139; 5,342,604; 5,274,119; 4,994,560; and 5,808,003; the contents of each of which are hereby incorporated by reference in its entirety).

Thus, the invention provides a diagnostic method of a disorder, which involves (a) assaying the expression level of a colon cancer polypeptide of the present invention in cells or body fluid of an individual, or more preferably, assaying the expression level of a colon cancer polypeptide of the present invention in colon cells or sera of an individual; and (b) comparing the assayed polypeptide expression level with a standard polypeptide expression level, whereby an increase or decrease in the assayed polypeptide expression level compared to the standard expression level is indicative of a disorder. With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or

aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Moreover, colon cancer antigen polypeptides of the present invention can be used to treat or prevent diseases or conditions such as, for example, neural disorders, immune system disorders, muscular disorders, reproductive disorders, gastrointestinal disorders, pulmonary disorders, cardiovascular disorders, renal disorders, proliferative disorders, and/or cancerous diseases and conditions, preferably proliferative disorders of the colon, and/or cancerous disease and conditions. For example, patients can be administered a polypeptide of the present invention in an effort to replace absent or decreased levels of the polypeptide (e.g., insulin), to supplement absent or decreased levels of a different polypeptide (e.g., hemoglobin S for hemoglobin B, SOD, catalase, DNA repair proteins), to inhibit the activity of a polypeptide (e.g., an oncogene or tumor suppressor), to activate the activity of a polypeptide (e.g., by binding to a receptor), to reduce the activity of a membrane bound receptor by competing with it for free ligand (e.g., soluble TNF receptors used in reducing inflammation), or to bring about a desired response (e.g., blood vessel growth inhibition, enhancement of the immune response to proliferative cells or tissues).

Similarly, antibodies directed to a polypeptide of the present invention can also be used to treat disease (as described supra, and elsewhere herein). For example, administration of an antibody directed to a polypeptide of the present invention can bind, and/or neutralize the polypeptide, and/or reduce overproduction of the polypeptide. Similarly, administration of an antibody can activate the polypeptide, such as by binding to a polypeptide bound to a membrane (receptor).

At the very least, the polypeptides of the present invention can be used as molecular weight markers on SDS-PAGE gels or on molecular sieve gel filtration columns using methods well known to those of skill in the art. Polypeptides can also be used to raise antibodies, which in turn are used to measure protein expression from a recombinant cell, as a way of assessing transformation of the host cell. Moreover, the polypeptides of the present invention can be used to test the following biological activities.

Gene Therapy Methods

Another aspect of the present invention is to gene therapy methods for treating or preventing disorders, diseases and conditions. The gene therapy methods relate to the introduction of nucleic acid (DNA, RNA and antisense DNA or RNA) sequences into an animal to achieve expression of the polypeptide of the present invention. This method requires a polynucleotide which codes for a polypeptide of the present invention operatively linked to a promoter and any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques are known in the art, see, for example, WO90/11092, which is herein incorporated by reference.

Thus, for example, cells from a patient may be engineered with a polynucleotide (DNA or RNA) comprising a promoter operably linked to a polynucleotide of the present invention ex vivo, with the engineered cells then being provided to a patient to be treated with the polypeptide of the present invention. Such methods are well-known in the art. For example, see Belldgrun, A., et al., J. Natl. Cancer Inst. 85: 207-216 (1993); Ferrantini, M. et al., Cancer Research 53: 1107-1112 (1993); Ferrantini, M. et al., J. Immunology 153: 4604-4615 (1994); Kaido, T., et al., Int. J. Cancer 60: 221-229 (1995); Ogura, H., et al., Cancer Research 50: 5102-5106 (1990); Santodonato, L., et al., Human Gene Therapy 7:1-10 (1996); Santodonato, L., et al., Gene Therapy 4:1246-1255 (1997); and Zhang, J.-F. et al., Cancer Gene Therapy 3: 31-38 (1996)), which are herein incorporated by reference. In one embodiment, the cells which are engineered are arterial cells. The arterial cells may be reintroduced into the patient through direct injection to the artery, the tissues surrounding the artery, or through catheter injection.

As discussed in more detail below, the polynucleotide constructs can be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of tissues (heart, muscle, skin, lung, liver, and the like). The polynucleotide constructs may be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

In one embodiment, the polynucleotide of the present invention is delivered as a naked polynucleotide. The term "naked" polynucleotide, DNA or RNA refers to sequences that are free from any delivery vehicle that acts to assist, promote or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating agents and the like. However, the polynucleotide of the present invention can

also be delivered in liposome formulations and lipofectin formulations and the like can be prepared by methods well known to those skilled in the art. Such methods are described, for example, in U.S. Patent Nos. 5,593,972, 5,589,466, and 5,580,859, which are herein incorporated by reference.

5 The polynucleotide vector constructs used in the gene therapy method are preferably constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Appropriate vectors include pWLNEO, pSV2CAT, pOG44, pXT1 and pSG available from Stratagene; pSVK3, pBPV, pMSG and pSVL available from Pharmacia; and pEF1/V5, pcDNA3.1, and pRc/CMV2 available from Invitrogen. Other suitable vectors
10 will be readily apparent to the skilled artisan.

Any strong promoter known to those skilled in the art can be used for driving the expression of the polynucleotide sequence. Suitable promoters include adenoviral promoters, such as the adenoviral major late promoter; or heterologous promoters, such as the cytomegalovirus (CMV) promoter; the respiratory syncytial virus (RSV) promoter; inducible
15 promoters, such as the MMT promoter, the metallothionein promoter; heat shock promoters; the albumin promoter; the ApoA1 promoter; human globin promoters; viral thymidine kinase promoters, such as the Herpes Simplex thymidine kinase promoter; retroviral LTRs; the b-actin promoter; and human growth hormone promoters. The promoter also may be the native promoter for the polynucleotide of the present invention.

20 Unlike other gene therapy techniques, one major advantage of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

The polynucleotide construct can be delivered to the interstitial space of tissues within
25 the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular, fluid, mucopolysaccharide matrix among the reticular fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of
30 fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is

preferred for the reasons discussed below. They may be conveniently delivered by injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, 5 stem cells of blood or skin fibroblasts. In vivo muscle cells are particularly competent in their ability to take up and express polynucleotides.

For the naked nucleic acid sequence injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 mg/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more 10 preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration.

The preferred route of administration is by the parenteral route of injection into the 15 interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous membranes of the nose. In addition, naked DNA constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The naked polynucleotides are delivered by any method known in the art, including, 20 but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, and so-called "gene guns". These delivery methods are known in the art.

The constructs may also be delivered with delivery vehicles such as viral sequences, viral particles, liposome formulations, lipofectin, precipitating agents, etc. Such methods of 25 delivery are known in the art.

In certain embodiments, the polynucleotide constructs are complexed in a liposome preparation. Liposomal preparations for use in the instant invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. However, cationic liposomes are particularly preferred because a tight charge complex can be formed 30 between the cationic liposome and the polyanionic nucleic acid. Cationic liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner et al., Proc. Natl. Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference); mRNA

(Malone et al., Proc. Natl. Acad. Sci. USA (1989) 86:6077-6081, which is herein incorporated by reference); and purified transcription factors (Debs et al., J. Biol. Chem. (1990) 265:10189-10192, which is herein incorporated by reference), in functional form.

Cationic liposomes are readily available. For example,
5 N[1-2,3-dioleoyloxy]propyl]-N,N,N-triethylammonium (DOTMA) liposomes are particularly useful and are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, N.Y. (See, also, Felgner et al., Proc. Natl Acad. Sci. USA (1987) 84:7413-7416, which is herein incorporated by reference). Other commercially available liposomes include transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer).

10 Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g. PCT Publication No. WO 90/11092 (which is herein incorporated by reference) for a description of the synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes. Preparation of DOTMA liposomes is explained in the literature, see, e.g., P. Felgner et al., Proc. Natl. Acad. Sci. USA
15 84:7413-7417, which is herein incorporated by reference. Similar methods can be used to prepare liposomes from other cationic lipid materials.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, Ala.), or can be easily prepared using readily available materials. Such materials include phosphatidyl, choline, cholesterol, phosphatidyl ethanolamine,
20 dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

For example, commercially dioleoylphosphatidyl choline (DOPC),
25 dioleoylphosphatidyl glycerol (DOPG), and dioleoylphosphatidyl ethanolamine (DOPE) can be used in various combinations to make conventional liposomes, with or without the addition of cholesterol. Thus, for example, DOPG/DOPC vesicles can be prepared by drying 50 mg each of DOPG and DOPC under a stream of nitrogen gas into a sonication vial. The sample is placed under a vacuum pump overnight and is hydrated the following day with
30 deionized water. The sample is then sonicated for 2 hours in a capped vial, using a Heat Systems model 350 sonicator equipped with an inverted cup (bath type) probe at the maximum setting while the bath is circulated at 15EC. Alternatively, negatively charged

vesicles can be prepared without sonication to produce multilamellar vesicles or by extrusion through nucleopore membranes to produce unilamellar vesicles of discrete size. Other methods are known and available to those of skill in the art.

The liposomes can comprise multilamellar vesicles (MLVs), small unilamellar vesicles (SUVs), or large unilamellar vesicles (LUVs), with SUVs being preferred. The various liposome-nucleic acid complexes are prepared using methods well known in the art. See, e.g., Straubinger et al., *Methods of Immunology* (1983), 101:512-527, which is herein incorporated by reference. For example, MLVs containing nucleic acid can be prepared by depositing a thin film of phospholipid on the walls of a glass tube and subsequently hydrating with a solution of the material to be encapsulated. SUVs are prepared by extended sonication of MLVs to produce a homogeneous population of unilamellar liposomes. The material to be entrapped is added to a suspension of preformed MLVs and then sonicated. When using liposomes containing cationic lipids, the dried lipid film is resuspended in an appropriate solution such as sterile water or an isotonic buffer solution such as 10 mM Tris/NaCl, sonicated, and then the preformed liposomes are mixed directly with the DNA. The liposome and DNA form a very stable complex due to binding of the positively charged liposomes to the cationic DNA. SUVs find use with small nucleic acid fragments. LUVs are prepared by a number of methods, well known in the art. Commonly used methods include Ca^{2+} -EDTA chelation (Papahadjopoulos et al., *Biochim. Biophys. Acta* (1975) 394:483; Wilson et al., *Cell* (1979) 17:77); ether injection (Deamer, D. and Bangham, A., *Biochim. Biophys. Acta* (1976) 443:629; Ostro et al., *Biochem. Biophys. Res. Commun.* (1977) 76:836; Fraley et al., *Proc. Natl. Acad. Sci. USA* (1979) 76:3348); detergent dialysis (Enoch, H. and Strittmatter, P., *Proc. Natl. Acad. Sci. USA* (1979) 76:145); and reverse-phase evaporation (REV) (Fraley et al., *J. Biol. Chem.* (1980) 255:10431; Szoka, F. and Papahadjopoulos, D., *Proc. Natl. Acad. Sci. USA* (1978) 75:145; Schaefer-Ridder et al., *Science* (1982) 215:166), which are herein incorporated by reference.

Generally, the ratio of DNA to liposomes will be from about 10:1 to about 1:10. Preferably, the ration will be from about 5:1 to about 1:5. More preferably, the ration will be about 3:1 to about 1:3. Still more preferably, the ratio will be about 1:1.

U.S. Patent No. 5,676,954 (which is herein incorporated by reference) reports on the injection of genetic material, complexed with cationic liposomes carriers, into mice. U.S. Patent Nos. 4,897,355, 4,946,787, 5,049,386, 5,459,127, 5,589,466, 5,693,622, 5,580,859,

5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide cationic lipids for use in transfecting DNA into cells and mammals. U.S. Patent Nos. 5,589,466, 5,693,622, 5,580,859, 5,703,055, and international publication no. WO 94/9469 (which are herein incorporated by reference) provide methods for delivering
5 DNA-cationic lipid complexes to mammals.

In certain embodiments, cells are engineered, *ex vivo* or *in vivo*, using a retroviral particle containing RNA which comprises a sequence encoding a polypeptide of the present invention. Retroviruses from which the retroviral plasmid vectors may be derived include, but are not limited to, Moloney Murine Leukemia Virus, spleen necrosis virus, Rous sarcoma
10 Virus, Harvey Sarcoma Virus, avian leukosis virus, gibbon ape leukemia virus, human immunodeficiency virus, Myeloproliferative Sarcoma Virus, and mammary tumor virus.

The retroviral plasmid vector is employed to transduce packaging cell lines to form producer cell lines. Examples of packaging cells which may be transfected include, but are not limited to, the PE501, PA317, R-2, R-AM, PA12, T19-14X, VT-19-17-H2, RCRE,
15 RCRIP, GP+E-86, GP+envAm12, and DAN cell lines as described in Miller, Human Gene Therapy 1:5-14 (1990), which is incorporated herein by reference in its entirety. The vector may transduce the packaging cells through any means known in the art. Such means include, but are not limited to, electroporation, the use of liposomes, and CaPO_4 precipitation. In one alternative, the retroviral plasmid vector may be encapsulated into a liposome, or coupled to a
20 lipid, and then administered to a host.

The producer cell line generates infectious retroviral vector particles which include polynucleotide encoding a polypeptide of the present invention. Such retroviral vector particles then may be employed, to transduce eukaryotic cells, either *in vitro* or *in vivo*. The transduced eukaryotic cells will express a polypeptide of the present invention.

25 In certain other embodiments, cells are engineered, *ex vivo* or *in vivo*, with polynucleotide contained in an adenovirus vector. Adenovirus can be manipulated such that it encodes and expresses a polypeptide of the present invention, and at the same time is inactivated in terms of its ability to replicate in a normal lytic viral life cycle. Adenovirus expression is achieved without integration of the viral DNA into the host cell chromosome,
30 thereby alleviating concerns about insertional mutagenesis. Furthermore, adenoviruses have been used as live enteric vaccines for many years with an excellent safety profile (Schwartz, A. R. et al. (1974) Am. Rev. Respir. Dis.109:233-238). Finally, adenovirus mediated gene

transfer has been demonstrated in a number of instances including transfer of alpha-1-antitrypsin and CFTR to the lungs of cotton rats (Rosenfeld, M. A. et al. (1991) Science 252:431-434; Rosenfeld et al., (1992) Cell 68:143-155). Furthermore, extensive studies to attempt to establish adenovirus as a causative agent in human cancer were uniformly negative (Green, M. et al. (1979) Proc. Natl. Acad. Sci. USA 76:6606).

Suitable adenoviral vectors useful in the present invention are described, for example, in Kozarsky and Wilson, Curr. Opin. Genet. Devel. 3:499-503 (1993); Rosenfeld et al., Cell 68:143-155 (1992); Engelhardt et al., Human Genet. Ther. 4:759-769 (1993); Yang et al., Nature Genet. 7:362-369 (1994); Wilson et al., Nature 365:691-692 (1993); and U.S. Patent No. 5,652,224, which are herein incorporated by reference. For example, the adenovirus vector Ad2 is useful and can be grown in human 293 cells. These cells contain the E1 region of adenovirus and constitutively express E1a and E1b, which complement the defective adenoviruses by providing the products of the genes deleted from the vector. In addition to Ad2, other varieties of adenovirus (e.g., Ad3, Ad5, and Ad7) are also useful in the present invention.

Preferably, the adenoviruses used in the present invention are replication deficient. Replication deficient adenoviruses require the aid of a helper virus and/or packaging cell line to form infectious particles. The resulting virus is capable of infecting cells and can express a polynucleotide of interest which is operably linked to a promoter, but cannot replicate in most cells. Replication deficient adenoviruses may be deleted in one or more of all or a portion of the following genes: E1a, E1b, E3, E4, E2a, or L1 through L5.

In certain other embodiments, the cells are engineered, ex vivo or in vivo, using an adeno-associated virus (AAV). AAVs are naturally occurring defective viruses that require helper viruses to produce infectious particles (Muzyczka, N., Curr. Topics in Microbiol. Immunol. 158:97 (1992)). It is also one of the few viruses that may integrate its DNA into non-dividing cells. Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate, but space for exogenous DNA is limited to about 4.5 kb. Methods for producing and using such AAVs are known in the art. See, for example, U.S. Patent Nos. 5,139,941, 5,173,414, 5,354,678, 5,436,146, 5,474,935, 5,478,745, and 5,589,377.

For example, an appropriate AAV vector for use in the present invention will include all the sequences necessary for DNA replication, encapsidation, and host-cell integration. The polynucleotide construct is inserted into the AAV vector using standard cloning

methods, such as those found in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989). The recombinant AAV vector is then transfected into packaging cells which are infected with a helper virus, using any standard technique, including lipofection, electroporation, calcium phosphate precipitation, etc. Appropriate
5 helper viruses include adenoviruses, cytomegaloviruses, vaccinia viruses, or herpes viruses. Once the packaging cells are transfected and infected, they will produce infectious AAV viral particles which contain the polynucleotide construct. These viral particles are then used to transduce eukaryotic cells, either ex vivo or in vivo. The transduced cells will contain the polynucleotide construct integrated into its genome, and will express a polypeptide of the
10 invention.

Another method of gene therapy involves operably associating heterologous control regions and endogenous polynucleotide sequences (e.g. encoding a polypeptide of the present invention) via homologous recombination (see, e.g., U.S. Patent No. 5,641,670, issued June 24, 1997; International Publication No. WO 96/29411, published September 26, 1996;
15 International Publication No. WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra et al., *Nature* 342:435-438 (1989). This method involves the activation of a gene which is present in the target cells, but which is not normally expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made, using standard techniques known in the art,
20 which contain the promoter with targeting sequences flanking the promoter. Suitable promoters are described herein. The targeting sequence is sufficiently complementary to an endogenous sequence to permit homologous recombination of the promoter-targeting sequence with the endogenous sequence. The targeting sequence will be sufficiently near the 5' end of the desired endogenous polynucleotide sequence so the promoter will be operably
25 linked to the endogenous sequence upon homologous recombination.

The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence
30 contains the same restriction site as the 3' end of the amplified promoter. The amplified promoter and targeting sequences are digested and ligated together.

The promoter-targeting sequence construct is delivered to the cells, either as naked polynucleotide, or in conjunction with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, whole viruses, lipofection, precipitating agents, etc., described in more detail above. The P promoter-targeting sequence can be delivered by any method, 5 included direct needle injection, intravenous injection, topical administration, catheter infusion, particle accelerators, etc. The methods are described in more detail below.

The promoter-targeting sequence construct is taken up by cells. Homologous recombination between the construct and the endogenous sequence takes place, such that an endogenous sequence is placed under the control of the promoter. The promoter then drives 10 the expression of the endogenous sequence.

Preferably, the polynucleotide encoding a polypeptide of the present invention contains a secretory signal sequence that facilitates secretion of the protein. Typically, the signal sequence is positioned in the coding region of the polynucleotide to be expressed 15 towards or at the 5' end of the coding region. The signal sequence may be homologous or heterologous to the polynucleotide of interest and may be homologous or heterologous to the cells to be transfected. Additionally, the signal sequence may be chemically synthesized using methods known in the art.

Any mode of administration of any of the above-described polynucleotides constructs can be used so long as the mode results in the expression of one or more molecules in an 20 amount sufficient to provide a therapeutic effect. This includes direct needle injection, systemic injection, catheter infusion, biolistic injectors, particle accelerators (i.e., "gene guns"), gel foam sponge depots, other commercially available depot materials, osmotic pumps (e.g., Alza minipumps), oral or suppository solid (tablet or pill) pharmaceutical formulations, and decanting or topical applications during surgery. For example, direct 25 injection of naked calcium phosphate-precipitated plasmid into rat liver and rat spleen or a protein-coated plasmid into the portal vein has resulted in gene expression of the foreign gene in the rat livers (Kaneda et al., Science 243:375 (1989)).

A preferred method of local administration is by direct injection. Preferably, a recombinant molecule of the present invention complexed with a delivery vehicle is 30 administered by direct injection into or locally within the area of arteries. Administration of a composition locally within the area of arteries refers to injecting the composition centimeters and preferably, millimeters within arteries.

Another method of local administration is to contact a polynucleotide construct of the present invention in or around a surgical wound. For example, a patient can undergo surgery and the polynucleotide construct can be coated on the surface of tissue inside the wound or the construct can be injected into areas of tissue inside the wound.

5 Therapeutic compositions useful in systemic administration, include recombinant molecules of the present invention complexed to a targeted delivery vehicle of the present invention. Suitable delivery vehicles for use with systemic administration comprise liposomes comprising ligands for targeting the vehicle to a particular site.

Preferred methods of systemic administration, include intravenous injection, aerosol,
10 oral and percutaneous (topical) delivery. Intravenous injections can be performed using methods standard in the art. Aerosol delivery can also be performed using methods standard in the art (see, for example, Stribling et al., Proc. Natl. Acad. Sci. USA 189:11277-11281, 1992, which is incorporated herein by reference). Oral delivery can be performed by complexing a polynucleotide construct of the present invention to a carrier capable of
15 withstanding degradation by digestive enzymes in the gut of an animal. Examples of such carriers, include plastic capsules or tablets, such as those known in the art. Topical delivery can be performed by mixing a polynucleotide construct of the present invention with a lipophilic reagent (e.g., DMSO) that is capable of passing into the skin.

Determining an effective amount of substance to be delivered can depend upon a
20 number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the animal, the precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and
25 timing of doses will be determined by the attending physician or veterinarian.

Therapeutic compositions of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and pigs, with humans being particularly preferred.

30 Biological Activities

Polynucleotides or polypeptides, or agonists or antagonists of the present invention, can be used in assays to test for one or more biological activities. If these polynucleotides or

polypeptides, or agonists or antagonists of the present invention, do exhibit activity in a particular assay, it is likely that these molecules may be involved in the diseases associated with the biological activity. Thus, the polynucleotides and polypeptides, and agonists or antagonists could be used to treat the associated disease.

5

Immune Activity

A polypeptide or polynucleotide, or agonists or antagonists of the present invention may be useful in treating deficiencies or disorders of the immune system, by activating or inhibiting the proliferation, differentiation, or mobilization (chemotaxis) of immune cells.

10 Immune cells develop through a process called hematopoiesis, producing myeloid (platelets, red blood cells, neutrophils, and macrophages) and lymphoid (B and T lymphocytes) cells from pluripotent stem cells. The etiology of these immune deficiencies or disorders may be genetic, somatic, such as cancer or some autoimmune disorders, acquired (e.g., by chemotherapy or toxins), or infectious. Moreover, polynucleotides or polypeptides, or
15 agonists or antagonists of the present invention can be used as a marker or detector of a particular immune system disease or disorder.

Polynucleotides or polypeptides, or agonists or antagonists of the present invention may be useful in treating or detecting deficiencies or disorders of hematopoietic cells. Polynucleotides or polypeptides, or agonists or antagonists of the present invention could be
20 used to increase differentiation and proliferation of hematopoietic cells, including the pluripotent stem cells, in an effort to treat those disorders associated with a decrease in certain (or many) types hematopoietic cells. Examples of immunologic deficiency syndromes include, but are not limited to: blood protein disorders (e.g. agammaglobulinemia, dysgammaglobulinemia), ataxia telangiectasia, common variable
25 immunodeficiency, Digeorge Syndrome, HIV infection, HTLV-BLV infection, leukocyte adhesion deficiency syndrome, lymphopenia, phagocyte bactericidal dysfunction, severe combined immunodeficiency (SCIDs), Wiskott-Aldrich Disorder, anemia, thrombocytopenia, or hemoglobinuria.

Moreover, polynucleotides or polypeptides, or agonists or antagonists of the present
30 invention could also be used to modulate hemostatic (the stopping of bleeding) or thrombolytic activity (clot formation). For example, by increasing hemostatic or thrombolytic activity, polynucleotides or polypeptides, or agonists or antagonists of the

present invention could be used to treat blood coagulation disorders (e.g., afibrinogenemia, factor deficiencies), blood platelet disorders (e.g. thrombocytopenia), or wounds resulting from trauma, surgery, or other causes. Alternatively, polynucleotides or polypeptides, or agonists or antagonists of the present invention that can decrease hemostatic or thrombolytic activity could be used to inhibit or dissolve clotting. These molecules could be important in the treatment of heart attacks (infarction), strokes, or scarring.

Polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be useful in treating or detecting autoimmune disorders. Many autoimmune disorders result from inappropriate recognition of self as foreign material by immune cells. This inappropriate recognition results in an immune response leading to the destruction of the host tissue. Therefore, the administration of polynucleotides or polypeptides, or agonists or antagonists of the present invention that can inhibit an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing autoimmune disorders.

Examples of autoimmune disorders that can be treated or detected include, but are not limited to: Addison's Disease, hemolytic anemia, antiphospholipid syndrome, rheumatoid arthritis, dermatitis, allergic encephalomyelitis, glomerulonephritis, Goodpasture's Syndrome, Graves' Disease, Multiple Sclerosis, Myasthenia Gravis, Neuritis, Ophthalmia, Bullous Pemphigoid, Pemphigus, Polyendocrinopathies, Purpura, Reiter's Disease, Stiff-Man Syndrome, Autoimmune Thyroiditis, Systemic Lupus Erythematosus, Autoimmune Pulmonary Inflammation, Guillain-Barre Syndrome, insulin dependent diabetes mellitus, and autoimmune inflammatory eye disease.

Similarly, allergic reactions and conditions, such as asthma (particularly allergic asthma) or other respiratory problems, may also be treated by polynucleotides or polypeptides, or agonists or antagonists of the present invention. Moreover, these molecules can be used to treat anaphylaxis, hypersensitivity to an antigenic molecule, or blood group incompatibility.

Polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to treat and/or prevent organ rejection or graft-versus-host disease (GVHD). Organ rejection occurs by host immune cell destruction of the transplanted tissue through an immune response. Similarly, an immune response is also involved in GVHD, but, in this case, the foreign transplanted immune cells destroy the host tissues. The administration of

polynucleotides or polypeptides, or agonists or antagonists of the present invention that inhibits an immune response, particularly the proliferation, differentiation, or chemotaxis of T-cells, may be an effective therapy in preventing organ rejection or GVHD.

Similarly, polynucleotides or polypeptides, or agonists or antagonists of the present invention may also be used to modulate inflammation. For example, polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation and differentiation of cells involved in an inflammatory response. These molecules can be used to treat inflammatory conditions, both chronic and acute conditions, including chronic prostatitis, granulomatous prostatitis and malacoplakia, inflammation associated with infection (e.g., septic shock, sepsis, or systemic inflammatory response syndrome (SIRS)), ischemia-reperfusion injury, endotoxin lethality, arthritis, complement-mediated hyperacute rejection, nephritis, cytokine or chemokine induced lung injury, inflammatory bowel disease, Crohn's disease, or resulting from over production of cytokines (e.g., TNF or IL-1.)

15 **Hyperproliferative Disorders**

Polynucleotides or polypeptides, or agonists or antagonists of the present invention can be used to treat or detect hyperproliferative disorders, including neoplasms. Polynucleotides or polypeptides, or agonists or antagonists of the present invention may inhibit the proliferation of the disorder through direct or indirect interactions. Alternatively, Polynucleotides or polypeptides, or agonists or antagonists of the present invention may proliferate other cells which can inhibit the hyperproliferative disorder.

For example, by increasing an immune response, particularly increasing antigenic qualities of the hyperproliferative disorder or by proliferating, differentiating, or mobilizing T-cells, hyperproliferative disorders can be treated. This immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response. Alternatively, decreasing an immune response may also be a method of treating hyperproliferative disorders, such as a chemotherapeutic agent.

Examples of hyperproliferative disorders that can be treated or detected by Polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to neoplasms located in the: colon, abdomen, bone, breast, digestive system, liver, pancreas, peritoneum, endocrine glands (adrenal, parathyroid, pituitary,

testicles, ovary, thymus, thyroid), eye, head and neck, nervous (central and peripheral), lymphatic system, pelvic, skin, soft tissue, spleen, thoracic, and urogenital.

Similarly, other hyperproliferative disorders can also be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention.

5 Examples of such hyperproliferative disorders include, but are not limited to: hypergammaglobulinemia, lymphoproliferative disorders, paraproteinemias, purpura, sarcoidosis, Sezary Syndrome, Waldenström's Macroglobulinemia, Gaucher's Disease, histiocytosis, and any other hyperproliferative disease, besides neoplasia, located in an organ system listed above.

10 One preferred embodiment utilizes polynucleotides of the present invention to inhibit aberrant cellular division, by gene therapy using the present invention, and/or protein fusions or fragments thereof.

Thus, the present invention provides a method for treating cell proliferative disorders by inserting into an abnormally proliferating cell a polynucleotide of the present invention,
15 wherein said polynucleotide represses said expression.

Another embodiment of the present invention provides a method of treating cell-proliferative disorders in individuals comprising administration of one or more active gene copies of the present invention to an abnormally proliferating cell or cells. In a preferred embodiment, polynucleotides of the present invention is a DNA construct comprising a
20 recombinant expression vector effective in expressing a DNA sequence encoding said polynucleotides. In another preferred embodiment of the present invention, the DNA construct encoding the polynucleotides of the present invention is inserted into cells to be treated utilizing a retrovirus, or more preferably an adenoviral vector (See G J. Nabel, et. al., PNAS 1999 96: 324-326, which is hereby incorporated by reference). In a most preferred
25 embodiment, the viral vector is defective and will not transform non-proliferating cells, only proliferating cells. Moreover, in a preferred embodiment, the polynucleotides of the present invention inserted into proliferating cells either alone, or in combination with or fused to other polynucleotides, can then be modulated via an external stimulus (i.e. magnetic, specific small molecule, chemical, or drug administration, etc.), which acts upon the promoter
30 upstream of said polynucleotides to induce expression of the encoded protein product. As such the beneficial therapeutic affect of the present invention may be expressly modulated

(i.e. to increase, decrease, or inhibit expression of the present invention) based upon said external stimulus.

Polynucleotides of the present invention may be useful in repressing expression of oncogenic genes or antigens. By "repressing expression of the oncogenic genes " is intended
5 the suppression of the transcription of the gene, the degradation of the gene transcript (pre-message RNA), the inhibition of splicing, the destruction of the messenger RNA, the prevention of the post-translational modifications of the protein, the destruction of the protein, or the inhibition of the normal function of the protein.

For local administration to abnormally proliferating cells, polynucleotides of the
10 present invention may be administered by any method known to those of skill in the art including, but not limited to transfection, electroporation, microinjection of cells, or in vehicles such as liposomes, lipofectin, or as naked polynucleotides, or any other method described throughout the specification. The polynucleotide of the present invention may be delivered by known gene delivery systems such as, but not limited to, retroviral vectors
15 (Gilboa, J. Virology 44:845 (1982); Hocke, Nature 320:275 (1986); Wilson, et al., Proc. Natl. Acad. Sci. U.S.A. 85:3014), vaccinia virus system (Chakrabarty et al., Mol. Cell Biol. 5:3403 (1985) or other efficient DNA delivery systems (Yates et al., Nature 313:812 (1985)) known to those skilled in the art. These references are exemplary only and are hereby incorporated by reference. In order to specifically deliver or transfect cells which are abnormally
20 proliferating and spare non-dividing cells, it is preferable to utilize a retrovirus, or adenoviral (as described in the art and elsewhere herein) delivery system known to those of skill in the art. Since host DNA replication is required for retroviral DNA to integrate and the retrovirus will be unable to self replicate due to the lack of the retrovirus genes needed for its life cycle. Utilizing such a retroviral delivery system for polynucleotides of the present invention will
25 target said gene and constructs to abnormally proliferating cells and will spare the non-dividing normal cells.

The polynucleotides of the present invention may be delivered directly to cell proliferative disorder/disease sites in internal organs, body cavities and the like by use of imaging devices used to guide an injecting needle directly to the disease site. The
30 polynucleotides of the present invention may also be administered to disease sites at the time of surgical intervention.

By "cell proliferative disease" is meant any human or animal disease or disorder, affecting any one or any combination of organs, cavities, or body parts, which is characterized by single or multiple local abnormal proliferations of cells, groups of cells, or tissues, whether benign or malignant.

- 5 Any amount of the polynucleotides of the present invention may be administered as long as it has a biologically inhibiting effect on the proliferation of the treated cells. Moreover, it is possible to administer more than one of the polynucleotide of the present invention simultaneously to the same site. By "biologically inhibiting" is meant partial or total growth inhibition as well as decreases in the rate of proliferation or growth of the cells.
- 10 The biologically inhibitory dose may be determined by assessing the effects of the polynucleotides of the present invention on target malignant or abnormally proliferating cell growth in tissue culture, tumor growth in animals and cell cultures, or any other method known to one of ordinary skill in the art.

- The present invention is further directed to antibody-based therapies which involve
- 15 administering of anti-polypeptides and anti-polynucleotide antibodies to a mammalian, preferably human, patient for treating one or more of the described disorders. Methods for producing anti-polypeptides and anti-polynucleotide antibodies polyclonal and monoclonal antibodies are described in detail elsewhere herein. Such antibodies may be provided in pharmaceutically acceptable compositions as known in the art or as described herein.

- 20 A summary of the ways in which the antibodies of the present invention may be used therapeutically includes binding polynucleotides or polypeptides of the present invention locally or systemically in the body or by direct cytotoxicity of the antibody, e.g. as mediated by complement (CDC) or by effector cells (ADCC). Some of these approaches are described in more detail below. Armed with the teachings provided herein, one of ordinary skill in the
- 25 art will know how to use the antibodies of the present invention for diagnostic, monitoring or therapeutic purposes without undue experimentation.

- In particular, the antibodies, fragments and derivatives of the present invention are useful for treating a subject having or developing cell proliferative and/or differentiation disorders as described herein. Such treatment comprises administering a single or multiple
- 30 doses of the antibody, or a fragment, derivative, or a conjugate thereof.

The antibodies of this invention may be advantageously utilized in combination with other monoclonal or chimeric antibodies, or with lymphokines or hematopoietic growth

factors, for example., which serve to increase the number or activity of effector cells which interact with the antibodies.

It is preferred to use high affinity and/or potent in vivo inhibiting and/or neutralizing antibodies against polypeptides or polynucleotides of the present invention. fragments or regions thereof, for both immunoassays directed to and therapy of disorders related to polynucleotides or polypeptides, including fragments thereof, of the present invention. Such antibodies, fragments, or regions, will preferably have an affinity for polynucleotides or polypeptides, including fragments thereof. Preferred binding affinities include those with a dissociation constant or K_d less than $5 \times 10^{-6}M$, $10^{-6}M$, $5 \times 10^{-7}M$, $10^{-7}M$, $5 \times 10^{-8}M$, $10^{-8}M$, $5 \times 10^{-9}M$, $10^{-9}M$, $5 \times 10^{-10}M$, $10^{-10}M$, $5 \times 10^{-11}M$, $10^{-11}M$, $5 \times 10^{-12}M$, $10^{-12}M$, $5 \times 10^{-13}M$, $10^{-13}M$, $5 \times 10^{-14}M$, $10^{-14}M$, $5 \times 10^{-15}M$, and $10^{-15}M$.

Moreover, polypeptides of the present invention are useful in inhibiting the angiogenesis of proliferative cells or tissues, either alone, as a protein fusion, or in combination with other polypeptides directly or indirectly, as described elsewhere herein. In a most preferred embodiment, said anti-angiogenesis effect may be achieved indirectly, for example, through the inhibition of hematopoietic, tumor-specific cells, such as tumor-associated macrophages (See Joseph IB, et al. J Natl Cancer Inst, 90(21):1648-53 (1998), which is hereby incorporated by reference). Antibodies directed to polypeptides or polynucleotides of the present invention may also result in inhibition of angiogenesis directly, or indirectly (See Witte L, et al., Cancer Metastasis Rev. 17(2):155-61 (1998), which is hereby incorporated by reference)).

Polypeptides, including protein fusions, of the present invention, or fragments thereof may be useful in inhibiting proliferative cells or tissues through the induction of apoptosis. Said polypeptides may act either directly, or indirectly to induce apoptosis of proliferative cells and tissues, for example in the activation of a death-domain receptor, such as tumor necrosis factor (TNF) receptor-1, CD95 (Fas/APO-1), TNF-receptor-related apoptosis-mediated protein (TRAMP) and TNF-related apoptosis-inducing ligand (TRAIL) receptor-1 and -2 (See Schulze-Osthoff K, et.al., Eur J Biochem 254(3):439-59 (1998), which is hereby incorporated by reference). Moreover, in another preferred embodiment of the present invention, said polypeptides may induce apoptosis through other mechanisms, such as in the activation of other proteins which will activate apoptosis, or through stimulating the expression of said proteins, either alone or in combination with small molecule drugs or

adjuvants, such as apoptonin, galectins, thioredoxins, antiinflammatory proteins (See for example, Mutat Res 400(1-2):447-55 (1998), Med Hypotheses.50(5):423-33 (1998), Chem Biol Interact. Apr 24;111-112:23-34 (1998), J Mol Med.76(6):402-12 (1998), Int J Tissue React;20(1):3-15 (1998), which are all hereby incorporated by reference).

5 Polypeptides, including protein fusions to, or fragments thereof, of the present invention are useful in inhibiting the metastasis of proliferative cells or tissues. Inhibition may occur as a direct result of administering polypeptides, or antibodies directed to said polypeptides as described elsewhere herein, or indirectly, such as activating the expression of proteins known to inhibit metastasis, for example alpha 4 integrins, (See, e.g., Curr Top
10 Microbiol Immunol 1998;231:125-41, which is hereby incorporated by reference). Such therapeutic affects of the present invention may be achieved either alone, or in combination with small molecule drugs or adjuvants.

In another embodiment, the invention provides a method of delivering compositions containing the polypeptides of the invention (e.g., compositions containing polypeptides or
15 polypeptide antibodies associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs) to targeted cells expressing the polypeptide of the present invention. Polypeptides or polypeptide antibodies of the invention may be associated with with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. Polypeptides, protein fusions to, or
20 fragments thereof, of the present invention are useful in enhancing the immunogenicity and/or antigenicity of proliferating cells or tissues, either directly, such as would occur if the polypeptides of the present invention 'vaccinated' the immune response to respond to proliferative antigens and immunogens, or indirectly, such as in activating the expression of proteins known to enhance the immune response (e.g. chemokines), to said antigens and
25 immunogens.

Cardiovascular Disorders

Polynucleotides or polypeptides, or agonists or antagonists of the present invention, may be used to treat cardiovascular disorders, including peripheral artery disease, such as
30 limb ischemia.

Cardiovascular disorders include cardiovascular abnormalities, such as arterio-arterial fistula, arteriovenous fistula, cerebral arteriovenous malformations, congenital heart defects,

pulmonary atresia, and Scimitar Syndrome. Congenital heart defects include aortic coarctation, cor triatriatum, coronary vessel anomalies, crisscross heart, dextrocardia, patent ductus arteriosus, Ebstein's anomaly, Eisenmenger complex, hypoplastic left heart syndrome, levocardia, tetralogy of fallot, transposition of great vessels, double outlet right ventricle, 5 tricuspid atresia, persistent truncus arteriosus, and heart septal defects, such as aortopulmonary septal defect, endocardial cushion defects, Lutembacher's Syndrome, trilog of Fallot, ventricular heart septal defects.

Cardiovascular disorders also include heart disease, such as arrhythmias, carcinoid heart disease, high cardiac output, low cardiac output, cardiac tamponade, endocarditis 10 (including bacterial), heart aneurysm, cardiac arrest, congestive heart failure, congestive cardiomyopathy, paroxysmal dyspnea, cardiac edema, heart hypertrophy, congestive cardiomyopathy, left ventricular hypertrophy, right ventricular hypertrophy, post-infarction heart rupture, ventricular septal rupture, heart valve diseases, myocardial diseases, myocardial ischemia, pericardial effusion, pericarditis (including constrictive and 15 tuberculous), pneumopericardium, postpericardiotomy syndrome, pulmonary heart disease, rheumatic heart disease, ventricular dysfunction, hyperemia, cardiovascular pregnancy complications, Scimitar Syndrome, cardiovascular syphilis, and cardiovascular tuberculosis.

Arrhythmias include sinus arrhythmia, atrial fibrillation, atrial flutter, bradycardia, extrasystole, Adams-Stokes Syndrome, bundle-branch block, sinoatrial block, long QT 20 syndrome, parasystole, Lown-Ganong-Levine Syndrome, Mahaim-type pre-excitation syndrome, Wolff-Parkinson-White syndrome, sick sinus syndrome, tachycardias, and ventricular fibrillation. Tachycardias include paroxysmal tachycardia, supraventricular tachycardia, accelerated idioventricular rhythm, atrioventricular nodal reentry tachycardia, ectopic atrial tachycardia, ectopic junctional tachycardia, sinoatrial nodal reentry tachycardia, 25 sinus tachycardia, Torsades de Pointes, and ventricular tachycardia.

Heart valve disease include aortic valve insufficiency, aortic valve stenosis, hear murmurs, aortic valve prolapse, mitral valve prolapse, tricuspid valve prolapse, mitral valve insufficiency, mitral valve stenosis, pulmonary atresia, pulmonary valve insufficiency, pulmonary valve stenosis, tricuspid atresia, tricuspid valve insufficiency, and tricuspid valve 30 stenosis.

Myocardial diseases include alcoholic cardiomyopathy, congestive cardiomyopathy, hypertrophic cardiomyopathy, aortic subvalvular stenosis, pulmonary subvalvular stenosis,

restrictive cardiomyopathy, Chagas cardiomyopathy, endocardial fibroelastosis, endomyocardial fibrosis, Kearns Syndrome, myocardial reperfusion injury, and myocarditis.

Myocardial ischemias include coronary disease, such as angina pectoris, coronary aneurysm, coronary arteriosclerosis, coronary thrombosis, coronary vasospasm, myocardial infarction and myocardial stunning.

Cardiovascular diseases also include vascular diseases such as aneurysms, angiodyplasia, angiomas, bacillary angiomas, Hippiel-Lindau Disease, Klippel-Trenaunay-Weber Syndrome, Sturge-Weber Syndrome, angioneurotic edema, aortic diseases, Takayasu's Arteritis, aortitis, Leriche's Syndrome, arterial occlusive diseases, arteritis, 10 enarteritis, polyarteritis nodosa, cerebrovascular disorders, diabetic angiopathies, diabetic retinopathy, embolisms, thrombosis, erythromelalgia, hemorrhoids, hepatic veno-occlusive disease, hypertension, hypotension, ischemia, peripheral vascular diseases, phlebitis, pulmonary veno-occlusive disease, Raynaud's disease, CREST syndrome, retinal vein occlusion, Scimitar syndrome, superior vena cava syndrome, telangiectasia, atacia 15 telangiectasia, hereditary hemorrhagic telangiectasia, varicocele, varicose veins, varicose ulcer, vasculitis, and venous insufficiency.

Aneurysms include dissecting aneurysms, false aneurysms, infected aneurysms, ruptured aneurysms, aortic aneurysms, cerebral aneurysms, coronary aneurysms, heart aneurysms, and iliac aneurysms.

Arterial occlusive diseases include arteriosclerosis, intermittent claudication, carotid stenosis, fibromuscular dysplasias, mesenteric vascular occlusion, Moyamoya disease, renal artery obstruction, retinal artery occlusion, and thromboangiitis obliterans.

Cerebrovascular disorders include carotid artery diseases, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformation, cerebral artery diseases, cerebral embolism and thrombosis, 25 carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, cerebral hemorrhage, epidural hematoma, subdural hematoma, subarachnoid hemorrhage, cerebral infarction, cerebral ischemia (including transient), subclavian steal syndrome, periventricular leukomalacia, vascular headache, cluster headache, migraine, and vertebrobasilar 30 insufficiency.

Embolisms include air embolisms, amniotic fluid embolisms, cholesterol embolisms, blue toe syndrome, fat embolisms, pulmonary embolisms, and thromboembolisms.

Thrombosis include coronary thrombosis, hepatic vein thrombosis, retinal vein occlusion, carotid artery thrombosis, sinus thrombosis, Wallenberg's syndrome, and thrombophlebitis.

Ischemia includes cerebral ischemia, ischemic colitis, compartment syndromes, anterior compartment syndrome, myocardial ischemia, reperfusion injuries, and peripheral limb ischemia. Vasculitis includes aortitis, arteritis, Behcet's Syndrome, Churg-Strauss Syndrome, mucocutaneous lymph node syndrome, thromboangiitis obliterans, hypersensitivity vasculitis, Schoenlein-Henoch purpura, allergic cutaneous vasculitis, and Wegener's granulomatosis.

Polynucleotides or polypeptides, or agonists or antagonists of the present invention, are especially effective for the treatment of critical limb ischemia and coronary disease.

Polypeptides may be administered using any method known in the art, including, but not limited to, direct needle injection at the delivery site, intravenous injection, topical administration, catheter infusion, biolistic injectors, particle accelerators, gelfoam sponge depots, other commercially available depot materials, osmotic pumps, oral or suppository solid pharmaceutical formulations, decanting or topical applications during surgery, aerosol delivery. Such methods are known in the art. Polypeptides may be administered as part of a Therapeutic, described in more detail below. Methods of delivering polynucleotides are described in more detail herein.

Anti-Angiogenesis Activity

The naturally occurring balance between endogenous stimulators and inhibitors of angiogenesis is one in which inhibitory influences predominate. Rastinejad *et al.*, *Cell* 56:345-355 (1989). In those rare instances in which neovascularization occurs under normal physiological conditions, such as wound healing, organ regeneration, embryonic development, and female reproductive processes, angiogenesis is stringently regulated and spatially and temporally delimited. Under conditions of pathological angiogenesis such as that characterizing solid tumor growth, these regulatory controls fail. Unregulated angiogenesis becomes pathologic and sustains progression of many neoplastic and non-neoplastic diseases. A number of serious diseases are dominated by abnormal neovascularization including solid tumor growth and metastases, arthritis, some types of eye disorders, and psoriasis. See, e.g., reviews by Moses *et al.*, *Biotech.* 9:630-634 (1991); Folkman *et al.*, *N. Engl. J. Med.*, 333:1757-1763 (1995); Auerbach *et al.*, *J. Microvasc. Res.*

29:401-411 (1985); Folkman, *Advances in Cancer Research*, eds. Klein and Weinhouse, Academic Press, New York, pp. 175-203 (1985); Patz, *Am. J. Ophthalmol.* 94:715-743 (1982); and Folkman *et al.*, *Science* 221:719-725 (1983). In a number of pathological conditions, the process of angiogenesis contributes to the disease state. For example, 5 significant data have accumulated which suggest that the growth of solid tumors is dependent on angiogenesis. Folkman and Klagsbrun, *Science* 235:442-447 (1987).

The polynucleotides encoding a polypeptide of the present invention may be administered along with other polynucleotides encoding an angiogenic protein. Examples of angiogenic proteins include, but are not limited to, acidic and basic fibroblast growth factors. 10 VEGF-1, VEGF-2, VEGF-3, epidermal growth factor alpha and beta, platelet-derived endothelial cell growth factor, platelet-derived growth factor, tumor necrosis factor alpha, hepatocyte growth factor, insulin like growth factor, colony stimulating factor, macrophage colony stimulating factor, granulocyte/macrophage colony stimulating factor, and nitric oxide synthase.

15 The present invention provides for treatment of diseases or disorders associated with neovascularization by administration of the polynucleotides and/or polypeptides of the invention, as well as agonists or antagonists of the present invention. Malignant and metastatic conditions which can be treated with the polynucleotides and polypeptides, or agonists or antagonists of the invention include, but are not limited to, malignancies, solid 20 tumors, and cancers described herein and otherwise known in the art (for a review of such disorders, see Fishman *et al.*, *Medicine*, 2d Ed., J. B. Lippincott Co., Philadelphia (1985)). Thus, the present invention provides a method of treating an angiogenesis-related disease and/or disorder, comprising administering to an individual in need thereof a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist 25 of the invention. For example, polynucleotides, polypeptides, antagonists and/or agonists may be utilized in a variety of additional methods in order to therapeutically treat a cancer or tumor. Cancers which may be treated with polynucleotides, polypeptides, antagonists and/or agonists include, but are not limited to solid tumors, including colon, rectum, prostate, lung, breast, ovarian, stomach, pancreas, larynx, esophagus, testes, liver, parotid, biliary tract, 30 cervix, uterus, endometrium, kidney, bladder, thyroid cancer; primary tumors and metastases; melanomas; glioblastoma; Kaposi's sarcoma; leiomyosarcoma; non-small cell lung cancer; colorectal cancer; advanced malignancies; and blood born tumors such as leukemias. For

example, polynucleotides, polypeptides, antagonists and/or agonists may be delivered topically, in order to treat cancers such as skin cancer, head and neck tumors, breast tumors, and Kaposi's sarcoma.

Within yet other aspects, polynucleotides, polypeptides, antagonists and/or agonists
5 may be utilized to treat superficial forms of bladder cancer by, for example, intravesical administration. Polynucleotides, polypeptides, antagonists and/or agonists may be delivered directly into the tumor, or near the tumor site, via injection or a catheter. Of course, as the artisan of ordinary skill will appreciate, the appropriate mode of administration will vary according to the cancer to be treated. Other modes of delivery are discussed herein.

10 Polynucleotides, polypeptides, antagonists and/or agonists may be useful in treating other disorders, besides cancers, which involve angiogenesis. These disorders include, but are not limited to: benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; arteriosclerotic plaques; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular
15 degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures; scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous
20 malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis.

For example, within one aspect of the present invention methods are provided for treating hypertrophic scars and keloids, comprising the step of administering a
25 polynucleotide, polypeptide, antagonist and/or agonist of the invention to a hypertrophic scar or keloid.

Within one embodiment of the present invention polynucleotides, polypeptides, antagonists and/or agonists are directly injected into a hypertrophic scar or keloid, in order to prevent the progression of these lesions. This therapy is of particular value in the
30 prophylactic treatment of conditions which are known to result in the development of hypertrophic scars and keloids (e.g., burns), and is preferably initiated after the proliferative phase has had time to progress (approximately 14 days after the initial injury), but before

hypertrophic scar or keloid development. As noted above, the present invention also provides methods for treating neovascular diseases of the eye, including for example, corneal neovascularization, neovascular glaucoma, proliferative diabetic retinopathy, retrolental fibroplasia and macular degeneration.

5 Moreover, Ocular disorders associated with neovascularization which can be treated with the polynucleotides and polypeptides of the present invention (including agonists and/or antagonists) include, but are not limited to: neovascular glaucoma, diabetic retinopathy, retinoblastoma, retrolental fibroplasia, uveitis, retinopathy of prematurity macular degeneration, corneal graft neovascularization, as well as other eye inflammatory diseases,
10 ocular tumors and diseases associated with choroidal or iris neovascularization. See, e.g., reviews by Waltman *et al.*, *Am. J. Ophthalmol.* 85:704-710 (1978) and Gartner *et al.*, *Surv. Ophthalmol.* 22:291-312 (1978).

 Thus, within one aspect of the present invention methods are provided for treating neovascular diseases of the eye such as corneal neovascularization (including corneal graft
15 neovascularization), comprising the step of administering to a patient a therapeutically effective amount of a compound (as described above) to the cornea, such that the formation of blood vessels is inhibited. Briefly, the cornea is a tissue which normally lacks blood vessels. In certain pathological conditions however, capillaries may extend into the cornea from the pericorneal vascular plexus of the limbus. When the cornea becomes vascularized,
20 it also becomes clouded, resulting in a decline in the patient's visual acuity. Visual loss may become complete if the cornea completely opacitates. A wide variety of disorders can result in corneal neovascularization, including for example, corneal infections (e.g., trachoma, herpes simplex keratitis, leishmaniasis and onchocerciasis), immunological processes (e.g., graft rejection and Stevens-Johnson's syndrome), alkali burns, trauma, inflammation (of any
25 cause), toxic and nutritional deficiency states, and as a complication of wearing contact lenses.

 Within particularly preferred embodiments of the invention, may be prepared for topical administration in saline (combined with any of the preservatives and antimicrobial agents commonly used in ocular preparations), and administered in eyedrop form. The
30 solution or suspension may be prepared in its pure form and administered several times daily. Alternatively, anti-angiogenic compositions, prepared as described above, may also be administered directly to the cornea. Within preferred embodiments, the anti-angiogenic

composition is prepared with a muco-adhesive polymer which binds to cornea. Within further embodiments, the anti-angiogenic factors or anti-angiogenic compositions may be utilized as an adjunct to conventional steroid therapy. Topical therapy may also be useful prophylactically in corneal lesions which are known to have a high probability of inducing an angiogenic response (such as chemical burns). In these instances the treatment, likely in combination with steroids, may be instituted immediately to help prevent subsequent complications.

Within other embodiments, the compounds described above may be injected directly into the corneal stroma by an ophthalmologist under microscopic guidance. The preferred site of injection may vary with the morphology of the individual lesion, but the goal of the administration would be to place the composition at the advancing front of the vasculature (i.e., interspersed between the blood vessels and the normal cornea). In most cases this would involve perilimbic corneal injection to "protect" the cornea from the advancing blood vessels. This method may also be utilized shortly after a corneal insult in order to prophylactically prevent corneal neovascularization. In this situation the material could be injected in the perilimbic cornea interspersed between the corneal lesion and its undesired potential limbic blood supply. Such methods may also be utilized in a similar fashion to prevent capillary invasion of transplanted corneas. In a sustained-release form injections might only be required 2-3 times per year. A steroid could also be added to the injection solution to reduce inflammation resulting from the injection itself.

Within another aspect of the present invention, methods are provided for treating neovascular glaucoma, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. In one embodiment, the compound may be administered topically to the eye in order to treat early forms of neovascular glaucoma. Within other embodiments, the compound may be implanted by injection into the region of the anterior chamber angle. Within other embodiments, the compound may also be placed in any location such that the compound is continuously released into the aqueous humor. Within another aspect of the present invention, methods are provided for treating proliferative diabetic retinopathy, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eyes, such that the formation of blood vessels is inhibited.

Within particularly preferred embodiments of the invention, proliferative diabetic retinopathy may be treated by injection into the aqueous humor or the vitreous, in order to increase the local concentration of the polynucleotide, polypeptide, antagonist and/or agonist in the retina. Preferably, this treatment should be initiated prior to the acquisition of severe disease requiring photocoagulation.

Within another aspect of the present invention, methods are provided for treating retrolental fibroplasia, comprising the step of administering to a patient a therapeutically effective amount of a polynucleotide, polypeptide, antagonist and/or agonist to the eye, such that the formation of blood vessels is inhibited. The compound may be administered topically, via intravitreal injection and/or via intraocular implants.

Additionally, disorders which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, hemangioma, arthritis, psoriasis, angiofibroma, atherosclerotic plaques, delayed wound healing, granulations, hemophilic joints, hypertrophic scars, nonunion fractures, Osler-Weber syndrome, pyogenic granuloma, scleroderma, trachoma, and vascular adhesions.

Moreover, disorders and/or states, which can be treated with the polynucleotides, polypeptides, agonists and/or antagonists include, but are not limited to, solid tumors, blood born tumors such as leukemias, tumor metastasis, Kaposi's sarcoma, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas, rheumatoid arthritis, psoriasis, ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, and uveitis, delayed wound healing, endometriosis, vasculogenesis, granulations, hypertrophic scars (keloids), nonunion fractures, scleroderma, trachoma, vascular adhesions, myocardial angiogenesis, coronary collaterals, cerebral collaterals, arteriovenous malformations, ischemic limb angiogenesis, Osler-Webber Syndrome, plaque neovascularization, telangiectasia, hemophilic joints, angiofibroma fibromuscular dysplasia, wound granulation, Crohn's disease, atherosclerosis, birth control agent by preventing vascularization required for embryo implantation controlling menstruation, diseases that have angiogenesis as a pathologic consequence such as cat scratch disease (Rochelie minalia quintosa), ulcers (*Helicobacter pylori*), Bartonellosis and bacillary angiomatosis.

In one aspect of the birth control method, an amount of the compound sufficient to block embryo implantation is administered before or after intercourse and fertilization have occurred, thus providing an effective method of birth control, possibly a "morning after" method. Polynucleotides, polypeptides, agonists and/or agonists may also be used in
5 controlling menstruation or administered as either a peritoneal lavage fluid or for peritoneal implantation in the treatment of endometriosis.

Polynucleotides, polypeptides, agonists and/or agonists of the present invention may be incorporated into surgical sutures in order to prevent stitch granulomas.

Polynucleotides, polypeptides, agonists and/or agonists may be utilized in a wide
10 variety of surgical procedures. For example, within one aspect of the present invention a compositions (in the form of, for example, a spray or film) may be utilized to coat or spray an area prior to removal of a tumor, in order to isolate normal surrounding tissues from malignant tissue, and/or to prevent the spread of disease to surrounding tissues. Within other aspects of the present invention, compositions (e.g., in the form of a spray) may be delivered
15 via endoscopic procedures in order to coat tumors, or inhibit angiogenesis in a desired locale. Within yet other aspects of the present invention, surgical meshes which have been coated with anti- angiogenic compositions of the present invention may be utilized in any procedure wherein a surgical mesh might be utilized. For example, within one embodiment of the invention a surgical mesh laden with an anti-angiogenic composition may be utilized during
20 abdominal cancer resection surgery (e.g., subsequent to colon resection) in order to provide support to the structure, and to release an amount of the anti-angiogenic factor.

Within further aspects of the present invention, methods are provided for treating tumor excision sites, comprising administering a polynucleotide, polypeptide, agonist and/or agonist to the resection margins of a tumor subsequent to excision, such that the local
25 recurrence of cancer and the formation of new blood vessels at the site is inhibited. Within one embodiment of the invention, the anti-angiogenic compound is administered directly to the tumor excision site (e.g., applied by swabbing, brushing or otherwise coating the resection margins of the tumor with the anti-angiogenic compound). Alternatively, the anti-angiogenic compounds may be incorporated into known surgical pastes prior to
30 administration. Within particularly preferred embodiments of the invention, the anti-angiogenic compounds are applied after hepatic resections for malignancy, and after neurosurgical operations.

Within one aspect of the present invention, polynucleotides, polypeptides, agonists and/or agonists may be administered to the resection margin of a wide variety of tumors, including for example, breast, colon, brain and hepatic tumors. For example, within one embodiment of the invention, anti-angiogenic compounds may be administered to the site of a neurological tumor subsequent to excision, such that the formation of new blood vessels at the site are inhibited.

The polynucleotides, polypeptides, agonists and/or agonists of the present invention may also be administered along with other anti-angiogenic factors. Representative examples of other anti-angiogenic factors include: Anti-Invasive Factor, retinoic acid and derivatives thereof, paclitaxel, Suramin, Tissue Inhibitor of Metalloproteinase-1, Tissue Inhibitor of Metalloproteinase-2, Plasminogen Activator Inhibitor-1, Plasminogen Activator Inhibitor-2, and various forms of the lighter "d group" transition metals.

Lighter "d group" transition metals include, for example, vanadium, molybdenum, tungsten, titanium, niobium, and tantalum species. Such transition metal species may form transition metal complexes. Suitable complexes of the above-mentioned transition metal species include oxo transition metal complexes.

Representative examples of vanadium complexes include oxo vanadium complexes such as vanadate and vanadyl complexes. Suitable vanadate complexes include metavanadate and orthovanadate complexes such as, for example, ammonium metavanadate, sodium metavanadate, and sodium orthovanadate. Suitable vanadyl complexes include, for example, vanadyl acetylacetonate and vanadyl sulfate including vanadyl sulfate hydrates such as vanadyl sulfate mono- and trihydrates.

Representative examples of tungsten and molybdenum complexes also include oxo complexes. Suitable oxo tungsten complexes include tungstate and tungsten oxide complexes. Suitable tungstate complexes include ammonium tungstate, calcium tungstate, sodium tungstate dihydrate, and tungstic acid. Suitable tungsten oxides include tungsten (IV) oxide and tungsten (VI) oxide. Suitable oxo molybdenum complexes include molybdate, molybdenum oxide, and molybdenyl complexes. Suitable molybdate complexes include ammonium molybdate and its hydrates, sodium molybdate and its hydrates, and potassium molybdate and its hydrates. Suitable molybdenum oxides include molybdenum (VI) oxide, molybdenum (VI) oxide, and molybdic acid. Suitable molybdenyl complexes include, for

example. molybdenyl acetylacetonate. Other suitable tungsten and molybdenum complexes include hydroxo derivatives derived from, for example, glycerol, tartaric acid, and sugars.

A wide variety of other anti-angiogenic factors may also be utilized within the context of the present invention. Representative examples include platelet factor 4; protamine sulphate; sulphated chitin derivatives (prepared from queen crab shells), (Murata et al., Cancer Res. 51:22-26, 1991); Sulphated Polysaccharide Peptidoglycan Complex (SP- PG) (the function of this compound may be enhanced by the presence of steroids such as estrogen, and tamoxifen citrate); Staurosporine; modulators of matrix metabolism, including for example, proline analogs, cishydroxyproline, d,L-3,4-dehydroproline, Thiaproline, 10 alpha,alpha-dipyridyl, aminopropionitrile fumarate; 4-propyl-5-(4-pyridinyl)-2(3H)-oxazolone; Methotrexate; Mitoxantrone; Heparin; Interferons; 2 Macroglobulin-serum; ChIMP-3 (Pavloff et al., J. Bio. Chem. 267:17321-17326, 1992); Chymostatin (Tomkinson et al., Biochem J. 286:475-480, 1992); Cyclodextrin Tetradecasulfate; Eponemycin; Camptothecin; Fumagillin (Ingber et al., Nature 348:555-557, 1990); Gold Sodium 15 Thiomaleate ("GST"; Matsubara and Ziff, J. Clin. Invest. 79:1440-1446, 1987); anticollagenase-serum; alpha2-antiplasmin (Holmes et al., J. Biol. Chem. 262(4):1659-1664, 1987); Bisantrene (National Cancer Institute); Lobenzarit disodium (N-(2)-carboxyphenyl-4-chloroanthronilic acid disodium or "CCA"; Takeuchi et al., Agents Actions 36:312-316, 1992); Thalidomide; Angostatic steroid; AGM-1470; carboxynaminolimidazole; and 20 metalloproteinase inhibitors such as BB94.

Diseases at the Cellular Level

Diseases associated with increased cell survival or the inhibition of apoptosis that could be treated or detected by polynucleotides or polypeptides, as well as antagonists or 25 agonists of the present invention, include cancers (such as follicular lymphomas, carcinomas with p53 mutations, and hormone-dependent tumors, including, but not limited to colon cancer, cardiac tumors, pancreatic cancer, melanoma, retinoblastoma, glioblastoma, lung cancer, intestinal cancer, testicular cancer, stomach cancer, neuroblastoma, myxoma, myoma, lymphoma, endothelioma, osteoblastoma, osteoclastoma, osteosarcoma, chondrosarcoma, adenoma, breast cancer, prostate cancer, Kaposi's sarcoma and ovarian cancer); autoimmune 30 disorders (such as, multiple sclerosis, Sjogren's syndrome, Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and

immune-related glomerulonephritis and rheumatoid arthritis) and viral infections (such as herpes viruses, pox viruses and adenoviruses), inflammation, graft v. host disease, acute graft rejection, and chronic graft rejection. In preferred embodiments, polynucleotides, polypeptides, and/or antagonists of the invention are used to inhibit growth, progression, and/or metasis of cancers, in particular those listed above.

Additional diseases or conditions associated with increased cell survival that could be treated or detected by polynucleotides or polypeptides, or agonists or antagonists of the present invention include, but are not limited to, progression, and/or metastases of malignancies and related disorders such as leukemia (including acute leukemias (e.g., acute lymphocytic leukemia, acute myelocytic leukemia (including myeloblastic, promyelocytic, myelomonocytic, monocytic, and erythroleukemia)) and chronic leukemias (e.g., chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia)), polycythemia vera, lymphomas (e.g., Hodgkin's disease and non-Hodgkin's disease), multiple myeloma, Waldenstrom's macroglobulinemia, heavy chain disease, and solid tumors including, but not limited to, sarcomas and carcinomas such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilm's tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, menangioma, melanoma, neuroblastoma, and retinoblastoma.

Diseases associated with increased apoptosis that could be treated or detected by polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, include AIDS; neurodegenerative disorders (such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, Retinitis pigmentosa, Cerebellar degeneration and brain tumor or prior associated disease); autoimmune disorders (such as, multiple sclerosis, Sjogren's

syndrome. Hashimoto's thyroiditis, biliary cirrhosis, Behcet's disease, Crohn's disease, polymyositis, systemic lupus erythematosus and immune-related glomerulonephritis and rheumatoid arthritis) myelodysplastic syndromes (such as aplastic anemia), graft v. host disease, ischemic injury (such as that caused by myocardial infarction, stroke and reperfusion injury), liver injury (e.g., hepatitis related liver injury, ischemia/reperfusion injury, cholestosis (bile duct injury) and liver cancer); toxin-induced liver disease (such as that caused by alcohol), septic shock, cachexia and anorexia.

Wound Healing and Epithelial Cell Proliferation

10 In accordance with yet a further aspect of the present invention, there is provided a process for utilizing polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, for therapeutic purposes, for example, to stimulate epithelial cell proliferation and basal keratinocytes for the purpose of wound healing, and to stimulate hair follicle production and healing of dermal wounds. Polynucleotides or polypeptides, as well as
15 agonists or antagonists of the present invention, may be clinically useful in stimulating wound healing including surgical wounds, excisional wounds, deep wounds involving damage of the dermis and epidermis, eye tissue wounds, dental tissue wounds, oral cavity wounds, diabetic ulcers, dermal ulcers, cubitus ulcers, arterial ulcers, venous stasis ulcers, burns resulting from heat exposure or chemicals, and other abnormal wound healing
20 conditions such as uremia, malnutrition, vitamin deficiencies and complications associated with systemic treatment with steroids, radiation therapy and antineoplastic drugs and antimetabolites. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to promote dermal reestablishment subsequent to dermal loss

25 Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to increase the adherence of skin grafts to a wound bed and to stimulate re-epithelialization from the wound bed. The following are types of grafts that polynucleotides or polypeptides, agonists or antagonists of the present invention, could be used to increase adherence to a wound bed: autografts, artificial skin, allografts, autodermic
30 graft, autoepidermic grafts, avascular grafts, Blair-Brown grafts, bone graft, brephoplastic grafts, cutis graft, delayed graft, dermic graft, epidermic graft, fascia graft, full thickness graft, heterologous graft, xenograft, homologous graft, hyperplastic graft, lamellar graft,

mesh graft, mucosal graft, Ollier-Thiersch graft, omentum graft, patch graft, pedicle graft, penetrating graft, split skin graft, thick split graft. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, can be used to promote skin strength and to improve the appearance of aged skin.

5 It is believed that polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, will also produce changes in hepatocyte proliferation, and epithelial cell proliferation in the lung, breast, pancreas, stomach, small intestine, and large intestine. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could promote proliferation of epithelial cells such as sebocytes, hair follicles, hepatocytes,
10 type II pneumocytes, mucin-producing goblet cells, and other epithelial cells and their progenitors contained within the skin, lung, liver, and gastrointestinal tract. Polynucleotides or polypeptides, agonists or antagonists of the present invention, may promote proliferation of endothelial cells, keratinocytes, and basal keratinocytes.

Polynucleotides or polypeptides, as well as agonists or antagonists of the present
15 invention, could also be used to reduce the side effects of gut toxicity that result from radiation, chemotherapy treatments or viral infections. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may have a cytoprotective effect on the small intestine mucosa. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, may also stimulate healing of mucositis (mouth ulcers)
20 that result from chemotherapy and viral infections.

Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could further be used in full regeneration of skin in full and partial thickness skin defects, including burns, (i.e., repopulation of hair follicles, sweat glands, and sebaceous glands), treatment of other skin defects such as psoriasis. Polynucleotides or polypeptides, as
25 well as agonists or antagonists of the present invention, could be used to treat epidermolysis bullosa, a defect in adherence of the epidermis to the underlying dermis which results in frequent, open and painful blisters by accelerating reepithelialization of these lesions. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could also be used to treat gastric and duodenal ulcers and help heal by scar formation of the
30 mucosal lining and regeneration of glandular mucosa and duodenal mucosal lining more rapidly. Inflammatory bowel diseases, such as Crohn's disease and ulcerative colitis, are diseases which result in destruction of the mucosal surface of the small or large intestine,

respectively. Thus, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to promote the resurfacing of the mucosal surface to aid more rapid healing and to prevent progression of inflammatory bowel disease. Treatment with polynucleotides or polypeptides, agonists or antagonists of the present invention, is expected to have a significant effect on the production of mucus throughout the gastrointestinal tract and could be used to protect the intestinal mucosa from injurious substances that are ingested or following surgery. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat diseases associated with the under expression.

Moreover, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to prevent and heal damage to the lungs due to various pathological states. Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, which could stimulate proliferation and differentiation and promote the repair of alveoli and bronchiolar epithelium to prevent or treat acute or chronic lung damage. For example, emphysema, which results in the progressive loss of alveoli, and inhalation injuries, i.e., resulting from smoke inhalation and burns, that cause necrosis of the bronchiolar epithelium and alveoli could be effectively treated using polynucleotides or polypeptides, agonists or antagonists of the present invention. Also, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to stimulate the proliferation of and differentiation of type II pneumocytes, which may help treat or prevent disease such as hyaline membrane diseases, such as infant respiratory distress syndrome and bronchopulmonary dysplasia, in premature infants.

Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could stimulate the proliferation and differentiation of hepatocytes and, thus, could be used to alleviate or treat liver diseases and pathologies such as fulminant liver failure caused by cirrhosis, liver damage caused by viral hepatitis and toxic substances (i.e., acetaminophen, carbon tetrachloride and other hepatotoxins known in the art).

In addition, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to treat or prevent the onset of diabetes mellitus. In patients with newly diagnosed Types I and II diabetes, where some islet cell function remains, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used to maintain the islet function so as to alleviate, delay or prevent permanent

manifestation of the disease. Also, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, could be used as an auxiliary in islet cell transplantation to improve or promote islet cell function.

5 Neurological Diseases

In accordance with yet a further aspect of the present invention, there is provided a process for utilizing polynucleotides or polypeptides, as well as agonists or antagonists of the present invention, for therapeutic purposes, for example, to stimulate neurological cell
10 proliferation and/or differentiation. Therefore, polynucleotides, polypeptides, agonists and/or antagonists of the invention may be used to treat and/or detect neurologic diseases. Moreover, polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used as a marker or detector of a particular nervous system disease or disorder.

Examples of neurologic diseases which can be treated or detected with
15 polynucleotides, polypeptides, agonists, and/or antagonists of the present invention include brain diseases, such as metabolic brain diseases which includes phenylketonuria such as maternal phenylketonuria, pyruvate carboxylase deficiency, pyruvate dehydrogenase complex deficiency, Wernicke's Encephalopathy, brain edema, brain neoplasms such as cerebellar neoplasms which include infratentorial neoplasms, cerebral ventricle neoplasms
20 such as choroid plexus neoplasms, hypothalamic neoplasms, supratentorial neoplasms, canavan disease, cerebellar diseases such as cerebellar ataxia which include spinocerebellar degeneration such as ataxia telangiectasia, cerebellar dyssynergia, Friederich's Ataxia, Machado-Joseph Disease, olivopontocerebellar atrophy, cerebellar neoplasms such as infratentorial neoplasms, diffuse cerebral sclerosis such as encephalitis periaxialis, globoid
25 cell leukodystrophy, metachromatic leukodystrophy and subacute sclerosing panencephalitis, cerebrovascular disorders (such as carotid artery diseases which include carotid artery thrombosis, carotid stenosis and Moyamoya Disease, cerebral amyloid angiopathy, cerebral aneurysm, cerebral anoxia, cerebral arteriosclerosis, cerebral arteriovenous malformations, cerebral artery diseases, cerebral embolism and thrombosis such as carotid artery thrombosis,
30 sinus thrombosis and Wallenberg's Syndrome, cerebral hemorrhage such as epidural hematoma, subdural hematoma and subarachnoid hemorrhage, cerebral infarction, cerebral ischemia such as transient cerebral ischemia, Subclavian Steal Syndrome and vertebrobasilar insufficiency, vascular dementia such as multi-infarct dementia, periventricular leukomalacia,

vascular headache such as cluster headache, migraine, dementia such as AIDS Dementia Complex, presenile dementia such as Alzheimer's Disease and Creutzfeldt-Jakob Syndrome, senile dementia such as Alzheimer's Disease and progressive supranuclear palsy, vascular dementia such as multi-infarct dementia, encephalitis which include encephalitis periaxialis, 5 viral encephalitis such as epidemic encephalitis, Japanese Encephalitis, St. Louis Encephalitis, tick-borne encephalitis and West Nile Fever, acute disseminated encephalomyelitis, meningoencephalitis such as uveomeningoencephalitic syndrome, Postencephalitic Parkinson Disease and subacute sclerosing panencephalitis, encephalomalacia such as periventricular leukomalacia, epilepsy such as generalized epilepsy 10 which includes infantile spasms, absence epilepsy, myoclonic epilepsy which includes MERRF Syndrome, tonic-clonic epilepsy, partial epilepsy such as complex partial epilepsy, frontal lobe epilepsy and temporal lobe epilepsy, post-traumatic epilepsy, status epilepticus such as Epilepsia Partialis Continua, Hallervorden-Spatz Syndrome, hydrocephalus such as Dandy-Walker Syndrome and normal pressure hydrocephalus, hypothalamic diseases such as 15 hypothalamic neoplasms, cerebral malaria, narcolepsy which includes cataplexy, bulbar poliomyelitis, cerebri pseudotumor, Rett Syndrome, Reye's Syndrome, thalamic diseases, cerebral toxoplasmosis, intracranial tuberculoma and Zellweger Syndrome, central nervous system infections such as AIDS Dementia Complex, Brain Abscess, subdural empyema, encephalomyelitis such as Equine Encephalomyelitis, Venezuelan Equine Encephalomyelitis, 20 Necrotizing Hemorrhagic Encephalomyelitis, Visna, cerebral malaria, meningitis such as arachnoiditis, aseptic meningitis such as viral meningitis which includes lymphocytic choriomeningitis. Bacterial meningitis which includes Haemophilus Meningitis, Listeria Meningitis. Meningococcal Meningitis such as Waterhouse-Friderichsen Syndrome, Pneumococcal Meningitis and meningeal tuberculosis, fungal meningitis such as 25 Cryptococcal Meningitis, subdural effusion, meningoencephalitis such as uveomeningoencephalitic syndrome, myelitis such as transverse myelitis, neurosyphilis such as tabes dorsalis, poliomyelitis which includes bulbar poliomyelitis and postpoliomyelitis syndrome, prion diseases (such as Creutzfeldt-Jakob Syndrome, Bovine Spongiform Encephalopathy, Gerstmann-Straussler Syndrome, Kuru, Scrapie) cerebral toxoplasmosis, 30 central nervous system neoplasms such as brain neoplasms that include cerebellar neoplasms such as infratentorial neoplasms, cerebral ventricle neoplasms such as choroid plexus neoplasms, hypothalamic neoplasms and supratentorial neoplasms, meningeal

neoplasms, spinal cord neoplasms which include epidural neoplasms, demyelinating diseases such as Canavan Diseases, diffuse cerebral sclerolosis which includes adrenoleukodystrophy, encephalitis periaxialis, globoid cell leukodystrophy, diffuse cerebral sclerosis such as metachromatic leukodystrophy, allergic encephalomyelitis, necrotizing hemorrhagic
5 encephalomyelitis, progressive multifocal leukoencephalopathy, multiple sclerosis, central pontine myelinolysis, transverse myelitis, neuromyelitis optica, Scrapie, Swayback, Chronic Fatigue Syndrome, Visna, High Pressure Nervous Syndrome, Meningism, spinal cord diseases such as amyotonia congenita, amyotrophic lateral sclerosis, spinal muscular atrophy such as Werdnig-Hoffmann Disease, spinal cord compression, spinal cord neoplasms such as
10 epidural neoplasms, syringomyelia, Tabes Dorsalis, Stiff-Man Syndrome, mental retardation such as Angelman Syndrome, Cri-du-Chat Syndrome, De Lange's Syndrome, Down Syndrome, Gangliosidoses such as gangliosidoses G(M1), Sandhoff Disease, Tay-Sachs Disease, Hartnup Disease, homocystinuria, Laurence-Moon- Biedl Syndrome, Lesch-Nyhan Syndrome, Maple Syrup Urine Disease, mucopolipidosis such as fucosidosis, neuronal ceroid-
15 lipofuscinosis, oculocerebrorenal syndrome, phenylketonuria such as maternal phenylketonuria, Prader-Willi Syndrome, Rett Syndrome, Rubinstein-Taybi Syndrome, Tuberous Sclerosis, WAGR Syndrome, nervous system abnormalities such as holoprosencephaly, neural tube defects such as anencephaly which includes hydrangencephaly, Arnold-Chairi Deformity, encephalocele, meningocele,
20 meningomyelocele, spinal dysraphism such as spina bifida cystica and spina bifida occulta, hereditary motor and sensory neuropathies which include Charcot-Marie Disease, Hereditary optic atrophy, Refsum's Disease, hereditary spastic paraplegia, Werdnig-Hoffmann Disease, Hereditary Sensory and Autonomic Neuropathies such as Congenital Analgesia and Familial Dysautonomia, Neurologic manifestations (such as agnosia that include Gerstmann's
25 Syndrome, Amnesia such as retrograde amnesia, apraxia, neurogenic bladder, cataplexy, communicative disorders such as hearing disorders that includes deafness, partial hearing loss, loudness recruitment and tinnitus, language disorders such as aphasia which include agraphia, anomia, broca aphasia, and Wernicke Aphasia, Dyslexia such as Acquired Dyslexia, language development disorders, speech disorders such as aphasia which includes
30 anomia, broca aphasia and Wernicke Aphasia. articulation disorders, communicative disorders such as speech disorders which include dysarthria, echolalia, mutism and stuttering, voice disorders such as aphonia and hoarseness, decerebrate state, delirium, fasciculation,

hallucinations. meningism. movement disorders such as angelman syndrome, ataxia, athetosis. chorea. dystonia, hypokinesia, muscle hypotonia, myoclonus. tic, torticollis and tremor, muscle hypertonia such as muscle rigidity such as stiff-man syndrome, muscle spasticity, paralysis such as facial paralysis which includes Herpes Zoster Oticus, 5 Gastroparesis. Hemiplegia, ophthalmoplegia such as diplopia, Duane's Syndrome, Horner's Syndrome, Chronic progressive external ophthalmoplegia such as Kearns Syndrome, Bulbar Paralysis, Tropical Spastic Paraparesis. Paraplegia such as Brown-Sequard Syndrome, quadriplegia, respiratory paralysis and vocal cord paralysis, paresis, phantom limb, taste disorders such as ageusia and dysgeusia, vision disorders such as amblyopia, blindness, color 10 vision defects, diplopia, hemianopsia. scotoma and subnormal vision, sleep disorders such as hypersomnia which includes Kleine-Levin Syndrome, insomnia, and somnambulism, spasm such as trismus. unconsciousness such as coma, persistent vegetative state and syncope and vertigo, neuromuscular diseases such as amyotonia congenita, amyotrophic lateral sclerosis, Lambert-Eaton Myasthenic Syndrome, motor neuron disease, muscular atrophy such as 15 spinal muscular atrophy, Charcot-Marie Disease and Werdnig-Hoffmann Disease, Postpoliomyelitis Syndrome, Muscular Dystrophy, Myasthenia Gravis, Myotonia Atrophica, Myotonia Confenita, Nemaline Myopathy, Familial Periodic Paralysis, Multiplex Paramyoclonus, Tropical Spastic Paraparesis and Stiff-Man Syndrome, peripheral nervous system diseases such as acrodynia, amyloid neuropathies, autonomic nervous system diseases 20 such as Adie's Syndrome, Barre-Lieou Syndrome, Familial Dysautonomia, Horner's Syndrome, Reflex Sympathetic Dystrophy and Shy-Drager Syndrome, Cranial Nerve Diseases such as Acoustic Nerve Diseases such as Acoustic Neuroma which includes Neurofibromatosis 2, Facial Nerve Diseases such as Facial Neuralgia, Melkersson-Rosenthal Syndrome, ocular motility disorders which includes amblyopia, nystagmus, oculomotor nerve 25 paralysis, ophthalmoplegia such as Duane's Syndrome, Horner's Syndrome, Chronic Progressive External Ophthalmoplegia which includes Kearns Syndrome, Strabismus such as Esotropia and Exotropia, Oculomotor Nerve Paralysis, Optic Nerve Diseases such as Optic Atrophy which includes Hereditary Optic Atrophy, Optic Disk Drusen. Optic Neuritis such as Neuromyelitis Optica, Papilledema, Trigeminal Neuralgia, Vocal Cord Paralysis, 30 Demyelinating Diseases such as Neuromyelitis Optica and Swayback. Diabetic neuropathies such as diabetic foot, nerve compression syndromes such as carpal tunnel syndrome, tarsal tunnel syndrome, thoracic outlet syndrome such as cervical rib syndrome, ulnar nerve

- compression syndrome, neuralgia such as causalgia, cervico-brachial neuralgia, facial neuralgia and trigeminal neuralgia, neuritis such as experimental allergic neuritis, optic neuritis, polyneuritis, polyradiculoneuritis and radiculities such as polyradiculitis, hereditary motor and sensory neuropathies such as Charcot-Marie Disease. Hereditary Optic Atrophy, Refsum's Disease, Hereditary Spastic Paraplegia and Werdnig-Hoffmann Disease. Hereditary Sensory and Autonomic Neuropathies which include Congenital Analgesia and Familial Dysautonomia, POEMS Syndrome, Sciatica, Gustatory Sweating and Tetany).

Infectious Disease

- 10 Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention can be used to treat or detect infectious agents. For example, by increasing the immune response, particularly increasing the proliferation and differentiation of B and/or T cells, infectious diseases may be treated. The immune response may be increased by either enhancing an existing immune response, or by initiating a new immune response.
- 15 Alternatively, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may also directly inhibit the infectious agent, without necessarily eliciting an immune response.

- Viruses are one example of an infectious agent that can cause disease or symptoms that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention. Examples of viruses, include, but are not limited to Examples of viruses, include, but are not limited to the following DNA and RNA viruses and viral families: Arbovirus, Adenoviridae, Arenaviridae, Arterivirus, Birnaviridae, Bunyaviridae, Caliciviridae, Circoviridae, Coronaviridae, Dengue, EBV, HIV, Flaviviridae, Hepadnaviridae (Hepatitis), Herpesviridae (such as, Cytomegalovirus, Herpes Simplex, Herpes Zoster), Mononegavirus (e.g., Paramyxoviridae, Morbillivirus, Rhabdoviridae), Orthomyxoviridae (e.g., Influenza A, Influenza B, and parainfluenza), Papiloma virus, Papovaviridae, Parvoviridae, Picornaviridae, Poxviridae (such as Smallpox or Vaccinia), Reoviridae (e.g., Rotavirus), Retroviridae (HTLV-I, HTLV-II, Lentivirus), and Togaviridae (e.g., Rubivirus). Viruses falling within these families can cause a variety of diseases or symptoms, including, but not limited to: arthritis, bronchiolitis, respiratory syncytial virus, encephalitis, eye infections (e.g., conjunctivitis, keratitis), chronic fatigue syndrome, hepatitis (A, B, C, E, Chronic Active, Delta), Japanese B encephalitis, Junin, Chikungunya, Rift Valley fever,

yellow fever, meningitis, opportunistic infections (e.g., AIDS), pneumonia, Burkitt's Lymphoma, chickenpox, hemorrhagic fever, Measles, Mumps, Parainfluenza, Rabies, the common cold, Polio, leukemia, Rubella, sexually transmitted diseases, skin diseases (e.g., Kaposi's, warts), and viremia. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat: meningitis, Dengue, EBV, and/or hepatitis (e.g., hepatitis B). In an additional specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat patients nonresponsive to one or more other commercially available hepatitis vaccines. In a further specific embodiment polynucleotides, polypeptides, or agonists or antagonists of the invention are used to treat AIDS.

Similarly, bacterial or fungal agents that can cause disease or symptoms and that can be treated or detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, include, but not limited to, the following Gram-Negative and Gram-positive bacteria and bacterial families and fungi: Actinomycetales (e.g., Corynebacterium, Mycobacterium, Norcardia), Cryptococcus neoformans, Aspergillosis, Bacillaceae (e.g., Anthrax, Clostridium), Bacteroidaceae, Blastomycosis, Bordetella, Borrelia (e.g., Borrelia burgdorferi, Brucellosis, Candidiasis, Campylobacter, Coccidioidomycosis, Cryptococcosis, Dermatocycoses, E. coli (e.g., Enterotoxigenic E. coli and Enterohemorrhagic E. coli), Enterobacteriaceae (Klebsiella, Salmonella (e.g., Salmonella typhi, and Salmonella paratyphi), Serratia, Yersinia), Erysipelothrix, Helicobacter, Legionellosis, Leptospirosis, Listeria, Mycoplasmatales, Mycobacterium leprae, Vibrio cholerae, Neisseriaceae (e.g., Acinetobacter, Gonorrhea, Meningococcal), Meisseria meningitidis, Pasteurellaceae Infections (e.g., Actinobacillus, Heamophilus (e.g., Heamophilus influenza type B), Pasteurella), Pseudomonas, Rickettsiaceae, Chlamydiaceae, Syphilis, Shigella spp., Staphylococcal, Meningiococcal, Pneumococcal and Streptococcal (e.g., Streptococcus pneumoniae and Group B Streptococcus). These bacterial or fungal families can cause the following diseases or symptoms, including, but not limited to: bacteremia, endocarditis, eye infections (conjunctivitis, tuberculosis, uveitis), gingivitis, opportunistic infections (e.g., AIDS related infections), paronychia, prosthesis-related infections, Reiter's Disease, respiratory tract infections, such as Whooping Cough or Empyema, sepsis, Lyme Disease, Cat-Scratch Disease, Dysentery, Paratyphoid Fever, food poisoning, Typhoid,

pneumonia, Gonorrhea, meningitis (e.g., meningitis types A and B), Chlamydia, Syphilis, Diphtheria, Leprosy, Paratuberculosis, Tuberculosis, Lupus, Botulism, gangrene, tetanus, impetigo, Rheumatic Fever, Scarlet Fever, sexually transmitted diseases, skin diseases (e.g., cellulitis, dermatocycoses), toxemia, urinary tract infections, wound infections.

- 5 Polynucleotides or polypeptides, agonists or antagonists of the invention, can be used to treat or detect any of these symptoms or diseases. In specific embodiments, Polynucleotides, polypeptides, agonists or antagonists of the invention are used to treat: tetanus, Diphtheria, botulism, and/or meningitis type B.

- Moreover, parasitic agents causing disease or symptoms that can be treated or
10 detected by a polynucleotide or polypeptide and/or agonist or antagonist of the present invention include, but not limited to, the following families or class: Amebiasis, Babesiosis, Coccidiosis, Cryptosporidiosis, Dientamoebiasis, Dourine, Ectoparasitic, Giardiasis, Helminthiasis, Leishmaniasis, Theileriasis, Toxoplasmosis, Trypanosomiasis, and Trichomonas and Sporozoans (e.g., Plasmodium virax, Plasmodium falciparum, Plasmodium
15 malariae and Plasmodium ovale). These parasites can cause a variety of diseases or symptoms, including, but not limited to: Scabies, Trombiculiasis, eye infections, intestinal disease (e.g., dysentery, giardiasis), liver disease, lung disease, opportunistic infections (e.g., AIDS related), malaria, pregnancy complications, and toxoplasmosis. polynucleotides or polypeptides, or agonists or antagonists of the invention, can be used to treat or detect any of
20 these symptoms or diseases.

- Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention of the present invention could either be by administering an effective amount of a polypeptide to the patient, or by removing cells from the patient, supplying the cells with a polynucleotide of the present invention, and returning the engineered cells to the patient (ex
25 vivo therapy). Moreover, the polypeptide or polynucleotide of the present invention can be used as an antigen in a vaccine to raise an immune response against infectious disease.

Regeneration

- Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention can be used to differentiate, proliferate, and attract cells, leading to the regeneration
30 of tissues. (See, Science 276:59-87 (1997).) The regeneration of tissues could be used to repair, replace, or protect tissue damaged by congenital defects, trauma (wounds, burns,

incisions, or ulcers). age, disease (e.g. osteoporosis, osteoarthritis, periodontal disease, liver failure), surgery, including cosmetic plastic surgery, fibrosis, reperfusion injury, or systemic cytokine damage.

Tissues that could be regenerated using the present invention include organs (e.g.,
5 pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal or cardiac),
vasculature (including vascular and lymphatics), nervous, hematopoietic, and skeletal (bone,
cartilage, tendon, and ligament) tissue. Preferably, regeneration occurs without or decreased
scarring. Regeneration also may include angiogenesis.

Moreover, polynucleotides or polypeptides, as well as agonists or antagonists of the
10 present invention, may increase regeneration of tissues difficult to heal. For example,
increased tendon/ligament regeneration would quicken recovery time after damage.
Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention
could also be used prophylactically in an effort to avoid damage. Specific diseases that could
be treated include of tendinitis, carpal tunnel syndrome, and other tendon or ligament defects.
15 A further example of tissue regeneration of non-healing wounds includes pressure ulcers,
ulcers associated with vascular insufficiency, surgical, and traumatic wounds.

Similarly, nerve and brain tissue could also be regenerated by using polynucleotides
or polypeptides, as well as agonists or antagonists of the present invention, to proliferate and
differentiate nerve cells. Diseases that could be treated using this method include central and
20 peripheral nervous system diseases, neuropathies, or mechanical and traumatic disorders
(e.g., spinal cord disorders, head trauma, cerebrovascular disease, and stroke). Specifically,
diseases associated with peripheral nerve injuries, peripheral neuropathy (e.g., resulting from
chemotherapy or other medical therapies), localized neuropathies, and central nervous system
diseases (e.g., Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic
25 lateral sclerosis, and Shy-Drager syndrome), could all be treated using the polynucleotides or
polypeptides, as well as agonists or antagonists of the present invention.

Chemotaxis

Polynucleotides or polypeptides, as well as agonists or antagonists of the present
30 invention may have chemotaxis activity. A chemotactic molecule attracts or mobilizes cells
(e.g., monocytes, fibroblasts, neutrophils, T-cells, mast cells, eosinophils, epithelial and/or
endothelial cells) to a particular site in the body, such as inflammation, infection, or site of

hyperproliferation. The mobilized cells can then fight off and/or heal the particular trauma or abnormality.

Polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may increase chemotactic activity of particular cells. These chemotactic molecules can then be used to treat inflammation, infection, hyperproliferative disorders, or any immune system disorder by increasing the number of cells targeted to a particular location in the body. For example, chemotactic molecules can be used to treat wounds and other trauma to tissues by attracting immune cells to the injured location. Chemotactic molecules of the present invention can also attract fibroblasts, which can be used to treat wounds.

It is also contemplated that polynucleotides or polypeptides, as well as agonists or antagonists of the present invention may inhibit chemotactic activity. These molecules could also be used to treat disorders. Thus, polynucleotides or polypeptides, as well as agonists or antagonists of the present invention could be used as an inhibitor of chemotaxis.

15 **Binding Activity**

A polypeptide of the present invention may be used to screen for molecules that bind to the polypeptide or for molecules to which the polypeptide binds. The binding of the polypeptide and the molecule may activate (agonist), increase, inhibit (antagonist), or decrease activity of the polypeptide or the molecule bound. Examples of such molecules include antibodies, oligonucleotides, proteins (e.g., receptors), or small molecules.

Preferably, the molecule is closely related to the natural ligand of the polypeptide, e.g., a fragment of the ligand, or a natural substrate, a ligand, a structural or functional mimetic. (See, Coligan et al., Current Protocols in Immunology 1(2):Chapter 5 (1991).) Similarly, the molecule can be closely related to the natural receptor to which the polypeptide binds, or at least, a fragment of the receptor capable of being bound by the polypeptide (e.g., active site). In either case, the molecule can be rationally designed using known techniques.

Preferably, the screening for these molecules involves producing appropriate cells which express the polypeptide. Preferred cells include cells from mammals, yeast, *Drosophila*, or *E. coli*. Cells expressing the polypeptide (or cell membrane containing the expressed polypeptide) are then preferably contacted with a test compound potentially containing the molecule to observe binding, stimulation, or inhibition of activity of either the polypeptide or the molecule.

The assay may simply test binding of a candidate compound to the polypeptide, wherein binding is detected by a label, or in an assay involving competition with a labeled competitor. Further, the assay may test whether the candidate compound results in a signal generated by binding to the polypeptide.

5 Alternatively, the assay can be carried out using cell-free preparations, polypeptide/molecule affixed to a solid support, chemical libraries, or natural product mixtures. The assay may also simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide, measuring polypeptide/molecule activity or binding, and comparing the polypeptide/molecule activity or binding to a standard.

10 Preferably, an ELISA assay can measure polypeptide level or activity in a sample (e.g., biological sample) using a monoclonal or polyclonal antibody. The antibody can measure polypeptide level or activity by either binding, directly or indirectly, to the polypeptide or by competing with the polypeptide for a substrate.

15 Additionally, the receptor to which the polypeptide of the present invention binds can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting (Coligan, et al., Current Protocols in Immun., 1(2), Chapter 5, (1991)). For example, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the polypeptides, for example, NIH3T3 cells which are known to contain multiple receptors for the FGF family proteins, and SC-3 cells, and a cDNA
20 library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the polypeptides. Transfected cells which are grown on glass slides are exposed to the polypeptide of the present invention, after they have been labelled. The polypeptides can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase.

25 Following fixation and incubation, the slides are subjected to auto-radiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an iterative sub-pooling and re-screening process, eventually yielding a single clones that encodes the putative receptor.

30 As an alternative approach for receptor identification, the labeled polypeptides can be photoaffinity linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE analysis and exposed to X-ray film. The labeled complex containing the receptors of the polypeptides can be excised, resolved

into peptide fragments, and subjected to protein microsequencing. The amino acid sequence obtained from microsequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the genes encoding the putative receptors.

Moreover, the techniques of gene-shuffling, motif-shuffling, exon-shuffling, and/or
5 codon-shuffling (collectively referred to as "DNA shuffling") may be employed to modulate the activities of the polypeptide of the present invention thereby effectively generating agonists and antagonists of the polypeptide of the present invention. *See generally*, U.S. Patent Nos. 5,605,793, 5,811,238, 5,830,721, 5,834,252, and 5,837,458, and Patten, P. A., *et al.*, *Curr. Opinion Biotechnol.* 8:724-33 (1997); Harayama, S. *Trends Biotechnol.* 16(2):76-
10 82 (1998); Hansson, L. O., *et al.*, *J. Mol. Biol.* 287:265-76 (1999); and Lorenzo, M. M. and Blasco, R. *Biotechniques* 24(2):308-13 (1998) (each of these patents and publications are hereby incorporated by reference). In one embodiment, alteration of polynucleotides and corresponding polypeptides may be achieved by DNA shuffling. DNA shuffling involves the assembly of two or more DNA segments into a desired molecule by homologous, or site-
15 specific, recombination. In another embodiment, polynucleotides and corresponding polypeptides may be altered by being subjected to random mutagenesis by error-prone PCR, random nucleotide insertion or other methods prior to recombination. In another embodiment, one or more components, motifs, sections, parts, domains, fragments, etc., of the polypeptide of the present invention may be recombined with one or more components,
20 motifs, sections, parts, domains, fragments, etc. of one or more heterologous molecules. In preferred embodiments, the heterologous molecules are family members. In further preferred embodiments, the heterologous molecule is a growth factor such as, for example, platelet-derived growth factor (PDGF), insulin-like growth factor (IGF-I), transforming growth factor (TGF)-alpha, epidermal growth factor (EGF), fibroblast growth factor (FGF),
25 TGF-beta, bone morphogenetic protein (BMP)-2, BMP-4, BMP-5, BMP-6, BMP-7, activins A and B, decapentaplegic(dpp), 60A, OP-2, dorsalin, growth differentiation factors (GDFs), nodal, MIS, inhibin-alpha, TGF-beta1, TGF-beta2, TGF-beta3, TGF-beta5, and glial-derived neurotrophic factor (GDNF).

Other preferred fragments are biologically active fragments of the polypeptide of the
30 present invention. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The

biological activity of the fragments may include an improved desired activity, or a decreased undesirable activity.

Additionally, this invention provides a method of screening compounds to identify those which modulate the action of the polypeptide of the present invention. An example of
5 such an assay comprises combining a mammalian fibroblast cell, a the polypeptide of the present invention, the compound to be screened and $^3\text{[H]}$ thymidine under cell culture conditions where the fibroblast cell would normally proliferate. A control assay may be performed in the absence of the compound to be screened and compared to the amount of fibroblast proliferation in the presence of the compound to determine if the compound
10 stimulates proliferation by determining the uptake of $^3\text{[H]}$ thymidine in each case. The amount of fibroblast cell proliferation is measured by liquid scintillation chromatography which measures the incorporation of $^3\text{[H]}$ thymidine. Both agonist and antagonist compounds may be identified by this procedure.

In another method, a mammalian cell or membrane preparation expressing a receptor
15 for a polypeptide of the present invention is incubated with a labeled polypeptide of the present invention in the presence of the compound. The ability of the compound to enhance or block this interaction could then be measured. Alternatively, the response of a known second messenger system following interaction of a compound to be screened and the receptor is measured and the ability of the compound to bind to the receptor and elicit a
20 second messenger response is measured to determine if the compound is a potential agonist or antagonist. Such second messenger systems include but are not limited to, cAMP guanylate cyclase, ion channels or phosphoinositide hydrolysis.

All of these above assays can be used as diagnostic or prognostic markers. The molecules discovered using these assays can be used to treat disease or to bring about a
25 particular result in a patient (e.g., blood vessel growth) by activating or inhibiting the polypeptide/molecule. Moreover, the assays can discover agents which may inhibit or enhance the production of the polypeptides of the invention from suitably manipulated cells or tissues.

Therefore, the invention includes a method of identifying compounds which bind to a
30 polypeptide of the invention comprising the steps of: (a) incubating a candidate binding compound with a polypeptide of the present invention; and (b) determining if binding has

occurred. Moreover, the invention includes a method of identifying agonists/antagonists comprising the steps of: (a) incubating a candidate compound with a polypeptide of the present invention, (b) assaying a biological activity, and (b) determining if a biological activity of the polypeptide has been altered.

5

Targeted Delivery

In another embodiment, the invention provides a method of delivering compositions to targeted cells expressing a receptor for a polypeptide of the invention, or cells expressing a cell bound form of a polypeptide of the invention.

10 As discussed herein, polypeptides or antibodies of the invention may be associated with heterologous polypeptides, heterologous nucleic acids, toxins, or prodrugs via hydrophobic, hydrophilic, ionic and/or covalent interactions. In one embodiment, the invention provides a method for the specific delivery of compositions of the invention to cells by administering polypeptides of the invention (including antibodies) that are associated with
15 heterologous polypeptides or nucleic acids. In one example, the invention provides a method for delivering a therapeutic protein into the targeted cell. In another example, the invention provides a method for delivering a single stranded nucleic acid (e.g., antisense or ribozymes) or double stranded nucleic acid (e.g., DNA that can integrate into the cell's genome or replicate episomally and that can be transcribed) into the targeted cell.

20 In another embodiment, the invention provides a method for the specific destruction of cells (e.g., the destruction of tumor cells) by administering polypeptides of the invention (e.g., polypeptides of the invention or antibodies of the invention) in association with toxins or cytotoxic prodrugs.

By "toxin" is meant compounds that bind and activate endogenous cytotoxic effector
25 systems, radioisotopes, holotoxins, modified toxins, catalytic subunits of toxins, or any molecules or enzymes not normally present in or on the surface of a cell that under defined conditions cause the cell's death. Toxins that may be used according to the methods of the invention include, but are not limited to, radioisotopes known in the art, compounds such as, for example, antibodies (or complement fixing containing portions thereof) that bind an
30 inherent or induced endogenous cytotoxic effector system, thymidine kinase, endonuclease, RNase, alpha toxin, ricin, abrin, *Pseudomonas* exotoxin A, diphtheria toxin, saporin,

momordin, gelonin, pokeweed antiviral protein, alpha-sarcin and cholera toxin. By "cytotoxic prodrug" is meant a non-toxic compound that is converted by an enzyme, normally present in the cell, into a cytotoxic compound. Cytotoxic prodrugs that may be used according to the methods of the invention include, but are not limited to, glutamyl
5 derivatives of benzoic acid mustard alkylating agent, phosphate derivatives of etoposide or mitomycin C, cytosine arabinoside, daunorubisin, and phenoxycetamide derivatives of doxorubicin.

Drug Screening

10 Further contemplated is the use of the polypeptides of the present invention, or the polynucleotides encoding these polypeptides, to screen for molecules which modify the activities of the polypeptides of the present invention. Such a method would include contacting the polypeptide of the present invention with a selected compound(s) suspected of having antagonist or agonist activity, and assaying the activity of these polypeptides
15 following binding.

This invention is particularly useful for screening therapeutic compounds by using the polypeptides of the present invention, or binding fragments thereof, in any of a variety of drug screening techniques. The polypeptide or fragment employed in such a test may be affixed to a solid support, expressed on a cell surface, free in solution, or located
20 intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. One may measure, for example, the formulation of complexes between the agent being tested and a polypeptide of the present invention.

25 Thus, the present invention provides methods of screening for drugs or any other agents which affect activities mediated by the polypeptides of the present invention. These methods comprise contacting such an agent with a polypeptide of the present invention or a fragment thereof and assaying for the presence of a complex between the agent and the polypeptide or a fragment thereof, by methods well known in the art. In such a competitive
30 binding assay, the agents to screen are typically labeled. Following incubation, free agent is separated from that present in bound form, and the amount of free or uncomplexed label is a

measure of the ability of a particular agent to bind to the polypeptides of the present invention.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to the polypeptides of the present invention, and is described in great detail in European Patent Application 84/03564, published on September 13, 1984, which is incorporated herein by reference herein. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The peptide test compounds are reacted with polypeptides of the present invention and washed. Bound polypeptides are then detected by methods well known in the art. Purified polypeptides are coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies may be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding polypeptides of the present invention specifically compete with a test compound for binding to the polypeptides or fragments thereof. In this manner, the antibodies are used to detect the presence of any peptide which shares one or more antigenic epitopes with a polypeptide of the invention.

Antisense And Ribozyme (Antagonists)

In specific embodiments, antagonists according to the present invention are nucleic acids corresponding to the sequences contained in SEQ ID NO:X, or the complementary strand thereof, and/or to nucleotide sequences contained in the cDNA contained in the related cDNA clone identified in Table 1. In one embodiment, antisense sequence is generated internally, by the organism, in another embodiment, the antisense sequence is separately administered (see, for example, O'Connor, J., Neurochem. 56:560 (1991). Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Antisense technology can be used to control gene expression through antisense DNA or RNA, or through triple-helix formation. Antisense techniques are discussed for example, in Okano, J., Neurochem. 56:560 (1991); Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988). Triple helix formation is discussed in, for instance, Lee et al., Nucleic Acids Research 6:3073 (1979); Cooney et al.,

Science 241:456 (1988); and Dervan et al., Science 251:1300 (1991). The methods are based on binding of a polynucleotide to a complementary DNA or RNA.

For example, the use of c-myc and c-myb antisense RNA constructs to inhibit the growth of the non-lymphocytic leukemia cell line HL-60 and other cell lines was previously described. (Wickstrom et al. (1988); Anfossi et al. (1989)). These experiments were performed in vitro by incubating cells with the oligoribonucleotide. A similar procedure for in vivo use is described in WO 91/15580. Briefly, a pair of oligonucleotides for a given antisense RNA is produced as follows: A sequence complimentary to the first 15 bases of the open reading frame is flanked by an EcoRI site on the 5' end and a HindIII site on the 3' end. Next, the pair of oligonucleotides is heated at 90°C for one minute and then annealed in 2X ligation buffer (20mM TRIS HCl pH 7.5, 10mM MgCl₂, 10mM dithiothreitol (DTT) and 0.2 mM ATP) and then ligated to the EcoRI/Hind III site of the retroviral vector PMV7 (WO 91/15580).

For example, the 5' coding portion of a polynucleotide that encodes the polypeptide of the present invention may be used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription thereby preventing transcription and the production of the receptor. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into receptor polypeptide.

In one embodiment, the antisense nucleic acid of the invention is produced intracellularly by transcription from an exogenous sequence. For example, a vector or a portion thereof, is transcribed, producing an antisense nucleic acid (RNA) of the invention. Such a vector would contain a sequence encoding the antisense nucleic acid. Such a vector can remain episomal or become chromosomally integrated, as long as it can be transcribed to produce the desired antisense RNA. Such vectors can be constructed by recombinant DNA technology methods standard in the art. Vectors can be plasmid, viral, or others known in the art, used for replication and expression in vertebrate cells. Expression of the sequence encoding the polypeptide of the present invention or fragments thereof, can be by any promoter known in the art to act in vertebrate, preferably human cells. Such promoters can be inducible or constitutive. Such promoters include, but are not limited to, the SV40 early promoter region (Bernoist and Chambon, Nature 29:304-310 (1981)), the promoter contained in the 3' long terminal repeat of Rous sarcoma virus (Yamamoto et al., Cell 22:787-797

(1980), the herpes thymidine promoter (Wagner et al., Proc. Natl. Acad. Sci. U.S.A. 78:1441-1445 (1981), the regulatory sequences of the metallothionein gene (Brinster, et al., Nature 296:39-42 (1982)), etc.

The antisense nucleic acids of the invention comprise a sequence complementary to at least a portion of an RNA transcript of a gene of the present invention. However, absolute complementarity, although preferred, is not required. A sequence "complementary to at least a portion of an RNA," referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the larger the hybridizing nucleic acid, the more base mismatches with a RNA it may contain and still form a stable duplex (or triplex as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the message, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have been shown to be effective at inhibiting translation of mRNAs as well. See generally, Wagner, R., 1994, Nature 372:333-335. Thus, oligonucleotides complementary to either the 5'- or 3'- non- translated, non-coding regions of polynucleotide sequences described herein could be used in an antisense approach to inhibit translation of endogenous mRNA. Oligonucleotides complementary to the 5' untranslated region of the mRNA should include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could be used in accordance with the invention. Whether designed to hybridize to the 5', 3'- or coding region of mRNA of the present invention, antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

The polynucleotides of the invention can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The

oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as peptides (e.g., for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al., 5 1989, Proc. Natl. Acad. Sci. U.S.A. 86:6553-6556; Lemaitre et al., 1987, Proc. Natl. Acad. Sci. 84:648-652; PCT Publication No. WO88/09810, published December 15, 1988) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134, published April 25, 1988), hybridization-triggered cleavage agents. (See, e.g., Krol et al., 1988, BioTechniques 6:958-976) or intercalating agents. (See, e.g., Zon, 1988, Pharm. Res. 5:539-549). To this end, the 10 oligonucleotide may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

The antisense oligonucleotide may comprise at least one modified base moiety which is selected from the group including, but not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 15 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta- 20 D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine.

25 The antisense oligonucleotide may also comprise at least one modified sugar moiety selected from the group including, but not limited to, arabinose, 2-fluoroarabinose, xylulose, and hexose.

In yet another embodiment, the antisense oligonucleotide comprises at least one modified phosphate backbone selected from the group including, but not limited to, a 30 phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In yet another embodiment, the antisense oligonucleotide is an a-anomeric oligonucleotide. An a-anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual b-units, the strands run parallel to each other (Gautier et al., 1987, Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al., 1987, Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al., 1987, FEBS Lett. 215:327-330).

Polynucleotides of the invention may be synthesized by standard methods known in the art, e.g. by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988, Nucl. Acids Res. 16:3209), methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al., 1988, Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc.

While antisense nucleotides complementary to the coding region sequence could be used, those complementary to the transcribed untranslated region are most preferred.

Potential antagonists according to the invention also include catalytic RNA, or a ribozyme (See, e.g., PCT International Publication WO 90/11364, published October 4, 1990; Sarver et al, Science 247:1222-1225 (1990). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach, Nature 334:585-591 (1988). There are numerous potential hammerhead ribozyme cleavage sites within the nucleotide sequence of SEQ ID NO:X. Preferably, the ribozyme is engineered so that the cleavage recognition site is located near the 5' end of the mRNA; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

As in the antisense approach, the ribozymes of the invention can be composed of modified oligonucleotides (e.g. for improved stability, targeting, etc.) and should be delivered to cells which express in vivo. DNA constructs encoding the ribozyme may be introduced into the cell in the same manner as described above for the introduction of antisense encoding DNA. A preferred method of delivery involves using a DNA construct "encoding" the

ribozyme under the control of a strong constitutive promoter, such as, for example, pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous messages and inhibit translation. Since ribozymes unlike antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

- 5 Antagonist/agonist compounds may be employed to inhibit the cell growth and proliferation effects of the polypeptides of the present invention on neoplastic cells and tissues, i.e. stimulation of angiogenesis of tumors, and, therefore, retard or prevent abnormal cellular growth and proliferation, for example, in tumor formation or growth.

- 10 The antagonist/agonist may also be employed to prevent hyper-vascular diseases, and prevent the proliferation of epithelial lens cells after extracapsular cataract surgery. Prevention of the mitogenic activity of the polypeptides of the present invention may also be desirable in cases such as restenosis after balloon angioplasty.

 The antagonist/agonist may also be employed to prevent the growth of scar tissue during wound healing.

- 15 The antagonist/agonist may also be employed to treat the diseases described herein.

- Thus, the invention provides a method of treating disorders or diseases, including but not limited to the disorders or diseases listed throughout this application, associated with overexpression of a polynucleotide of the present invention by administering to a patient (a) an antisense molecule directed to the polynucleotide of the present invention, and/or (b) a
20 ribozyme directed to the polynucleotide of the present invention.

Other Activities

- A polypeptide, polynucleotide, agonist, or antagonist of the present invention, as a result of the ability to stimulate vascular endothelial cell growth, may be employed in
25 treatment for stimulating re-vascularization of ischemic tissues due to various disease conditions such as thrombosis, arteriosclerosis, and other cardiovascular conditions. The polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed to stimulate angiogenesis and limb regeneration, as discussed above.

- 30 A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for treating wounds due to injuries, burns, post-operative tissue repair, and ulcers since they are mitogenic to various cells of different origins, such as fibroblast cells

and skeletal muscle cells, and therefore, facilitate the repair or replacement of damaged or diseased tissue.

5 A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed stimulate neuronal growth and to treat and prevent neuronal damage which occurs in certain neuronal disorders or neuro-degenerative conditions such as Alzheimer's disease, Parkinson's disease, and AIDS-related complex. A polypeptide, polynucleotide, agonist, or antagonist of the present invention may have the ability to stimulate chondrocyte growth, therefore, they may be employed to enhance bone and periodontal regeneration and aid in tissue transplants or bone grafts.

10 A polypeptide, polynucleotide, agonist, or antagonist of the present invention may be also be employed to prevent skin aging due to sunburn by stimulating keratinocyte growth.

A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed for preventing hair loss, since FGF family members activate hair-forming cells and promotes melanocyte growth. Along the same lines, a polypeptide, polynucleotide, 15 agonist, or antagonist of the present invention may be employed to stimulate growth and differentiation of hematopoietic cells and bone marrow cells when used in combination with other cytokines.

A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be employed to maintain organs before transplantation or for supporting cell culture of primary tissues. A polypeptide, polynucleotide, agonist, or antagonist of the present 20 invention may also be employed for inducing tissue of mesodermal origin to differentiate in early embryos.

A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also increase or decrease the differentiation or proliferation of embryonic stem cells, besides, 25 as discussed above, hematopoietic lineage.

A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be used to modulate mammalian characteristics. such as body height, weight, hair color, eye color, skin, percentage of adipose tissue, pigmentation, size, and shape (e.g., cosmetic surgery). Similarly, a polypeptide, polynucleotide, agonist, or antagonist of the present 30 invention may be used to modulate mammalian metabolism affecting catabolism, anabolism, processing, utilization, and storage of energy.

A polypeptide, polynucleotide, agonist, or antagonist of the present invention may be used to change a mammal's mental state or physical state by influencing biorhythms, circadian rhythms, depression (including depressive disorders), tendency for violence, tolerance for pain, reproductive capabilities (preferably by Activin or Inhibin-like activity),
5 hormonal or endocrine levels, appetite, libido, memory, stress, or other cognitive qualities.

A polypeptide, polynucleotide, agonist, or antagonist of the present invention may also be used as a food additive or preservative, such as to increase or decrease storage capabilities, fat content, lipid, protein, carbohydrate, vitamins, minerals, cofactors or other nutritional components.

10 The above-recited applications have uses in a wide variety of hosts. Such hosts include, but are not limited to, human, murine, rabbit, goat, guinea pig, camel, horse, mouse, rat, hamster, pig, micro-pig, chicken, goat, cow, sheep, dog, cat, non-human primate, and human. In specific embodiments, the host is a mouse, rabbit, goat, guinea pig, chicken, rat, hamster, pig, sheep, dog or cat. In preferred embodiments, the host is a mammal. In most
15 preferred embodiments, the host is a human.

Other Preferred Embodiments

Other preferred embodiments of the claimed invention include an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence
20 of at least about 50 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, and/or the cDNA in the related cDNA clone contained in the deposit.

Also preferred is a nucleic acid molecule wherein said sequence of contiguous nucleotides is included in the nucleotide sequence of SEQ ID NO:X in the range of positions
25 identified as "Start" and "End" in columns 7 and 8 as defined for SEQ ID NO:X in Table 1.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 150 contiguous nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, and/or the cDNA in the related cDNA clone contained in the deposit.

30 Further preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least about 500 contiguous

nucleotides in the nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, and/or the cDNA in the related cDNA clone contained in the deposit.

5 A further preferred embodiment is a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the nucleotide sequence of SEQ ID NO:X in the range of positions identified as "Start" and "End" in columns 7 and 8 as defined for SEQ ID NO:X in Table 1.

10 A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, and/or the cDNA in the related cDNA clone contained in the deposit.

Also preferred is an isolated nucleic acid molecule which hybridizes under stringent hybridization conditions to a nucleic acid molecule comprising a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto, and/or the cDNA in the related cDNA clone contained in the deposit, wherein said nucleic acid molecule which hybridizes does not
15 hybridize under stringent hybridization conditions to a nucleic acid molecule having a nucleotide sequence consisting of only A residues or of only T residues.

Also preferred is a composition of matter comprising a DNA molecule which comprises a cDNA clone contained in the deposit.

20 Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in the nucleotide sequence of the cDNA in the related cDNA clone contained in the deposit.

Also preferred is an isolated nucleic acid molecule, wherein said sequence of at least 50 contiguous nucleotides is included in the nucleotide sequence of an open reading frame sequence encoded by the cDNA in the related cDNA clone contained in the deposit.

25 Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 150 contiguous nucleotides in the nucleotide sequence encoded by the cDNA in the related cDNA clone contained in the deposit.

30 A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to sequence of at least 500 contiguous nucleotides in the nucleotide sequence encoded by the cDNA in the related cDNA clone contained in the deposit.

A further preferred embodiment is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to the complete nucleotide sequence encoded by the cDNA in the related cDNA clone contained in the deposit.

5 A further preferred embodiment is a method for detecting in a biological sample a nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; and a nucleotide sequence encoded by the cDNA in the related cDNA clone contained in the deposit; which method comprises a step of comparing a nucleotide sequence of at least one
10 nucleic acid molecule in said sample with a sequence selected from said group and determining whether the sequence of said nucleic acid molecule in said sample is at least 95% identical to said selected sequence.

Also preferred is the above method wherein said step of comparing sequences comprises determining the extent of nucleic acid hybridization between nucleic acid
15 molecules in said sample and a nucleic acid molecule comprising said sequence selected from said group. Similarly, also preferred is the above method wherein said step of comparing sequences is performed by comparing the nucleotide sequence determined from a nucleic acid molecule in said sample with said sequence selected from said group. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

20 A further preferred embodiment is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting nucleic acid molecules in said sample, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand
25 thereto; and a nucleotide sequence encoded by the cDNA in the related cDNA clone contained in the deposit.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample which comprises a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one
30 sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a nucleotide sequence of SEQ ID NO:X; or the cDNA contained in the related cDNA clone referenced in Table I which encodes a protein, wherein the method comprises a step of detecting in a biological sample obtained
5 from said subject nucleic acid molecules, if any, comprising a nucleotide sequence that is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; and a nucleotide sequence of the cDNA in the related cDNA clone contained in the deposit.

10 Also preferred is the above method for diagnosing a pathological condition which comprises a step of detecting nucleic acid molecules comprising a nucleotide sequence in a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from said group.

15 Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a panel of at least two nucleotide sequences, wherein at least one sequence in said panel is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand
20 thereto; and a nucleotide sequence encoded by the cDNA in the related cDNA clone contained in the deposit. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is a composition of matter comprising isolated nucleic acid molecules wherein the nucleotide sequences of said nucleic acid molecules comprise a DNA microarray or "chip" of at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 40, 50, 100, 150, 200, 250, 300,
25 500, 1000, 2000, 3000 or 4000 nucleotide sequences, wherein at least one sequence in said DNA microarray or "chip" is at least 95% identical to a sequence of at least 50 contiguous nucleotides in a sequence selected from the group consisting of: a nucleotide sequence of SEQ ID NO:X or the complementary strand thereto; and a nucleotide sequence encoded by
30 the cDNA in the cDNA clone referenced in Table I. The nucleic acid molecules can comprise DNA molecules or RNA molecules.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and/or a polypeptide encoded by the cDNA in the related cDNA clone contained in the deposit.

5 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and/or a polypeptide encoded by the cDNA in the related cDNA clone contained in the deposit.

Further preferred is an isolated polypeptide comprising an amino acid sequence at
10 least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and/or a polypeptide encoded by the cDNA in the related cDNA clone contained in the deposit.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the complete amino acid sequence of SEQ ID NO:Y; a polypeptide
15 encoded by SEQ ID NO:X; and/or a polypeptide encoded by the cDNA in the related cDNA clone contained in the deposit.

Further preferred is an isolated polypeptide comprising an amino acid sequence at least 90% identical to a sequence of at least about 10 contiguous amino acids in the complete amino acid sequence of a polypeptide encoded by the cDNA clone referenced in Table 1.

20 Also preferred is a polypeptide wherein said sequence of contiguous amino acids is included in the amino acid sequence of a portion of said polypeptide encoded by the cDNA clone referenced in Table 1; a polypeptide encoded by SEQ ID NO:X; and/or the polypeptide sequence of SEQ ID NO:Y.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least
25 95% identical to a sequence of at least about 30 contiguous amino acids in the amino acid sequence of a polypeptide encoded by the cDNA clone referenced in Table 1.

Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to a sequence of at least about 100 contiguous amino acids in the amino acid sequence of a polypeptide encoded by the cDNA clone referenced in Table 1.

30 Also preferred is an isolated polypeptide comprising an amino acid sequence at least 95% identical to the amino acid sequence of a polypeptide encoded by the cDNA clone referenced in Table 1.

Further preferred is an isolated antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide
5 encoded by the cDNA in the related cDNA clone contained in the deposit.

Further preferred is a method for detecting in a biological sample a polypeptide comprising an amino acid sequence which is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide
10 encoded by the cDNA in the related cDNA clone referenced in Table 1; which method comprises a step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group and determining whether the sequence of said polypeptide molecule in said sample is at least 90% identical to said sequence of at least 10 contiguous amino acids.

Also preferred is the above method wherein said step of comparing an amino acid sequence of at least one polypeptide molecule in said sample with a sequence selected from said group comprises determining the extent of specific binding of polypeptides in said sample to an antibody which binds specifically to a polypeptide comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a
15 sequence selected from the group consisting of: a polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide encoded by the cDNA in the related cDNA clone referenced in Table 1.

Also preferred is the above method wherein said step of comparing sequences is performed by comparing the amino acid sequence determined from a polypeptide molecule in
20 said sample with said sequence selected from said group.

Also preferred is a method for identifying the species, tissue or cell type of a biological sample which method comprises a step of detecting polypeptide molecules in said sample, if any, comprising an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of:
25 polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide encoded by the cDNA in the related cDNA clone referenced in Table 1.

Also preferred is the above method for identifying the species, tissue or cell type of a biological sample, which method comprises a step of detecting polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the above group.

Also preferred is a method for diagnosing in a subject a pathological condition associated with abnormal structure or expression of a nucleic acid sequence identified in Table 1 encoding a polypeptide, which method comprises a step of detecting in a biological sample obtained from said subject polypeptide molecules comprising an amino acid sequence in a panel of at least two amino acid sequences, wherein at least one sequence in said panel is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide encoded by the cDNA in the related cDNA clone referenced in Table 1.

In any of these methods, the step of detecting said polypeptide molecules includes using an antibody.

Also preferred is an isolated nucleic acid molecule comprising a nucleotide sequence which is at least 95% identical to a nucleotide sequence encoding a polypeptide wherein said polypeptide comprises an amino acid sequence that is at least 90% identical to a sequence of at least 10 contiguous amino acids in a sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide encoded by the cDNA in the related cDNA clone referenced in Table 1.

Also preferred is an isolated nucleic acid molecule, wherein said nucleotide sequence encoding a polypeptide has been optimized for expression of said polypeptide in a prokaryotic host.

Also preferred is an isolated nucleic acid molecule, wherein said polypeptide comprises an amino acid sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide encoded by the cDNA in the related cDNA clone referenced in Table 1.

Further preferred is a method of making a recombinant vector comprising inserting any of the above isolated nucleic acid molecule into a vector. Also preferred is the recombinant vector produced by this method. Also preferred is a method of making a

recombinant host cell comprising introducing the vector into a host cell. as well as the recombinant host cell produced by this method.

Also preferred is a method of making an isolated polypeptide comprising culturing this recombinant host cell under conditions such that said polypeptide is expressed and recovering said polypeptide. Also preferred is this method of making an isolated polypeptide, wherein said recombinant host cell is a eukaryotic cell and said polypeptide is a human protein comprising an amino acid sequence selected from the group consisting of: polypeptide sequence of SEQ ID NO:Y; a polypeptide encoded by SEQ ID NO:X; and a polypeptide encoded by the cDNA in the related cDNA clone referenced in Table 1. The isolated polypeptide produced by this method is also preferred.

Also preferred is a method of treatment of an individual in need of an increased level of a protein activity, which method comprises administering to such an individual a Therapeutic comprising an amount of an isolated polypeptide, polynucleotide, immunogenic fragment or analogue thereof, binding agent, antibody, or antigen binding fragment of the claimed invention effective to increase the level of said protein activity in said individual.

Also preferred is a method of treatment of an individual in need of a decreased level of a protein activity, which method comprised administering to such an individual a Therapeutic comprising an amount of an isolated polypeptide, polynucleotide, immunogenic fragment or analogue thereof, binding agent, antibody, or antigen binding fragment of the claimed invention effective to decrease the level of said protein activity in said individual.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

*Examples**Example 1: Isolation of a Selected cDNA Clone From the Deposited Sample*

5

Each deposited cDNA clone is contained in a plasmid vector. Table 5 identifies the vectors used to construct the cDNA library from which each clone was isolated. In many cases, the vector used to construct the library is a phage vector from which a plasmid has been excised. The following correlates the related plasmid for each phage vector used in constructing the cDNA library. For example, where a particular clone is identified in Table 5 as being isolated in the vector "Lambda Zap," the corresponding deposited clone is in "pBluescript."

	<u>Vector Used to Construct Library</u>	<u>Corresponding Deposited Plasmid</u>
	Lambda Zap	pBluescript (pBS)
15	Uni-Zap XR	pBluescript (pBS)
	Zap Express	pBK
	lafrmid BA	plafmid BA
	pSport1	pSport1
	pCMVSPORT 2.0	pCMVSPORT 2.0
20	pCMVSPORT 3.0	pCMVSPORT 3.0
	pCR [®] 2.1	pCR [®] 2.1

Vectors Lambda Zap (U.S. Patent Nos. 5,128,256 and 5,286,636), Uni-Zap XR (U.S. Patent Nos. 5,128,256 and 5,286,636), Zap Express (U.S. Patent Nos. 5,128,256 and 5,286,636), pBluescript (pBS) (Short, J. M. et al., Nucleic Acids Res. 16:7583-7600 (1988); Alting-Mees, M. A. and Short, J. M., Nucleic Acids Res. 17:9494 (1989)) and pBK (Alting-Mees, M. A. et al., Strategies 5:58-61 (1992)) are commercially available from Stratagene Cloning Systems, Inc., 11011 N. Torrey Pines Road, La Jolla, CA, 92037. pBS contains an ampicillin resistance gene and pBK contains a neomycin resistance gene. Both can be transformed into E. coli strain XL-1 Blue, also available from Stratagene. pBS comes in 4 forms SK+, SK-, KS+ and KS-. The S and K refers to the orientation of the polylinker to the T7 and T3 primer sequences which flank the polylinker region ("S" is for SacI and "K" is for KpnI which are the first sites on each respective end of the linker). "+" or "-" refer to the

orientation of the fl origin of replication ("ori"), such that in one orientation, single stranded rescue initiated from the fl ori generates sense strand DNA and in the other, antisense.

Vectors pSport1, pCMVSPORT 2.0 and pCMVSPORT 3.0, were obtained from Life Technologies, Inc., P. O. Box 6009, Gaithersburg, MD 20897. All Sport vectors contain an
5 ampicillin resistance gene and may be transformed into E. coli strain DH10B, also available from Life Technologies. (See, for instance, Gruber, C. E., et al., Focus 15:59 (1993).) Vector lafmid BA (Bento Soares, Columbia University, NY) contains an ampicillin resistance gene and can be transformed into E. coli strain XL-1 Blue. Vector pCR[®]2.1, which is available from Invitrogen, 1600 Faraday Avenue, Carlsbad, CA 92008, contains an
10 ampicillin resistance gene and may be transformed into E. coli strain DH10B, available from Life Technologies. (See, for instance, Clark, J. M., Nuc. Acids Res. 16:9677-9686 (1988) and Mead, D. et al., Bio/Technology 9: (1991).) Preferably, a polynucleotide of the present invention does not comprise the phage vector sequences identified for the particular clone in Table 5, as well as the corresponding plasmid vector sequences designated above.

15 The deposited material in the sample assigned the ATCC Deposit Number cited by reference to Table 2 and 5 for any given cDNA clone also may contain one or more additional plasmids, each comprising a cDNA clone different from that given clone. Thus, deposits sharing the same ATCC Deposit Number contain at least a plasmid for each cDNA clone referenced in Table 1.

TABLE 5

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HUKA HUKB HUKC HUKD HUKF HUKF HUKG	Human Uterine Cancer	Lambda ZAP II	LP01
HCNA HCNB	Human Colon	Lambda Zap II	LP01
HFFA	Human Fetal Brain, random primed	Lambda Zap II	LP01
HTWA	Resting T-Cell	Lambda ZAP II	LP01
HBQA	Early Stage Human Brain, random primed	Lambda ZAP II	LP01
HLMB HLMF HLMG HLMH HLMJ HLMJ HLMN HLMN	breast lymph node CDNA library	Lambda ZAP II	LP01
HCQA HCOB	human colon cancer	Lambda ZAP II	LP01
HMEA HMEC HMEF HMEF HMEF HMEG HMEI HMEJ HMEK HMEK HMEK HMEK	Human Microvascular Endothelial Cells, fract. A	Lambda ZAP II	LP01
HUSA HUSC	Human Umbilical Vein Endothelial Cells, fract. A	Lambda ZAP II	LP01
HLQA HLQB	Hepatocellular Tumor	Lambda ZAP II	LP01
HHGA HHGB HHGC HHGD	Hemangiopericytoma	Lambda ZAP II	LP01
HSDM	Human Striatum Depression, re-rescue	Lambda ZAP II	LP01
HUSH	H Umbilical Vein Endothelial Cells, fract. A, re-excision	Lambda ZAP II	LP01
HSGS	Salivary gland, subtracted	Lambda ZAP II	LP01
HFXA HFXB HFXC HFXD HFXE HFXF HFXG HFXH	Brain frontal cortex	Lambda ZAP II	LP01
HPQA HPQB HPQC	PERM TF274	Lambda ZAP II	LP01
HFXJ HFXK	Brain Frontal Cortex, re-excision	Lambda ZAP II	LP01
HCWA HCWB HCWC HCWD HCWE HCWF HCWG HCWH HCWI HCWJ HCWK	CD34 positive cells (Cord Blood)	ZAP Express	LP02
HCUA HCUB HCUC	CD34 depleted Buffy Coat (Cord Blood)	ZAP Express	LP02
HRSM	A-14 cell line	ZAP Express	LP02
HRSA	A1-CELL LINE	ZAP Express	LP02
HCUD HCUE HCUF HCUG HCUH HCUH HCUH	CD34 depleted Buffy Coat (Cord Blood), re-excision	ZAP Express	LP02
HBXE HBXF HBXG	H. Whole Brain #2, re-excision	ZAP Express	LP02
HRLM	L8 cell line	ZAP Express	LP02
HBXA HBXB HBXC HBXD	Human Whole Brain #2 - Oligo dT > 1.5Kb	ZAP Express	LP02
HUDA HUDB HUDC	Testes	ZAP Express	LP02
HHTM HHTN HHTO	H. hypothalamus, fract. A, re-excision	ZAP Express	LP02
HHTL	H. hypothalamus, fract. A	ZAP Express	LP02
HASA HASD	Human Adult Spleen	Uni-ZAP XR	LP03
HFKC HFKD HFKF HFKF HFKG	Human Fetal Kidney	Uni-ZAP XR	LP03
HE8A HE8B HE8C HE8D HE8E HE8F HE8M HE8N	Human 8 Week Whole Embryo	Uni-ZAP XR	LP03
HGBA HGBD HGBE HGBF HGBG HGBH HGBI	Human Gall Bladder	Uni-ZAP XR	LP03
HLHA HLHB HLHC HLHD HLHE	Human Fetal Lung III	Uni-ZAP XR	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HLHF HLHG HLHH HLHQ			
HPMA HPMB HPMC HPMD HPME HPMF HPMG HPMH	Human Placenta	Uni-ZAP XR	LP03
HPRA HPRB HPRC HPRD	Human Prostate	Uni-ZAP XR	LP03
HSIA HSIC HSID HSIE	Human Adult Small Intestine	Uni-ZAP XR	LP03
HTEA HTEB HTEC HTED HTEE HTEF HTEG HTEH HTEI HTEJ HTEK	Human Testes	Uni-ZAP XR	LP03
HTPA HTPB HTPC HTPD HTPF	Human Pancreas Tumor	Uni-ZAP XR	LP03
HTTA HTTB HTTC HTTD HTTE HTTF	Human Testes Tumor	Uni-ZAP XR	LP03
HAPA HAPB HAPC HAPM	Human Adult Pulmonary	Uni-ZAP XR	LP03
HETA HETB HETC HETD HETE HETF HETG HETH HETI	Human Endometrial Tumor	Uni-ZAP XR	LP03
HHFB HHFC HHFD HHFE HHFF HHFG HHFH HHFI	Human Fetal Heart	Uni-ZAP XR	LP03
HHPB HHPC HHPD HHPE HHPF HHPG HHPH	Human Hippocampus	Uni-ZAP XR	LP03
HCE1 HCE2 HCE3 HCE4 HCE5 HCEB HCEC HCED HCEE HCEF HCEG	Human Cerebellum	Uni-ZAP XR	LP03
HUVB HUVC HUVD HUVE	Human Umbilical Vein. Endo. remake	Uni-ZAP XR	LP03
HSTA HSTB HSTC HSTD	Human Skin Tumor	Uni-ZAP XR	LP03
HTAA HTAB HTAC HTAD HTAE	Human Activated T-Cells	Uni-ZAP XR	LP03
HFEA HFEB HFEC	Human Fetal Epithelium (Skin)	Uni-ZAP XR	LP03
HJPA HJPB HJPC HJPD	HUMAN JURKAT MEMBRANE BOUND POLYSOMES	Uni-ZAP XR	LP03
HESA	Human epithelioid sarcoma	Uni-Zap XR	LP03
HLTA HLTB HLTC HLTD HLTE HLTF	Human T-Cell Lymphoma	Uni-ZAP XR	LP03
HFTA HFTB HFTC HFTD	Human Fetal Dura Mater	Uni-ZAP XR	LP03
HRDA HRDB HRDC HRDD HRDE HRDF	Human Rhabdomyosarcoma	Uni-ZAP XR	LP03
HCAA HCAB HCAC	Cem cells cyclohexamide treated	Uni-ZAP XR	LP03
HRGA HRGB HRGC HRGD	Raji Cells. cyclohexamide treated	Uni-ZAP XR	LP03
HSUA HSUB HSUC HSUM	Supt Cells. cyclohexamide treated	Uni-ZAP XR	LP03
HT4A HT4C HT4D	Activated T-Cells. 12 hrs.	Uni-ZAP XR	LP03
HE9A HE9B HE9C HE9D HE9E HE9F HE9G HE9H HE9M HE9N	Nine Week Old Early Stage Human	Uni-ZAP XR	LP03
HATA HATB HATC HATD HATE	Human Adrenal Gland Tumor	Uni-ZAP XR	LP03
HT5A	Activated T-Cells. 24 hrs.	Uni-ZAP XR	LP03
HFGA HFGM	Human Fetal Brain	Uni-ZAP XR	LP03
HNEA HNEB HNEC HNED HNEE	Human Neutrophil	Uni-ZAP XR	LP03
HBGB HBGD	Human Primary Breast Cancer	Uni-ZAP XR	LP03
HBNA HBNB	Human Normal Breast	Uni-ZAP XR	LP03
HCAS	Cem Cells. cyclohexamide treated. subtra	Uni-ZAP XR	LP03
HHPS	Human Hippocampus. subtracted	pBS	LP03
HKCS HKCU	Human Colon Cancer. subtracted	pBS	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HRGS	Raji cells, cyclohexamide treated, subtracted	pBS	LP03
HSUT	Supt cells, cyclohexamide treated, differentially expressed	pBS	LP03
HT4S	Activated T-Cells, 12 hrs. subtracted	Uni-ZAP XR	LP03
HCDA HCDB HCDC HCDD HCDE	Human Chondrosarcoma	Uni-ZAP XR	LP03
HOAA HOAB HOAC	Human Osteosarcoma	Uni-ZAP XR	LP03
HTLA HTLB HTLC HTLD HTLE HTLF	Human adult testis, large inserts	Uni-ZAP XR	LP03
HLMA HLMB HLMD	Breast Lymph node cDNA library	Uni-ZAP XR	LP03
H6EA H6EB H6EC	HL-60, PMA 4H	Uni-ZAP XR	LP03
HTXA HTXB HTXC HTXD HTXE HTXF HTXG HTXH	Activated T-Cell (12hs)/Thiouridine labelledEco	Uni-ZAP XR	LP03
HNFA HNFB HNFC HNFD HNFE HNFF HNFG HNFH HNFJ	Human Neutrophil, Activated	Uni-ZAP XR	LP03
HTOB HTOC	HUMAN TONSILS, FRACTION 2	Uni-ZAP XR	LP03
HMGB	Human OB MG63 control fraction I	Uni-ZAP XR	LP03
HOPB	Human OB HOS control fraction I	Uni-ZAP XR	LP03
HORB	Human OB HOS treated (10 nM E2) fraction I	Uni-ZAP XR	LP03
HSVA HSVB HSVC	Human Chronic Synovitis	Uni-ZAP XR	LP03
HROA	HUMAN STOMACH	Uni-ZAP XR	LP03
HBJA HBJB HBJC HBJD HBJE HBJF HBJG HBJH HBJI HBJJ HBJK	HUMAN B CELL LYMPHOMA	Uni-ZAP XR	LP03
HCRA HCRB HCRC	human corpus colosum	Uni-ZAP XR	LP03
HODA HODB HODC HODD	human ovarian cancer	Uni-ZAP XR	LP03
HDSA	Dermatofibrosarcoma Protuberance	Uni-ZAP XR	LP03
HMWA HMWB HMWC HMWD HMWE HMWF HMWG HMWH HMWI HMWJ	Bone Marrow Cell Line (RS4;11)	Uni-ZAP XR	LP03
HSOA	stomach cancer (human)	Uni-ZAP XR	LP03
HERA	SKIN	Uni-ZAP XR	LP03
HMDA	Brain-medulloblastoma	Uni-ZAP XR	LP03
HGLA HGLB HGLD	Glioblastoma	Uni-ZAP XR	LP03
HEAA	H. Atrophic Endometrium	Uni-ZAP XR	LP03
HBCA HBCB	H. Lymph node breast Cancer	Uni-ZAP XR	LP03
HPWT	Human Prostate BPH, re-excision	Uni-ZAP XR	LP03
HFVG HFVH HFVI	Fetal Liver, subtraction II	pBS	LP03
HNFI	Human Neutrophils, Activated, re-excision	pBS	LP03
HBMB HBMC HBMD	Human Bone Marrow, re-excision	pBS	LP03
HKML HKMM HKMN	H. Kidney Medulla, re-excision	pBS	LP03
HKIX HKIY	H. Kidney Cortex, subtracted	pBS	LP03
HADT	H. Amygdala Depression, subtracted	pBS	LP03
H6AS	HL-60, untreated, subtracted	Uni-ZAP XR	LP03
H6ES	HL-60, PMA 4H, subtracted	Uni-ZAP XR	LP03
H6BS	HL-60, RA 4h, Subtracted	Uni-ZAP XR	LP03

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
H6CS	HL-60, PMA 1d. subtracted	Uni-ZAP XR	LP03
HTXJ HTXK	Activated T-cell(12h)/Thiouridine-re-excision	Uni-ZAP XR	LP03
HMSA HMSB HMSC HMSE HMSF HMSG HMSH HMSI HMSJ HMSK	Monocyte activated	Uni-ZAP XR	LP03
HAGA HAGB HAGC HAGD HAGE HAGF	Human Amygdala	Uni-ZAP XR	LP03
HSRA HSRB HSRE	STROMAL -OSTEOCLASTOMA	Uni-ZAP XR	LP03
HSRD HSRF HSRG HSRH	Human Osteoclastoma Stromal Cells - unamplified	Uni-ZAP XR	LP03
HSQA HSQB HSQC HSQD HSQE HSQF HSQG	Stromal cell TF274	Uni-ZAP XR	LP03
HSKA HSKB HSKC HSKD HSKE HSKF HSKZ	Smooth muscle, serum treated	Uni-ZAP XR	LP03
HSLA HSLB HSLC HSLD HSLE HSLF HSLG	Smooth muscle control	Uni-ZAP XR	LP03
HSDA HSDD HSDE HSDF HSDG HSDH	Spinal cord	Uni-ZAP XR	LP03
HPWS	Prostate-BPH subtracted II	pBS	LP03
HSKV HSKX HSKY	Smooth Muscle- HASTE normalized	pBS	LP03
HFPB HFPC HFPD	H. Frontal cortex.epileptic:re-excision	Uni-ZAP XR	LP03
HSDI HSDJ HSDK	Spinal Cord, re-excision	Uni-ZAP XR	LP03
HSKN HSKO	Smooth Muscle Serum Treated, Norm	pBS	LP03
HSKG HSKH HSKI	Smooth muscle, serum induced.re-exc	pBS	LP03
HFCA HFCB HFCC HFCD HFCE HFCF	Human Fetal Brain	Uni-ZAP XR	LP04
HPTA HPTB HPTD	Human Pituitary	Uni-ZAP XR	LP04
HTHB HTHC HTHD	Human Thymus	Uni-ZAP XR	LP04
HE6B HE6C HE6D HE6E HE6F HE6G HE6S	Human Whole Six Week Old Embryo	Uni-ZAP XR	LP04
HSSA HSSB HSSC HSSD HSSE HSSF HSSG HSSH HSSI HSSJ HSSK HE7T	Human Synovial Sarcoma	Uni-ZAP XR	LP04
HEPA HEPB HEPC	7 Week Old Early Stage Human, subtracted	Uni-ZAP XR	LP04
HSNA HSNB HSNB HSNM HSNM	Human Epididymus	Uni-ZAP XR	LP04
HPFB HPFC HPFD HPFE	Human Synovium	Uni-ZAP XR	LP04
HE2A HE2D HE2E HE2H HE2I HE2M HE2N HE2O	Human Prostate Cancer, Stage C fraction	Uni-ZAP XR	LP04
HE2B HE2C HE2F HE2G HE2P HE2Q	12 Week Old Early Stage Human	Uni-ZAP XR	LP04
HPTS HPTT HPTU	12 Week Old Early Stage Human, II	Uni-ZAP XR	LP04
HAUA HAUB HAUC	Human Pituitary, subtracted	Uni-ZAP XR	LP04
HAQA HAQB HAQC HAQD	Amniotic Cells - TNF induced	Uni-ZAP XR	LP04
HWTA HWTB HWTC	Amniotic Cells - Primary Culture	Uni-ZAP XR	LP04
HBSD	wilm's tumor	Uni-ZAP XR	LP04
HSGB	Bone Cancer, re-excision	Uni-ZAP XR	LP04
HSJA HSJB HSJC	Salivary gland, re-excision	Uni-ZAP XR	LP04
HSXA HSXB HSXC HSXD	Smooth muscle-ILb induced	Uni-ZAP XR	LP04
	Human Substantia Nigra	Uni-ZAP XR	LP04

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HSHA HSHB HSHC	Smooth muscle, IL1b induced	Uni-ZAP XR	LP04
HOUA HOUB HOU C HOUD HOUE	Adipocytes	Uni-ZAP XR	LP04
HPWA HPWB HPWC HPWD HPWE	Prostate BPH	Uni-ZAP XR	LP04
HELA HELB HELC HELD HELE HELF HELG HELH	Endothelial cells-control	Uni-ZAP XR	LP04
HEMA HEMB HEMC HEMD HEME HEMF HEMG HEMH	Endothelial-induced	Uni-ZAP XR	LP04
HBIA HBIB HBIC	Human Brain, Striatum	Uni-ZAP XR	LP04
HSHA HSHB HHSC HHSD HHSE	Human Hypothalamus, Schizophrenia	Uni-ZAP XR	LP04
HNGA HNGB HNGC HNGD HNGE HNGF HNGG HNGH HNGI HNGJ	neutrophils control	Uni-ZAP XR	LP04
HNHA HNHB HNHC HNHD HNHE HNHF HNHG HNHH HNHI HNHI	Neutrophils IL-1 and LPS induced	Uni-ZAP XR	LP04
HSDB HSDC	STRIATUM DEPRESSION	Uni-ZAP XR	LP04
HHPT	Hypothalamus	Uni-ZAP XR	LP04
HSAT HSAU HSAV HSAW HSAX HSAY HSAZ	Anergic T-cell	Uni-ZAP XR	LP04
HBMS HBMT HBMU HBMV HBMW HBMX	Bone marrow	Uni-ZAP XR	LP04
HOEA HOEB HOEC HOED HOEE HOEF HOEJ	Osteoblasts	Uni-ZAP XR	LP04
HAIA HAIB HAIC HAID HAIE HAIF	Epithelial-TNF α and INF induced	Uni-ZAP XR	LP04
HTGA HTGB HTGC HTGD	Apoptotic T-cell	Uni-ZAP XR	LP04
HMCA HMCB HMCC HMCD HMCE	Macrophage-oxLDL	Uni-ZAP XR	LP04
HMAA HMA B HMA C HMA D HMAE HMAF HMA G	Macrophage (GM-CSF treated)	Uni-ZAP XR	LP04
HPHA	Normal Prostate	Uni-ZAP XR	LP04
HPIA HPIB HPIC	LNCAP prostate cell line	Uni-ZAP XR	LP04
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP04
HOSE HOSF HOSG	Human Osteoclastoma, re-excision	Uni-ZAP XR	LP04
HTGE HTGF	Apoptotic T-cell, re-excision	Uni-ZAP XR	LP04
HMAJ HMAK	H Macrophage (GM-CSF treated), re-excision	Uni-ZAP XR	LP04
HACB HACC HACD	Human Adipose Tissue, re-excision	Uni-ZAP XR	LP04
HFPA	H. Frontal Cortex, Epileptic	Uni-ZAP XR	LP04
HFAA HFAB HFAC HFAD HFAE	Alzheimers, spongy change	Uni-ZAP XR	LP04
HFAM	Frontal Lobe, Dementia	Uni-ZAP XR	LP04
HMIA HMIB HMIC	Human Manic Depression Tissue	Uni-ZAP XR	LP04
HTSA HTSE HTSF HTSG HTSH	Human Thymus	pBS	LP05
HPBA HPBB HPBC HPBD HPBE	Human Pineal Gland	pBS	LP05
HSAA HSAB HSAC	HSA 172 Cells	pBS	LP05
HSBA HSBB HSBC HSBM	HSC172 cells	pBS	LP05
HJAA HJAB HJAC HJAD	Jurkat T-cell G1 phase	pBS	LP05
HJBA HJBB HJBC HJBD	Jurkat T-Cell, S phase	pBS	LP05
HAFA HAFB	Aorta endothelial cells + TNF- α	pBS	LP05
HAWA HAWB HAWC	Human White Adipose	pBS	LP05
HTNA HTNB	Human Thyroid	pBS	LP05

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HONA	Normal Ovary, Premenopausal	pBS	LP05
HARA HARB	Human Adult Retina	pBS	LP05
HLJA HLJB	Human Lung	pCMVSPORT 1	LP06
HOFM HOFN HOFO	H. Ovarian Tumor, II, OV5232	pCMVSPORT 2.0	LP07
HOGA HOGB HOGC	OV 10-3-95	pCMVSPORT 2.0	LP07
HCGL	CD34+cells, II	pCMVSPORT 2.0	LP07
HDLA	Hodgkin's Lymphoma I	pCMVSPORT 2.0	LP07
HDTA HDTB HDTC HDTD HDTE	Hodgkin's Lymphoma II	pCMVSPORT 2.0	LP07
HKAA HKAB HKAC HKAD HKAE HKAF HKAG HKAH	Keratinocyte	pCMVSPORT2.0	LP07
HCIM	CAPFINDER, Crohn's Disease, lib 2	pCMVSPORT 2.0	LP07
HKAL	Keratinocyte, lib 2	pCMVSPORT2.0	LP07
HKAT	Keratinocyte, lib 3	pCMVSPORT2.0	LP07
HNDA	Nasal polyps	pCMVSPORT2.0	LP07
HDRA	H. Primary Dendritic Cells.lib 3	pCMVSPORT2.0	LP07
HOHA HOHB HOHC	Human Osteoblasts II	pCMVSPORT2.0	LP07
HLDA HLDB HLDC	Liver, Hepatoma	pCMVSPORT3.0	LP08
HLDN HLDO HLDP	Human Liver, normal	pCMVSPORT3.0	LP08
HMTA	pBMC stimulated w/ poly I/C	pCMVSPORT3.0	LP08
HNTA	NTERA2, control	pCMVSPORT3.0	LP08
HDP A HDPB HDPC HDPD HDPE HDPG HDPI HDPI HDPI HDPI HDPI	Primary Dendritic Cells, lib 1	pCMVSPORT3.0	LP08
HDPM HDPN HDPO HDPP	Primary Dendritic cells.frac 2	pCMVSPORT3.0	LP08
HMUA HMUB HMUC	Myeloid Progenitor Cell Line	pCMVSPORT3.0	LP08
HHEA HHEB HHEC HHED	T Cell helper I	pCMVSPORT3.0	LP08
HHEM HHEN HHEO HHEP	T cell helper II	pCMVSPORT3.0	LP08
HEQA HEQB HEQC	Human endometrial stromal cells	pCMVSPORT3.0	LP08
HJMA HJMB	Human endometrial stromal cells-treated with progesterone	pCMVSPORT3.0	LP08
HSWA HSWB HSWC	Human endometrial stromal cells-treated with estradiol	pCMVSPORT3.0	LP08
HSYA HSYB HSYC	Human Thymus Stromal Cells	pCMVSPORT3.0	LP08
HLWA HLWB HLWC	Human Placenta	pCMVSPORT3.0	LP08
HRAA HRAB HRAC	Rejected Kidney, lib 4	pCMVSPORT3.0	LP08
HMTM	PCR, pBMC I/C treated	PCR11	LP09
HMJA	H. Meningioma, M6	pSport 1	LP10
HMKA HMKB HMKC HMKD HMKE	H. Meningioma, M1	pSport 1	LP10
HUSG HUSI	Human umbilical vein endothelial cells, IL-4 induced	pSport 1	LP10
HUSX HUSY	Human Umbilical Vein Endothelial Cells, uninduced	pSport 1	LP10
HOFA	Ovarian Tumor I, OV5232	pSport 1	LP10
HCFA HCFB HCFC HCFC	T-Cell PHA 16 hrs	pSport 1	LP10
HCFL HCFM HCFN HCFO	T-Cell PHA 24 hrs	pSport 1	LP10
HADA HADC HADD HADE HADF HADG	Human Adipose	pSport 1	LP10

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HOVA HOVB HOVC	Human Ovary	pSport 1	LP10
HTWB HTWC HTWD HTWE HTWF	Resting T-Cell Library.II	pSport 1	LP10
HMMA	Spleen metastatic melanoma	pSport 1	LP10
HLYA HLYB HLYC HLYD HLYE	Spleen, Chronic lymphocytic leukemia	pSport 1	LP10
HCGA	CD34+ cell. I	pSport 1	LP10
HEOM HEON	Human Eosinophils	pSport 1	LP10
HTDA	Human Tonsil, Lib 3	pSport 1	LP10
HSPA	Salivary Gland, Lib 2	pSport 1	LP10
HCHA HCHB HCHC	Breast Cancer cell line, MDA 36	pSport 1	LP10
HCHM HCHN	Breast Cancer Cell line, angiogenic	pSport 1	LP10
HCIA	Crohn's Disease	pSport 1	LP10
HDAA HDAB HDAC	HEL cell line	pSport 1	LP10
HABA	Human Astrocyte	pSport 1	LP10
HUFA HUFB HUFC	Ulcerative Colitis	pSport 1	LP10
HNTM	NTERA2 + retinoic acid, 14 days	pSport 1	LP10
HDQA	Primary Dendritic cells.CapFinder2, frac 1	pSport 1	LP10
HDQM	Primary Dendritic Cells, CapFinder, frac 2	pSport 1	LP10
HLDX	Human Liver, normal.CapFinder	pSport 1	LP10
HULA HULB HULC	Human Dermal Endothelial Cells,untreated	pSport 1	LP10
HUMA	Human Dermal Endothelial cells,treated	pSport 1	LP10
HCJA	Human Stromal Endometrial fibroblasts, untreated	pSport 1	LP10
HCJM	Human Stromal endometrial fibroblasts, treated w/ estradiol	pSport 1	LP10
HEDA	Human Stromal endometrial fibroblasts, treated with progesterone	pSport 1	LP10
HFNA	Human ovary tumor cell OV350721	pSport 1	LP10
HKGA HKGB HKGC HKGD	Merkel Cells	pSport 1	LP10
HISA HISB HISC	Pancreas Islet Cell Tumor	pSport 1	LP10
HLSA	Skin, burned	pSport 1	LP10
HBZA	Prostate,BPH, Lib 2	pSport 1	LP10
HBZS	Prostate BPH,Lib 2, subtracted	pSport 1	LP10
HFIA HFIB HFIC	Synovial Fibroblasts (control)	pSport 1	LP10
HFIH HFII HFIJ	Synovial hypoxia	pSport 1	LP10
HFIT HFIU HFIV	Synovial IL-1/TNF stimulated	pSport 1	LP10
HGCA	Mesangial cell, frac 1	pSport 1	LP10
HMVA HMVB HMVC	Bone Marrow Stromal Cell, untreated	pSport 1	LP10
HFIX HFY HFIZ	Synovial Fibroblasts (III/TNF), subt	pSport 1	LP10
HFOX HFOY HFOZ	Synovial hypoxia-RSF subtracted	pSport 1	LP10
HMQA HMQB HMQC HMQD	Human Activated Monocytes	Uni-ZAP XR	LP11
HLIA HLIB HLIC	Human Liver	pCMVSPORT 1	LP012
HHBA HHBB HHBC HHBD HHBE	Human Heart	pCMVSPORT 1	LP012
HBBA HBBB	Human Brain	pCMVSPORT 1	LP012

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HLJA HLJB HLJC HLJD HLJE	Human Lung	pCMVSPORT 1	LP012
HOGA HOGB HOGC	Ovarian Tumor	pCMVSPORT 2.0	LP012
HTJM	Human Tonsils. Lib 2	pCMVSPORT 2.0	LP012
HAMF HAMG	KMH2	pCMVSPORT 3.0	LP012
HAJA HAJB HAJC	L428	pCMVSPORT 3.0	LP012
HWBA HWBB HWBC HWBD HWBE	Dendritic cells. pooled	pCMVSPORT 3.0	LP012
HWAA HWAB HWAC HWAD HWAE	Human Bone Marrow. treated	pCMVSPORT 3.0	LP012
HYAA HYAB HYAC	B Cell lymphoma	pCMVSPORT 3.0	LP012
HWHG HWHH HWHI	Healing groin wound, 6.5 hours post incision	pCMVSPORT 3.0	LP012
HWHP HWHQ HWHR	Healing groin wound: 7.5 hours post incision	pCMVSPORT 3.0	LP012
HARM	Healing groin wound - zero hr post-incision (control)	pCMVSPORT 3.0	LP012
HBIM	Olfactory epithelium: nasal cavity	pCMVSPORT 3.0	LP012
HWDA	Healing Abdomen wound: 70&90 min post incision	pCMVSPORT 3.0	LP012
HWEA	Healing Abdomen Wound:15 days post incision	pCMVSPORT 3.0	LP012
HWJA	Healing Abdomen Wound:21&29 days	pCMVSPORT 3.0	LP012
HNAL	Human Tongue. frac 2	pSport1	LP012
HMJA	H. Meningioma, M6	pSport1	LP012
HMKA HMKB HMKC HMKE	H. Meningioma, M1	pSport1	LP012
HOFA	Ovarian Tumor I. OV5232	pSport1	LP012
HCFA HCFB HCFC HCFD	T-Cell PHA 16 hrs	pSport1	LP012
HCFL HCFM HCFN HCFO	T-Cell PHA 24 hrs	pSport1	LP012
HMMA HMMB HMMC	Spleen metastatic melanoma	pSport1	LP012
HTDA	Human Tonsil. Lib 3	pSport1	LP012
HDBA	Human Fetal Thymus	pSport1	LP012
HDLA	Pericardium	pSport1	LP012
HBZA	Prostate.BPH. Lib 2	pSport1	LP012
HWCA	Larynx tumor	pSport1	LP012
HWKA	Normal lung	pSport1	LP012
HSMB	Bone marrow stroma.treated	pSport1	LP012
HBHM	Normal trachea	pSport1	LP012
HLFC	Human Larynx	pSport1	LP012
HLRB	Siebben Polyposis	pSport1	LP012
HNIA	Mammary Gland	pSport1	LP012
HNJB	Palate carcinoma	pSport1	LP012
HNKA	Palate normal	pSport1	LP012
HMZA	Pharynx carcinoma	pSport1	LP012
HABG	Cheek Carcinoma	pSport1	LP012
HMZM	Pharynx Carcinoma	pSport1	LP012
HDRM	Larynx Carcinoma	pSport1	LP012
HVAA	Pancreas normal PCA4 No	pSport1	LP012
HICA	Tongue carcinoma	pSport1	LP012

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HUKA HUKB HUKC HUKD HUK E	Human Uterine Cancer	Lambda ZAP II	LP013
HFFA	Human Fetal Brain, random primed	Lambda ZAP II	LP013
HTUA	Activated T-cell labeled with 4-thiour	Lambda ZAP II	LP013
HBQA	Early Stage Human Brain, random primed	Lambda ZAP II	LP013
HMEB	Human microvascular Endothelial cells, fract. B	Lambda ZAP II	LP013
HUSH	Human Umbilical Vein Endothelial cells, fract. A, re-excision	Lambda ZAP II	LP013
HLOC HLOD	Hepatocellular tumor, re-excision	Lambda ZAP II	LP013
HTWJ HTWK HTWL	Resting T-cell, re-excision	Lambda ZAP II	LP013
HF6S	Human Whole 6 week Old Embryo (II), subt	pBluescript	LP013
HHPS	Human Hippocampus, subtracted	pBluescript	LP013
HLIS	LNAP, differential expression	pBluescript	LP013
HLHS HLHT	Early Stage Human Lung, Subtracted	pBluescript	LP013
HSUS	Supt cells, cyclohexamide treated, subtracted	pBluescript	LP013
HSUT	Supt cells, cyclohexamide treated, differentially expressed	pBluescript	LP013
HSDS	H. Striatum Depression, subtracted	pBluescript	LP013
HPTZ	Human Pituitary, Subtracted VII	pBluescript	LP013
HSDX	H. Striatum Depression, subt II	pBluescript	LP013
HSDZ	H. Striatum Depression, subt	pBluescript	LP013
HPBA HPBB HPBC HPBD HPBE	Human Pineal Gland	pBluescript SK-	LP013
HRTA	Colorectal Tumor	pBluescript SK-	LP013
HSBA HSBB HSBC HSBM	HSC172 cells	pBluescript SK-	LP013
HJAA HJAB HJAC HJAD	Jurkat T-cell G1 phase	pBluescript SK-	LP013
HJBA HJBB HJBC HJBD	Jurkat T-cell, S1 phase	pBluescript SK-	LP013
HTNA HTNB	Human Thyroid	pBluescript SK-	LP013
HAHA HAHB	Human Adult Heart	Uni-ZAP XR	LP013
HE6A	Whole 6 week Old Embryo	Uni-ZAP XR	LP013
HFCA HFCB HFCC HFCD HFCE	Human Fetal Brain	Uni-ZAP XR	LP013
HFKE HFKE HFKE HFKE HFKE	Human Fetal Kidney	Uni-ZAP XR	LP013
HGBA HGBD HGBE HGBF HGBG	Human Gall Bladder	Uni-ZAP XR	LP013
HPRA HPRB HPRC HPRD	Human Prostate	Uni-ZAP XR	LP013
HTEA HTEB HTEC HTED HTEE	Human Testes	Uni-ZAP XR	LP013
HTTA HTTB HTTC HTTD HTTE	Human Testes Tumor	Uni-ZAP XR	LP013
HYBA HYBB	Human Fetal Bone	Uni-ZAP XR	LP013
HFLA	Human Fetal Liver	Uni-ZAP XR	LP013
HHFB HHFC HHFD HHFE HHFF	Human Fetal Heart	Uni-ZAP XR	LP013
HUVB HUVB HUVB HUVB	Human Umbilical Vein, End. remake	Uni-ZAP XR	LP013
HTHB HTHC HTHD	Human Thymus	Uni-ZAP XR	LP013
HSTA HSTB HSTC HSTD	Human Skin Tumor	Uni-ZAP XR	LP013
HTAA HTAB HTAC HTAD HTAE	Human Activated T-cells	Uni-ZAP XR	LP013
HFEA HFEB HFEC	Human Fetal Epithelium (skin)	Uni-ZAP XR	LP013
HJPA HJPB HJPC HJPD	Human Jurkat Membrane Bound Polysomes	Uni-ZAP XR	LP013
HESA	Human Epithelioid Sarcoma	Uni-ZAP XR	LP013

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HALS	Human Adult Liver. Subtracted	Uni-ZAP XR	LP013
HFTA HFTB HFTC HFTD	Human Fetal Dura Mater	Uni-ZAP XR	LP013
HCAA HCAB HCAC	Cem cells. cyclohexamide treated	Uni-ZAP XR	LP013
HRGA HRGB HRGC HRGD	Raji Cells. cyclohexamide treated	Uni-ZAP XR	LP013
HE9A HE9B HE9C HE9D HE9E	Nine Week Old Early Stage Human	Uni-ZAP XR	LP013
HSFA	Human Fibrosarcoma	Uni-ZAP XR	LP013
HATA HATB HATC HATD HATE	Human Adrenal Gland Tumor	Uni-ZAP XR	LP013
HTRA	Human Trachea Tumor	Uni-ZAP XR	LP013
HE2A HE2D HE2E HE2H HE2I	12 Week Old Early Stage Human	Uni-ZAP XR	LP013
HE2B HE2C HE2F HE2G HE2P	12 Week Old Early Stage Human. II	Uni-ZAP XR	LP013
HNEA HNEB HNEC HNED HNEE	Human Neutrophil	Uni-ZAP XR	LP013
HBGA	Human Primary Breast Cancer	Uni-ZAP XR	LP013
HPTS HPTT HPTU	Human Pituitary. subtracted	Uni-ZAP XR	LP013
HMQA HMQB HMQC HMQD	Human Activated Monocytes	Uni-ZAP XR	LP013
HOAA HOAB HOAC	Human Osteosarcoma	Uni-ZAP XR	LP013
HTOA HTOD HTOE HTOF HTOG	human tonsils	Uni-ZAP XR	LP013
HMGB	Human OB MG63 control fraction I	Uni-ZAP XR	LP013
HOPB	Human OB HOS control fraction I	Uni-ZAP XR	LP013
HOQB	Human OB HOS treated (1 nM E2) fraction I	Uni-ZAP XR	LP013
HAUA HAUB HAUC	Amniotic Cells - TNF induced	Uni-ZAP XR	LP013
HAQA HAQB HAQC HAQD	Amniotic Cells - Primary Culture	Uni-ZAP XR	LP013
HROA HROC	HUMAN STOMACH	Uni-ZAP XR	LP013
HBJA HBJB HBJC HBJD HBJE	HUMAN B CELL LYMPHOMA	Uni-ZAP XR	LP013
HODA HODB HODC HODD	human ovarian cancer	Uni-ZAP XR	LP013
HCPA	Corpus Callosum	Uni-ZAP XR	LP013
HSOA	stomach cancer (human)	Uni-ZAP XR	LP013
HERA	SKIN	Uni-ZAP XR	LP013
HMDA	Brain-medulloblastoma	Uni-ZAP XR	LP013
HGLA HGLB HGLD	Glioblastoma	Uni-ZAP XR	LP013
HWTA HWTB HWTC	wilm's tumor	Uni-ZAP XR	LP013
HEAA	H. Atrophic Endometrium	Uni-ZAP XR	LP013
HAPN HAPO HAPP HAQO HAPR	Human Adult Pulmonary:re-excision	Uni-ZAP XR	LP013
HLTG HLTH	Human T-cell lymphoma:re-excision	Uni-ZAP XR	LP013
HAHC HAHD HAHE	Human Adult Heart:re-excision	Uni-ZAP XR	LP013
HAGA HAGB HAGC HAGD HAGE	Human Amygdala	Uni-ZAP XR	LP013
HSJA HSJB HSJC	Smooth muscle-ILb induced	Uni-ZAP XR	LP013
HSJA HSHB HSHC	Smooth muscle, IL1b induced	Uni-ZAP XR	LP013
HPWA HPWB HPWC HPWD HPWE	Prostate BPH	Uni-ZAP XR	LP013
HPJA HPJB HPJC	LNCAP prostate cell line	Uni-ZAP XR	LP013
HPJA HPJB HPJC	PC3 Prostate cell line	Uni-ZAP XR	LP013
HBTA	Bone Marrow Stroma. TNF&LPS ind	Uni-ZAP XR	LP013
HMCJ HMCG HMCH HMCJ HMCJ	Macrophage-oxLDL: re-excision	Uni-ZAP XR	LP013
HAGG HAGH HAGI	Human Amygdala:re-excision	Uni-ZAP XR	LP013
HACA	H. Adipose Tissue	Uni-ZAP XR	LP013
HKFB	K562 + PMA (36 hrs).re-excision	ZAP Express	LP013
HCWT HCWU HCWV	CD34 positive cells (cord blood).re-ex	ZAP Express	LP013

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HBWA	Whole brain	ZAP Express	LP013
HBXA HBXB HBXC HBXD	Human Whole Brain #2 - Oligo dT > 1.5Kb	ZAP Express	LP013
HAVM	Temporal cortex-Alzheimer	pT-Adv	LP014
HAVT	Hippocampus, Alzheimer Subtracted	pT-Adv	LP014
HHAS	CHME Cell Line	Uni-ZAP XR	LP014
HAJR	Larynx normal	pSport 1	LP014
HWLE HWLF HWLG HWLH	Colon Normal	pSport 1	LP014
HCRM HCRN HCRO	Colon Carcinoma	pSport 1	LP014
HWLI HWLJ HWLK	Colon Normal	pSport 1	LP014
HWLQ HWLR HWLS HWLT	Colon Tumor	pSport 1	LP014
HBFM	Gastrocnemius Muscle	pSport 1	LP014
HBOD HBOE	Quadriceps Muscle	pSport 1	LP014
HBKD HBKE	Soleus Muscle	pSport 1	LP014
HCCM	Pancreatic Langerhans	pSport 1	LP014
HWGA	Larynx carcinoma	pSport 1	LP014
HWGM HWGN	Larynx carcinoma	pSport 1	LP014
HWLA HWLB HWLC	Normal colon	pSport 1	LP014
HWLM HWLN	Colon Tumor	pSport 1	LP014
HVAM HVAN HVAO	Pancreas Tumor	pSport 1	LP014
HWGO	Larynx carcinoma	pSport 1	LP014
HAQM HAQN	Salivary Gland	pSport 1	LP014
HASM	Stomach; normal	pSport 1	LP014
HBCM	Uterus; normal	pSport 1	LP014
HCDM	Testis; normal	pSport 1	LP014
HDJM	Brain; normal	pSport 1	LP014
HEFM	Adrenal Gland; normal	pSport 1	LP014
HBAA	Rectum normal	pSport 1	LP014
HFDM	Rectum tumour	pSport 1	LP014
HGAM	Colon, normal	pSport 1	LP014
HHMM	Colon, tumour	pSport 1	LP014
HCLB HCLC	Human Lung Cancer	Lambda Zap II	LP015
HRLA	L1 Cell line	ZAP Express	LP015
HHAM	Hypothalamus, Alzheimer's	pCMVSPORT 3.0	LP015
HKBA	Ku 812F Basophils Line	pSport 1	LP015
HS2S	Saos2, Dexamethosone Treated	pSport 1	LP016
HA5A	Lung Carcinoma A549 TNFalpha activated	pSport 1	LP016
HTFM	TF-1 Cell Line GM-CSF Treated	pSport 1	LP016
HYAS	Thyroid Tumour	pSport 1	LP016
HUTS	Larynx Normal	pSport 1	LP016
HXOA	Larynx Tumor	pSport 1	LP016
HEAH	Ea.hy.926 cell line	pSport 1	LP016
HINA	Adenocarcinoma Human	pSport 1	LP016
HRMA	Lung Mesothelium	pSport 1	LP016
HLCL	Human Pre-Differentiated Adipocytes	Uni-Zap XR	LP017
HS2A	Saos2 Cells	pSport 1	LP020

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HS2I	Saos2 Cells: Vitamin D3 Treated	pSport 1	LP020
HUCM	CHME Cell Line, untreated	pSport 1	LP020
HEPN	Aryepiglottis Normal	pSport 1	LP020
HPSN	Sinus Piniformis Tumour	pSport 1	LP020
HNSA	Stomach Normal	pSport 1	LP020
HNSM	Stomach Tumour	pSport 1	LP020
HNLA	Liver Normal Met5No	pSport 1	LP020
HUTA	Liver Tumour Met 5 Tu	pSport 1	LP020
HOCN	Colon Normal	pSport 1	LP020
HOCT	Colon Tumor	pSport 1	LP020
HTNT	Tongue Tumour	pSport 1	LP020
HLXN	Larynx Normal	pSport 1	LP020
HLXT	Larynx Tumour	pSport 1	LP020
HTYN	Thymus	pSport 1	LP020
HPLN	Placenta	pSport 1	LP020
HTNG	Tongue Normal	pSport 1	LP020
HZAA	Thyroid Normal (SDCA2 No)	pSport 1	LP020
HWES	Thyroid Thyroiditis	pSport 1	LP020
HFHD	Ficoll Human Stromal Cells, 5Fu treated	pTriplEx2	LP021
HFHM, HFHN	Ficoll Human Stromal Cells, Untreated	pTriplEx2	LP021
HPCI	Hep G2 Cells, lambda library	lambda Zap-CMV XR	LP021
HBCA, HBCB, HBCC	H. Lymph node breast Cancer	Uni-ZAP XR	LP021
HCOK	Chondrocytes	pSPORT1	LP022
HDCA, HDCB, HDCC	Dendritic Cells From CD34 Cells	pSPORT1	LP022
HDMA, HDMB	CD40 activated monocyte dendritic cells	pSPORT1	LP022
HDDM, HDDN, HDDO	LPS activated derived dendritic cells	pSPORT1	LP022
HPCR	Hep G2 Cells, PCR library	lambda Zap-CMV XR	LP022
HAAA, HAAB, HAAC	Lung, Cancer (4005313 A3): Invasive Poorly Differentiated Lung Adenocarcinoma	pSPORT1	LP022
HIPB, HIPB, HIPC	Lung, Cancer (4005163 B7): Invasive, Poorly Diff. Adenocarcinoma, Metastatic	pSPORT1	LP022
HOOH, HOOI	Ovary, Cancer: (4004562 B6) Papillary Serous Cystic Neoplasm, Low Malignant Pot	pSPORT1	LP022
HIDA	Lung, Normal: (4005313 B1)	pSPORT1	LP022
HUJA, HUJB, HUJC, HUJD, HUJE	B-Cells	pCMVSPORT 3.0	LP022
HNOA, HNOB, HNOG, HNOD	Ovary, Normal: (9805C040R)	pSPORT1	LP022
HNLM	Lung, Normal: (4005313 B1)	pSPORT1	LP022
HSCL	Stromal Cells	pSPORT1	LP022
HAAX	Lung, Cancer: (4005313 A3) Invasive Poorly-differentiated Metastatic lung adenocarcinoma	pSPORT1	LP022
HUUA, HUUB, HUUC, HUUD	B-cells (unstimulated)	pTriplEx2	LP022
HWWA, HWWB, HWWC, HWWD, HWE, HWWF, HWWG	B-cells (stimulated)	pSPORT1	LP022
HCCC	Colon, Cancer: (9808C064R)	pCMVSPORT 3.0	LP023

Libraries owned by Catalog	Catalog Description	Vector	ATCC Deposit
HPDO HPDP HPDQ HPDR HPD	Ovary, Cancer (9809C332): Poorly differentiated adenocarcinoma	pSport 1	LP023
HPCO HPCP HPCQ HPCT	Ovary, Cancer (15395A1F): Grade II Papillary Carcinoma	pSport 1	LP023
HOCM HOCO HOCQ HOCQ	Ovary, Cancer: (15799A1F) Poorly differentiated carcinoma	pSport 1	LP023
HCBM HCBN HCBO	Breast, Cancer: (4004943 A5)	pSport 1	LP023
HNBT HNBU HNBV	Breast, Normal: (4005522B2)	pSport 1	LP023
HBCP HBCQ	Breast, Cancer: (4005522 A2)	pSport 1	LP023
HBCJ	Breast, Cancer: (9806C012R)	pSport 1	LP023
HSAM HSAN	Stromal cells 3.88	pSport 1	LP023
HVCA HV'CB HVCC HVCD	Ovary, Cancer: (4004332 A2)	pSport 1	LP023
HSCK HSEN HSEO	Stromal cells (HBM3.18)	pSport 1	LP023
HSCP HSCQ	stromal cell clone 2.5	pSport 1	LP023
HUXA	Breast Cancer: (4005385 A2)	pSport 1	LP023
HCOM HCON HCOO HCOP HCOQ	Ovary, Cancer (4004650 A3): Well-Differentiated Micropapillary Serous Carcinoma	pSport 1	LP023
HBNM	Breast, Cancer: (9802C020E)	pSport 1	LP023
HVVA HV'VB HVVC HVVD HVVE	Human Bone Marrow, treated	pSport 1	LP023

Two approaches can be used to isolate a particular clone from the deposited sample of plasmid DNAs cited for that clone in Table 5. First, a plasmid is directly isolated by screening the clones using a polynucleotide probe corresponding to the nucleotide sequence of SEQ ID NO:X.

5 Particularly, a specific polynucleotide with 30-40 nucleotides is synthesized using an Applied Biosystems DNA synthesizer according to the sequence reported. The oligonucleotide is labeled, for instance, with ^{32}P - γ -ATP using T4 polynucleotide kinase and purified according to routine methods. (E.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring, NY (1982).) The plasmid
10 mixture is transformed into a suitable host, as indicated above (such as XL-1 Blue (Stratagene)) using techniques known to those of skill in the art, such as those provided by the vector supplier or in related publications or patents cited above. The transformants are plated on 1.5% agar plates (containing the appropriate selection agent, e.g., ampicillin) to a density of about 150 transformants (colonies) per plate. These plates are screened using
15 Nylon membranes according to routine methods for bacterial colony screening (e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Edit., (1989), Cold Spring Harbor Laboratory Press, pages 1.93 to 1.104), or other techniques known to those of skill in the art.

 Alternatively, two primers of 17-20 nucleotides derived from both ends of the
20 nucleotide sequence of SEQ ID NO:X are synthesized and used to amplify the desired cDNA using the deposited cDNA plasmid as a template. The polymerase chain reaction is carried out under routine conditions, for instance, in 25 μl of reaction mixture with 0.5 μg of the above cDNA template. A convenient reaction mixture is 1.5-5 mM MgCl_2 , 0.01% (w/v) gelatin, 20 μM each of dATP, dCTP, dGTP, dTTP, 25 pmol of each primer and 0.25 Unit of
25 Taq polymerase. Thirty five cycles of PCR (denaturation at 94°C for 1 min; annealing at 55°C for 1 min; elongation at 72°C for 1 min) are performed with a Perkin-Elmer Cetus automated thermal cycler. The amplified product is analyzed by agarose gel electrophoresis and the DNA band with expected molecular weight is excised and purified. The PCR product is verified to be the selected sequence by subcloning and sequencing the DNA
30 product.

Several methods are available for the identification of the 5' or 3' non-coding portions of a gene which may not be present in the deposited clone. These methods include but are not

limited to. filter probing, clone enrichment using specific probes, and protocols similar or identical to 5' and 3' "RACE" protocols which are well known in the art. For instance, a method similar to 5' RACE is available for generating the missing 5' end of a desired full-length transcript. (Fromont-Racine et al., Nucleic Acids Res. 21(7):1683-1684 (1993).)

5 Briefly, a specific RNA oligonucleotide is ligated to the 5' ends of a population of RNA presumably containing full-length gene RNA transcripts. A primer set containing a primer specific to the ligated RNA oligonucleotide and a primer specific to a known sequence of the gene of interest is used to PCR amplify the 5' portion of the desired full-length gene. This amplified product may then be sequenced and used to generate the full
10 length gene.

This above method starts with total RNA isolated from the desired source, although poly-A+ RNA can be used. The RNA preparation can then be treated with phosphatase if necessary to eliminate 5' phosphate groups on degraded or damaged RNA which may interfere with the later RNA ligase step. The phosphatase should then be inactivated and the
15 RNA treated with tobacco acid pyrophosphatase in order to remove the cap structure present at the 5' ends of messenger RNAs. This reaction leaves a 5' phosphate group at the 5' end of the cap cleaved RNA which can then be ligated to an RNA oligonucleotide using T4 RNA ligase.

This modified RNA preparation is used as a template for first strand cDNA synthesis
20 using a gene specific oligonucleotide. The first strand synthesis reaction is used as a template for PCR amplification of the desired 5' end using a primer specific to the ligated RNA oligonucleotide and a primer specific to the known sequence of the gene of interest. The resultant product is then sequenced and analyzed to confirm that the 5' end sequence belongs to the desired gene.

25

Example 2: Isolation of Genomic Clones Corresponding to a Polynucleotide

A human genomic P1 library (Genomic Systems, Inc.) is screened by PCR using primers selected for the sequence corresponding to SEQ ID NO:X, according to the method
30 described in Example 1. (See also, Sambrook.)

Example 3: Tissue specific expression analysis

The Human Genome Sciences, Inc. (HGS) database is derived from sequencing tissue specific cDNA libraries. Libraries generated from a particular tissue are selected and the specific tissue expression pattern of EST groups or assembled contigs within these libraries is determined by comparison of the expression patterns of those groups or contigs within the entire database. ESTs which show tissue specific expression are selected.

The original clone from which the specific EST sequence was generated, is obtained from the catalogued library of clones and the insert amplified by PCR using methods known in the art. The PCR product is denatured then transferred in 96 well format to a nylon membrane (Schleicher and Scheull) generating an array filter of tissue specific clones. Housekeeping genes, maize genes, and known tissue specific genes are included on the filters. These targets can be used in signal normalization and to validate assay sensitivity. Additional targets are included to monitor probe length and specificity of hybridization.

Radioactively labeled hybridization probes are generated by first strand cDNA synthesis per the manufacturer's instructions (Life Technologies) from mRNA/RNA samples prepared from the specific tissue being analyzed. The hybridization probes are purified by gel exclusion chromatography, quantitated, and hybridized with the array filters in hybridization bottles at 65°C overnight. The filters are washed under stringent conditions and signals are captured using a Fuji phosphorimager.

Data is extracted using AIS software and following background subtraction, signal normalization is performed. This includes a normalization of filter-wide expression levels between different experimental runs. Genes that are differentially expressed in the tissue of interest are identified and the full length sequence of these clones is generated.

Example 4: Chromosomal Mapping of the Polynucleotides

An oligonucleotide primer set is designed according to the sequence at the 5' end of SEQ ID NO:X. This primer preferably spans about 100 nucleotides. This primer set is then used in a polymerase chain reaction under the following set of conditions : 30 seconds, 95°C; 1 minute, 56°C; 1 minute, 70°C. This cycle is repeated 32 times followed by one 5 minute

cycle at 70°C. Human, mouse, and hamster DNA is used as template in addition to a somatic cell hybrid panel containing individual chromosomes or chromosome fragments (Bios, Inc). The reactions is analyzed on either 8% polyacrylamide gels or 3.5 % agarose gels. Chromosome mapping is determined by the presence of an approximately 100 bp PCR
5 fragment in the particular somatic cell hybrid.

Example 5: Bacterial Expression of a Polypeptide

A polynucleotide encoding a polypeptide of the present invention is amplified using
10 PCR oligonucleotide primers corresponding to the 5' and 3' ends of the DNA sequence, as outlined in Example 1, to synthesize insertion fragments. The primers used to amplify the cDNA insert should preferably contain restriction sites, such as BamHI and XbaI, at the 5' end of the primers in order to clone the amplified product into the expression vector. For example, BamHI and XbaI correspond to the restriction enzyme sites on the bacterial
15 expression vector pQE-9. (Qiagen, Inc., Chatsworth, CA). This plasmid vector encodes antibiotic resistance (Amp^r), a bacterial origin of replication (ori), an IPTG-regulatable promoter/operator (P/O), a ribosome binding site (RBS), a 6-histidine tag (6-His), and restriction enzyme cloning sites.

The pQE-9 vector is digested with BamHI and XbaI and the amplified fragment is
20 ligated into the pQE-9 vector maintaining the reading frame initiated at the bacterial RBS. The ligation mixture is then used to transform the E. coli strain M15/rep4 (Qiagen, Inc.) which contains multiple copies of the plasmid pREP4, which expresses the lacI repressor and also confers kanamycin resistance (Kan^r). Transformants are identified by their ability to grow on LB plates and ampicillin/kanamycin resistant colonies are selected. Plasmid DNA is
25 isolated and confirmed by restriction analysis.

Clones containing the desired constructs are grown overnight (O/N) in liquid culture in LB media supplemented with both Amp (100 ug/ml) and Kan (25 ug/ml). The O/N culture is used to inoculate a large culture at a ratio of 1:100 to 1:250. The cells are grown to an optical density 600 (O.D.⁶⁰⁰) of between 0.4 and 0.6. IPTG (Isopropyl-B-D-thiogalacto
30 pyranoside) is then added to a final concentration of 1 mM. IPTG induces by inactivating the lacI repressor, clearing the P/O leading to increased gene expression.

Cells are grown for an extra 3 to 4 hours. Cells are then harvested by centrifugation (20 mins at 6000Xg). The cell pellet is solubilized in the chaotropic agent 6 Molar Guanidine HCl by stirring for 3-4 hours at 4°C. The cell debris is removed by centrifugation, and the supernatant containing the polypeptide is loaded onto a nickel-nitrilo-tri-acetic acid ("Ni-NTA") affinity resin column (available from QIAGEN, Inc., *supra*). Proteins with a 6 x His tag bind to the Ni-NTA resin with high affinity and can be purified in a simple one-step procedure (for details see: The QIAexpressionist (1995) QIAGEN, Inc., *supra*).

Briefly, the supernatant is loaded onto the column in 6 M guanidine-HCl, pH 8. the column is first washed with 10 volumes of 6 M guanidine-HCl, pH 8. then washed with 10 volumes of 6 M guanidine-HCl pH 6, and finally the polypeptide is eluted with 6 M guanidine-HCl, pH 5.

The purified protein is then renatured by dialyzing it against phosphate-buffered saline (PBS) or 50 mM Na-acetate, pH 6 buffer plus 200 mM NaCl. Alternatively, the protein can be successfully refolded while immobilized on the Ni-NTA column. The recommended conditions are as follows: renature using a linear 6M-1M urea gradient in 500 mM NaCl, 20% glycerol, 20 mM Tris/HCl pH 7.4, containing protease inhibitors. The renaturation should be performed over a period of 1.5 hours or more. After renaturation the proteins are eluted by the addition of 250 mM imidazole. Imidazole is removed by a final dialyzing step against PBS or 50 mM sodium acetate pH 6 buffer plus 200 mM NaCl.

The purified protein is stored at 4°C or frozen at -80°C.

In addition to the above expression vector, the present invention further includes an expression vector comprising phage operator and promoter elements operatively linked to a polynucleotide of the present invention, called pHE4a. (ATCC Accession Number 209645, deposited on February 25, 1998.) This vector contains: 1) a neomycinphosphotransferase gene as a selection marker, 2) an E. coli origin of replication, 3) a T5 phage promoter sequence, 4) two lac operator sequences, 5) a Shine-Delgarno sequence, and 6) the lactose operon repressor gene (*lacIq*). The origin of replication (*oriC*) is derived from pUC19 (LTI, Gaithersburg, MD). The promoter sequence and operator sequences are made synthetically.

DNA can be inserted into the pHEa by restricting the vector with NdeI and XbaI, BamHI, XhoI, or Asp718, running the restricted product on a gel, and isolating the larger fragment (the stuffer fragment should be about 310 base pairs). The DNA insert is generated according to the PCR protocol described in Example 1, using PCR primers having restriction

sites for NdeI (5' primer) and XbaI, BamHI, XhoI, or Asp718 (3' primer). The PCR insert is gel purified and restricted with compatible enzymes. The insert and vector are ligated according to standard protocols.

The engineered vector could easily be substituted in the above protocol to express
5 protein in a bacterial system.

Example 6: Purification of a Polypeptide from an Inclusion Body

The following alternative method can be used to purify a polypeptide expressed in *E*
10 *coli* when it is present in the form of inclusion bodies. Unless otherwise specified, all of the following steps are conducted at 4-10°C.

Upon completion of the production phase of the *E. coli* fermentation, the cell culture is cooled to 4-10°C and the cells harvested by continuous centrifugation at 15,000 rpm (Heraeus Sepatech). On the basis of the expected yield of protein per unit weight of cell
15 paste and the amount of purified protein required, an appropriate amount of cell paste, by weight, is suspended in a buffer solution containing 100 mM Tris, 50 mM EDTA, pH 7.4. The cells are dispersed to a homogeneous suspension using a high shear mixer.

The cells are then lysed by passing the solution through a microfluidizer (Microfluidics, Corp. or APV Gaulin, Inc.) twice at 4000-6000 psi. The homogenate is then
20 mixed with NaCl solution to a final concentration of 0.5 M NaCl, followed by centrifugation at 7000 xg for 15 min. The resultant pellet is washed again using 0.5M NaCl, 100 mM Tris, 50 mM EDTA, pH 7.4.

The resulting washed inclusion bodies are solubilized with 1.5 M guanidine hydrochloride (GuHCl) for 2-4 hours. After 7000 xg centrifugation for 15 min., the pellet is
25 discarded and the polypeptide containing supernatant is incubated at 4°C overnight to allow further GuHCl extraction.

Following high speed centrifugation (30,000 xg) to remove insoluble particles, the GuHCl solubilized protein is refolded by quickly mixing the GuHCl extract with 20 volumes of buffer containing 50 mM sodium, pH 4.5, 150 mM NaCl, 2 mM EDTA by vigorous
30 stirring. The refolded diluted protein solution is kept at 4°C without mixing for 12 hours prior to further purification steps.

To clarify the refolded polypeptide solution, a previously prepared tangential filtration unit equipped with 0.16 μm membrane filter with appropriate surface area (e.g., Filtron), equilibrated with 40 mM sodium acetate, pH 6.0 is employed. The filtered sample is loaded onto a cation exchange resin (e.g., Poros HS-50, Perseptive Biosystems). The column is washed with 40 mM sodium acetate, pH 6.0 and eluted with 250 mM, 500 mM, 1000 mM, and 1500 mM NaCl in the same buffer, in a stepwise manner. The absorbance at 280 nm of the effluent is continuously monitored. Fractions are collected and further analyzed by SDS-PAGE.

Fractions containing the polypeptide are then pooled and mixed with 4 volumes of water. The diluted sample is then loaded onto a previously prepared set of tandem columns of strong anion (Poros HQ-50, Perseptive Biosystems) and weak anion (Poros CM-20, Perseptive Biosystems) exchange resins. The columns are equilibrated with 40 mM sodium acetate, pH 6.0. Both columns are washed with 40 mM sodium acetate, pH 6.0, 200 mM NaCl. The CM-20 column is then eluted using a 10 column volume linear gradient ranging from 0.2 M NaCl, 50 mM sodium acetate, pH 6.0 to 1.0 M NaCl, 50 mM sodium acetate, pH 6.5. Fractions are collected under constant A_{280} monitoring of the effluent. Fractions containing the polypeptide (determined, for instance, by 16% SDS-PAGE) are then pooled.

The resultant polypeptide should exhibit greater than 95% purity after the above refolding and purification steps. No major contaminant bands should be observed from Commassie blue stained 16% SDS-PAGE gel when 5 μg of purified protein is loaded. The purified protein can also be tested for endotoxin/LPS contamination, and typically the LPS content is less than 0.1 ng/ml according to LAL assays.

Example 7: Cloning and Expression of a Polypeptide in a Baculovirus Expression System

25

In this example, the plasmid shuttle vector pA2 is used to insert a polynucleotide into a baculovirus to express a polypeptide. This expression vector contains the strong polyhedrin promoter of the *Autographa californica* nuclear polyhedrosis virus (AcMNPV) followed by convenient restriction sites such as BamHI, Xba I and Asp718. The polyadenylation site of the simian virus 40 ("SV40") is used for efficient polyadenylation. For easy selection of recombinant virus, the plasmid contains the beta-galactosidase gene from *E. coli* under control of a weak *Drosophila* promoter in the same orientation, followed by the

polyadenylation signal of the polyhedrin gene. The inserted genes are flanked on both sides by viral sequences for cell-mediated homologous recombination with wild-type viral DNA to generate a viable virus that express the cloned polynucleotide.

Many other baculovirus vectors can be used in place of the vector above, such as pAc373, pVL941, and pAcIM1, as one skilled in the art would readily appreciate, as long as the construct provides appropriately located signals for transcription, translation, secretion and the like, including a signal peptide and an in-frame AUG as required. Such vectors are described, for instance, in Luckow et al., *Virology* 170:31-39 (1989).

Specifically, the cDNA sequence contained in the deposited clone, including the AUG initiation codon, is amplified using the PCR protocol described in Example 1. If a naturally occurring signal sequence is used to produce the polypeptide of the present invention, the pA2 vector does not need a second signal peptide. Alternatively, the vector can be modified (pA2 GP) to include a baculovirus leader sequence, using the standard methods described in Summers et al., "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures," Texas Agricultural Experimental Station Bulletin No. 1555 (1987).

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

The plasmid is digested with the corresponding restriction enzymes and optionally, can be dephosphorylated using calf intestinal phosphatase, using routine procedures known in the art. The DNA is then isolated from a 1% agarose gel using a commercially available kit ("GeneClean" BIO 101 Inc., La Jolla, Ca.).

The fragment and the dephosphorylated plasmid are ligated together with T4 DNA ligase. *E. coli* HB101 or other suitable *E. coli* hosts such as XL-1 Blue (Stratagene Cloning Systems, La Jolla, CA) cells are transformed with the ligation mixture and spread on culture plates. Bacteria containing the plasmid are identified by digesting DNA from individual colonies and analyzing the digestion product by gel electrophoresis. The sequence of the cloned fragment is confirmed by DNA sequencing.

Five µg of a plasmid containing the polynucleotide is co-transfected with 1.0 µg of a commercially available linearized baculovirus DNA ("BaculoGold™ baculovirus DNA", Pharmingen, San Diego, CA), using the lipofection method described by Felgner et al., *Proc.*

Natl. Acad. Sci. USA 84:7413-7417 (1987). One μg of BaculoGold™ virus DNA and 5 μg of the plasmid are mixed in a sterile well of a microtiter plate containing 50 μl of serum-free Grace's medium (Life Technologies Inc., Gaithersburg, MD). Afterwards, 10 μl Lipofectin plus 90 μl Grace's medium are added, mixed and incubated for 15 minutes at room temperature. Then the transfection mixture is added drop-wise to Sf9 insect cells (ATCC CRL 1711) seeded in a 35 mm tissue culture plate with 1 ml Grace's medium without serum. The plate is then incubated for 5 hours at 27° C. The transfection solution is then removed from the plate and 1 ml of Grace's insect medium supplemented with 10% fetal calf serum is added. Cultivation is then continued at 27° C for four days.

After four days the supernatant is collected and a plaque assay is performed, as described by Summers and Smith, *supra*. An agarose gel with "Blue Gal" (Life Technologies Inc., Gaithersburg) is used to allow easy identification and isolation of gal-expressing clones, which produce blue-stained plaques. (A detailed description of a "plaque assay" of this type can also be found in the user's guide for insect cell culture and baculovirology distributed by Life Technologies Inc., Gaithersburg, page 9-10.) After appropriate incubation, blue stained plaques are picked with the tip of a micropipettor (e.g., Eppendorf). The agar containing the recombinant viruses is then resuspended in a microcentrifuge tube containing 200 μl of Grace's medium and the suspension containing the recombinant baculovirus is used to infect Sf9 cells seeded in 35 mm dishes. Four days later the supernatants of these culture dishes are harvested and then they are stored at 4° C.

To verify the expression of the polypeptide, Sf9 cells are grown in Grace's medium supplemented with 10% heat-inactivated FBS. The cells are infected with the recombinant baculovirus containing the polynucleotide at a multiplicity of infection ("MOI") of about 2. If radiolabeled proteins are desired, 6 hours later the medium is removed and is replaced with SF900 II medium minus methionine and cysteine (available from Life Technologies Inc., Rockville, MD). After 42 hours, 5 μCi of ^{35}S -methionine and 5 μCi ^{35}S -cysteine (available from Amersham) are added. The cells are further incubated for 16 hours and then are harvested by centrifugation. The proteins in the supernatant as well as the intracellular proteins are analyzed by SDS-PAGE followed by autoradiography (if radiolabeled).

Microsequencing of the amino acid sequence of the amino terminus of purified protein may be used to determine the amino terminal sequence of the produced protein.

Example 8: Expression of a Polypeptide in Mammalian Cells

The polypeptide of the present invention can be expressed in a mammalian cell. A typical mammalian expression vector contains a promoter element, which mediates the initiation of transcription of mRNA, a protein coding sequence, and signals required for the termination of transcription and polyadenylation of the transcript. Additional elements include enhancers, Kozak sequences and intervening sequences flanked by donor and acceptor sites for RNA splicing. Highly efficient transcription is achieved with the early and late promoters from SV40, the long terminal repeats (LTRs) from Retroviruses, e.g., RSV, HTLV, HIV and the early promoter of the cytomegalovirus (CMV). However, cellular elements can also be used (e.g., the human actin promoter).

Suitable expression vectors for use in practicing the present invention include, for example, vectors such as pSVL and pMSG (Pharmacia, Uppsala, Sweden), pRSVcat (ATCC 37152), pSV2dhfr (ATCC 37146), pBC12MI (ATCC 67109), pCMVSPORT 2.0, and pCMVSPORT 3.0. Mammalian host cells that could be used include, human Hela, 293, H9 and Jurkat cells, mouse NIH3T3 and C127 cells, Cos 1, Cos 7 and CV1, quail QC1-3 cells, mouse L cells and Chinese hamster ovary (CHO) cells.

Alternatively, the polypeptide can be expressed in stable cell lines containing the polynucleotide integrated into a chromosome. The co-transfection with a selectable marker such as DHFR, gpt, neomycin, hygromycin allows the identification and isolation of the transfected cells.

The transfected gene can also be amplified to express large amounts of the encoded protein. The DHFR (dihydrofolate reductase) marker is useful in developing cell lines that carry several hundred or even several thousand copies of the gene of interest. (See, e.g., Alt, F. W., et al., J. Biol. Chem. 253:1357-1370 (1978); Hamlin, J. L. and Ma, C., Biochem. et Biophys. Acta, 1097:107-143 (1990); Page, M. J. and Sydenham, M. A., Biotechnology 9:64-68 (1991).) Another useful selection marker is the enzyme glutamine synthase (GS) (Murphy et al., Biochem J. 227:277-279 (1991); Bebbington et al., Bio/Technology 10:169-175 (1992). Using these markers, the mammalian cells are grown in selective medium and the cells with the highest resistance are selected. These cell lines contain the amplified gene(s) integrated into a chromosome. Chinese hamster ovary (CHO) and NSO cells are often used for the production of proteins.

Derivatives of the plasmid pSV2-dhfr (ATCC Accession No. 37146), the expression vectors pC4 (ATCC Accession No. 209646) and pC6 (ATCC Accession No. 209647) contain the strong promoter (LTR) of the Rous Sarcoma Virus (Cullen et al., Molecular and Cellular Biology, 438-447 (March, 1985)) plus a fragment of the CMV-enhancer (Boshart et al., Cell 5 41:521-530 (1985).) Multiple cloning sites, e.g., with the restriction enzyme cleavage sites BamHI, XbaI and Asp718, facilitate the cloning of the gene of interest. The vectors also contain the 3' intron, the polyadenylation and termination signal of the rat preproinsulin gene, and the mouse DHFR gene under control of the SV40 early promoter.

Specifically, the plasmid pC6, for example, is digested with appropriate restriction 10 enzymes and then dephosphorylated using calf intestinal phosphates by procedures known in the art. The vector is then isolated from a 1% agarose gel.

A polynucleotide of the present invention is amplified according to the protocol outlined in Example 1. If a naturally occurring signal sequence is used to produce the polypeptide of the present invention, the vector does not need a second signal peptide. 15 Alternatively, if a naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

The amplified fragment is isolated from a 1% agarose gel using a commercially available kit ("GeneClean," BIO 101 Inc., La Jolla, Ca.). The fragment then is digested with appropriate restriction enzymes and again purified on a 1% agarose gel.

20 The amplified fragment is then digested with the same restriction enzyme and purified on a 1% agarose gel. The isolated fragment and the dephosphorylated vector are then ligated with T4 DNA ligase. *E. coli* HB101 or XL-1 Blue cells are then transformed and bacteria are identified that contain the fragment inserted into plasmid pC6 using, for instance, restriction enzyme analysis.

25 Chinese hamster ovary cells lacking an active DHFR gene is used for transfection. Five μ g of the expression plasmid pC6 or pC4 is cotransfected with 0.5 μ g of the plasmid pSVneo using lipofectin (Felgner et al., *supra*). The plasmid pSV2-neo contains a dominant selectable marker, the *neo* gene from Tn5 encoding an enzyme that confers resistance to a group of antibiotics including G418. The cells are seeded in alpha minus MEM 30 supplemented with 1 mg/ml G418. After 2 days, the cells are trypsinized and seeded in hybridoma cloning plates (Greiner, Germany) in alpha minus MEM supplemented with 10, 25, or 50 ng/ml of methotrexate plus 1 mg/ml G418. After about 10-14 days single clones

are trypsinized and then seeded in 6-well petri dishes or 10 ml flasks using different concentrations of methotrexate (50 nM, 100 nM, 200 nM, 400 nM, 800 nM). Clones growing at the highest concentrations of methotrexate are then transferred to new 6-well plates containing even higher concentrations of methotrexate (1 μ M, 2 μ M, 5 μ M, 10 mM, 20 mM). The same procedure is repeated until clones are obtained which grow at a concentration of 100 - 200 μ M. Expression of the desired gene product is analyzed, for instance, by SDS-PAGE and Western blot or by reversed phase HPLC analysis.

Example 9: Protein Fusions

10

The polypeptides of the present invention are preferably fused to other proteins. These fusion proteins can be used for a variety of applications. For example, fusion of the present polypeptides to His-tag, HA-tag, protein A, IgG domains, and maltose binding protein facilitates purification. (See Example 5; see also EP A 394,827; Traunecker, et al., Nature 331:84-86 (1988).) Similarly, fusion to IgG-1, IgG-3, and albumin increases the half-life time in vivo. Nuclear localization signals fused to the polypeptides of the present invention can target the protein to a specific subcellular localization, while covalent heterodimer or homodimers can increase or decrease the activity of a fusion protein. Fusion proteins can also create chimeric molecules having more than one function. Finally, fusion proteins can increase solubility and/or stability of the fused protein compared to the non-fused protein. All of the types of fusion proteins described above can be made by modifying the following protocol, which outlines the fusion of a polypeptide to an IgG molecule, or the protocol described in Example 5.

25 Briefly, the human Fc portion of the IgG molecule can be PCR amplified, using primers that span the 5' and 3' ends of the sequence described below. These primers also should have convenient restriction enzyme sites that will facilitate cloning into an expression vector, preferably a mammalian expression vector.

30 For example, if pC4 (Accession No. 209646) is used, the human Fc portion can be ligated into the BamHI cloning site. Note that the 3' BamHI site should be destroyed. Next, the vector containing the human Fc portion is re-restricted with BamHI, linearizing the vector, and a polynucleotide of the present invention, isolated by the PCR protocol described in Example 1, is ligated into this BamHI site. Note that the polynucleotide is cloned without

a stop codon, otherwise a fusion protein will not be produced.

If the naturally occurring signal sequence is used to produce the polypeptide of the present invention, pC4 does not need a second signal peptide. Alternatively, if the naturally occurring signal sequence is not used, the vector can be modified to include a heterologous signal sequence. (See, e.g., WO 96/34891.)

Human IgG Fc region:

```
GGGATCCGGAGCCCAAATCTTCTGACAAACTCACACATGCCCACCGTGCCCAG
CACCTGAATTCGAGGGTGACCGTCAGTCTTCTCTTCCCCCAAAACCCAAGGA
10 CACCTCATGATCTCCCGGACTCCTGAGGTCACATGCGTGGTGGTGGACGTAAGC
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCAT
AATGCCAAGACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTC
AGCGTCCTCACCCTGTCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGC
AAGGTCTCCAACAAAGCCCTCCCAACCCCCATCGAGAAAACCATCTCCAAAGCC
15 AAAGGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAG
CTGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAAGC
GACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGAC
CACGCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACC
GTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGCAT
20 GAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAT
GAGTGCGACGGCCGCGACTCTAGAGGAT (SEQ ID NO:1547)
```

Example 10: Production of an Antibody from a Polypeptide

25 a) Hybridoma Technology

The antibodies of the present invention can be prepared by a variety of methods. (See, Current Protocols, Chapter 2.) As one example of such methods, cells expressing polypeptide of the present invention are administered to an animal to induce the production of sera containing polyclonal antibodies. In a preferred method, a preparation of polypeptide of the present invention is prepared and purified to render it substantially free of natural contaminants. Such a preparation is then introduced into an animal in order to produce polyclonal antisera of greater specific activity.

Monoclonal antibodies specific for polypeptide of the present invention are prepared using hybridoma technology. (Kohler et al., Nature 256:495 (1975); Kohler et al., Eur. J. Immunol. 6:511 (1976); Kohler et al., Eur. J. Immunol. 6:292 (1976); Hammerling et al., in: Monoclonal Antibodies and T-Cell Hybridomas, Elsevier, N.Y., pp. 563-681 (1981)). In
5 general, an animal (preferably a mouse) is immunized with polypeptide of the present invention or, more preferably, with a secreted polypeptide of the present invention-expressing cell. Such polypeptide-expressing cells are cultured in any suitable tissue culture medium, preferably in Earle's modified Eagle's medium supplemented with 10% fetal bovine serum (inactivated at about 56°C), and supplemented with about 10 g/l of nonessential amino
10 acids, about 1,000 U/ml of penicillin, and about 100 µg/ml of streptomycin.

The splenocytes of such mice are extracted and fused with a suitable myeloma cell line. Any suitable myeloma cell line may be employed in accordance with the present invention; however, it is preferable to employ the parent myeloma cell line (SP2O), available from the ATCC. After fusion, the resulting hybridoma cells are selectively maintained in
15 HAT medium, and then cloned by limiting dilution as described by Wands et al. (Gastroenterology 80:225-232 (1981)). The hybridoma cells obtained through such a selection are then assayed to identify clones which secrete antibodies capable of binding the polypeptide of the present invention.

Alternatively, additional antibodies capable of binding to polypeptide of the present
20 invention can be produced in a two-step procedure using anti-idiotypic antibodies. Such a method makes use of the fact that antibodies are themselves antigens, and therefore, it is possible to obtain an antibody which binds to a second antibody. In accordance with this method, protein specific antibodies are used to immunize an animal, preferably a mouse. The splenocytes of such an animal are then used to produce hybridoma cells, and the hybridoma
25 cells are screened to identify clones which produce an antibody whose ability to bind to the polypeptide of the present invention-specific antibody can be blocked by polypeptide of the present invention. Such antibodies comprise anti-idiotypic antibodies to the polypeptide of the present invention-specific antibody and are used to immunize an animal to induce formation of further polypeptide of the present invention-specific antibodies.

30 For in vivo use of antibodies in humans, an antibody is "humanized". Such antibodies can be produced using genetic constructs derived from hybridoma cells producing the monoclonal antibodies described above. Methods for producing chimeric and humanized

antibodies are known in the art and are discussed herein. (See, for review, Morrison. Science 229:1202 (1985); Oi et al., BioTechniques 4:214 (1986); Cabilly et al., U.S. Patent No. 4,816,567; Taniguchi et al., EP 171496; Morrison et al., EP 173494; Neuberger et al., WO 8601533; Robinson et al., WO 8702671; Boulianne et al., Nature 312:643 (1984); Neuberger et al., Nature 314:268 (1985).)

b) Isolation Of Antibody Fragments Directed Against Polypeptide of the Present Invention From A Library Of scFvs

Naturally occurring V-genes isolated from human PBLs are constructed into a library of antibody fragments which contain reactivities against polypeptide of the present invention to which the donor may or may not have been exposed (see e.g., U.S. Patent 5,885,793 incorporated herein by reference in its entirety).

Rescue of the Library. A library of scFvs is constructed from the RNA of human PBLs as described in PCT publication WO 92/01047. To rescue phage displaying antibody fragments, approximately 10⁹ E. coli harboring the phagemid are used to inoculate 50 ml of 2xTY containing 1% glucose and 100 µg/ml of ampicillin (2xTY-AMP-GLU) and grown to an O.D. of 0.8 with shaking. Five ml of this culture is used to inoculate 50 ml of 2xTY-AMP-GLU, 2 x 10⁸ TU of delta gene 3 helper (M13 delta gene III, see PCT publication WO 92/01047) are added and the culture incubated at 37°C for 45 minutes without shaking and then at 37°C for 45 minutes with shaking. The culture is centrifuged at 4000 r.p.m. for 10 min. and the pellet resuspended in 2 liters of 2xTY containing 100 µg/ml ampicillin and 50 µg/ml kanamycin and grown overnight. Phage are prepared as described in PCT publication WO 92/01047.

M13 delta gene III is prepared as follows: M13 delta gene III helper phage does not encode gene III protein, hence the phage(mid) displaying antibody fragments have a greater avidity of binding to antigen. Infectious M13 delta gene III particles are made by growing the helper phage in cells harboring a pUC19 derivative supplying the wild type gene III protein during phage morphogenesis. The culture is incubated for 1 hour at 37° C without shaking and then for a further hour at 37°C with shaking. Cells are spun down (IEC-Centra 8,400 r.p.m. for 10 min), resuspended in 300 ml 2xTY broth containing 100 µg ampicillin/ml and 25 µg kanamycin/ml (2xTY-AMP-KAN) and grown overnight, shaking at 37°C. Phage particles are purified and concentrated from the culture medium by two PEG-precipitations

(Sambrook et al., 1990), resuspended in 2 ml PBS and passed through a 0.45 μ m filter (Minisart NML: Sartorius) to give a final concentration of approximately 10¹³ transducing units/ml (ampicillin-resistant clones).

Panning of the Library. Immunotubes (Nunc) are coated overnight in PBS with 4 ml of either 100 μ g/ml or 10 μ g/ml of a polypeptide of the present invention. Tubes are blocked with 2% Marvel-PBS for 2 hours at 37°C and then washed 3 times in PBS. Approximately 10¹³ TU of phage is applied to the tube and incubated for 30 minutes at room temperature tumbling on an over and under turntable and then left to stand for another 1.5 hours. Tubes are washed 10 times with PBS 0.1% Tween-20 and 10 times with PBS. Phage are eluted by adding 1 ml of 100 mM triethylamine and rotating 15 minutes on an under and over turntable after which the solution is immediately neutralized with 0.5 ml of 1.0M Tris-HCl, pH 7.4. Phage are then used to infect 10 ml of mid-log E. coli TG1 by incubating eluted phage with bacteria for 30 minutes at 37°C. The E. coli are then plated on TYE plates containing 1% glucose and 100 μ g/ml ampicillin. The resulting bacterial library is then rescued with delta gene 3 helper phage as described above to prepare phage for a subsequent round of selection. This process is then repeated for a total of 4 rounds of affinity purification with tube-washing increased to 20 times with PBS, 0.1% Tween-20 and 20 times with PBS for rounds 3 and 4.

Characterization of Binders. Eluted phage from the 3rd and 4th rounds of selection are used to infect E. coli HB 2151 and soluble scFv is produced (Marks, et al., 1991) from single colonies for assay. ELISAs are performed with microtitre plates coated with either 10 μ g/ml of the polypeptide of the present invention in 50 mM bicarbonate pH 9.6. Clones positive in ELISA are further characterized by PCR fingerprinting (see, e.g., PCT publication WO 92/01047) and then by sequencing. These ELISA positive clones may also be further characterized by techniques known in the art, such as, for example, epitope mapping, binding affinity, receptor signal transduction, ability to block or competitively inhibit antibody/antigen binding, and competitive agonistic or antagonistic activity.

Example 11: Method of Determining Alterations in a Gene Corresponding to a Polynucleotide

30

RNA isolated from entire families or individual patients presenting with a phenotype of interest (such as a disease) is isolated. cDNA is then generated from these RNA

samples using protocols known in the art. (See, Sambrook.) The cDNA is then used as a template for PCR, employing primers surrounding regions of interest in SEQ ID NO:X; and/or the nucleotide sequence of the related cDNA in the cDNA clone contained in a deposited library. Suggested PCR conditions consist of 35 cycles at 95 degrees C for 30
5 seconds; 60-120 seconds at 52-58 degrees C; and 60-120 seconds at 70 degrees C, using buffer solutions described in Sidransky et al., Science 252:706 (1991).

PCR products are then sequenced using primers labeled at their 5' end with T4 polynucleotide kinase, employing SequiTherm Polymerase. (Epicentre Technologies). The intron-exon borders of selected exons is also determined and genomic PCR products
10 analyzed to confirm the results. PCR products harboring suspected mutations is then cloned and sequenced to validate the results of the direct sequencing.

PCR products is cloned into T-tailed vectors as described in Holton et al., Nucleic Acids Research, 19:1156 (1991) and sequenced with T7 polymerase (United States Biochemical). Affected individuals are identified by mutations not present in unaffected
15 individuals.

Genomic rearrangements are also observed as a method of determining alterations in a gene corresponding to a polynucleotide. Genomic clones isolated according to Example 2 are nick-translated with digoxigenin deoxy-uridine 5'-triphosphate (Boehringer Mannheim), and FISH performed as described in Johnson et al., Methods Cell Biol. 35:73-99 (1991).
20 Hybridization with the labeled probe is carried out using a vast excess of human cot-1 DNA for specific hybridization to the corresponding genomic locus.

Chromosomes are counterstained with 4,6-diamino-2-phenylidole and propidium iodide, producing a combination of C- and R-bands. Aligned images for precise mapping are obtained using a triple-band filter set (Chroma Technology, Brattleboro, VT) in combination
25 with a cooled charge-coupled device camera (Photometrics, Tucson, AZ) and variable excitation wavelength filters. (Johnson et al., Genet. Anal. Tech. Appl., 8:75 (1991).) Image collection, analysis and chromosomal fractional length measurements are performed using the ISee Graphical Program System. (Inovision Corporation, Durham, NC.) Chromosome alterations of the genomic region hybridized by the probe are identified as insertions,
30 deletions, and translocations. These alterations are used as a diagnostic marker for an associated disease.

Example 12: Method of Detecting Abnormal Levels of a Polypeptide in a Biological Sample

A polypeptide of the present invention can be detected in a biological sample, and if an increased or decreased level of the polypeptide is detected, this polypeptide is a marker for a particular phenotype. Methods of detection are numerous, and thus, it is understood that one skilled in the art can modify the following assay to fit their particular needs.

For example, antibody-sandwich ELISAs are used to detect polypeptides in a sample, preferably a biological sample. Wells of a microtiter plate are coated with specific antibodies, at a final concentration of 0.2 to 10 ug/ml. The antibodies are either monoclonal or polyclonal and are produced by the method described in Example 10. The wells are blocked so that non-specific binding of the polypeptide to the well is reduced.

The coated wells are then incubated for > 2 hours at RT with a sample containing the polypeptide. Preferably, serial dilutions of the sample should be used to validate results. The plates are then washed three times with deionized or distilled water to remove unbounded polypeptide.

Next, 50 ul of specific antibody-alkaline phosphatase conjugate, at a concentration of 25-400 ng, is added and incubated for 2 hours at room temperature. The plates are again washed three times with deionized or distilled water to remove unbounded conjugate.

Add 75 ul of 4-methylumbelliferyl phosphate (MUP) or p-nitrophenyl phosphate (NPP) substrate solution to each well and incubate 1 hour at room temperature. Measure the reaction by a microtiter plate reader. Prepare a standard curve, using serial dilutions of a control sample, and plot polypeptide concentration on the X-axis (log scale) and fluorescence or absorbance of the Y-axis (linear scale). Interpolate the concentration of the polypeptide in the sample using the standard curve.

Example 13: Formulation

The invention also provides methods of treatment and/or prevention of diseases or disorders (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of a Therapeutic. By therapeutic is meant a polynucleotides or polypeptides of the invention (including fragments and variants), agonists or antagonists thereof, and/or antibodies thereto, in combination with

a pharmaceutically acceptable carrier type (e.g., a sterile carrier).

The Therapeutic will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the Therapeutic alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

As a general proposition, the total pharmaceutically effective amount of the Therapeutic administered parenterally per dose will be in the range of about 1 µg/kg/day to 10 mg/kg/day of patient body weight, although, as noted above, this will be subject to therapeutic discretion. More preferably, this dose is at least 0.01 mg/kg/day, and most preferably for humans between about 0.01 and 1 mg/kg/day for the hormone. If given continuously, the Therapeutic is typically administered at a dose rate of about 1 µg/kg/hour to about 50 µg/kg/hour, either by 1-4 injections per day or by continuous subcutaneous infusions, for example, using a mini-pump. An intravenous bag solution may also be employed. The length of treatment needed to observe changes and the interval following treatment for responses to occur appears to vary depending on the desired effect.

Therapeutics can be administered orally, rectally, parenterally, intracistemally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics are administered orally, rectally, parenterally, intracistemally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. The term "parenteral" as used herein refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

Therapeutics of the invention are also suitably administered by sustained-release systems. Suitable examples of sustained-release Therapeutics include suitable polymeric materials (such as, for example, semi-permeable polymer matrices in the form of shaped articles, e.g., films, or microcapsules), suitable hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, and sparingly soluble derivatives (such as, for example, a sparingly soluble salt).

Sustained-release matrices include polylactides (U.S. Pat. No. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma-ethyl-L-glutamate (Sidman et al., Biopolymers 22:547-556 (1983)), poly (2- hydroxyethyl methacrylate) (Langer et al., J. Biomed. Mater. Res. 15:167-277 (1981), and Langer, Chem. Tech. 12:98-105 (1982)), ethylene vinyl acetate (Langer et al., Id.) or poly-D- (-)-3-hydroxybutyric acid (EP 133,988).

Sustained-release Therapeutics also include liposomally entrapped Therapeutics of the invention (*see* generally, Langer, *Science* 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 317 -327 and 353-365 (1989)). Liposomes containing the Therapeutic are prepared by methods known per se: DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. (USA) 82:3688-3692 (1985); Hwang et al., Proc. Natl. Acad. Sci.(USA) 77:4030-4034 (1980); EP 52,322; EP 36,676; EP 88,046; EP 143,949; EP 142,641; Japanese Pat. Appl. 83-118008; U.S. Pat. Nos. 4,485,045 and 4,544,545; and EP 102,324. Ordinarily, the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol. percent cholesterol, the selected proportion being adjusted for the optimal Therapeutic.

In yet an additional embodiment, the Therapeutics of the invention are delivered by way of a pump (*see* Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)).

Other controlled release systems are discussed in the review by Langer (*Science* 249:1527-1533 (1990)).

For parenteral administration, in one embodiment, the Therapeutic is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation preferably does not

include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

Generally, the formulations are prepared by contacting the Therapeutic uniformly and intimately with liquid carriers or finely divided solid carriers or both. Then, if necessary, the product is shaped into the desired formulation. Preferably the carrier is a parenteral carrier, more preferably a solution that is isotonic with the blood of the recipient. Examples of such carrier vehicles include water, saline, Ringer's solution, and dextrose solution. Non-aqueous vehicles such as fixed oils and ethyl oleate are also useful herein, as well as liposomes.

The carrier suitably contains minor amounts of additives such as substances that enhance isotonicity and chemical stability. Such materials are non-toxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, succinate, acetic acid, and other organic acids or their salts; antioxidants such as ascorbic acid; low molecular weight (less than about ten residues) polypeptides, e.g., polyarginine or tripeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids, such as glycine, glutamic acid, aspartic acid, or arginine; monosaccharides, disaccharides, and other carbohydrates including cellulose or its derivatives, glucose, manose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; counterions such as sodium; and/or nonionic surfactants such as polysorbates, poloxamers, or PEG.

The Therapeutic is typically formulated in such vehicles at a concentration of about 0.1 mg/ml to 100 mg/ml, preferably 1-10 mg/ml, at a pH of about 3 to 8. It will be understood that the use of certain of the foregoing excipients, carriers, or stabilizers will result in the formation of polypeptide salts.

Any pharmaceutical used for therapeutic administration can be sterile. Sterility is readily accomplished by filtration through sterile filtration membranes (e.g., 0.2 micron membranes). Therapeutics generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

Therapeutics ordinarily will be stored in unit or multi-dose containers, for example, sealed ampoules or vials, as an aqueous solution or as a lyophilized formulation for reconstitution. As an example of a lyophilized formulation, 10-ml vials are filled with 5 ml of sterile-filtered 1% (w/v) aqueous Therapeutic solution, and the resulting mixture is

lyophilized. The infusion solution is prepared by reconstituting the lyophilized Therapeutic using bacteriostatic Water-for-Injection.

The invention also provides a pharmaceutical pack or kit comprising one or more containers filled with one or more of the ingredients of the Therapeutics of the invention.

5 Associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of manufacture, use or sale for human administration. In addition, the Therapeutics may be employed in conjunction with other therapeutic compounds.

10 The Therapeutics of the invention may be administered alone or in combination with adjuvants. Adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, alum, alum plus deoxycholate (ImmunoAg), MTP-PE (Biocine Corp.), QS21 (Genentech, Inc.), BCG, and MPL. In a specific embodiment, Therapeutics of the invention are administered in combination with alum. In another specific
15 embodiment, Therapeutics of the invention are administered in combination with QS-21. Further adjuvants that may be administered with the Therapeutics of the invention include, but are not limited to, Monophosphoryl lipid immunomodulator, AdjuVax 100a, QS-21, QS-18, CRL1005, Aluminum salts, MF-59, and Virosomal adjuvant technology. Vaccines that may be administered with the Therapeutics of the invention include, but are not limited to,
20 vaccines directed toward protection against MMR (measles, mumps, rubella), polio, varicella, tetanus/diphtheria, hepatitis A, hepatitis B, haemophilus influenzae B, whooping cough, pneumonia, influenza, Lyme's Disease, rotavirus, cholera, yellow fever, Japanese encephalitis, poliomyelitis, rabies, typhoid fever, and pertussis. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or
25 concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

30 The Therapeutics of the invention may be administered alone or in combination with other therapeutic agents. Therapeutic agents that may be administered in combination with the Therapeutics of the invention, include but not limited to, other members of the TNF

family, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, cytokines and/or growth factors. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

In one embodiment, the Therapeutics of the invention are administered in combination with members of the TNF family. TNF, TNF-related or TNF-like molecules that may be administered with the Therapeutics of the invention include, but are not limited to, soluble forms of TNF-alpha, lymphotoxin-alpha (LT-alpha, also known as TNF-beta), LT-beta (found in complex heterotrimer LT-alpha2-beta), OPGL, FasL, CD27L, CD30L, CD40L, 4-1BBL, DcR3, OX40L, TNF-gamma (International Publication No. WO 96/14328), AIM-I (International Publication No. WO 97/33899), endokine-alpha (International Publication No. WO 98/07880), TR6 (International Publication No. WO 98/30694), OPG, and neutrokin-alpha (International Publication No. WO 98/18921, OX40, and nerve growth factor (NGF), and soluble forms of Fas, CD30, CD27, CD40 and 4-1BB, TR2 (International Publication No. WO 96/34095), DR3 (International Publication No. WO 97/33904), DR4 (International Publication No. WO 98/32856), TR5 (International Publication No. WO 98/30693), TR6 (International Publication No. WO 98/30694), TR7 (International Publication No. WO 98/41629), TRANK, TR9 (International Publication No. WO 98/56892), TR10 (International Publication No. WO 98/54202), 312C2 (International Publication No. WO 98/06842), and TR12, and soluble forms CD154, CD70, and CD153.

In certain embodiments, Therapeutics of the invention are administered in combination with antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors. Nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, RETROVIR™ (zidovudine/AZT), VIDEX™ (didanosine/ddI), HIVID™ (zalcitabine/ddC), ZERIT™ (stavudine/d4T), EPIVIR™ (lamivudine/3TC), and COMBIVIR™ (zidovudine/lamivudine). Non-nucleoside reverse transcriptase inhibitors that may be administered in combination with the Therapeutics of the

invention. include, but are not limited to, VIRAMUNE™ (nevirapine), RESCRIPTOR™ (delavirdine), and SUSTIVA™ (efavirenz). Protease inhibitors that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, CRIXIVAN™ (indinavir), NORVIR™ (ritonavir), INVIRASE™ (saquinavir), and
5 VIRACEPT™ (nelfinavir). In a specific embodiment, antiretroviral agents, nucleoside reverse transcriptase inhibitors, non-nucleoside reverse transcriptase inhibitors, and/or protease inhibitors may be used in any combination with Therapeutics of the invention to treat AIDS and/or to prevent or treat HIV infection.

In other embodiments, Therapeutics of the invention may be administered in
10 combination with anti-opportunistic infection agents. Anti-opportunistic agents that may be administered in combination with the Therapeutics of the invention, include, but are not limited to, TRIMETHOPRIM-SULFAMETHOXAZOLE™, DAPSONE™, PENTAMIDINE™, ATOVAQUONE™, ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, ETHAMBUTOL™, RIFABUTIN™, CLARITHROMYCIN™, AZITHROMYCIN™,
15 GANCICLOVIR™, FOSCARNET™, CIDOFOVIR™, FLUCONAZOLE™, ITRACONAZOLE™, KETOCONAZOLE™, ACYCLOVIR™, FAMCICOLVIR™, PYRIMETHAMINE™, LEUCOVORIN™, NEUPOGEN™ (filgrastim/G-CSF), and LEUKINE™ (sargramostim/GM-CSF). In a specific embodiment, Therapeutics of the invention are used in any combination with TRIMETHOPRIM-SULFAMETHOXAZOLE™,
20 DAPSONE™, PENTAMIDINE™, and/or ATOVAQUONE™ to prophylactically treat or prevent an opportunistic *Pneumocystis carinii* pneumonia infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ISONIAZID™, RIFAMPIN™, PYRAZINAMIDE™, and/or ETHAMBUTOL™ to prophylactically treat or prevent an opportunistic *Mycobacterium avium* complex infection. In another specific
25 embodiment, Therapeutics of the invention are used in any combination with RIFABUTIN™, CLARITHROMYCIN™, and/or AZITHROMYCIN™ to prophylactically treat or prevent an opportunistic *Mycobacterium tuberculosis* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with GANCICLOVIR™, FOSCARNET™, and/or CIDOFOVIR™ to prophylactically treat or prevent an opportunistic
30 cytomegalovirus infection. In another specific embodiment, Therapeutics of the invention are used in any combination with FLUCONAZOLE™, ITRACONAZOLE™, and/or

KETOCONAZOLE™ to prophylactically treat or prevent an opportunistic fungal infection. In another specific embodiment, Therapeutics of the invention are used in any combination with ACYCLOVIR™ and/or FAMCICOLVIR™ to prophylactically treat or prevent an opportunistic herpes simplex virus type I and/or type II infection. In another specific
5 embodiment, Therapeutics of the invention are used in any combination with PYRIMETHAMINE™ and/or LEUCOVORIN™ to prophylactically treat or prevent an opportunistic *Toxoplasma gondii* infection. In another specific embodiment, Therapeutics of the invention are used in any combination with LEUCOVORIN™ and/or NEUPOGEN™ to prophylactically treat or prevent an opportunistic bacterial infection.

10 In a further embodiment, the Therapeutics of the invention are administered in combination with an antiviral agent. Antiviral agents that may be administered with the Therapeutics of the invention include, but are not limited to, acyclovir, ribavirin, amantadine, and remantidine.

In a further embodiment, the Therapeutics of the invention are administered in
15 combination with an antibiotic agent. Antibiotic agents that may be administered with the Therapeutics of the invention include, but are not limited to, amoxicillin, beta-lactamases, aminoglycosides, beta-lactam (glycopeptide), beta-lactamases, Clindamycin, chloramphenicol, cephalosporins, ciprofloxacin, ciprofloxacin, erythromycin, fluoroquinolones, macrolides, metronidazole, penicillins, quinolones, rifampin, streptomycin,
20 sulfonamide, tetracyclines, trimethoprim, trimethoprim-sulfamthoxazole, and vancomycin.

Conventional nonspecific immunosuppressive agents, that may be administered in combination with the Therapeutics of the invention include, but are not limited to, steroids, cyclosporine, cyclosporine analogs, cyclophosphamide methylprednisone, prednisone, azathioprine, FK-506, 15-deoxyspergualin, and other immunosuppressive agents that act by
25 suppressing the function of responding T cells.

In specific embodiments, Therapeutics of the invention are administered in combination with immunosuppressants. Immunosuppressants preparations that may be administered with the Therapeutics of the invention include, but are not limited to, ORTHOCLONE™ (OKT3), SANDIMMUNE™/NEORAL™/SANGDYA™ (cyclosporin),
30 PROGRAF™ (tacrolimus), CELLCEPT™ (mycophenolate), Azathioprine, glucocorticosteroids, and RAPAMUNE™ (sirolimus). In a specific embodiment, immunosuppressants may be

used to prevent rejection of organ or bone marrow transplantation.

In an additional embodiment, Therapeutics of the invention are administered alone or in combination with one or more intravenous immune globulin preparations. Intravenous immune globulin preparations that may be administered with the Therapeutics of the invention include, but not limited to, GAMMAR™, IVEEGAM™, SANDOGLOBULIN™, 5 GAMMAGARD S/D™, and GAMIMUNE™. In a specific embodiment, Therapeutics of the invention are administered in combination with intravenous immune globulin preparations in transplantation therapy (e.g., bone marrow transplant).

In an additional embodiment, the Therapeutics of the invention are administered alone 10 or in combination with an anti-inflammatory agent. Anti-inflammatory agents that may be administered with the Therapeutics of the invention include, but are not limited to, glucocorticoids and the nonsteroidal anti-inflammatories, aminoarylcarboxylic acid derivatives, arylacetic acid derivatives, arylbutyric acid derivatives, arylcarboxylic acids, arylpropionic acid derivatives, pyrazoles, pyrazolones, salicylic acid derivatives, 15 thiazinecarboxamides, e-acetamidocaproic acid, S-adenosylmethionine, 3-amino-4-hydroxybutyric acid, amixetrine, bendazac, benzydamine, bucolome, difenpiramide, ditazol, emorfazone, guaiazulene, nabumetone, nimesulide, orgotein, oxaceprol, paranyline, perisoxal, pifoxime, proquazone, proxazole, and tenidap.

In another embodiment, compositions of the invention are administered in 20 combination with a chemotherapeutic agent. Chemotherapeutic agents that may be administered with the Therapeutics of the invention include, but are not limited to, antibiotic derivatives (e.g., doxorubicin, bleomycin, daunorubicin, and dactinomycin); antiestrogens (e.g., tamoxifen); antimetabolites (e.g., fluorouracil, 5-FU, methotrexate, floxuridine, interferon alpha-2b, glutamic acid, plicamycin, mercaptopurine, and 6-thioguanine); 25 cytotoxic agents (e.g., carmustine, BCNU, lomustine, CCNU, cytosine arabinoside, cyclophosphamide, estramustine, hydroxyurea, procarbazine, mitomycin, busulfan, cis-platin, and vincristine sulfate); hormones (e.g., medroxyprogesterone, estramustine phosphate sodium, ethinyl estradiol, estradiol, megestrol acetate, methyltestosterone, diethylstilbestrol diphosphate, chlorotrianisene, and testolactone); nitrogen mustard derivatives (e.g., 30 mephallen, chorambucil, mechlorethamine (nitrogen mustard) and thiotepa); steroids and combinations (e.g., bethamethasone sodium phosphate); and others (e.g., dicarbazine, asparaginase, mitotane, vincristine sulfate, vinblastine sulfate, and etoposide).

In a specific embodiment, Therapeutics of the invention are administered in combination with CHOP (cyclophosphamide, doxorubicin, vincristine, and prednisone) or any combination of the components of CHOP. In another embodiment, Therapeutics of the invention are administered in combination with Rituximab. In a further embodiment, Therapeutics of the invention are administered with Rituximab and CHOP, or Rituximab and any combination of the components of CHOP.

In an additional embodiment, the Therapeutics of the invention are administered in combination with cytokines. Cytokines that may be administered with the Therapeutics of the invention include, but are not limited to, IL2, IL3, IL4, IL5, IL6, IL7, IL10, IL12, IL13, IL15, anti-CD40, CD40L, IFN-gamma and TNF-alpha. In another embodiment, Therapeutics of the invention may be administered with any interleukin, including, but not limited to, IL-1alpha, IL-1beta, IL-2, IL-3, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IL-16, IL-17, IL-18, IL-19, IL-20, and IL-21.

In an additional embodiment, the Therapeutics of the invention are administered in combination with angiogenic proteins. Angiogenic proteins that may be administered with the Therapeutics of the invention include, but are not limited to, Glioma Derived Growth Factor (GDGF), as disclosed in European Patent Number EP-399816; Platelet Derived Growth Factor-A (PDGF-A), as disclosed in European Patent Number EP-682110; Platelet Derived Growth Factor-B (PDGF-B), as disclosed in European Patent Number EP-282317; Placental Growth Factor (PIGF), as disclosed in International Publication Number WO 92/06194; Placental Growth Factor-2 (PIGF-2), as disclosed in Hauser et al., Growth Factors, 4:259-268 (1993); Vascular Endothelial Growth Factor (VEGF), as disclosed in International Publication Number WO 90/13649; Vascular Endothelial Growth Factor-A (VEGF-A), as disclosed in European Patent Number EP-506477; Vascular Endothelial Growth Factor-2 (VEGF-2), as disclosed in International Publication Number WO 96/39515; Vascular Endothelial Growth Factor B (VEGF-3); Vascular Endothelial Growth Factor B-186 (VEGF-B186), as disclosed in International Publication Number WO 96/26736; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/02543; Vascular Endothelial Growth Factor-D (VEGF-D), as disclosed in International Publication Number WO 98/07832; and Vascular Endothelial Growth Factor-E (VEGF-E), as disclosed in German Patent Number DE19639601. The above mentioned references are incorporated herein by reference herein.

In an additional embodiment, the Therapeutics of the invention are administered in combination with hematopoietic growth factors. Hematopoietic growth factors that may be administered with the Therapeutics of the invention include, but are not limited to, LEUKINE™ (SARGRAMOSTIM™) and NEUPOGEN™ (FILGRASTIM™).

5 In an additional embodiment, the Therapeutics of the invention are administered in combination with Fibroblast Growth Factors. Fibroblast Growth Factors that may be administered with the Therapeutics of the invention include, but are not limited to, FGF-1, FGF-2, FGF-3, FGF-4, FGF-5, FGF-6, FGF-7, FGF-8, FGF-9, FGF-10, FGF-11, FGF-12, FGF-13, FGF-14, and FGF-15.

10 In additional embodiments, the Therapeutics of the invention are administered in combination with other therapeutic or prophylactic regimens, such as, for example, radiation therapy.

Example 14: Method of Treating Decreased Levels of the Polypeptide

15

The present invention relates to a method for treating an individual in need of an increased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an agonist of the invention (including polypeptides of the invention). Moreover, it will be appreciated that conditions caused by a decrease in the standard or normal expression level of a polypeptide of the present invention in an individual can be treated by administering the agonist or antagonist of the present invention. Thus, the invention also provides a method of treatment of an individual in need of an increased level of the polypeptide comprising administering to such an individual a Therapeutic comprising an amount of the agonist or antagonist to increase the activity level of the polypeptide in such an individual.

20 For example, a patient with decreased levels of a polypeptide receives a daily dose 0.1-100 ug/kg of the agonist or antagonist for six consecutive days. The exact details of the dosing scheme, based on administration and formulation, are provided in Example 13.

30 *Example 15: Method of Treating Increased Levels of the Polypeptide*

The present invention also relates to a method of treating an individual in need of a

decreased level of a polypeptide of the invention in the body comprising administering to such an individual a composition comprising a therapeutically effective amount of an antagonist of the invention (including polypeptides and antibodies of the invention).

In one example, antisense technology is used to inhibit production of a polypeptide of the present invention. This technology is one example of a method of decreasing levels of a polypeptide, due to a variety of etiologies, such as cancer.

For example, a patient diagnosed with abnormally increased levels of a polypeptide is administered intravenously antisense polynucleotides at 0.5, 1.0, 1.5, 2.0 and 3.0 mg/kg day for 21 days. This treatment is repeated after a 7-day rest period if the treatment was well tolerated. The formulation of the antisense polynucleotide is provided in Example 13.

Example 16: Method of Treatment Using Gene Therapy-Ex Vivo

One method of gene therapy transplants fibroblasts, which are capable of expressing a polypeptide, onto a patient. Generally, fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in tissue-culture medium and separated into small pieces. Small chunks of the tissue are placed on a wet surface of a tissue culture flask, approximately ten pieces are placed in each flask. The flask is turned upside down, closed tight and left at room temperature over night. After 24 hours at room temperature, the flask is inverted and the chunks of tissue remain fixed to the bottom of the flask and fresh media (e.g., Ham's F12 media, with 10% FBS, penicillin and streptomycin) is added. The flasks are then incubated at 37 degree C for approximately one week.

At this time, fresh media is added and subsequently changed every several days. After an additional two weeks in culture, a monolayer of fibroblasts emerge. The monolayer is trypsinized and scaled into larger flasks.

pMV-7 (Kirschmeier, P.T. et al., DNA, 7:219-25 (1988)), flanked by the long terminal repeats of the Moloney murine sarcoma virus, is digested with EcoRI and HindIII and subsequently treated with calf intestinal phosphatase. The linear vector is fractionated on agarose gel and purified, using glass beads.

The cDNA encoding a polypeptide of the present invention can be amplified using PCR primers which correspond to the 5' and 3' end sequences respectively as set forth in Example 1 using primers and having appropriate restriction sites and initiation/stop codons, if

necessary. Preferably, the 5' primer contains an EcoRI site and the 3' primer includes a HindIII site. Equal quantities of the Moloney murine sarcoma virus linear backbone and the amplified EcoRI and HindIII fragment are added together, in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The ligation mixture is then used to transform bacteria HB101, which are then plated onto agar containing kanamycin for the purpose of confirming that the vector has the gene of interest properly inserted.

The amphotropic pA317 or GP+am12 packaging cells are grown in tissue culture to confluent density in Dulbecco's Modified Eagles Medium (DMEM) with 10% calf serum (CS), penicillin and streptomycin. The MSV vector containing the gene is then added to the media and the packaging cells transduced with the vector. The packaging cells now produce infectious viral particles containing the gene (the packaging cells are now referred to as producer cells).

Fresh media is added to the transduced producer cells, and subsequently, the media is harvested from a 10 cm plate of confluent producer cells. The spent media, containing the infectious viral particles, is filtered through a millipore filter to remove detached producer cells and this media is then used to infect fibroblast cells. Media is removed from a sub-confluent plate of fibroblasts and quickly replaced with the media from the producer cells. This media is removed and replaced with fresh media. If the titer of virus is high, then virtually all fibroblasts will be infected and no selection is required. If the titer is very low, then it is necessary to use a retroviral vector that has a selectable marker, such as neo or his. Once the fibroblasts have been efficiently infected, the fibroblasts are analyzed to determine whether protein is produced.

The engineered fibroblasts are then transplanted onto the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads.

Example 17: Gene Therapy Using Endogenous Genes Corresponding To Polynucleotides of the Invention

Another method of gene therapy according to the present invention involves operably associating the endogenous polynucleotide sequence of the invention with a promoter via homologous recombination as described, for example, in U.S. Patent NO: 5,641,670, issued

June 24, 1997; International Publication NO: WO 96/29411, published September 26, 1996; International Publication NO: WO 94/12650, published August 4, 1994; Koller et al., *Proc. Natl. Acad. Sci. USA*, 86:8932-8935 (1989); and Zijlstra et al., *Nature*, 342:435-438 (1989).

This method involves the activation of a gene which is present in the target cells, but which is not expressed in the cells, or is expressed at a lower level than desired.

Polynucleotide constructs are made which contain a promoter and targeting sequences, which are homologous to the 5' non-coding sequence of endogenous polynucleotide sequence, flanking the promoter. The targeting sequence will be sufficiently near the 5' end of the polynucleotide sequence so the promoter will be operably linked to the endogenous sequence upon homologous recombination. The promoter and the targeting sequences can be amplified using PCR. Preferably, the amplified promoter contains distinct restriction enzyme sites on the 5' and 3' ends. Preferably, the 3' end of the first targeting sequence contains the same restriction enzyme site as the 5' end of the amplified promoter and the 5' end of the second targeting sequence contains the same restriction site as the 3' end of the amplified promoter.

The amplified promoter and the amplified targeting sequences are digested with the appropriate restriction enzymes and subsequently treated with calf intestinal phosphatase. The digested promoter and digested targeting sequences are added together in the presence of T4 DNA ligase. The resulting mixture is maintained under conditions appropriate for ligation of the two fragments. The construct is size fractionated on an agarose gel then purified by phenol extraction and ethanol precipitation.

In this Example, the polynucleotide constructs are administered as naked polynucleotides via electroporation. However, the polynucleotide constructs may also be administered with transfection-facilitating agents, such as liposomes, viral sequences, viral particles, precipitating agents, etc. Such methods of delivery are known in the art.

Once the cells are transfected, homologous recombination will take place which results in the promoter being operably linked to the endogenous polynucleotide sequence. This results in the expression of polynucleotide corresponding to the polynucleotide in the cell. Expression may be detected by immunological staining, or any other method known in the art.

Fibroblasts are obtained from a subject by skin biopsy. The resulting tissue is placed in DMEM + 10% fetal calf serum. Exponentially growing or early stationary phase

fibroblasts are trypsinized and rinsed from the plastic surface with nutrient medium. An aliquot of the cell suspension is removed for counting, and the remaining cells are subjected to centrifugation. The supernatant is aspirated and the pellet is resuspended in 5 ml of electroporation buffer (20 mM HEPES pH 7.3, 137 mM NaCl, 5 mM KCl, 0.7 mM Na₂HPO₄, 6 mM dextrose). The cells are recentrifuged, the supernatant aspirated, and the cells resuspended in electroporation buffer containing 1 mg/ml acetylated bovine serum albumin. The final cell suspension contains approximately 3×10^6 cells/ml. Electroporation should be performed immediately following resuspension.

Plasmid DNA is prepared according to standard techniques. For example, to construct a plasmid for targeting to the locus corresponding to the polynucleotide of the invention, plasmid pUC18 (MBI Fermentas, Amherst, NY) is digested with HindIII. The CMV promoter is amplified by PCR with an XbaI site on the 5' end and a BamHI site on the 3' end. Two non-coding sequences are amplified via PCR: one non-coding sequence (fragment 1) is amplified with a HindIII site at the 5' end and an Xba site at the 3' end; the other non-coding sequence (fragment 2) is amplified with a BamHI site at the 5' end and a HindIII site at the 3' end. The CMV promoter and the fragments (1 and 2) are digested with the appropriate enzymes (CMV promoter - XbaI and BamHI; fragment 1 - XbaI; fragment 2 - BamHI) and ligated together. The resulting ligation product is digested with HindIII, and ligated with the HindIII-digested pUC18 plasmid.

Plasmid DNA is added to a sterile cuvette with a 0.4 cm electrode gap (Bio-Rad). The final DNA concentration is generally at least 120 µg/ml. 0.5 ml of the cell suspension (containing approximately 1.5×10^6 cells) is then added to the cuvette, and the cell suspension and DNA solutions are gently mixed. Electroporation is performed with a Gene-Pulser apparatus (Bio-Rad). Capacitance and voltage are set at 960 µF and 250-300 V, respectively. As voltage increases, cell survival decreases, but the percentage of surviving cells that stably incorporate the introduced DNA into their genome increases dramatically. Given these parameters, a pulse time of approximately 14-20 mSec should be observed.

Electroporated cells are maintained at room temperature for approximately 5 min, and the contents of the cuvette are then gently removed with a sterile transfer pipette. The cells are added directly to 10 ml of prewarmed nutrient media (DMEM with 15% calf serum) in a 10 cm dish and incubated at 37 degree C. The following day, the media is aspirated and replaced with 10 ml of fresh media and incubated for a further 16-24 hours.

The engineered fibroblasts are then injected into the host, either alone or after having been grown to confluence on cytodex 3 microcarrier beads. The fibroblasts now produce the protein product. The fibroblasts can then be introduced into a patient as described above.

5 *Example 18: Method of Treatment Using Gene Therapy - In Vivo*

Another aspect of the present invention is using *in vivo* gene therapy methods to treat disorders, diseases and conditions. The gene therapy method relates to the introduction of naked nucleic acid (DNA, RNA, and antisense DNA or RNA) sequences into an animal to
10 increase or decrease the expression of the polypeptide. The polynucleotide of the present invention may be operatively linked to a promoter or any other genetic elements necessary for the expression of the polypeptide by the target tissue. Such gene therapy and delivery techniques and methods are known in the art, see, for example, WO90/11092, WO98/11779; U.S. Patent NO. 5693622, 5705151, 5580859; Tabata et al., *Cardiovasc. Res.* 35(3):470-479
15 (1997); Chao et al., *Pharmacol. Res.* 35(6):517-522 (1997); Wolff, *Neuromuscul. Disord.* 7(5):314-318 (1997); Schwartz et al., *Gene Ther.* 3(5):405-411 (1996); Tsurumi et al., *Circulation* 94(12):3281-3290 (1996) (incorporated herein by reference).

The polynucleotide constructs may be delivered by any method that delivers injectable materials to the cells of an animal, such as, injection into the interstitial space of
20 tissues (heart, muscle, skin, lung, liver, intestine and the like). The polynucleotide constructs can be delivered in a pharmaceutically acceptable liquid or aqueous carrier.

The term "naked" polynucleotide, DNA or RNA, refers to sequences that are free from any delivery vehicle that acts to assist, promote, or facilitate entry into the cell, including viral sequences, viral particles, liposome formulations, lipofectin or precipitating
25 agents and the like. However, the polynucleotides of the present invention may also be delivered in liposome formulations (such as those taught in Felgner P.L. et al. (1995) *Ann. NY Acad. Sci.* 772:126-139 and Abdallah B. et al. (1995) *Biol. Cell* 85(1):1-7) which can be prepared by methods well known to those skilled in the art.

The polynucleotide vector constructs used in the gene therapy method are preferably
30 constructs that will not integrate into the host genome nor will they contain sequences that allow for replication. Any strong promoter known to those skilled in the art can be used for driving the expression of DNA. Unlike other gene therapies techniques, one major advantage

of introducing naked nucleic acid sequences into target cells is the transitory nature of the polynucleotide synthesis in the cells. Studies have shown that non-replicating DNA sequences can be introduced into cells to provide production of the desired polypeptide for periods of up to six months.

5 The polynucleotide construct can be delivered to the interstitial space of tissues within the an animal, including of muscle, skin, brain, lung, liver, spleen, bone marrow, thymus, heart, lymph, blood, bone, cartilage, pancreas, kidney, gall bladder, stomach, intestine, testis, ovary, uterus, rectum, nervous system, eye, gland, and connective tissue. Interstitial space of the tissues comprises the intercellular fluid, mucopolysaccharide matrix among the reticular
10 fibers of organ tissues, elastic fibers in the walls of vessels or chambers, collagen fibers of fibrous tissues, or that same matrix within connective tissue ensheathing muscle cells or in the lacunae of bone. It is similarly the space occupied by the plasma of the circulation and the lymph fluid of the lymphatic channels. Delivery to the interstitial space of muscle tissue is preferred for the reasons discussed below. They may be conveniently delivered by
15 injection into the tissues comprising these cells. They are preferably delivered to and expressed in persistent, non-dividing cells which are differentiated, although delivery and expression may be achieved in non-differentiated or less completely differentiated cells, such as, for example, stem cells of blood or skin fibroblasts. *In vivo* muscle cells are particularly competent in their ability to take up and express polynucleotides.

20 For the naked polynucleotide injection, an effective dosage amount of DNA or RNA will be in the range of from about 0.05 g/kg body weight to about 50 mg/kg body weight. Preferably the dosage will be from about 0.005 mg/kg to about 20 mg/kg and more preferably from about 0.05 mg/kg to about 5 mg/kg. Of course, as the artisan of ordinary skill will appreciate, this dosage will vary according to the tissue site of injection. The appropriate and
25 effective dosage of nucleic acid sequence can readily be determined by those of ordinary skill in the art and may depend on the condition being treated and the route of administration. The preferred route of administration is by the parenteral route of injection into the interstitial space of tissues. However, other parenteral routes may also be used, such as, inhalation of an aerosol formulation particularly for delivery to lungs or bronchial tissues, throat or mucous
30 membranes of the nose. In addition, naked polynucleotide constructs can be delivered to arteries during angioplasty by the catheter used in the procedure.

The dose response effects of injected polynucleotide in muscle *in vivo* is determined

as follows. Suitable template DNA for production of mRNA coding for polypeptide of the present invention is prepared in accordance with a standard recombinant DNA methodology. The template DNA, which may be either circular or linear, is either used as naked DNA or complexed with liposomes. The quadriceps muscles of mice are then injected with various
5 amounts of the template DNA.

Five to six week old female and male Balb/C mice are anesthetized by intraperitoneal injection with 0.3 ml of 2.5% Avertin. A 1.5 cm incision is made on the anterior thigh, and the quadriceps muscle is directly visualized. The template DNA is injected in 0.1 ml of carrier in a 1 cc syringe through a 27 gauge needle over one minute, approximately 0.5 cm
10 from the distal insertion site of the muscle into the knee and about 0.2 cm deep. A suture is placed over the injection site for future localization, and the skin is closed with stainless steel clips.

After an appropriate incubation time (e.g., 7 days) muscle extracts are prepared by excising the entire quadriceps. Every fifth 15 um cross-section of the individual quadriceps
15 muscles is histochemically stained for protein expression. A time course for protein expression may be done in a similar fashion except that quadriceps from different mice are harvested at different times. Persistence of DNA in muscle following injection may be determined by Southern blot analysis after preparing total cellular DNA and HIRT supernatants from injected and control mice. The results of the above experimentation in
20 mice can be use to extrapolate proper dosages and other treatment parameters in humans and other animals using naked DNA.

Example 19: Transgenic Animals

25 The polypeptides of the invention can also be expressed in transgenic animals. Animals of any species, including, but not limited to, mice, rats, rabbits, hamsters, guinea pigs, pigs, micro-pigs, goats, sheep, cows and non-human primates, e.g., baboons, monkeys, and chimpanzees may be used to generate transgenic animals. In a specific embodiment, techniques described herein or otherwise known in the art, are used to express polypeptides
30 of the invention in humans. as part of a gene therapy protocol.

Any technique known in the art may be used to introduce the transgene (i.e., polynucleotides of the invention) into animals to produce the founder lines of transgenic

animals. Such techniques include, but are not limited to, pronuclear microinjection (Paterson et al., Appl. Microbiol. Biotechnol. 40:691-698 (1994); Carver et al., Biotechnology (NY) 11:1263-1270 (1993); Wright et al., Biotechnology (NY) 9:830-834 (1991); and Hoppe et al., U.S. Pat. No. 4,873,191 (1989)); retrovirus mediated gene transfer into germ lines (Van der Putten et al., Proc. Natl. Acad. Sci., USA 82:6148-6152 (1985)), blastocysts or embryos; gene targeting in embryonic stem cells (Thompson et al., Cell 56:313-321 (1989)); electroporation of cells or embryos (Lo, 1983, Mol Cell. Biol. 3:1803-1814 (1983)); introduction of the polynucleotides of the invention using a gene gun (see, e.g., Ulmer et al., Science 259:1745 (1993)); introducing nucleic acid constructs into embryonic pluripotent stem cells and transferring the stem cells back into the blastocyst; and sperm-mediated gene transfer (Lavitrano et al., Cell 57:717-723 (1989); etc. For a review of such techniques, see Gordon, "Transgenic Animals," Intl. Rev. Cytol. 115:171-229 (1989), which is incorporated by reference herein in its entirety.

Any technique known in the art may be used to produce transgenic clones containing polynucleotides of the invention, for example, nuclear transfer into enucleated oocytes of nuclei from cultured embryonic, fetal, or adult cells induced to quiescence (Campell et al., Nature 380:64-66 (1996); Wilmut et al., Nature 385:810-813 (1997)).

The present invention provides for transgenic animals that carry the transgene in all their cells, as well as animals which carry the transgene in some, but not all their cells, *i.e.*, mosaic animals or chimeric. The transgene may be integrated as a single transgene or as multiple copies such as in concatamers, *e.g.*, head-to-head tandems or head-to-tail tandems. The transgene may also be selectively introduced into and activated in a particular cell type by following, for example, the teaching of Lasko et al. (Lasko et al., Proc. Natl. Acad. Sci. USA 89:6232-6236 (1992)). The regulatory sequences required for such a cell-type specific activation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art. When it is desired that the polynucleotide transgene be integrated into the chromosomal site of the endogenous gene, gene targeting is preferred. Briefly, when such a technique is to be utilized, vectors containing some nucleotide sequences homologous to the endogenous gene are designed for the purpose of integrating, via homologous recombination with chromosomal sequences, into and disrupting the function of the nucleotide sequence of the endogenous gene. The transgene may also be selectively introduced into a particular cell type, thus inactivating the endogenous gene in only that cell type, by following, for example,

the teaching of Gu et al. (Gu et al., Science 265:103-106 (1994)). The regulatory sequences required for such a cell-type specific inactivation will depend upon the particular cell type of interest, and will be apparent to those of skill in the art.

Once transgenic animals have been generated, the expression of the recombinant gene may be assayed utilizing standard techniques. Initial screening may be accomplished by Southern blot analysis or PCR techniques to analyze animal tissues to verify that integration of the transgene has taken place. The level of mRNA expression of the transgene in the tissues of the transgenic animals may also be assessed using techniques which include, but are not limited to, Northern blot analysis of tissue samples obtained from the animal, *in situ* hybridization analysis, and reverse transcriptase-PCR (rt-PCR). Samples of transgenic gene-expressing tissue may also be evaluated immunocytochemically or immunohistochemically using antibodies specific for the transgene product.

Once the founder animals are produced, they may be bred, inbred, outbred, or crossbred to produce colonies of the particular animal. Examples of such breeding strategies include, but are not limited to: outbreeding of founder animals with more than one integration site in order to establish separate lines; inbreeding of separate lines in order to produce compound transgenics that express the transgene at higher levels because of the effects of additive expression of each transgene; crossing of heterozygous transgenic animals to produce animals homozygous for a given integration site in order to both augment expression and eliminate the need for screening of animals by DNA analysis; crossing of separate homozygous lines to produce compound heterozygous or homozygous lines; and breeding to place the transgene on a distinct background that is appropriate for an experimental model of interest.

Transgenic animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

Example 20: Knock-Out Animals

30

Endogenous gene expression can also be reduced by inactivating or "knocking out" the gene and/or its promoter using targeted homologous recombination. (E.g., see Smithies

et al., Nature 317:230-234 (1985); Thomas & Capecchi, Cell 51:503-512 (1987); Thompson et al., Cell 5:313-321 (1989); each of which is incorporated by reference herein in its entirety). For example, a mutant, non-functional polynucleotide of the invention (or a completely unrelated DNA sequence) flanked by DNA homologous to the endogenous polynucleotide sequence (either the coding regions or regulatory regions of the gene) can be used, with or without a selectable marker and/or a negative selectable marker, to transfect cells that express polypeptides of the invention *in vivo*. In another embodiment, techniques known in the art are used to generate knockouts in cells that contain, but do not express the gene of interest. Insertion of the DNA construct, via targeted homologous recombination, results in inactivation of the targeted gene. Such approaches are particularly suited in research and agricultural fields where modifications to embryonic stem cells can be used to generate animal offspring with an inactive targeted gene (*e.g.*, see Thomas & Capecchi 1987 and Thompson 1989, *supra*). However this approach can be routinely adapted for use in humans provided the recombinant DNA constructs are directly administered or targeted to the required site *in vivo* using appropriate viral vectors that will be apparent to those of skill in the art.

In further embodiments of the invention, cells that are genetically engineered to express the polypeptides of the invention, or alternatively, that are genetically engineered not to express the polypeptides of the invention (*e.g.*, knockouts) are administered to a patient *in vivo*. Such cells may be obtained from the patient (*i.e.*, animal, including human) or an MHC compatible donor and can include, but are not limited to fibroblasts, bone marrow cells, blood cells (*e.g.*, lymphocytes), adipocytes, muscle cells, endothelial cells etc. The cells are genetically engineered *in vitro* using recombinant DNA techniques to introduce the coding sequence of polypeptides of the invention into the cells, or alternatively, to disrupt the coding sequence and/or endogenous regulatory sequence associated with the polypeptides of the invention, *e.g.*, by transduction (using viral vectors, and preferably vectors that integrate the transgene into the cell genome) or transfection procedures, including, but not limited to, the use of plasmids, cosmids, YACs, naked DNA, electroporation, liposomes, etc. The coding sequence of the polypeptides of the invention can be placed under the control of a strong constitutive or inducible promoter or promoter/enhancer to achieve expression, and preferably secretion, of the polypeptides of the invention. The engineered cells which express and preferably secrete the polypeptides of the invention can be introduced into the

patient systemically. e.g., in the circulation, or intraperitoneally.

Alternatively, the cells can be incorporated into a matrix and implanted in the body, e.g., genetically engineered fibroblasts can be implanted as part of a skin graft; genetically engineered endothelial cells can be implanted as part of a lymphatic or vascular graft. (See, 5 for example, Anderson et al. U.S. Patent No. 5,399,349; and Mulligan & Wilson, U.S. Patent No. 5,460,959 each of which is incorporated by reference herein in its entirety).

When the cells to be administered are non-autologous or non-MHC compatible cells, they can be administered using well known techniques which prevent the development of a host immune response against the introduced cells. For example, the cells may be introduced 10 in an encapsulated form which, while allowing for an exchange of components with the immediate extracellular environment, does not allow the introduced cells to be recognized by the host immune system.

Transgenic and "knock-out" animals of the invention have uses which include, but are not limited to, animal model systems useful in elaborating the biological function of 15 polypeptides of the present invention, studying conditions and/or disorders associated with aberrant expression, and in screening for compounds effective in ameliorating such conditions and/or disorders.

20 *Example 22: Assays Detecting Stimulation or Inhibition of B cell Proliferation and Differentiation*

Generation of functional humoral immune responses requires both soluble and cognate signaling between B-lineage cells and their microenvironment. Signals may impart a positive stimulus that allows a B-lineage cell to continue its programmed development, or a 25 negative stimulus that instructs the cell to arrest its current developmental pathway. To date, numerous stimulatory and inhibitory signals have been found to influence B cell responsiveness including IL-2, IL-4, IL-5, IL-6, IL-7, IL10, IL-13, IL-14 and IL-15. Interestingly, these signals are by themselves weak effectors but can, in combination with various co-stimulatory proteins, induce activation, proliferation, differentiation, homing, 30 tolerance and death among B cell populations.

One of the best studied classes of B-cell co-stimulatory proteins is the TNF-superfamily. Within this family CD40, CD27, and CD30 along with their respective ligands

CD154, CD70, and CD153 have been found to regulate a variety of immune responses. Assays which allow for the detection and/or observation of the proliferation and differentiation of these B-cell populations and their precursors are valuable tools in determining the effects various proteins may have on these B-cell populations in terms of proliferation and differentiation. Listed below are two assays designed to allow for the detection of the differentiation, proliferation, or inhibition of B-cell populations and their precursors.

In Vitro Assay- Agonists or antagonists of the invention can be assessed for its ability to induce activation, proliferation, differentiation or inhibition and/or death in B-cell populations and their precursors. The activity of the agonists or antagonists of the invention on purified human tonsillar B cells, measured qualitatively over the dose range from 0.1 to 10,000 ng/mL, is assessed in a standard B-lymphocyte co-stimulation assay in which purified tonsillar B cells are cultured in the presence of either formalin-fixed *Staphylococcus aureus* Cowan I (SAC) or immobilized anti-human IgM antibody as the priming agent. Second signals such as IL-2 and IL-15 synergize with SAC and IgM crosslinking to elicit B cell proliferation as measured by tritiated-thymidine incorporation. Novel synergizing agents can be readily identified using this assay. The assay involves isolating human tonsillar B cells by magnetic bead (MACS) depletion of CD3-positive cells. The resulting cell population is greater than 95% B cells as assessed by expression of CD45R(B220).

Various dilutions of each sample are placed into individual wells of a 96-well plate to which are added 10^5 B-cells suspended in culture medium (RPMI 1640 containing 10% FBS, 5×10^{-5} M 2ME, 100U/ml penicillin, 10ug/ml streptomycin, and 10^{-5} dilution of SAC) in a total volume of 150ul. Proliferation or inhibition is quantitated by a 20h pulse (1uCi/well) with 3 H-thymidine (6.7 Ci/mM) beginning 72h post factor addition. The positive and negative controls are IL2 and medium respectively.

In Vivo Assay- BALB/c mice are injected (i.p.) twice per day with buffer only, or 2 mg/Kg of agonists or antagonists of the invention, or truncated forms thereof. Mice receive this treatment for 4 consecutive days, at which time they are sacrificed and various tissues and serum collected for analyses. Comparison of H&E sections from normal spleens and spleens treated with agonists or antagonists of the invention identify the results of the activity of the agonists or antagonists on spleen cells, such as the diffusion of peri-arterial lymphatic sheaths, and/or significant increases in the nucleated cellularity of the red pulp regions, which

may indicate the activation of the differentiation and proliferation of B-cell populations. Immunohistochemical studies using a B cell marker, anti-CD45R(B220), are used to determine whether any physiological changes to splenic cells, such as splenic disorganization, are due to increased B-cell representation within loosely defined B-cell zones that infiltrate established T-cell regions.

Flow cytometric analyses of the spleens from mice treated with agonist or antagonist is used to indicate whether the agonists or antagonists specifically increases the proportion of ThB+, CD45R(B220)dull B cells over that which is observed in control mice. Likewise, a predicted consequence of increased mature B-cell representation in vivo is a relative increase in serum Ig titers. Accordingly, serum IgM and IgA levels are compared between buffer and agonists or antagonists-treated mice.

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

15

Example 23: T Cell Proliferation Assay

A CD3-induced proliferation assay is performed on PBMCs and is measured by the uptake of ³H-thymidine. The assay is performed as follows. Ninety-six well plates are coated with 100 µl/well of mAb to CD3 (HIT3a, Pharmingen) or isotype-matched control mAb (B33.1) overnight at 4 degrees C (1 µg/ml in .05M bicarbonate buffer, pH 9.5), then washed three times with PBS. PBMC are isolated by F/H gradient centrifugation from human peripheral blood and added to quadruplicate wells (5 x 10⁴/well) of mAb coated plates in RPMI containing 10% FCS and P/S in the presence of varying concentrations of agonists or antagonists of the invention (total volume 200 µl). Relevant protein buffer and medium alone are controls. After 48 hr. culture at 37 degrees C, plates are spun for 2 min. at 1000 rpm and 100 µl of supernatant is removed and stored -20 degrees C for measurement of IL-2 (or other cytokines) if effect on proliferation is observed. Wells are supplemented with 100 µl of medium containing 0.5 uCi of ³H-thymidine and cultured at 37 degrees C for 18-24 hr. Wells are harvested and incorporation of ³H-thymidine used as a measure of proliferation. Anti-CD3 alone is the positive control for proliferation. IL-2 (100 U/ml) is also used as a control which enhances proliferation. Control antibody which does not induce proliferation

of T cells is used as the negative controls for the effects of agonists or antagonists of the invention.

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 24: Effect of Agonists or Antagonists of the Invention on the Expression of MHC Class II, Costimulatory and Adhesion Molecules and Cell Differentiation of Monocytes and Monocyte-Derived Human Dendritic Cells

10

Dendritic cells are generated by the expansion of proliferating precursors found in the peripheral blood: adherent PBMC or elutriated monocytic fractions are cultured for 7-10 days with GM-CSF (50 ng/ml) and IL-4 (20 ng/ml). These dendritic cells have the characteristic phenotype of immature cells (expression of CD1, CD80, CD86, CD40 and MHC class II antigens). Treatment with activating factors, such as TNF- α , causes a rapid change in surface phenotype (increased expression of MHC class I and II, costimulatory and adhesion molecules, downregulation of FC γ RII, upregulation of CD83). These changes correlate with increased antigen-presenting capacity and with functional maturation of the dendritic cells.

15

FACS analysis of surface antigens is performed as follows. Cells are treated 1-3 days with increasing concentrations of agonist or antagonist of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degrees C. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

20

25

Effect on the production of cytokines. Cytokines generated by dendritic cells, in particular IL-12, are important in the initiation of T-cell dependent immune responses. IL-12 strongly influences the development of Th1 helper T-cell immune response, and induces cytotoxic T and NK cell function. An ELISA is used to measure the IL-12 release as follows. Dendritic cells (10^6 /ml) are treated with increasing concentrations of agonists or antagonists of the invention for 24 hours. LPS (100 ng/ml) is added to the cell culture as positive control. Supernatants from the cell cultures are then collected and analyzed for IL-12 content using

30

commercial ELISA kit (e.g., R & D Systems (Minneapolis, MN)). The standard protocols provided with the kits are used.

Effect on the expression of MHC Class II, costimulatory and adhesion molecules. Three major families of cell surface antigens can be identified on monocytes: adhesion molecules, molecules involved in antigen presentation, and Fc receptor. Modulation of the expression of MHC class II antigens and other costimulatory molecules, such as B7 and ICAM-1, may result in changes in the antigen presenting capacity of monocytes and ability to induce T cell activation. Increase expression of Fc receptors may correlate with improved monocyte cytotoxic activity, cytokine release and phagocytosis.

FACS analysis is used to examine the surface antigens as follows. Monocytes are treated 1-5 days with increasing concentrations of agonists or antagonists of the invention or LPS (positive control), washed with PBS containing 1% BSA and 0.02 mM sodium azide, and then incubated with 1:20 dilution of appropriate FITC- or PE-labeled monoclonal antibodies for 30 minutes at 4 degreesC. After an additional wash, the labeled cells are analyzed by flow cytometry on a FACScan (Becton Dickinson).

Monocyte activation and/or increased survival. Assays for molecules that activate (or alternatively, inactivate) monocytes and/or increase monocyte survival (or alternatively, decrease monocyte survival) are known in the art and may routinely be applied to determine whether a molecule of the invention functions as an inhibitor or activator of monocytes. Agonists or antagonists of the invention can be screened using the three assays described below. For each of these assays, Peripheral blood mononuclear cells (PBMC) are purified from single donor leukopacks (American Red Cross, Baltimore, MD) by centrifugation through a Histopaque gradient (Sigma). Monocytes are isolated from PBMC by counterflow centrifugal elutriation.

Monocyte Survival Assay. Human peripheral blood monocytes progressively lose viability when cultured in absence of serum or other stimuli. Their death results from internally regulated process (apoptosis). Addition to the culture of activating factors, such as TNF-alpha dramatically improves cell survival and prevents DNA fragmentation. Propidium iodide (PI) staining is used to measure apoptosis as follows. Monocytes are cultured for 48 hours in

polypropylene tubes in serum-free medium (positive control), in the presence of 100 ng/ml TNF-alpha (negative control), and in the presence of varying concentrations of the compound to be tested. Cells are suspended at a concentration of 2×10^6 /ml in PBS containing PI at a final concentration of 5 μ g/ml, and then incubated at room temperature for 5 minutes before

5 FACSscan analysis. PI uptake has been demonstrated to correlate with DNA fragmentation in this experimental paradigm.

Effect on cytokine release. An important function of monocytes/macrophages is their regulatory activity on other cellular populations of the immune system through the release of

10 cytokines after stimulation. An ELISA to measure cytokine release is performed as follows. Human monocytes are incubated at a density of 5×10^5 cells/ml with increasing concentrations of agonists or antagonists of the invention and under the same conditions, but in the absence of agonists or antagonists. For IL-12 production, the cells are primed overnight with IFN (100 U/ml) in presence of agonist or antagonist of the invention. LPS (10

15 ng/ml) is then added. Conditioned media are collected after 24h and kept frozen until use. Measurement of TNF-alpha, IL-10, MCP-1 and IL-8 is then performed using a commercially available ELISA kit (e. g, R & D Systems (Minneapolis, MN)) and applying the standard protocols provided with the kit.

20 Oxidative burst. Purified monocytes are plated in 96-w plate at 2×10^5 cell/well. Increasing concentrations of agonists or antagonists of the invention are added to the wells in a total volume of 0.2 ml culture medium (RPMI 1640 + 10% FCS, glutamine and antibiotics). After 3 days incubation, the plates are centrifuged and the medium is removed from the wells. To the macrophage monolayers, 0.2 ml per well of phenol red solution (140 mM NaCl, 10 mM

25 potassium phosphate buffer pH 7.0, 5.5 mM dextrose, 0.56 mM phenol red and 19 U/ml of HRPO) is added, together with the stimulant (200 nM PMA). The plates are incubated at 37°C for 2 hours and the reaction is stopped by adding 20 μ l 1N NaOH per well. The absorbance is read at 610 nm. To calculate the amount of H_2O_2 produced by the macrophages, a standard curve of a H_2O_2 solution of known molarity is performed for each

30 experiment.

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test

the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 25: Biological Effects of Agonists or Antagonists of the Invention

5

Astrocyte and Neuronal Assays.

Agonists or antagonists of the invention, expressed in *Escherichia coli* and purified as described above, can be tested for activity in promoting the survival, neurite outgrowth, or phenotypic differentiation of cortical neuronal cells and for inducing the proliferation of glial fibrillary acidic protein immunopositive cells, astrocytes. The selection of cortical cells for the bioassay is based on the prevalent expression of FGF-1 and FGF-2 in cortical structures and on the previously reported enhancement of cortical neuronal survival resulting from FGF-2 treatment. A thymidine incorporation assay, for example, can be used to elucidate an agonist or antagonist of the invention's activity on these cells.

Moreover, previous reports describing the biological effects of FGF-2 (basic FGF) on cortical or hippocampal neurons *in vitro* have demonstrated increases in both neuron survival and neurite outgrowth (Walicke et al., "Fibroblast growth factor promotes survival of dissociated hippocampal neurons and enhances neurite extension." *Proc. Natl. Acad. Sci. USA* 83:3012-3016. (1986), assay herein incorporated by reference in its entirety). However, reports from experiments done on PC-12 cells suggest that these two responses are not necessarily synonymous and may depend on not only which FGF is being tested but also on which receptor(s) are expressed on the target cells. Using the primary cortical neuronal culture paradigm, the ability of an agonist or antagonist of the invention to induce neurite outgrowth can be compared to the response achieved with FGF-2 using, for example, a thymidine incorporation assay.

Fibroblast and endothelial cell assays.

Human lung fibroblasts are obtained from Clonetics (San Diego, CA) and maintained in growth media from Clonetics. Dermal microvascular endothelial cells are obtained from Cell Applications (San Diego, CA). For proliferation assays, the human lung fibroblasts and

dermal microvascular endothelial cells can be cultured at 5,000 cells/well in a 96-well plate for one day in growth medium. The cells are then incubated for one day in 0.1% BSA basal medium. After replacing the medium with fresh 0.1% BSA medium, the cells are incubated with the test proteins for 3 days. Alamar Blue (Alamar Biosciences, Sacramento, CA) is added to each well to a final concentration of 10%. The cells are incubated for 4 hr. Cell viability is measured by reading in a CytoFluor fluorescence reader. For the PGE₂ assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or agonists or antagonists of the invention with or without IL-1 α for 24 hours. The supernatants are collected and assayed for PGE₂ by EIA kit (Cayman, Ann Arbor, MI). For the IL-6 assays, the human lung fibroblasts are cultured at 5,000 cells/well in a 96-well plate for one day. After a medium change to 0.1% BSA basal medium, the cells are incubated with FGF-2 or with or without agonists or antagonists of the invention IL-1 α for 24 hours. The supernatants are collected and assayed for IL-6 by ELISA kit (Endogen, Cambridge, MA).

Human lung fibroblasts are cultured with FGF-2 or agonists or antagonists of the invention for 3 days in basal medium before the addition of Alamar Blue to assess effects on growth of the fibroblasts. FGF-2 should show a stimulation at 10 - 2500 ng/ml which can be used to compare stimulation with agonists or antagonists of the invention.

20 Parkinson Models.

The loss of motor function in Parkinson's disease is attributed to a deficiency of striatal dopamine resulting from the degeneration of the nigrostriatal dopaminergic projection neurons. An animal model for Parkinson's that has been extensively characterized involves the systemic administration of 1-methyl-4 phenyl 1,2,3,6-tetrahydropyridine (MPTP). In the CNS, MPTP is taken-up by astrocytes and catabolized by monoamine oxidase B to 1-methyl-4-phenyl pyridine (MPP⁺) and released. Subsequently, MPP⁺ is actively accumulated in dopaminergic neurons by the high-affinity reuptake transporter for dopamine. MPP⁺ is then concentrated in mitochondria by the electrochemical gradient and selectively inhibits nicotinamide adenine disphosphate: ubiquinone oxidoreductionase (complex I), thereby interfering with electron transport and eventually generating oxygen radicals.

It has been demonstrated in tissue culture paradigms that FGF-2 (basic FGF) has

trophic activity towards nigral dopaminergic neurons (Ferrari et al., Dev. Biol. 1989). Recently, Dr. Unsicker's group has demonstrated that administering FGF-2 in gel foam implants in the striatum results in the near complete protection of nigral dopaminergic neurons from the toxicity associated with MPTP exposure (Otto and Unsicker, J. Neuroscience, 1990).

Based on the data with FGF-2, agonists or antagonists of the invention can be evaluated to determine whether it has an action similar to that of FGF-2 in enhancing dopaminergic neuronal survival *in vitro* and it can also be tested *in vivo* for protection of dopaminergic neurons in the striatum from the damage associated with MPTP treatment. The potential effect of an agonist or antagonist of the invention is first examined *in vitro* in a dopaminergic neuronal cell culture paradigm. The cultures are prepared by dissecting the midbrain floor plate from gestation day 14 Wistar rat embryos. The tissue is dissociated with trypsin and seeded at a density of 200,000 cells/cm² on polyorthinine-laminin coated glass coverslips. The cells are maintained in Dulbecco's Modified Eagle's medium and F12 medium containing hormonal supplements (N1). The cultures are fixed with paraformaldehyde after 8 days *in vitro* and are processed for tyrosine hydroxylase, a specific marker for dopaminergic neurons, immunohistochemical staining. Dissociated cell cultures are prepared from embryonic rats. The culture medium is changed every third day and the factors are also added at that time.

Since the dopaminergic neurons are isolated from animals at gestation day 14, a developmental time which is past the stage when the dopaminergic precursor cells are proliferating, an increase in the number of tyrosine hydroxylase immunopositive neurons would represent an increase in the number of dopaminergic neurons surviving *in vitro*. Therefore, if an agonist or antagonist of the invention acts to prolong the survival of dopaminergic neurons, it would suggest that the agonist or antagonist may be involved in Parkinson's Disease.

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 26: The Effect of Agonists or Antagonists of the Invention on the Growth of

Vascular Endothelial Cells

On day 1, human umbilical vein endothelial cells (HUVEC) are seeded at 2.5×10^4 cells/35 mm dish density in M199 medium containing 4% fetal bovine serum (FBS), 16 units/ml heparin, and 50 units/ml endothelial cell growth supplements (ECGS, Biotechnology, Inc.). On day 2, the medium is replaced with M199 containing 10% FBS, 8 units/ml heparin. An agonist or antagonist of the invention, and positive controls, such as VEGF and basic FGF (bFGF) are added, at varying concentrations. On days 4 and 6, the medium is replaced. On day 8, cell number is determined with a Coulter Counter.

10 An increase in the number of HUVEC cells indicates that the compound of the invention may proliferate vascular endothelial cells, while a decrease in the number of HUVEC cell indicates that the compound of the invention inhibits vascular endothelial cells.

The studies described in this example tested activity of a polypeptide of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity
15 of polynucleotides (e.g., gene therapy), agonists, and/or antagonists of the invention.

Example 27: Rat Corneal Wound Healing Model

This animal model shows the effect of an agonist or antagonist of the invention on
20 neovascularization. The experimental protocol includes:

- a) Making a 1-1.5 mm long incision from the center of cornea into the stromal layer.
- b) Inserting a spatula below the lip of the incision facing the outer corner of the eye.
- 25 c) Making a pocket (its base is 1-1.5 mm from the edge of the eye).
- d) Positioning a pellet, containing 50ng- 5ug of an agonist or antagonist of the invention, within the pocket.
- e) Treatment with an agonist or antagonist of the invention can also be applied topically to the corneal wounds in a dosage range of 20mg - 500mg (daily treatment for five
30 days).

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test

the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 28: Diabetic Mouse and Glucocorticoid-Impaired Wound Healing Models

5 A. *Diabetic db+/db+ Mouse Model.*

To demonstrate that an agonist or antagonist of the invention accelerates the healing process, the genetically diabetic mouse model of wound healing is used. The full thickness wound healing model in the db+/db+ mouse is a well characterized, clinically relevant and reproducible model of impaired wound healing. Healing of the diabetic wound is dependent
10 on formation of granulation tissue and re-epithelialization rather than contraction (Gartner, M.H. *et al.*, *J. Surg. Res.* 52:389 (1992); Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)).

The diabetic animals have many of the characteristic features observed in Type II diabetes mellitus. Homozygous (db+/db+) mice are obese in comparison to their normal
15 heterozygous (db+/+m) littermates. Mutant diabetic (db+/db+) mice have a single autosomal recessive mutation on chromosome 4 (db+) (Coleman *et al.* *Proc. Natl. Acad. Sci. USA* 77:283-293 (1982)). Animals show polyphagia, polydipsia and polyuria. Mutant diabetic mice (db+/db+) have elevated blood glucose, increased or normal insulin levels, and suppressed cell-mediated immunity (Mandel *et al.*, *J. Immunol.* 120:1375 (1978); Debray-
20 Sachs, M. *et al.*, *Clin. Exp. Immunol.* 51(1):1-7 (1983); Leiter *et al.*, *Am. J. of Pathol.* 114:46-55 (1985)). Peripheral neuropathy, myocardial complications, and microvascular lesions, basement membrane thickening and glomerular filtration abnormalities have been described in these animals (Norido, F. *et al.*, *Exp. Neurol.* 83(2):221-232 (1984); Robertson *et al.*, *Diabetes* 29(1):60-67 (1980); Giacomelli *et al.*, *Lab Invest.* 40(4):460-473 (1979); Coleman,
25 D.L., *Diabetes* 31 (Suppl):1-6 (1982)). These homozygous diabetic mice develop hyperglycemia that is resistant to insulin analogous to human type II diabetes (Mandel *et al.*, *J. Immunol.* 120:1375-1377 (1978)).

The characteristics observed in these animals suggests that healing in this model may be similar to the healing observed in human diabetes (Greenhalgh, *et al.*, *Am. J. of Pathol.*
30 136:1235-1246 (1990)).

Genetically diabetic female C57BL/KsJ (db+/db+) mice and their non-diabetic (db+/+m) heterozygous littermates are used in this study (Jackson Laboratories). The

animals are purchased at 6 weeks of age and are 8 weeks old at the beginning of the study. Animals are individually housed and received food and water ad libitum. All manipulations are performed using aseptic techniques. The experiments are conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

Wounding protocol is performed according to previously reported methods (Tsuboi, R. and Rifkin, D.B., *J. Exp. Med.* 172:245-251 (1990)). Briefly, on the day of wounding, animals are anesthetized with an intraperitoneal injection of Avertin (0.01 mg/mL), 2,2,2-tribromoethanol and 2-methyl-2-butanol dissolved in deionized water. The dorsal region of the animal is shaved and the skin washed with 70% ethanol solution and iodine. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is then created using a Keyes tissue punch. Immediately following wounding, the surrounding skin is gently stretched to eliminate wound expansion. The wounds are left open for the duration of the experiment. Application of the treatment is given topically for 5 consecutive days commencing on the day of wounding. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of surgery and at two day intervals thereafter. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

An agonist or antagonist of the invention is administered using at a range different doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for histology and immunohistochemistry. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes between biopsy sponges for further processing.

Three groups of 10 animals each (5 diabetic and 5 non-diabetic controls) are evaluated: 1) Vehicle placebo control, 2) untreated group, and 3) treated group.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis and obtaining the total square area of the wound. Contraction is then estimated by establishing

the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

5
$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned perpendicular to the wound surface (5mm) and cut using a Reichert-Jung microtome. Routine hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected
10 wounds. Histologic examination of the wounds are used to assess whether the healing process and the morphologic appearance of the repaired skin is altered by treatment with an agonist or antagonist of the invention. This assessment included verification of the presence of cell accumulation, inflammatory cells, capillaries, fibroblasts, re-epithelialization and epidermal maturity (Greenhalgh, D.G. *et al.*, *Am. J. Pathol.* 136:1235 (1990)). A calibrated
15 lens micrometer is used by a blinded observer.

Tissue sections are also stained immunohistochemically with a polyclonal rabbit anti-human keratin antibody using ABC Elite detection system. Human skin is used as a positive tissue control while non-immune IgG is used as a negative control. Keratinocyte growth is determined by evaluating the extent of reepithelialization of the wound using a calibrated
20 lens micrometer.

Proliferating cell nuclear antigen/cyclin (PCNA) in skin specimens is demonstrated by using anti-PCNA antibody (1:50) with an ABC Elite detection system. Human colon cancer served as a positive tissue control and human brain tissue is used as a negative tissue control. Each specimen included a section with omission of the primary antibody and
25 substitution with non-immune mouse IgG. Ranking of these sections is based on the extent of proliferation on a scale of 0-8, the lower side of the scale reflecting slight proliferation to the higher side reflecting intense proliferation.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

30

B. Steroid Impaired Rat Model

The inhibition of wound healing by steroids has been well documented in various *in vitro* and

in vivo systems (Wahl, Glucocorticoids and Wound healing. In: Anti-Inflammatory Steroid Action: Basic and Clinical Aspects. 280-302 (1989); Wahl *et al.*, *J. Immunol.* 115: 476-481 (1975); Werb *et al.*, *J. Exp. Med.* 147:1684-1694 (1978)). Glucocorticoids retard wound healing by inhibiting angiogenesis, decreasing vascular permeability (Ebert *et al.*, *An. Intern. Med.* 37:701-705 (1952)), fibroblast proliferation, and collagen synthesis (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978)) and producing a transient reduction of circulating monocytes (Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989)). The systemic administration of steroids to impaired wound healing is a well establish phenomenon in rats (Beck *et al.*, *Growth Factors.* 5: 295-304 (1991); Haynes *et al.*, *J. Clin. Invest.* 61: 703-797 (1978); Wahl, "Glucocorticoids and wound healing", In: Antiinflammatory Steroid Action: Basic and Clinical Aspects, Academic Press, New York, pp. 280-302 (1989); Pierce *et al.*, *Proc. Natl. Acad. Sci. USA* 86: 2229-2233 (1989)).

To demonstrate that an agonist or antagonist of the invention can accelerate the healing process, the effects of multiple topical applications of the agonist or antagonist on full thickness excisional skin wounds in rats in which healing has been impaired by the systemic administration of methylprednisolone is assessed.

Young adult male Sprague Dawley rats weighing 250-300 g (Charles River Laboratories) are used in this example. The animals are purchased at 8 weeks of age and are 9 weeks old at the beginning of the study. The healing response of rats is impaired by the systemic administration of methylprednisolone (17mg/kg/rat intramuscularly) at the time of wounding. Animals are individually housed and received food and water *ad libitum*. All manipulations are performed using aseptic techniques. This study is conducted according to the rules and guidelines of Human Genome Sciences, Inc. Institutional Animal Care and Use Committee and the Guidelines for the Care and Use of Laboratory Animals.

The wounding protocol is followed according to section A, above. On the day of wounding, animals are anesthetized with an intramuscular injection of ketamine (50 mg/kg) and xylazine (5 mg/kg). The dorsal region of the animal is shaved and the skin washed with 70% ethanol and iodine solutions. The surgical area is dried with sterile gauze prior to wounding. An 8 mm full-thickness wound is created using a Keyes tissue punch. The wounds are left open for the duration of the experiment. Applications of the testing materials

are given topically once a day for 7 consecutive days commencing on the day of wounding and subsequent to methylprednisolone administration. Prior to treatment, wounds are gently cleansed with sterile saline and gauze sponges.

Wounds are visually examined and photographed at a fixed distance at the day of
5 wounding and at the end of treatment. Wound closure is determined by daily measurement on days 1-5 and on day 8. Wounds are measured horizontally and vertically using a calibrated Jameson caliper. Wounds are considered healed if granulation tissue is no longer visible and the wound is covered by a continuous epithelium.

The agonist or antagonist of the invention is administered using at a range different
10 doses, from 4mg to 500mg per wound per day for 8 days in vehicle. Vehicle control groups received 50mL of vehicle solution.

Animals are euthanized on day 8 with an intraperitoneal injection of sodium
pentobarbital (300mg/kg). The wounds and surrounding skin are then harvested for
histology. Tissue specimens are placed in 10% neutral buffered formalin in tissue cassettes
15 between biopsy sponges for further processing.

Four groups of 10 animals each (5 with methylprednisolone and 5 without
glucocorticoid) are evaluated: 1) Untreated group 2) Vehicle placebo control 3) treated
groups.

Wound closure is analyzed by measuring the area in the vertical and horizontal axis
20 and obtaining the total area of the wound. Closure is then estimated by establishing the differences between the initial wound area (day 0) and that of post treatment (day 8). The wound area on day 1 is 64mm², the corresponding size of the dermal punch. Calculations are made using the following formula:

25
$$[\text{Open area on day 8}] - [\text{Open area on day 1}] / [\text{Open area on day 1}]$$

Specimens are fixed in 10% buffered formalin and paraffin embedded blocks are sectioned
perpendicular to the wound surface (5mm) and cut using an Olympus microtome. Routine
hematoxylin-eosin (H&E) staining is performed on cross-sections of bisected wounds.
30 Histologic examination of the wounds allows assessment of whether the healing process and the morphologic appearance of the repaired skin is improved by treatment with an agonist or antagonist of the invention. A calibrated lens micrometer is used by a blinded observer to

determine the distance of the wound gap.

Experimental data are analyzed using an unpaired t test. A p value of < 0.05 is considered significant.

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 29: Lymphadema Animal Model

The purpose of this experimental approach is to create an appropriate and consistent lymphedema model for testing the therapeutic effects of an agonist or antagonist of the invention in lymphangiogenesis and re-establishment of the lymphatic circulatory system in the rat hind limb. Effectiveness is measured by swelling volume of the affected limb, quantification of the amount of lymphatic vasculature, total blood plasma protein, and histopathology. Acute lymphedema is observed for 7-10 days. Perhaps more importantly, the chronic progress of the edema is followed for up to 3-4 weeks.

Prior to beginning surgery, blood sample is drawn for protein concentration analysis. Male rats weighing approximately ~350g are dosed with Pentobarbital. Subsequently, the right legs are shaved from knee to hip. The shaved area is swabbed with gauze soaked in 70% EtOH. Blood is drawn for serum total protein testing. Circumference and volumetric measurements are made prior to injecting dye into paws after marking 2 measurement levels (0.5 cm above heel, at mid-pt of dorsal paw). The intradermal dorsum of both right and left paws are injected with 0.05 ml of 1% Evan's Blue. Circumference and volumetric measurements are then made following injection of dye into paws.

Using the knee joint as a landmark, a mid-leg inguinal incision is made circumferentially allowing the femoral vessels to be located. Forceps and hemostats are used to dissect and separate the skin flaps. After locating the femoral vessels, the lymphatic vessel that runs along side and underneath the vessel(s) is located. The main lymphatic vessels in this area are then electrically coagulated or suture ligated.

Using a microscope, muscles in back of the leg (near the semitendinosus and adductors) are bluntly dissected. The popliteal lymph node is then located. The 2 proximal and 2 distal lymphatic vessels and distal blood supply of the popliteal node are then and

ligated by suturing. The popliteal lymph node, and any accompanying adipose tissue, is then removed by cutting connective tissues.

Care is taken to control any mild bleeding resulting from this procedure. After lymphatics are occluded, the skin flaps are sealed by using liquid skin (Vetbond) (AJ Buck).

- 5 The separated skin edges are sealed to the underlying muscle tissue while leaving a gap of ~0.5 cm around the leg. Skin also may be anchored by suturing to underlying muscle when necessary.

- To avoid infection, animals are housed individually with mesh (no bedding). Recovering animals are checked daily through the optimal edematous peak, which typically occurred by day 5-7. The plateau edematous peak are then observed. To evaluate the intensity of the lymphedema, the circumference and volumes of 2 designated places on each paw before operation and daily for 7 days are measured. The effect plasma proteins on lymphedema is determined and whether protein analysis is a useful testing perimeter is also investigated. The weights of both control and edematous limbs are evaluated at 2 places.
- 15 Analysis is performed in a blind manner.

Circumference Measurements: Under brief gas anesthetic to prevent limb movement, a cloth tape is used to measure limb circumference. Measurements are done at the ankle bone and dorsal paw by 2 different people then those 2 readings are averaged. Readings are taken from both control and edematous limbs.

- 20 Volumetric Measurements: On the day of surgery, animals are anesthetized with Pentobarbital and are tested prior to surgery. For daily volumetrics animals are under brief halothane anesthetic (rapid immobilization and quick recovery), both legs are shaved and equally marked using waterproof marker on legs. Legs are first dipped in water, then dipped into instrument to each marked level then measured by Buxco edema software(Chen/Victor).
- 25 Data is recorded by one person, while the other is dipping the limb to marked area.

Blood-plasma protein measurements: Blood is drawn, spun, and serum separated prior to surgery and then at conclusion for total protein and Ca²⁺ comparison.

- Limb Weight Comparison: After drawing blood, the animal is prepared for tissue collection. The limbs are amputated using a quillitine, then both experimental and control legs are cut at the ligature and weighed. A second weighing is done as the tibio-cacaneal joint is disarticulated and the foot is weighed.
- 30

Histological Preparations: The transverse muscle located behind the knee (popliteal)

area is dissected and arranged in a metal mold. filled with freezeGel, dipped into cold methylbutane, placed into labeled sample bags at - 80EC until sectioning. Upon sectioning, the muscle is observed under fluorescent microscopy for lymphatics..

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test
5 the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 30: Suppression of TNF alpha-induced adhesion molecule expression by a Agonist or Antagonist of the Invention

10

The recruitment of lymphocytes to areas of inflammation and angiogenesis involves specific receptor-ligand interactions between cell surface adhesion molecules (CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion
15 molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of
20 cytokines and growth factor participate in the modulation of the expression of these CAMs.

Tumor necrosis factor alpha (TNF-a), a potent proinflammatory cytokine, is a stimulator of all three CAMs on endothelial cells and may be involved in a wide variety of inflammatory responses, often resulting in a pathological outcome.

The potential of an agonist or antagonist of the invention to mediate a suppression of TNF-a
25 induced CAM expression can be examined. A modified ELISA assay which uses ECs as a solid phase absorbent is employed to measure the amount of CAM expression on TNF-a treated ECs when co-stimulated with a member of the FGF family of proteins.

To perform the experiment, human umbilical vein endothelial cell (HUVEC) cultures are obtained from pooled cord harvests and maintained in growth medium (EGM-2; Clonetics. San Diego, CA) supplemented with 10% FCS and 1% penicillin/streptomycin in a
30 37 degree C humidified incubator containing 5% CO₂. HUVECs are seeded in 96-well plates at concentrations of 1×10^4 cells/well in EGM medium at 37 degree C for 18-24 hrs or

until confluent. The monolayers are subsequently washed 3 times with a serum-free solution of RPMI-1640 supplemented with 100 U/ml penicillin and 100 mg/ml streptomycin, and treated with a given cytokine and/or growth factor(s) for 24 h at 37 degree C. Following incubation, the cells are then evaluated for CAM expression.

- 5 Human Umbilical Vein Endothelial cells (HUVECs) are grown in a standard 96 well plate to confluence. Growth medium is removed from the cells and replaced with 90 ul of 199 Medium (10% FBS). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 ul volumes). Plates are incubated at 37 degree C for either 5 h (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to
10 remove medium and 100 µl of 0.1% paraformaldehyde-PBS(with Ca++ and Mg++) is added to each well. Plates are held at 4°C for 30 min.

- Fixative is then removed from the wells and wells are washed 1X with PBS(+Ca,Mg)+0.5% BSA and drained. Do not allow the wells to dry. Add 10 µl of diluted primary antibody to the test and control wells. Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin
15 and Anti-E-selectin-Biotin are used at a concentration of 10 µg/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed X3 with PBS(+Ca,Mg)+0.5% BSA.

- Then add 20 µl of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution) to each well and incubated at 37°C for 30 min. Wells are washed X3 with PBS(+Ca,Mg)+0.5%
20 BSA. 1 tablet of p-Nitrophenol Phosphate pNPP is dissolved in 5 ml of glycine buffer (pH 10.4). 100 µl of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: 1:5,000 (10^0) > $10^{0.5}$ > 10^{-1} > $10^{-1.5}$. 5 µl of each dilution is added to triplicate wells and the resulting AP content in each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 µl
25 of pNPP reagent must then be added to each of the standard wells. The plate must be incubated at 37°C for 4h. A volume of 50 µl of 3M NaOH is added to all wells. The results are quantified on a plate reader at 405 nm. The background subtraction option is used on blank wells filled with glycine buffer only. The template is set up to indicate the concentration of AP-conjugate in each standard well [5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng].
30 Results are indicated as amount of bound AP-conjugate in each sample.

The studies described in this example tested activity of agonists or antagonists of the invention. However, one skilled in the art could easily modify the exemplified studies to test

the activity of polynucleotides or polypeptides of the invention (e.g., gene therapy).

Example 31: Production Of Polypeptide of the Invention For High-Throughput Screening Assays

5

The following protocol produces a supernatant containing polypeptide of the present invention to be tested. This supernatant can then be used in the Screening Assays described in Examples 33-42.

10 First, dilute Poly-D-Lysine (644 587 Boehringer-Mannheim) stock solution (1mg/ml in PBS) 1:20 in PBS (w/o calcium or magnesium 17-516F Biowhittaker) for a working solution of 50ug/ml. Add 200 ul of this solution to each well (24 well plates) and incubate at RT for 20 minutes. Be sure to distribute the solution over each well (note: a 12-channel pipetter may be used with tips on every other channel). Aspirate off the Poly-D-Lysine solution and rinse with 1ml PBS (Phosphate Buffered Saline). The PBS should remain in the well until just prior to plating the cells and plates may be poly-lysine coated in advance for up to two weeks.

20 Plate 293T cells (do not carry cells past P+20) at 2×10^5 cells/well in .5ml DMEM(Dulbecco's Modified Eagle Medium)(with 4.5 G/L glucose and L-glutamine (12-604F Biowhittaker))/10% heat inactivated FBS(14-503F Biowhittaker)/1x Penstrep(17-602E Biowhittaker). Let the cells grow overnight.

25 The next day, mix together in a sterile solution basin: 300 ul Lipofectamine (18324-012 Gibco/BRL) and 5ml Optimem I (31985070 Gibco/BRL)/96-well plate. With a small volume multi-channel pipetter, aliquot approximately 2ug of an expression vector containing a polynucleotide insert, produced by the methods described in Examples 8-10, into an appropriately labeled 96-well round bottom plate. With a multi-channel pipetter, add 50ul of the Lipofectamine/Optimem I mixture to each well. Pipette up and down gently to mix. Incubate at RT 15-45 minutes. After about 20 minutes, use a multi-channel pipetter to add 150ul Optimem I to each well. As a control, one plate of vector DNA lacking an insert should be transfected with each set of transfections.

30 Preferably, the transfection should be performed by tag-teaming the following tasks. By tag-teaming, hands on time is cut in half, and the cells do not spend too much time on PBS. First, person A aspirates off the media from four 24-well plates of cells, and then

person B rinses each well with .5-1ml PBS. Person A then aspirates off PBS rinse, and person B, using a12-channel pipetter with tips on every other channel, adds the 200ul of DNA/Lipofectamine/Optimem I complex to the odd wells first, then to the even wells, to each row on the 24-well plates. Incubate at 37 degree C for 6 hours.

- 5 While cells are incubating, prepare appropriate media, either 1%BSA in DMEM with 1x penstrep, or HGS CHO-5 media (116.6 mg/L of CaCl₂ (anhyd); 0.00130 mg/L CuSO₄-5H₂O; 0.050 mg/L of Fe(NO₃)₃-9H₂O; 0.417 mg/L of FeSO₄-7H₂O; 311.80 mg/L of Kcl; 28.64 mg/L of MgCl₂; 48.84 mg/L of MgSO₄; 6995.50 mg/L of NaCl; 2400.0 mg/L of NaHCO₃; 62.50 mg/L of NaH₂PO₄-H₂O; 71.02 mg/L of Na₂HPO₄; .4320 mg/L of ZnSO₄-10 7H₂O; .002 mg/L of Arachidonic Acid ; 1.022 mg/L of Cholesterol; .070 mg/L of DL-alpha-Tocopherol-Acetate; 0.0520 mg/L of Linoleic Acid; 0.010 mg/L of Linolenic Acid; 0.010 mg/L of Myristic Acid; 0.010 mg/L of Oleic Acid; 0.010 mg/L of Palmitric Acid; 0.010 mg/L of Palmitic Acid; 100 mg/L of Pluronic F-68; 0.010 mg/L of Stearic Acid; 2.20 mg/L of Tween 80; 4551 mg/L of D-Glucose; 130.85 mg/ml of L- Alanine; 147.50 mg/ml of L-Arginine-HCL; 7.50 mg/ml of L-Asparagine-H₂O; 6.65 mg/ml of L-Aspartic Acid; 29.56 15 mg/ml of L-Cystine-2HCL-H₂O; 31.29 mg/ml of L-Cystine-2HCL; 7.35 mg/ml of L-Glutamic Acid; 365.0 mg/ml of L-Glutamine; 18.75 mg/ml of Glycine; 52.48 mg/ml of L-Histidine-HCL-H₂O; 106.97 mg/ml of L-Isoleucine; 111.45 mg/ml of L-Leucine; 163.75 mg/ml of L-Lysine HCL; 32.34 mg/ml of L-Methionine; 68.48 mg/ml of L-Phenylalanine; 20 40.0 mg/ml of L-Proline; 26.25 mg/ml of L-Serine; 101.05 mg/ml of L-Threonine; 19.22 mg/ml of L-Tryptophan; 91.79 mg/ml of L-Tyrosine-2Na-2H₂O; and 99.65 mg/ml of L-Valine; 0.0035 mg/L of Biotin; 3.24 mg/L of D-Ca Pantothenate; 11.78 mg/L of Choline Chloride; 4.65 mg/L of Folic Acid; 15.60 mg/L of i-Inositol; 3.02 mg/L of Niacinamide; 3.00 mg/L of Pyridoxal HCL; 0.031 mg/L of Pyridoxine HCL; 0.319 mg/L of Riboflavin; 3.17 25 mg/L of Thiamine HCL; 0.365 mg/L of Thymidine; 0.680 mg/L of Vitamin B₁₂; 25 mM of HEPES Buffer; 2.39 mg/L of Na Hypoxanthine; 0.105 mg/L of Lipoic Acid; 0.081 mg/L of Sodium Putrescine-2HCL; 55.0 mg/L of Sodium Pyruvate; 0.0067 mg/L of Sodium Selenite; 20uM of Ethanolamine; 0.122 mg/L of Ferric Citrate; 41.70 mg/L of Methyl-B-Cyclodextrin complexed with Linoleic Acid; 33.33 mg/L of Methyl-B-Cyclodextrin complexed with Oleic 30 Acid; 10 mg/L of Methyl-B-Cyclodextrin complexed with Retinal Acetate. Adjust

osmolarity to 327 mOsm) with 2mm glutamine and 1x penstrep. (BSA (81-068-3 Bayer) 100gm dissolved in 1L DMEM for a 10% BSA stock solution). Filter the media and collect 50 ul for endotoxin assay in 15ml polystyrene conical.

5 The transfection reaction is terminated, preferably by tag-teaming, at the end of the incubation period. Person A aspirates off the transfection media, while person B adds 1.5ml appropriate media to each well. Incubate at 37 degree C for 45 or 72 hours depending on the media used: 1%BSA for 45 hours or CHO-5 for 72 hours.

10 On day four, using a 300ul multichannel pipetter, aliquot 600ul in one 1ml deep well plate and the remaining supernatant into a 2ml deep well. The supernatants from each well can then be used in the assays described in Examples 33-40.

It is specifically understood that when activity is obtained in any of the assays described below using a supernatant, the activity originates from either the polypeptide of the present invention directly (e.g., as a secreted protein) or by polypeptide of the present invention inducing expression of other proteins, which are then secreted into the supernatant.

15 Thus, the invention further provides a method of identifying the protein in the supernatant characterized by an activity in a particular assay.

Example 32: Construction of GAS Reporter Construct

20 One signal transduction pathway involved in the differentiation and proliferation of cells is called the Jaks-STATs pathway. Activated proteins in the Jaks-STATs pathway bind to gamma activation site "GAS" elements or interferon-sensitive responsive element ("ISRE"), located in the promoter of many genes. The binding of a protein to these elements alter the expression of the associated gene.

25 GAS and ISRE elements are recognized by a class of transcription factors called Signal Transducers and Activators of Transcription, or "STATs." There are six members of the STATs family. Stat1 and Stat3 are present in many cell types, as is Stat2 (as response to IFN-alpha is widespread). Stat4 is more restricted and is not in many cell types though it has been found in T helper class 1, cells after treatment with IL-12. Stat5 was originally called

30 mammary growth factor, but has been found at higher concentrations in other cells including myeloid cells. It can be activated in tissue culture cells by many cytokines.

The STATs are activated to translocate from the cytoplasm to the nucleus upon

tyrosine phosphorylation by a set of kinases known as the Janus Kinase ("Jaks") family. Jaks represent a distinct family of soluble tyrosine kinases and include Tyk2, Jak1, Jak2, and Jak3. These kinases display significant sequence similarity and are generally catalytically inactive in resting cells.

5 The Jaks are activated by a wide range of receptors summarized in the Table below. (Adapted from review by Schidler and Darnell, Ann. Rev. Biochem. 64:621-51 (1995).) A cytokine receptor family, capable of activating Jaks, is divided into two groups: (a) Class 1 includes receptors for IL-2, IL-3, IL-4, IL-6, IL-7, IL-9, IL-11, IL-12, IL-15, Epo, PRL, GH, G-CSF, GM-CSF, LIF, CNTF, and thrombopoietin; and (b) Class 2 includes IFN-a, IFN-g,
10 and IL-10. The Class 1 receptors share a conserved cysteine motif (a set of four conserved cysteines and one tryptophan) and a WSXWS motif (a membrane proximal region encoding Trp-Ser-Xxx-Trp-Ser (SEQ ID NO:1548)).

 Thus, on binding of a ligand to a receptor, Jaks are activated, which in turn activate STATs, which then translocate and bind to GAS elements. This entire process is
15 encompassed in the Jaks-STATs signal transduction pathway.

 Therefore, activation of the Jaks-STATs pathway, reflected by the binding of the GAS or the ISRE element, can be used to indicate proteins involved in the proliferation and differentiation of cells. For example, growth factors and cytokines are known to activate the Jaks-STATs pathway. (See Table below.) Thus, by using GAS elements linked to reporter
20 molecules, activators of the Jaks-STATs pathway can be identified.

	<u>Ligand</u>	<u>JAKs</u>				<u>STATS GAS(elements) or ISRE</u>	
		<u>tyk2</u>	<u>Jak1</u>	<u>Jak2</u>	<u>Jak3</u>		
	<u>IFN family</u>						
5	IFN-a/B	+	+	-	-	1,2,3	ISRE
	IFN-g		+	+	-	1	GAS
	(IRF1>Lys6>IFP)						
	IL-10	+	?	?	-	1,3	
10	<u>gp130 family</u>						
	IL-6 (Pleiotrohic)	+	+	+	?	1,3	GAS
	(IRF1>Lys6>IFP)						
	IL-11(Pleiotrohic)	?	+	?	?	1,3	
	OnM(Pleiotrohic)	?	+	+	?	1,3	
15	LIF(Pleiotrohic)	?	+	+	?	1,3	
	CNTF(Pleiotrohic)	-/+	+	+	?	1,3	
	G-CSF(Pleiotrohic)	?	+	?	?	1,3	
	IL-12(Pleiotrohic)	+	-	+	+	1,3	
20	<u>g-C family</u>						
	IL-2 (lymphocytes)	-	+	-	+	1,3,5	GAS
	IL-4 (lymph/myeloid)	-	+	-	+	6	GAS (IRF1 = IFP
	>>Ly6)(IgH)						
	IL-7 (lymphocytes)	-	+	-	+	5	GAS
25	IL-9 (lymphocytes)	-	+	-	+	5	GAS
	IL-13 (lymphocyte)	-	+	?	?	6	GAS
	IL-15	?	+	?	+	5	GAS
	<u>gp140 family</u>						
30	IL-3 (myeloid)	-	-	+	-	5	GAS
	(IRF1>IFP>>Ly6)						
	IL-5 (myeloid)	-	-	+	-	5	GAS
	GM-CSF (myeloid)	-	-	+	-	5	GAS

Growth hormone family

	GH	?	-	+	-	5	
	PRL	?	+/-	+	-	1,3,5	
5	EPO	?	-	+	-	5	GAS(B-
	CAS>IRF1=IFP>>Ly6)						

Receptor Tyrosine Kinases

	EGF	?	+	+	-	1,3	GAS (IRF1)
10	PDGF	?	+	+	-	1,3	
	CSF-1	?	+	+	-	1,3	GAS (not IRF1)

To construct a synthetic GAS containing promoter element, which is used in the Biological Assays described in Examples 33-34, a PCR based strategy is employed to generate a GAS-SV40 promoter sequence. The 5' primer contains four tandem copies of the GAS binding site found in the IRF1 promoter and previously demonstrated to bind STATs upon induction with a range of cytokines (Rothman et al., Immunity 1:457-468 (1994).), although other GAS or ISRE elements can be used instead. The 5' primer also contains 18bp of sequence complementary to the SV40 early promoter sequence and is flanked with an XhoI site. The sequence of the 5' primer is:

5' GCGCCTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAG:3' (SEQ ID NO:1549)

The downstream primer is complementary to the SV40 promoter and is flanked with a Hind III site: 5' GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO:1550)

PCR amplification is performed using the SV40 promoter template present in the B-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is digested with XhoI/Hind III and subcloned into BLSK2-. (Stratagene.) Sequencing with forward and reverse primers confirms that the insert contains the following sequence:

5' CTCTCGAGATTTCCTCCGAAATCTAGATTTCCTCCGAAATGATTTCCTCCGAAATGATTTCCTCCGAAATATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCATGGCTGACTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGCTTTTGGAGGCCTAGGCTTTGGCAAAGCTT:3' (SEQ ID NO:1551)

With this GAS promoter element linked to the SV40 promoter, a GAS:SEAP2 reporter construct is next engineered. Here, the reporter molecule is a secreted alkaline phosphatase, or "SEAP." Clearly, however, any reporter molecule can be instead of SEAP, in this or in any of the other Examples. Well known reporter molecules that can be used instead of SEAP include chloramphenicol

acetyltransferase (CAT), luciferase, alkaline phosphatase, B-galactosidase, green fluorescent protein (GFP), or any protein detectable by an antibody.

The above sequence confirmed synthetic GAS-SV40 promoter element is subcloned into the pSEAP-Promoter vector obtained from Clontech using HindIII and XhoI, effectively replacing the SV40 promoter with the amplified GAS:SV40 promoter element, to create the GAS-SEAP vector. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

Thus, in order to generate mammalian stable cell lines expressing the GAS-SEAP reporter, the GAS-SEAP cassette is removed from the GAS-SEAP vector using SalI and NotI, and inserted into a backbone vector containing the neomycin resistance gene, such as pGFP-1 (Clontech), using these restriction sites in the multiple cloning site, to create the GAS-SEAP/Neo vector. Once this vector is transfected into mammalian cells, this vector can then be used as a reporter molecule for GAS binding as described in Examples 33-34.

Other constructs can be made using the above description and replacing GAS with a different promoter sequence. For example, construction of reporter molecules containing NFK-B and EGR promoter sequences are described in Examples 35 and 36. However, many other promoters can be substituted using the protocols described in these Examples. For instance, SRE, IL-2, NFAT, or Osteocalcin promoters can be substituted, alone or in combination (e.g., GAS/NF-KB/EGR, GAS/NF-KB, IL-2/NFAT, or NF-KB/GAS). Similarly, other cell lines can be used to test reporter construct activity, such as HELA (epithelial), HUVEC (endothelial), Reh (B-cell), Saos-2 (osteoblast), HUVAC (aortic), or Cardiomyocyte.

Example 33: High-Throughput Screening Assay for T-cell Activity.

The following protocol is used to assess T-cell activity by identifying factors, and determining whether supernate containing a polypeptide of the invention proliferates and/or differentiates T-cells. T-cell activity is assessed using the

GAS/SEAP/Neo construct produced in Example 32. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The T-cell used in this assay is Jurkat T-cells (ATCC Accession No. TIB-152), although Molt-3 cells (ATCC Accession No. CRL-1552) and Molt-4 cells (ATCC
5 Accession No. CRL-1582) cells can also be used.

Jurkat T-cells are lymphoblastic CD4⁺ Th1 helper cells. In order to generate stable cell lines, approximately 2 million Jurkat cells are transfected with the GAS-SEAP/neo vector using DMRIE-C (Life Technologies)(transfection procedure described below). The transfected cells are seeded to a density of approximately
10 20,000 cells per well and transfectants resistant to 1 mg/ml gentamicin selected. Resistant colonies are expanded and then tested for their response to increasing concentrations of interferon gamma. The dose response of a selected clone is demonstrated.

Specifically, the following protocol will yield sufficient cells for 75 wells
15 containing 200 ul of cells. Thus, it is either scaled up, or performed in multiple to generate sufficient cells for multiple 96 well plates. Jurkat cells are maintained in RPMI + 10% serum with 1%Pen-Strep. Combine 2.5 mls of OPTI-MEM (Life Technologies) with 10 ug of plasmid DNA in a T25 flask. Add 2.5 ml OPTI-MEM containing 50 ul of DMRIE-C and incubate at room temperature for 15-45 mins.

20 During the incubation period, count cell concentration, spin down the required number of cells (10^7 per transfection), and resuspend in OPTI-MEM to a final concentration of 10^7 cells/ml. Then add 1ml of 1×10^7 cells in OPTI-MEM to T25 flask and incubate at 37 degree C for 6 hrs. After the incubation, add 10 ml of RPMI + 15% serum.

25 The Jurkat:GAS-SEAP stable reporter lines are maintained in RPMI + 10% serum, 1 mg/ml Gentamicin, and 1% Pen-Strep. These cells are treated with supernatants containing polypeptide of the present invention or polypeptide of the present invention induced polypeptides as produced by the protocol described in Example 31.

30 On the day of treatment with the supernatant, the cells should be washed and

resuspended in fresh RPMI + 10% serum to a density of 500,000 cells per ml. The exact number of cells required will depend on the number of supernatants being screened. For one 96 well plate, approximately 10 million cells (for 10 plates, 100 million cells) are required.

5 Transfer the cells to a triangular reservoir boat, in order to dispense the cells into a 96 well dish, using a 12 channel pipette. Using a 12 channel pipette, transfer 200 ul of cells into each well (therefore adding 100,000 cells per well).

 After all the plates have been seeded, 50 ul of the supernatants are transferred directly from the 96 well plate containing the supernatants into each well using a 12
10 channel pipette. In addition, a dose of exogenous interferon gamma (0.1, 1.0, 10 ng) is added to wells H9, H10, and H11 to serve as additional positive controls for the assay.

 The 96 well dishes containing Jurkat cells treated with supernatants are placed in an incubator for 48 hrs (note: this time is variable between 48-72 hrs). 35 ul
15 samples from each well are then transferred to an opaque 96 well plate using a 12 channel pipette. The opaque plates should be covered (using sellophene covers) and stored at -20 degree C until SEAP assays are performed according to Example 37. The plates containing the remaining treated cells are placed at 4 degree C and serve as a source of material for repeating the assay on a specific well if desired.

20 As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate Jurkat T cells. Over 30 fold induction is typically observed in the positive control wells.

 The above protocol may be used in the generation of both transient, as well as, stable transfected cells, which would be apparent to those of skill in the art.

25

Example 34: High-Throughput Screening Assay Identifying Myeloid Activity

 The following protocol is used to assess myeloid activity of polypeptide of the present invention by determining whether polypeptide of the present invention
30 proliferates and/or differentiates myeloid cells. Myeloid cell activity is assessed using

the GAS/SEAP/Neo construct produced in Example 32. Thus, factors that increase SEAP activity indicate the ability to activate the Jaks-STATS signal transduction pathway. The myeloid cell used in this assay is U937, a pre-monocyte cell line, although TF-1, HL60, or KG1 can be used.

- 5 To transiently transfect U937 cells with the GAS/SEAP/Neo construct produced in Example 32, a DEAE-Dextran method (Kharbanda et. al., 1994, Cell Growth & Differentiation, 5:259-265) is used. First, harvest 2×10^7 U937 cells and wash with PBS. The U937 cells are usually grown in RPMI 1640 medium containing 10% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml
10 penicillin and 100 mg/ml streptomycin.

Next, suspend the cells in 1 ml of 20 mM Tris-HCl (pH 7.4) buffer containing 0.5 mg/ml DEAE-Dextran, 8 ug GAS-SEAP2 plasmid DNA, 140 mM NaCl, 5 mM KCl, 375 uM $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$, 1 mM MgCl_2 , and 675 uM CaCl_2 . Incubate at 37 degrees C for 45 min.

- 15 Wash the cells with RPMI 1640 medium containing 10% FBS and then resuspend in 10 ml complete medium and incubate at 37 degree C for 36 hr.

The GAS-SEAP/U937 stable cells are obtained by growing the cells in 400 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 400 ug/ml G418 for couple of passages.

- 20 These cells are tested by harvesting 1×10^8 cells (this is enough for ten 96-well plates assay) and wash with PBS. Suspend the cells in 200 ml above described growth medium, with a final density of 5×10^5 cells/ml. Plate 200 ul cells per well in the 96-well plate (or 1×10^5 cells/well).

- Add 50 ul of the supernatant prepared by the protocol described in Example
25 31. Incubate at 37 degree C for 48 to 72 hr. As a positive control, 100 Unit/ml interferon gamma can be used which is known to activate U937 cells. Over 30 fold induction is typically observed in the positive control wells. SEAP assay the supernatant according to the protocol described in Example 37.

- 30 *Example 35: High-Throughput Screening Assay Identifying Neuronal Activity.*

When cells undergo differentiation and proliferation, a group of genes are activated through many different signal transduction pathways. One of these genes, EGR1 (early growth response gene 1), is induced in various tissues and cell types upon activation. The promoter of EGR1 is responsible for such induction. Using the EGR1 promoter linked to reporter molecules, activation of cells can be assessed by polypeptide of the present invention.

Particularly, the following protocol is used to assess neuronal activity in PC12 cell lines. PC12 cells (rat phenochromocytoma cells) are known to proliferate and/or differentiate by activation with a number of mitogens, such as TPA (tetradecanoyl phorbol acetate), NGF (nerve growth factor), and EGF (epidermal growth factor). The EGR1 gene expression is activated during this treatment. Thus, by stably transfecting PC12 cells with a construct containing an EGR promoter linked to SEAP reporter, activation of PC12 cells by polypeptide of the present invention can be assessed.

The EGR/SEAP reporter construct can be assembled by the following protocol. The EGR-1 promoter sequence (-633 to +1)(Sakamoto K et al., Oncogene 6:867-871 (1991)) can be PCR amplified from human genomic DNA using the following primers:

20 5' GCGCTCGAGGGATGACAGCGATAGAACCCCGG -3' (SEQ ID NO: 1552)

5' GCGAAGCTTCGCGACTCCCCGGATCCGCCTC-3' (SEQ ID NO: 1553)

Using the GAS:SEAP/Neo vector produced in Example 32, EGR1 amplified product can then be inserted into this vector. Linearize the GAS:SEAP/Neo vector using restriction enzymes XhoI/HindIII, removing the GAS/SV40 stuffer. Restrict the EGR1 amplified product with these same enzymes. Ligate the vector and the EGR1 promoter.

To prepare 96 well-plates for cell culture, two mls of a coating solution (1:30 dilution of collagen type I (Upstate Biotech Inc. Cat#08-115) in 30% ethanol (filter

sterilized)) is added per one 10 cm plate or 50 ml per well of the 96-well plate, and allowed to air dry for 2 hr.

PC12 cells are routinely grown in RPMI-1640 medium (Bio Whittaker) containing 10% horse serum (JRH BIOSCIENCES, Cat. # 12449-78P), 5% heat-inactivated fetal bovine serum (FBS) supplemented with 100 units/ml penicillin and 100 ug/ml streptomycin on a precoated 10 cm tissue culture dish. One to four split is done every three to four days. Cells are removed from the plates by scraping and resuspended with pipetting up and down for more than 15 times.

Transfect the EGR/SEAP/Neo construct into PC12 using the Lipofectamine protocol described in Example 31. EGR-SEAP/PC12 stable cells are obtained by growing the cells in 300 ug/ml G418. The G418-free medium is used for routine growth but every one to two months, the cells should be re-grown in 300 ug/ml G418 for couple of passages.

To assay for neuronal activity, a 10 cm plate with cells around 70 to 80% confluent is screened by removing the old medium. Wash the cells once with PBS (Phosphate buffered saline). Then starve the cells in low serum medium (RPMI-1640 containing 1% horse serum and 0.5% FBS with antibiotics) overnight.

The next morning, remove the medium and wash the cells with PBS. Scrape off the cells from the plate, suspend the cells well in 2 ml low serum medium. Count the cell number and add more low serum medium to reach final cell density as 5×10^5 cells/ml.

Add 200 ul of the cell suspension to each well of 96-well plate (equivalent to 1×10^5 cells/well). Add 50 ul supernatant produced by Example 31, 37 degree C for 48 to 72 hr. As a positive control, a growth factor known to activate PC12 cells through EGR can be used, such as 50 ng/ul of Neuronal Growth Factor (NGF). Over fifty-fold induction of SEAP is typically seen in the positive control wells. SEAP assay the supernatant according to Example 37.

Example 36: High-Throughput Screening Assay for T-cell Activity

NF-KB (Nuclear Factor KB) is a transcription factor activated by a wide variety of agents including the inflammatory cytokines IL-1 and TNF. CD30 and CD40, lymphotoxin-alpha and lymphotoxin-beta, by exposure to LPS or thrombin, and by expression of certain viral gene products. As a transcription factor, NF-KB regulates the expression of genes involved in immune cell activation, control of apoptosis (NF- KB appears to shield cells from apoptosis), B and T-cell development, anti-viral and antimicrobial responses, and multiple stress responses.

In non-stimulated conditions, NF- KB is retained in the cytoplasm with I-KB (Inhibitor KB). However, upon stimulation, I- KB is phosphorylated and degraded, causing NF- KB to shuttle to the nucleus, thereby activating transcription of target genes. Target genes activated by NF- KB include IL-2, IL-6, GM-CSF, ICAM-1 and class I MHC.

Due to its central role and ability to respond to a range of stimuli, reporter constructs utilizing the NF-KB promoter element are used to screen the supernatants produced in Example 31. Activators or inhibitors of NF-KB would be useful in treating, preventing, and/or diagnosing diseases. For example, inhibitors of NF-KB could be used to treat those diseases related to the acute or chronic activation of NF-KB, such as rheumatoid arthritis.

To construct a vector containing the NF-KB promoter element, a PCR based strategy is employed. The upstream primer contains four tandem copies of the NF-KB binding site (GGGGACTTTCCC) (SEQ ID NO:1554), 18 bp of sequence complementary to the 5' end of the SV40 early promoter sequence, and is flanked with an XhoI site:

5':GCGGCCTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGAC
TTTCCATCCTGCCATCTCAATTAG:3' (SEQ ID NO:1555)

The downstream primer is complementary to the 3' end of the SV40 promoter and is flanked with a Hind III site:

5':GCGGCAAGCTTTTGGCAAAGCCTAGGC:3' (SEQ ID NO:1550)

PCR amplification is performed using the SV40 promoter template present in the pB-gal:promoter plasmid obtained from Clontech. The resulting PCR fragment is

digested with XhoI and Hind III and subcloned into BLSK2-. (Stratagene) Sequencing with the T7 and T3 primers confirms the insert contains the following sequence:

5':CTCGAGGGGACTTTCCCGGGGACTTTCCGGGGACTTTCCGGGACTTTCC
 5 ATCTGCCATCTCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCC
 ATCCCGCCCCTAACTCCGCCCAGTTCGCCCCATTCTCCGCCCCATGGCTGA
 CTAATTTTTTTTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTA
 TTCCAGAAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAA
 GCTT:3' (SEQ ID NO:1556)

10 Next, replace the SV40 minimal promoter element present in the pSEAP2-promoter plasmid (Clontech) with this NF-KB/SV40 fragment using XhoI and HindIII. However, this vector does not contain a neomycin resistance gene, and therefore, is not preferred for mammalian expression systems.

In order to generate stable mammalian cell lines, the NF-KB/SV40/SEAP
 15 cassette is removed from the above NF-KB/SEAP vector using restriction enzymes Sall and NotI, and inserted into a vector containing neomycin resistance. Particularly, the NF-KB/SV40/SEAP cassette was inserted into pGFP-1 (Clontech), replacing the GFP gene, after restricting pGFP-1 with Sall and NotI.

Once NF-KB/SV40/SEAP/Neo vector is created, stable Jurkat T-cells are
 20 created and maintained according to the protocol described in Example 33. Similarly, the method for assaying supernatants with these stable Jurkat T-cells is also described in Example 33. As a positive control, exogenous TNF alpha (0.1, 1, 10 ng) is added to wells H9, H10, and H11, with a 5-10 fold activation typically observed.

25 *Example 37: Assay for SEAP Activity*

As a reporter molecule for the assays described in Examples 33-36, SEAP activity is assayed using the Tropix Phospho-light Kit (Cat. BP-400) according to the following general procedure. The Tropix Phospho-light Kit supplies the Dilution,
 30 Assay, and Reaction Buffers used below.

Prime a dispenser with the 2.5x Dilution Buffer and dispense 15 ul of 2.5x dilution buffer into Optiplates containing 35 ul of a supernatant. Seal the plates with a plastic sealer and incubate at 65 degree C for 30 min. Separate the Optiplates to avoid uneven heating.

- 5 Cool the samples to room temperature for 15 minutes. Empty the dispenser and prime with the Assay Buffer. Add 50 ml Assay Buffer and incubate at room temperature 5 min. Empty the dispenser and prime with the Reaction Buffer (see the table below). Add 50 ul Reaction Buffer and incubate at room temperature for 20 minutes. Since the intensity of the chemiluminescent signal is time dependent, and it
- 10 takes about 10 minutes to read 5 plates on luminometer, one should treat 5 plates at each time and start the second set 10 minutes later.

Read the relative light unit in the luminometer. Set H12 as blank, and print the results. An increase in chemiluminescence indicates reporter activity.

15 Reaction Buffer Formulation:

# of plates	Rxn buffer diluent (ml)	CSPD (ml)
10	60	3
11	65	3.25
12	70	3.5
13	75	3.75
14	80	4
15	85	4.25
16	90	4.5
17	95	4.75
18	100	5
19	105	5.25
20	110	5.5
21	115	5.75
22	120	6
23	125	6.25
24	130	6.5

384

25	135	6.75
26	140	7
27	145	7.25
28	150	7.5
29	155	7.75
30	160	8
31	165	8.25
32	170	8.5
33	175	8.75
34	180	9
35	185	9.25
36	190	9.5
37	195	9.75
38	200	10
39	205	10.25
40	210	10.5
41	215	10.75
42	220	11
43	225	11.25
44	230	11.5
45	235	11.75
46	240	12
47	245	12.25
48	250	12.5
49	255	12.75
50	260	13

Example 38: High-Throughput Screening Assay Identifying Changes in Small Molecule Concentration and Membrane Permeability

- 5 Binding of a ligand to a receptor is known to alter intracellular levels of small molecules, such as calcium, potassium, sodium, and pH, as well as alter membrane potential. These alterations can be measured in an assay to identify supernatants which bind to receptors of a particular cell. Although the following protocol

describes an assay for calcium, this protocol can easily be modified to detect changes in potassium, sodium, pH, membrane potential, or any other small molecule which is detectable by a fluorescent probe.

5 The following assay uses Fluorometric Imaging Plate Reader ("FLIPR") to measure changes in fluorescent molecules (Molecular Probes) that bind small molecules. Clearly, any fluorescent molecule detecting a small molecule can be used instead of the calcium fluorescent molecule. fluo-4 (Molecular Probes, Inc.; catalog no. F-14202), used here.

10 For adherent cells, seed the cells at 10,000 -20,000 cells/well in a Co-star black 96-well plate with clear bottom. The plate is incubated in a CO₂ incubator for 20 hours. The adherent cells are washed two times in Biotek washer with 200 ul of HBSS (Hank's Balanced Salt Solution) leaving 100 ul of buffer after the final wash.

15 A stock solution of 1 mg/ml fluo-4 is made in 10% pluronic acid DMSO. To load the cells with fluo-4, 50 ul of 12 ug/ml fluo-4 is added to each well. The plate is incubated at 37 degrees C in a CO₂ incubator for 60 min. The plate is washed four times in the Biotek washer with HBSS leaving 100 ul of buffer.

20 For non-adherent cells, the cells are spun down from culture media. Cells are re-suspended to 2.5×10^6 cells/ml with HBSS in a 50-ml conical tube. 4 ul of 1 mg/ml fluo-4 solution in 10% pluronic acid DMSO is added to each ml of cell suspension. The tube is then placed in a 37 degrees C water bath for 30-60 min. The cells are washed twice with HBSS, resuspended to 1×10^6 cells/ml, and dispensed into a microplate, 100 ul/well. The plate is centrifuged at 1000 rpm for 5 min. The plate is then washed once in Denley Cell Wash with 200 ul, followed by an aspiration step to 100 ul final volume.

25 For a non-cell based assay, each well contains a fluorescent molecule, such as fluo-4. The supernatant is added to the well, and a change in fluorescence is detected.

30 To measure the fluorescence of intracellular calcium, the FLIPR is set for the following parameters: (1) System gain is 300-800 mW; (2) Exposure time is 0.4 second; (3) Camera F/stop is F/2; (4) Excitation is 488 nm; (5) Emission is 530 nm;

and (6) Sample addition is 50 ul. Increased emission at 530 nm indicates an extracellular signaling event caused by the a molecule. either polypeptide of the present invention or a molecule induced by polypeptide of the present invention, which has resulted in an increase in the intracellular Ca^{++} concentration.

5

Example 40: High-Throughput Screening Assay Identifying Tyrosine Kinase Activity

The Protein Tyrosine Kinases (PTK) represent a diverse group of transmembrane and cytoplasmic kinases. Within the Receptor Protein Tyrosine
10 Kinase RPTK) group are receptors for a range of mitogenic and metabolic growth factors including the PDGF, FGF, EGF, NGF, HGF and Insulin receptor subfamilies. In addition there are a large family of RPTKs for which the corresponding ligand is unknown. Ligands for RPTKs include mainly secreted small proteins, but also membrane-bound and extracellular matrix proteins.

15 Activation of RPTK by ligands involves ligand-mediated receptor dimerization, resulting in transphosphorylation of the receptor subunits and activation of the cytoplasmic tyrosine kinases. The cytoplasmic tyrosine kinases include receptor associated tyrosine kinases of the src-family (e.g., src, yes, lck, lyn, fyn) and non-receptor linked and cytosolic protein tyrosine kinases, such as the Jak family,
20 members of which mediate signal transduction triggered by the cytokine superfamily of receptors (e.g., the Interleukins, Interferons, GM-CSF, and Leptin).

Because of the wide range of known factors capable of stimulating tyrosine kinase activity, identifying whether polypeptide of the present invention or a molecule induced by polypeptide of the present invention is capable of activating tyrosine
25 kinase signal transduction pathways is of interest. Therefore, the following protocol is designed to identify such molecules capable of activating the tyrosine kinase signal transduction pathways.

Seed target cells (e.g., primary keratinocytes) at a density of approximately 25,000 cells per well in a 96 well Loprodyne Silent Screen Plates purchased from
30 Nalge Nunc (Naperville, IL). The plates are sterilized with two 30 minute rinses with

100% ethanol, rinsed with water and dried overnight. Some plates are coated for 2 hr with 100 ml of cell culture grade type I collagen (50 mg/ml), gelatin (2%) or polylysine (50 mg/ml), all of which can be purchased from Sigma Chemicals (St. Louis, MO) or 10% Matrigel purchased from Becton Dickinson (Bedford, MA), or calf serum, rinsed with PBS and stored at 4 degree C. Cell growth on these plates is assayed by seeding 5,000 cells/well in growth medium and indirect quantitation of cell number through use of alamarBlue as described by the manufacturer Alamar Biosciences, Inc. (Sacramento, CA) after 48 hr. Falcon plate covers #3071 from Becton Dickinson (Bedford, MA) are used to cover the Loprodyne Silent Screen Plates. Falcon Microtest III cell culture plates can also be used in some proliferation experiments.

To prepare extracts, A431 cells are seeded onto the nylon membranes of Loprodyne plates (20,000/200ml/well) and cultured overnight in complete medium. Cells are quiesced by incubation in serum-free basal medium for 24 hr. After 5-20 minutes treatment with EGF (60ng/ml) or 50 ul of the supernatant produced in Example 31, the medium was removed and 100 ml of extraction buffer ((20 mM HEPES pH 7.5, 0.15 M NaCl, 1% Triton X-100, 0.1% SDS, 2 mM Na₃VO₄, 2 mM Na₄P₂O₇ and a cocktail of protease inhibitors (# 1836170) obtained from Boehringer Mannheim (Indianapolis, IN) is added to each well and the plate is shaken on a rotating shaker for 5 minutes at 4°C. The plate is then placed in a vacuum transfer manifold and the extract filtered through the 0.45 mm membrane bottoms of each well using house vacuum. Extracts are collected in a 96-well catch/assay plate in the bottom of the vacuum manifold and immediately placed on ice. To obtain extracts clarified by centrifugation, the content of each well, after detergent solubilization for 5 minutes, is removed and centrifuged for 15 minutes at 4 degree C at 16,000 x g.

Test the filtered extracts for levels of tyrosine kinase activity. Although many methods of detecting tyrosine kinase activity are known, one method is described here.

Generally, the tyrosine kinase activity of a supernatant is evaluated by

determining its ability to phosphorylate a tyrosine residue on a specific substrate (a biotinylated peptide). Biotinylated peptides that can be used for this purpose include PSK1 (corresponding to amino acids 6-20 of the cell division kinase cdc2-p34) and PSK2 (corresponding to amino acids 1-17 of gastrin). Both peptides are substrates for a range of tyrosine kinases and are available from Boehringer Mannheim.

The tyrosine kinase reaction is set up by adding the following components in order. First, add 10ul of 5uM Biotinylated Peptide, then 10ul ATP/Mg₂⁺ (5mM ATP/50mM MgCl₂), then 10ul of 5x Assay Buffer (40mM imidazole hydrochloride, pH7.3, 40 mM beta-glycerophosphate, 1mM EGTA, 100mM MgCl₂, 5 mM MnCl₂, 0.5 mg/ml BSA); then 5ul of Sodium Vanadate(1mM), and then 5ul of water. Mix the components gently and preincubate the reaction mix at 30 degree C for 2 min. Initial the reaction by adding 10ul of the control enzyme or the filtered supernatant.

The tyrosine kinase assay reaction is then terminated by adding 10 ul of 120mM EDTA and place the reactions on ice.

Tyrosine kinase activity is determined by transferring 50 ul aliquot of reaction mixture to a microtiter plate (MTP) module and incubating at 37 degree C for 20 min. This allows the streptavidin coated 96 well plate to associate with the biotinylated peptide. Wash the MTP module with 300ul/well of PBS four times. Next add 75 ul of anti-phosphotyrosine antibody conjugated to horse radish peroxidase(anti-P-Tyr-POD(0.5u/ml)) to each well and incubate at 37 degree C for one hour. Wash the well as above.

Next add 100ul of peroxidase substrate solution (Boehringer Mannheim) and incubate at room temperature for at least 5 mins (up to 30 min). Measure the absorbance of the sample at 405 nm by using ELISA reader. The level of bound peroxidase activity is quantitated using an ELISA reader and reflects the level of tyrosine kinase activity.

Example 41: High-Throughput Screening Assay Identifying Phosphorylation Activity

As a potential alternative and/or compliment to the assay of protein tyrosine

kinase activity described in Example 40. an assay which detects activation (phosphorylation) of major intracellular signal transduction intermediates can also be used. For example, as described below one particular assay can detect tyrosine phosphorylation of the Erk-1 and Erk-2 kinases. However, phosphorylation of other molecules, such as Raf, JNK, p38 MAP. Map kinase kinase (MEK), MEK kinase, Src, 5 Muscle specific kinase (MuSK), IRAK, Tec, and Janus, as well as any other phosphoserine, phosphotyrosine, or phosphothreonine molecule, can be detected by substituting these molecules for Erk-1 or Erk-2 in the following assay.

Specifically, assay plates are made by coating the wells of a 96-well ELISA 10 plate with 0.1ml of protein G (1ug/ml) for 2 hr at room temp, (RT). The plates are then rinsed with PBS and blocked with 3% BSA/PBS for 1 hr at RT. The protein G plates are then treated with 2 commercial monoclonal antibodies (100ng/well) against Erk-1 and Erk-2 (1 hr at RT) (Santa Cruz Biotechnology). (To detect other molecules, this step can easily be modified by substituting a monoclonal antibody 15 detecting any of the above described molecules.) After 3-5 rinses with PBS, the plates are stored at 4 degree C until use.

A431 cells are seeded at 20,000/well in a 96-well Loprodyne filterplate and cultured overnight in growth medium. The cells are then starved for 48 hr in basal medium (DMEM) and then treated with EGF (6ng/well) or 50 ul of the supernatants 20 obtained in Example 31 for 5-20 minutes. The cells are then solubilized and extracts filtered directly into the assay plate.

After incubation with the extract for 1 hr at RT, the wells are again rinsed. As a positive control, a commercial preparation of MAP kinase (10ng/well) is used in place of A431 extract. Plates are then treated with a commercial polyclonal (rabbit) 25 antibody (1ug/ml) which specifically recognizes the phosphorylated epitope of the Erk-1 and Erk-2 kinases (1 hr at RT). This antibody is biotinylated by standard procedures. The bound polyclonal antibody is then quantitated by successive incubations with Europium-streptavidin and Europium fluorescence enhancing reagent in the Wallac DELFIA instrument (time-resolved fluorescence). An increased 30 fluorescent signal over background indicates a phosphorylation by polypeptide of the

present invention or a molecule induced by polypeptide of the present invention.

Example 42: Assay for the Stimulation of Bone Marrow CD34+ Cell Proliferation

5 This assay is based on the ability of human CD34+ to proliferate in the presence of hematopoietic growth factors and evaluates the ability of isolated polypeptides expressed in mammalian cells to stimulate proliferation of CD34+ cells.

 It has been previously shown that most mature precursors will respond to only a single signal. More immature precursors require at least two signals to respond.
10 Therefore, to test the effect of polypeptides on hematopoietic activity of a wide range of progenitor cells, the assay contains a given polypeptide in the presence or absence of other hematopoietic growth factors. Isolated cells are cultured for 5 days in the presence of Stem Cell Factor (SCF) in combination with tested sample. SCF alone has a very limited effect on the proliferation of bone marrow (BM) cells, acting in
15 such conditions only as a "survival" factor. However, combined with any factor exhibiting stimulatory effect on these cells (e.g., IL-3), SCF will cause a synergistic effect. Therefore, if the tested polypeptide has a stimulatory effect on a hematopoietic progenitors, such activity can be easily detected. Since normal BM cells have a low level of cycling cells, it is likely that any inhibitory effect of a given polypeptide, or
20 agonists or antagonists thereof, might not be detected. Accordingly, assays for an inhibitory effect on progenitors is preferably tested in cells that are first subjected to *in vitro* stimulation with SCF+IL+3, and then contacted with the compound that is being evaluated for inhibition of such induced proliferation.

 Briefly, CD34+ cells are isolated using methods known in the art. The cells
25 are thawed and resuspended in medium (QBSF 60 serum-free medium with 1% L-glutamine (500ml) Quality Biological, Inc., Gaithersburg, MD Cat# 160-204-101). After several gentle centrifugation steps at 200 x g, cells are allowed to rest for one hour. The cell count is adjusted to 2.5×10^5 cells/ml. During this time, 100 μ l of sterile water is added to the peripheral wells of a 96-well plate. The cytokines that
30 can be tested with a given polypeptide in this assay is rhSCF (R&D Systems,

Minneapolis, MN, Cat# 255-SC) at 50 ng/ml alone and in combination with rhSCF and rhIL-3 (R&D Systems, Minneapolis, MN, Cat# 203-ML) at 30 ng/ml. After one hour, 10 μ l of prepared cytokines, 50 μ l of the supernatants prepared in Example 31 (supernatants at 1:2 dilution = 50 μ l) and 20 μ l of diluted cells are added to the media
5 which is already present in the wells to allow for a final total volume of 100 μ l. The plates are then placed in a 37°C/5% CO₂ incubator for five days.

Eighteen hours before the assay is harvested, 0.5 μ Ci/well of [3H] Thymidine is added in a 10 μ l volume to each well to determine the proliferation rate. The experiment is terminated by harvesting the cells from each 96-well plate to a filtermat
10 using the Tomtec Harvester 96. After harvesting, the filtermats are dried, trimmed and placed into OmniFilter assemblies consisting of one OmniFilter plate and one OmniFilter Tray. 60 μ l Microscint is added to each well and the plate sealed with TopSeal-A press-on sealing film. A bar code 15 sticker is affixed to the first plate for counting. The sealed plates is then loaded and the level of radioactivity determined
15 via the Packard Top Count and the printed data collected for analysis. The level of radioactivity reflects the amount of cell proliferation.

The studies described in this example test the activity of a given polypeptide to stimulate bone marrow CD34+ cell proliferation. One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene
20 therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof. As a nonlimiting example, potential antagonists tested in this assay would be expected to inhibit cell proliferation in the presence of cytokines and/or to increase the inhibition of cell proliferation in the presence of cytokines and a given polypeptide. In contrast, potential agonists tested in this assay would be expected to enhance cell
25 proliferation and/or to decrease the inhibition of cell proliferation in the presence of cytokines and a given polypeptide.

The ability of a gene to stimulate the proliferation of bone marrow CD34+ cells indicates that polynucleotides and polypeptides corresponding to the gene are useful for the diagnosis and treatment of disorders affecting the immune system and
30 hematopoiesis. Representative uses are described in the "Immune Activity" and

"Infectious Disease" sections above, and elsewhere herein.

Example 43: Assay for Extracellular Matrix Enhanced Cell Response (EMECCR)

5 The objective of the Extracellular Matrix Enhanced Cell Response (EMECCR) assay is to identify gene products (e.g., isolated polypeptides) that act on the hematopoietic stem cells in the context of the extracellular matrix (ECM) induced signal.

10 Cells respond to the regulatory factors in the context of signal(s) received from the surrounding microenvironment. For example, fibroblasts, and endothelial and epithelial stem cells fail to replicate in the absence of signals from the ECM. Hematopoietic stem cells can undergo self-renewal in the bone marrow, but not in *in vitro* suspension culture. The ability of stem cells to undergo self-renewal *in vitro* is dependent upon their interaction with the stromal cells and the ECM protein
15 fibronectin (fn). Adhesion of cells to fn is mediated by the $\alpha_3\beta_1$ and $\alpha_4\beta_1$ integrin receptors, which are expressed by human and mouse hematopoietic stem cells. The factor(s) which integrate with the ECM environment and responsible for stimulating stem cell self-renewal has not yet been identified. Discovery of such factors should be of great interest in gene therapy and bone marrow transplant applications

20 Briefly, polystyrene, non tissue culture treated, 96-well plates are coated with fn fragment at a coating concentration of $0.2 \mu\text{g}/\text{cm}^2$. Mouse bone marrow cells are plated (1,000 cells/well) in 0.2 ml of serum-free medium. Cells cultured in the presence of IL-3 (5 ng/ml) + SCF (50 ng/ml) would serve as the positive control, conditions under which little self-renewal but pronounced differentiation of the stem
25 cells is to be expected. Gene products of the invention (e.g., including, but not limited to, polynucleotides and polypeptides of the present invention, and supernatants produced in Example 31), are tested with appropriate negative controls in the presence and absence of SCF(5.0 ng/ml), where test factor supernates represent 10% of the total assay volume. The plated cells are then allowed to grow by incubating in
30 a low oxygen environment (5% CO_2 , 7% O_2 , and 88% N_2) tissue culture incubator

for 7 days. The number of proliferating cells within the wells is then quantitated by measuring thymidine incorporation into cellular DNA. Verification of the positive hits in the assay will require phenotypic characterization of the cells, which can be accomplished by scaling up of the culture system and using appropriate antibody
5 reagents against cell surface antigens and FACScan.

One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

If a particular polypeptide of the present invention is found to be a stimulator
10 of hematopoietic progenitors, polynucleotides and polypeptides corresponding to the gene encoding said polypeptide may be useful for the diagnosis and treatment of disorders affecting the immune system and hematopoiesis. Representative uses are described in the "Immune Activity" and "Infectious Disease" sections above, and elsewhere herein. The gene product may also be useful in the expansion of stem cells
15 and committed progenitors of various blood lineages, and in the differentiation and/or proliferation of various cell types.

Additionally, the polynucleotides and/or polypeptides of the gene of interest and/or agonists and/or antagonists thereof, may also be employed to inhibit the proliferation and differentiation of hematopoietic cells and therefore may be
20 employed to protect bone marrow stem cells from chemotherapeutic agents during chemotherapy. This antiproliferative effect may allow administration of higher doses of chemotherapeutic agents and, therefore, more effective chemotherapeutic treatment.

Moreover, polynucleotides and polypeptides corresponding to the gene of
25 interest may also be useful for the treatment and diagnosis of hematopoietic related disorders such as, for example, anemia, pancytopenia, leukopenia, thrombocytopenia or leukemia since stromal cells are important in the production of cells of hematopoietic lineages. The uses include bone marrow cell ex-vivo culture, bone marrow transplantation, bone marrow reconstitution, radiotherapy or chemotherapy of
30 neoplasia.

Example 44: Human Dermal Fibroblast and Aortic Smooth Muscle Cell Proliferation

- The polypeptide of interest is added to cultures of normal human dermal
- 5 fibroblasts (NHDF) and human aortic smooth muscle cells (AoSMC) and two co-assays are performed with each sample. The first assay examines the effect of the polypeptide of interest on the proliferation of normal human dermal fibroblasts (NHDF) or aortic smooth muscle cells (AoSMC). Aberrant growth of fibroblasts or smooth muscle cells is a part of several pathological processes, including fibrosis, and
- 10 restenosis. The second assay examines IL6 production by both NHDF and SMC. IL6 production is an indication of functional activation. Activated cells will have increased production of a number of cytokines and other factors, which can result in a proinflammatory or immunomodulatory outcome. Assays are run with and without co-TNF α stimulation, in order to check for costimulatory or inhibitory activity.
- 15 Briefly, on day 1, 96-well black plates are set up with 1000 cells/well (NHDF) or 2000 cells/well (AoSMC) in 100 μ l culture media. NHDF culture media contains: Clonetics FB basal media, 1mg/ml hFGF, 5mg/ml insulin, 50mg/ml gentamycin, 2%FBS, while AoSMC culture media contains Clonetics SM basal media, 0.5 μ g/ml hEGF, 5mg/ml insulin, 1 μ g/ml hFGF, 50mg/ml gentamycin, 50 μ g/ml Amphotericin
- 20 B, 5%FBS. After incubation at 37°C for at least 4-5 hours, culture media is aspirated and replaced with growth arrest media. Growth arrest media for NHDF contains fibroblast basal media, 50mg/ml gentamycin, 2% FBS, while growth arrest media for AoSMC contains SM basal media, 50mg/ml gentamycin, 50 μ g/ml Amphotericin B, 0.4% FBS. Incubate at 37°C until day 2.
- 25 On day 2, serial dilutions and templates of the polypeptide of interest are designed such that they always include media controls and known-protein controls. For both stimulation and inhibition experiments, proteins are diluted in growth arrest media. For inhibition experiments, TNF α is added to a final concentration of 2ng/ml (NHDF) or 5ng/ml (AoSMC). Add 1/3 vol media containing controls or polypeptides
- 30 of the present invention and incubate at 37°C/5% CO₂ until day 5.

Transfer 60 μ l from each well to another labeled 96-well plate. cover with a plate-sealer. and store at 4°C until Day 6 (for IL6 ELISA). To the remaining 100 μ l in the cell culture plate, aseptically add Alamar Blue in an amount equal to 10% of the culture volume (10 μ l). Return plates to incubator for 3 to 4 hours. Then measure
5 fluorescence with excitation at 530nm and emission at 590nm using the CytoFluor. This yields the growth stimulation/inhibition data.

On day 5, the IL6 ELISA is performed by coating a 96 well plate with 50-100 μ l/well of Anti-Human IL6 Monoclonal antibody diluted in PBS, pH 7.4, incubate ON at room temperature.

10 On day 6, empty the plates into the sink and blot on paper towels. Prepare Assay Buffer containing PBS with 4% BSA. Block the plates with 200 μ l/well of Pierce Super Block blocking buffer in PBS for 1-2 hr and then wash plates with wash buffer (PBS, 0.05% Tween-20). Blot plates on paper towels. Then add 50 μ l/well of diluted Anti-Human IL-6 Monoclonal, Biotin-labeled antibody at 0.50 mg/ml. Make
15 dilutions of IL-6 stock in media (30, 10, 3, 1, 0.3, 0 ng/ml). Add duplicate samples to top row of plate. Cover the plates and incubate for 2 hours at RT on shaker. Plates are washed with wash buffer and blotted on paper towels. Dilute EU-labeled Streptavidin 1:1000 in Assay buffer, and add 100 μ l/well. Cover the plate and incubate 1 h at RT. Plates are again washed with wash buffer and blotted on paper towels. Add 100
20 μ l/well of Enhancement Solution and shake for 5 minutes. Read the plate on the Wallac DELFIA Fluorometer. Readings from triplicate samples in each assay are tabulated and averaged.

A positive result in this assay suggests AoSMC cell proliferation and that the polypeptide of the present invention may be involved in dermal fibroblast
25 proliferation and/or smooth muscle cell proliferation. A positive result also suggests many potential uses of polypeptides, polynucleotides, agonists and/or antagonists of the polynucleotide/polypeptide of the present invention which gives a positive result. For example, inflammation and immune responses, wound healing, and angiogenesis, as detailed throughout this specification. Particularly, polypeptides of the present
30 invention and polynucleotides of the present invention may be used in wound healing

and dermal regeneration, as well as the promotion of vasculogenesis, both of the blood vessels and lymphatics. The growth of vessels can be used in the treatment of, for example, cardiovascular diseases. Additionally, antagonists of polypeptides and polynucleotides of the invention may be useful in treating diseases, disorders, and/or

5 conditions which involve angiogenesis by acting as an anti-vascular (e.g., anti-angiogenesis). These diseases, disorders, and/or conditions are known in the art and/or are described herein, such as, for example, malignancies, solid tumors, benign tumors, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; atherosclerotic plaques; ocular angiogenic diseases, for

10 example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis, retinoblastoma, uveitis and Pterygia (abnormal blood vessel growth) of the eye; rheumatoid arthritis; psoriasis; delayed wound healing; endometriosis; vasculogenesis; granulations; hypertrophic scars (keloids); nonunion fractures;

15 scleroderma; trachoma; vascular adhesions; myocardial angiogenesis; coronary collaterals; cerebral collaterals; arteriovenous malformations; ischemic limb angiogenesis; Osler-Webber Syndrome; plaque neovascularization; telangiectasia; hemophilic joints; angiofibroma; fibromuscular dysplasia; wound granulation; Crohn's disease; and atherosclerosis. Moreover, antagonists of polypeptides and

20 polynucleotides of the invention may be useful in treating anti-hyperproliferative diseases and/or anti-inflammatory known in the art and/or described herein.

One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

25

Example 45: Cellular Adhesion Molecule (CAM) Expression on Endothelial Cells

The recruitment of lymphocytes to areas of inflammation and angiogenesis

30 involves specific receptor-ligand interactions between cell surface adhesion molecules

(CAMs) on lymphocytes and the vascular endothelium. The adhesion process, in both normal and pathological settings, follows a multi-step cascade that involves intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and endothelial leukocyte adhesion molecule-1 (E-selectin) expression on endothelial cells (EC). The expression of these molecules and others on the vascular endothelium determines the efficiency with which leukocytes may adhere to the local vasculature and extravasate into the local tissue during the development of an inflammatory response. The local concentration of cytokines and growth factor participate in the modulation of the expression of these CAMs.

10 Briefly, endothelial cells (e.g., Human Umbilical Vein Endothelial cells (HUVECs)) are grown in a standard 96 well plate to confluence, growth medium is removed from the cells and replaced with 100 μ l of 199 Medium (10% fetal bovine serum (FBS)). Samples for testing and positive or negative controls are added to the plate in triplicate (in 10 μ l volumes). Plates are then incubated at 37°C for either 5 h
15 (selectin and integrin expression) or 24 h (integrin expression only). Plates are aspirated to remove medium and 100 μ l of 0.1% paraformaldehyde-PBS(with Ca++ and Mg++) is added to each well. Plates are held at 4°C for 30 min. Fixative is removed from the wells and wells are washed 1X with PBS(+Ca,Mg) + 0.5% BSA and drained. 10 μ l of diluted primary antibody is added to the test and control wells.
20 Anti-ICAM-1-Biotin, Anti-VCAM-1-Biotin and Anti-E-selectin-Biotin are used at a concentration of 10 μ g/ml (1:10 dilution of 0.1 mg/ml stock antibody). Cells are incubated at 37°C for 30 min. in a humidified environment. Wells are washed three times with PBS(+Ca,Mg) + 0.5% BSA. 20 μ l of diluted ExtrAvidin-Alkaline Phosphatase (1:5,000 dilution, referred to herein as the working dilution) are added to
25 each well and incubated at 37°C for 30 min. Wells are washed three times with PBS(+Ca,Mg)+0.5% BSA. Dissolve 1 tablet of p-Nitrophenol Phosphate pNPP per 5 ml of glycine buffer (pH 10.4). 100 μ l of pNPP substrate in glycine buffer is added to each test well. Standard wells in triplicate are prepared from the working dilution of the ExtrAvidin-Alkaline Phosphatase in glycine buffer: 1:5,000 (10^0) > $10^{-0.5}$ > 10^{-1} >
30 $10^{-1.5}$. 5 μ l of each dilution is added to triplicate wells and the resulting AP content in

each well is 5.50 ng, 1.74 ng, 0.55 ng, 0.18 ng. 100 µl of pNPN reagent is then added to each of the standard wells. The plate is incubated at 37°C for 4h. A volume of 50 µl of 3M NaOH is added to all wells. The plate is read on a plate reader at 405 nm using the background subtraction option on blank wells filled with glycine buffer only. Additionally, the template is set up to indicate the concentration of AP-conjugate in each standard well [5.50 ng; 1.74 ng; 0.55 ng; 0.18 ng]. Results are indicated as amount of bound AP-conjugate in each sample.

Example 46: Alamar Blue Endothelial Cells Proliferation Assay

10

This assay may be used to quantitatively determine protein mediated inhibition of bFGF-induced proliferation of Bovine Lymphatic Endothelial Cells (LECs), Bovine Aortic Endothelial Cells (BAECs) or Human Microvascular Uterine Myometrial Cells (UTMECs). This assay incorporates a fluorometric growth indicator based on detection of metabolic activity. A standard Alamar Blue Proliferation Assay is prepared in EGM-2MV with 10 ng /ml of bFGF added as a source of endothelial cell stimulation. This assay may be used with a variety of endothelial cells with slight changes in growth medium and cell concentration. Dilutions of the protein batches to be tested are diluted as appropriate. Serum-free medium (GIBCO SFM) without bFGF is used as a non-stimulated control and Angiostatin or TSP-1 are included as a known inhibitory controls.

Briefly, LEC, BAECs or UTMECs are seeded in growth media at a density of 5000 to 2000 cells/well in a 96 well plate and placed at 37-C overnight. After the overnight incubation of the cells, the growth media is removed and replaced with GIBCO EC-SFM. The cells are treated with the appropriate dilutions of the protein of interest or control protein sample(s) (prepared in SFM) in triplicate wells with additional bFGF to a concentration of 10 ng/ ml. Once the cells have been treated with the samples, the plate(s) is/are placed back in the 37° C incubator for three days. After three days 10 ml of stock alamar blue (Biosource Cat# DAL1100) is added to each well and the plate(s) is/are placed back in the 37°C incubator for four hours. The

plate(s) are then read at 530nm excitation and 590nm emission using the CytoFluor fluorescence reader. Direct output is recorded in relative fluorescence units.

Alamar blue is an oxidation-reduction indicator that both fluoresces and changes color in response to chemical reduction of growth medium resulting from cell growth. As cells grow in culture, innate metabolic activity results in a chemical reduction of the immediate surrounding environment. Reduction related to growth causes the indicator to change from oxidized (non-fluorescent blue) form to reduced (fluorescent red) form. i.e. stimulated proliferation will produce a stronger signal and inhibited proliferation will produce a weaker signal and the total signal is proportional to the total number of cells as well as their metabolic activity. The background level of activity is observed with the starvation medium alone. This is compared to the output observed from the positive control samples (bFGF in growth medium) and protein dilutions.

15 *Example 47: Detection of Inhibition of a Mixed Lymphocyte Reaction*

This assay can be used to detect and evaluate inhibition of a Mixed Lymphocyte Reaction (MLR) by gene products (e.g., isolated polypeptides). Inhibition of a MLR may be due to a direct effect on cell proliferation and viability, modulation of costimulatory molecules on interacting cells, modulation of adhesiveness between lymphocytes and accessory cells, or modulation of cytokine production by accessory cells. Multiple cells may be targeted by these polypeptides since the peripheral blood mononuclear fraction used in this assay includes T, B and natural killer lymphocytes, as well as monocytes and dendritic cells.

25 Polypeptides of interest found to inhibit the MLR may find application in diseases associated with lymphocyte and monocyte activation or proliferation. These include, but are not limited to, diseases such as asthma, arthritis, diabetes, inflammatory skin conditions, psoriasis, eczema, systemic lupus erythematosus, multiple sclerosis, glomerulonephritis, inflammatory bowel disease, crohn's disease, ulcerative colitis, arteriosclerosis, cirrhosis, graft vs. host disease, host vs. graft

disease, hepatitis, leukemia and lymphoma.

Briefly, PBMCs from human donors are purified by density gradient centrifugation using Lymphocyte Separation Medium (LSM[®], density 1.0770 g/ml, Organon Teknika Corporation, West Chester, PA). PBMCs from two donors are
5 adjusted to 2×10^6 cells/ml in RPMI-1640 (Life Technologies, Grand Island, NY) supplemented with 10% FCS and 2 mM glutamine. PBMCs from a third donor is adjusted to 2×10^5 cells/ml. Fifty microliters of PBMCs from each donor is added to wells of a 96-well round bottom microtiter plate. Dilutions of test materials (50 μ l) is added in triplicate to microtiter wells. Test samples (of the protein of interest) are
10 added for final dilution of 1:4; rhuIL-2 (R&D Systems, Minneapolis, MN, catalog number 202-IL) is added to a final concentration of 1 μ g/ml; anti-CD4 mAb (R&D Systems, clone 34930.11, catalog number MAB379) is added to a final concentration of 10 μ g/ml. Cells are cultured for 7-8 days at 37°C in 5% CO₂, and 1 μ C of [³H] thymidine is added to wells for the last 16 hrs of culture. Cells are harvested and
15 thymidine incorporation determined using a Packard TopCount. Data is expressed as the mean and standard deviation of triplicate determinations.

Samples of the protein of interest are screened in separate experiments and compared to the negative control treatment, anti-CD4 mAb, which inhibits proliferation of lymphocytes and the positive control treatment, IL-2 (either as
20 recombinant material or supernatant), which enhances proliferation of lymphocytes.

One skilled in the art could easily modify the exemplified studies to test the activity of polynucleotides (e.g., gene therapy), antibodies, agonists, and/or antagonists and fragments and variants thereof.

It will be clear that the invention may be practiced otherwise than as
25 particularly described in the foregoing description and examples. Numerous modifications and variations of the present invention are possible in light of the above teachings and, therefore, are within the scope of the appended claims.

The entire disclosure of each document cited (including patents, patent applications, journal articles, abstracts, laboratory manuals, books, or other
30 disclosures) in the Background of the Invention, Detailed Description, and Examples

is hereby incorporated herein by reference. Further, the hard copy of the sequence listing submitted herewith and the corresponding computer readable form are both incorporated herein by reference in their entireties. Moreover, the hard copy of and the corresponding computer readable form of the Sequence Listing of Serial No.

5 60/124,270 are also incorporated herein by reference in their entireties.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209059
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

ATCC Deposit No.: 209059

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209059

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209060
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<input type="checkbox"/> For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

<input type="checkbox"/> For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

ATCC Deposit No.: 209060

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209060

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209061
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209061

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209061**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209062
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209062

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209062

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209063
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209063

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209063**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209064
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209064

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209064

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>20 May 1997</u>	Accession Number <u>209065</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

ATCC Deposit No.: 209065

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209065

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209066
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209066

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209066

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>20 May 1997</u>	Accession Number <u>209067</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
<u>Europe</u> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

ATCC Deposit No.: 209067

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209067

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209068
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209068

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209068

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 20 May 1997	Accession Number 209069
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<p style="text-align: center;">For receiving Office use only</p> <p><input type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer</p>	<p style="text-align: center;">For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p>
---	--

ATCC Deposit No.: 209069

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209069

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

435

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 12 January 1998	Accession Number 209579
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 209579

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209579**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 12 January 1998	Accession Number 209578
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	
For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

ATCC Deposit No.: 209578**CANADA**

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 209578

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 16 July 1998	Accession Number 203067
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 203067

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203067

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

444

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> . line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 16 July 1998	Accession Number 203068
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

<p style="text-align: center;">For receiving Office use only</p> <p><input type="checkbox"/> This sheet was received with the international application</p> <p>Authorized officer</p>	<p style="text-align: center;">For International Bureau use only</p> <p><input type="checkbox"/> This sheet was received by the International Bureau on:</p> <p>Authorized officer</p>
---	--

ATCC Deposit No.: 203068

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203068

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 01 February 1999	Accession Number 203609
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 203609

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203609

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

450

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 01 February 1999	Accession Number 203610
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g. "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 203610

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203610**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>17 November 1998</u>	Accession Number <u>203485</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
<u>Europe</u> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: 203485

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: 203485

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> . line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 18 June 1999	Accession Number PTA-252
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE. (if the indications are not for all designated States) Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only
<input type="checkbox"/> This sheet was received with the international application
Authorized officer

For International Bureau use only
<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer

ATCC Deposit No.: PTA-252

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: PTA-252**DENMARK**

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
--	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u> .	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution American Type Culture Collection	
Address of depositary institution (including postal code and country) 10801 University Boulevard Manassas, Virginia 20110-2209 United States of America	
Date of deposit 18 June 1999	Accession Number PTA-253
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
Europe In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

For receiving Office use only	For International Bureau use only
<input type="checkbox"/> This sheet was received with the international application	<input type="checkbox"/> This sheet was received by the International Bureau on:
Authorized officer	Authorized officer

ATCC Deposit No.: PTA-253

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: PTA-253

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

Applicant's or agent's file reference number	PA102PCT	International application No.	UNASSIGNED
---	----------	-------------------------------	------------

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>91</u> , line <u>N/A</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution <u>American Type Culture Collection</u>	
Address of depositary institution (including postal code and country) <u>10801 University Boulevard</u> <u>Manassas, Virginia 20110-2209</u> <u>United States of America</u>	
Date of deposit <u>22 December 1999</u>	Accession Number <u>PTA-1081</u>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States) <u>Europe</u> In respect to those designations in which a European Patent is sought a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample (Rule 28 (4) EPC).	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable) The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit") 	
For receiving Office use only <input type="checkbox"/> This sheet was received with the international application Authorized officer	For International Bureau use only <input type="checkbox"/> This sheet was received by the International Bureau on: Authorized officer

ATCC Deposit No.: PTA-1081

CANADA

The applicant requests that, until either a Canadian patent has been issued on the basis of an application or the application has been refused, or is abandoned and no longer subject to reinstatement, or is withdrawn, the Commissioner of Patents only authorizes the furnishing of a sample of the deposited biological material referred to in the application to an independent expert nominated by the Commissioner, the applicant must, by a written statement, inform the International Bureau accordingly before completion of technical preparations for publication of the international application.

NORWAY

The applicant hereby requests that the application has been laid open to public inspection (by the Norwegian Patent Office), or has been finally decided upon by the Norwegian Patent Office without having been laid open inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Norwegian Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Norwegian Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on the list of recognized experts drawn up by the Norwegian Patent Office or any person approved by the applicant in the individual case.

AUSTRALIA

The applicant hereby gives notice that the furnishing of a sample of a microorganism shall only be effected prior to the grant of a patent, or prior to the lapsing, refusal or withdrawal of the application, to a person who is a skilled addressee without an interest in the invention (Regulation 3.25(3) of the Australian Patents Regulations).

FINLAND

The applicant hereby requests that, until the application has been laid open to public inspection (by the National Board of Patents and Regulations), or has been finally decided upon by the National Board of Patents and Registration without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art.

UNITED KINGDOM

The applicant hereby requests that the furnishing of a sample of a microorganism shall only be made available to an expert. The request to this effect must be filed by the applicant with the International Bureau before the completion of the technical preparations for the international publication of the application.

ATCC Deposit No.: PTA-1081

DENMARK

The applicant hereby requests that, until the application has been laid open to public inspection (by the Danish Patent Office), or has been finally decided upon by the Danish Patent office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the Danish Patent Office not later than at the time when the application is made available to the public under Sections 22 and 33(3) of the Danish Patents Act. If such a request has been filed by the applicant, any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Danish Patent Office or any person by the applicant in the individual case.

SWEDEN

The applicant hereby requests that, until the application has been laid open to public inspection (by the Swedish Patent Office), or has been finally decided upon by the Swedish Patent Office without having been laid open to public inspection, the furnishing of a sample shall only be effected to an expert in the art. The request to this effect shall be filed by the applicant with the International Bureau before the expiration of 16 months from the priority date (preferably on the Form PCT/RO/134 reproduced in annex Z of Volume I of the PCT Applicant's Guide). If such a request has been filed by the applicant any request made by a third party for the furnishing of a sample shall indicate the expert to be used. That expert may be any person entered on a list of recognized experts drawn up by the Swedish Patent Office or any person approved by a applicant in the individual case.

NETHERLANDS

The applicant hereby requests that until the date of a grant of a Netherlands patent or until the date on which the application is refused or withdrawn or lapsed, the microorganism shall be made available as provided in the 31F(1) of the Patent Rules only by the issue of a sample to an expert. The request to this effect must be furnished by the applicant with the Netherlands Industrial Property Office before the date on which the application is made available to the public under Section 22C or Section 25 of the Patents Act of the Kingdom of the Netherlands, whichever of the two dates occurs earlier.

What Is Claimed Is:

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 95% identical to a sequence selected from the group consisting of:
- 5 (a) a polynucleotide fragment of SEQ ID NO:X or a polynucleotide fragment of the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X;
- (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:Y or a polypeptide fragment encoded by the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X;
- 10 (c) a polynucleotide encoding a polypeptide fragment of a polypeptide encoded by SEQ ID NO:X or a polypeptide fragment encoded by the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X;
- 15 (d) a polynucleotide encoding a polypeptide domain of SEQ ID NO:Y or a polypeptide domain encoded by the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X;
- (e) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:Y or a polypeptide epitope encoded by the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X;
- 20 (f) a polynucleotide encoding a polypeptide of SEQ ID NO:Y or the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X, having biological activity;
- (g) a polynucleotide which is a variant of SEQ ID NO:X;
- 25 (h) a polynucleotide which is an allelic variant of SEQ ID NO:X;
- (i) a polynucleotide which encodes a species homologue of the SEQ ID NO:Y;
- (j) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(i), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide
- 30

sequence of only A residues or of only T residues.

2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a protein.

5

3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:Y or the polypeptide encoded by the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X.

10

4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:X or the cDNA sequence included in the related cDNA clone, which is hybridizable to SEQ ID NO:X.

15

5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.

20

6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.

25

7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.

8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.

30

9. A recombinant host cell produced by the method of claim 8.

10. The recombinant host cell of claim 9 comprising vector sequences.
11. An isolated polypeptide comprising an amino acid sequence at least
5 95% identical to a sequence selected from the group consisting of:
- (a) a polypeptide fragment of SEQ ID NO:Y or of the sequence encoded by the cDNA included in the related cDNA clone;
 - (b) a polypeptide fragment of SEQ ID NO:Y or of the sequence encoded by the cDNA included in the related cDNA clone, having biological activity;
 - 10 (c) a polypeptide domain of SEQ ID NO:Y or of the sequence encoded by the cDNA included in the related cDNA clone;
 - (d) a polypeptide epitope of SEQ ID NO:Y or of the sequence encoded by the cDNA included in the related cDNA clone;
 - (e) a full length protein of SEQ ID NO:Y or of the sequence encoded by the
15 cDNA included in the related cDNA clone;
 - (f) a variant of SEQ ID NO:Y;
 - (g) an allelic variant of SEQ ID NO:Y; or
 - (h) a species homologue of the SEQ ID NO:Y.
- 20 12. The isolated polypeptide of claim 11, wherein the full length protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.
- 25 13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.
14. A recombinant host cell that expresses the isolated polypeptide of claim 11.
- 30 15. A method of making an isolated polypeptide comprising:

(a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and

(b) recovering said polypeptide.

5 16. The polypeptide produced by claim 15.

 17. A method for preventing, treating, or ameliorating a medical condition, comprising administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 11 or the polynucleotide of claim 1.

10

 18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

 (a) determining the presence or absence of a mutation in the polynucleotide of claim 1; and

15 (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.

 19. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject comprising:

20 (a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample; and

 (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.

25 20. A method for identifying a binding partner to the polypeptide of claim 11 comprising:

 (a) contacting the polypeptide of claim 11 with a binding partner; and

 (b) determining whether the binding partner effects an activity of the polypeptide.

30

21. The gene corresponding to the cDNA sequence of SEQ ID NO:Y.
22. A method of identifying an activity in a biological assay, wherein the method comprises:
- 5 (a) expressing SEQ ID NO:X in a cell;
- (b) isolating the supernatant;
- (c) detecting an activity in a biological assay; and
- (d) identifying the protein in the supernatant having the activity.
- 10 23. The product produced by the method of claim 20.

SEQUENCE LISTING

<110> Craig Rosen,
Steve Ruben

<120> Human Colon Cancer Associated Gene Sequences and Polypeptides

<130> PA102PCT

<140> Unassigned

<141> 2000-03-08

<150> 60/124,270

<151> 1999-03-12

<160> 1556

<170> PatentIn Ver. 2.0

<210> 1

<211> 633

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (206)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (606)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (616)

<223> n equals a,t,g, or c

<400> 1

```
gcaggatgag ctacaggaag aatcagaaat gtcagaaaaa aagtcatgct cctcttctcc 60
caccctaaagt gagatatcca catcgctgcc tccagataga caaaggagaa aaagggagct 120
tcgcaccttt tcattttctg acgatgaaaa taaacctcct tcaccaaagg aaataaggat 180
cgaagtgtgt gaaggcttca cctggnacag taacctcttg aagtggagtg tggcagacgt 240
tgtgcggttc atcagatcca ctgactgtgc ctccattagc aagaatattc ctwgaccagg 300
aaattgatgg gcaggccctg ttgctcctta cccttccac tgttcaagaa tgcattggact 360
taaaattggg ccctgcatca aactttgcca tcacatagag aggrtcaagt ttgcttttta 420
tgagcagttt gccaaactgag aaggacaacc aaagtgagct gggatctttg aaggcacaaa 480
tgcaggcaaa tccttcaccc tgctttatta agtgggagct gggaatagtc ctggggggct 540
tctggggggt ttgcagggta ttcagctttg ctctcttttg gcacttttcg gggggaaggg 600
aggganttca cagtttaggg gaaggccaaa aac 633
```


<210> 2
<211> 295
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (147)
<223> n equals a,t,g, or c

<400> 2
agcggcacga gcggcacgag aagccctact cggatggggc acggactgtc caccttttct 60
aatgtgtgtt gtcagcctgt gctgtggcat agacatggat gcaaggacca ctttgagac 120
tgggtggcc tcaagagcac acagagnaag ggaagaagg gcatcacag gatgccagcc 180
cctgcctggg ttggggggcac tcagccacgg accagcccct tcctgggtat ttattctcta 240
tttattgggg gataggagaa gaggcattct rcctgggtgg gacaaacccc tttaa 295

<210> 3
<211> 442
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (378)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (422)
<223> n equals a,t,g, or c

<400> 3
aattcggcag agagagctgc tcagttagga cccagagga accatggaaa cccagcgca 60
ttkctyttcc tcctgtact ctggctccca gatattaccg cagaaattgt gttgacgcat 120
ttccagacac cctgtctttg tctcctgggg aaagagccac cctctcctgc aggaccagtc 180
agagtgttg cagcaacttc ttaacctggt atgaacaaaa atctggccag tytcccaggc 240
tcctcatgtt tggttaactcc cgcagccac tggcatccca gacagggta gtggcagtgg 300
gtctgggaca gayttcacty tcaccatcag cagactggag gytgaagatt ttgcagtgt 360
ttwctgttca acaatwtngg tttcgttcaa caatgtttca attttggggc agggggccma 420
gnrtggaaty aaaggactgt gg 442

<210> 4
<211> 754
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (446)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (662)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (700)

<223> n equals a,t,g, or c

<400> 4

```
tccgaagtcg gggaccctcc taccacctgt agagaagcgg gagtggatct aaaataaaat 60
ccaggawtct ggggggttcct agacggagcc agayttcggg aggggtgtcct gctactcctg 120
ctggggctcc tccaggacaa gggcacacaa ctgggtccgt taagcccctc tytcgctcag 180
acgccatgga gctggatctg tctccacctc atcttagcag ctctccggaa gacctttgcc 240
cagcccctgg aacccctcct ggaactcccc ggcccctga taccctctg cctgaggagg 300
taaagaggtc ccagcctctc ctcatcccaa ccaccggcag gaaacttcga gaggaggaga 360
ggcgtgccac ctccctcccc tctatcccca acccttccc tgagctctgc agtcctccct 420
cacagagccc cattctcggg ggcccntcca gtgcaagggg gctgctcccc gcgaatgcca 480
gccgccccca tgtagtaaag gtgtacagtg aggatggggc ctgcagtctg tggagggtgc 540
agcagggtgcc acagytgcc acgtgtgtga aatgctggtg cacgagctma sgcttgagcg 600
acgagacctg ggggtttgtg gagtgccacc ccaactagca tggagcgggt tttgaggacc 660
angatccctg ttgaagtgca gttgctggcc ctgggggagn tagccgttcg tttccggaaa 720
tttggaagt aggattttta aggttccaaa ttct 754
```

<210> 5

<211> 393

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (273)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (274)

<223> n equals a,t,g, or c

<400> 5

```
ttttctcttt gatgttctgg ccctgacttc tgaaccaggt gttctgggga gaggcatttg 60
agagtgaccc agggcctggg aaccggggct tttctcgtg gtcttaggcc tgttctgcaa 120
ccaaggcaag gccaggattt ccggaaatat gaggaaggtt ttgacccta ctctatgttc 180
accccagagc agatcatggg gaaggatgtc cggtcctac gcatcaagaa ggagggatcc 240
ttagacctgg ccctggaagg cgggtgtggc tcnnccattg ggaaggtggt cgtttctgct 300
gtgtatgagc ggggagctgc tgagcggcat ggtggcattg tgaaagggga cgagatcatg 360
gcaatcaacg gcaagattgt gacagattac acc 393
```

<210> 6
<211> 539
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (486)
<223> n equals a,t,g, or c

<400> 6
gctcgtgccg ctgaaatgac ttccacggct gggacgggaa ccttcacccc acagctatgc 60
ctctgattgg tgaatggtga aggtgcctgt ctaacttttc tgtaaaaaga accagctgcc 120
tccaggcagc magccctcaa gcatcactta caggaccaga gggacaagac atgactgtga 180
tgaggagctg ctttcgccaa tttaacacca agaagaattg aggctgcttg ggaggaaggc 240
caggaggaac acgagactga gagatgaatt ttcaacagag gctgcaaagc tgtggacttt 300
agccagaccc ttctgccctc ctttgctggc gacacttctt caaatgcaga tggttgtgct 360
cccttgccctg gggttttacc tgcttctytg gagccaggtg tcagggggccc agggccaaga 420
attccacttt gggccctgcc aagtgaaggg gggtgttccc cagaaamtkt ggggaagsctt 480
ctgggntgtg aaagacayta tgcaagytca ggwtacatc asgrgtsccc ggttgttgc 539

<210> 7
<211> 804
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (758)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (759)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (771)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (802)
<223> n equals a,t,g, or c

<400> 7
aattcggcac gagttcaaca ccaagtgtcc ctctggcagt tgtgtgatga atcagtatct 60
gagttcaaaa ttcccaaagg atttcagtac atcttgccgt gcacattttg aaagatacct 120
tttatctcag aaaccaaagt gcctgctgca agcacctatt cctacaaata taatgacaac 180
accagtgtgt gggaaccacc ttctagaagt gggagaagac tgtgattgtg gctctcctaa 240

ggagtgtacc aatctctgct gtgaagccct aacgtgtaaa ctgaagcctg gaactgattg 300
cggaggagat gctccaaacc ataccacaga gtgaatccaa aagtctgctt cactgagatg 360
ctaccttgcc aggacaagaa ccaagaacty taactgtccc aggaatcttg tgaattttca 420
cccataatgg tctttcactt gtcattctac tttctatatt gttatcagtc caggaaacag 480
gtaaacagat gtaattagag acattggctc tttgtttagg cctaactctt ctttttactt 540
tttttttctt ttttcttttt ttttaaagat catgaatttg tgacttagtt ctgcccyytg 600
gagaacaaaa gaaagcagtc ttccatcaaa tcaccttaaa atgcacggct aaactattca 660
gagttaacac tccagaattg ttaaattaca agtactatgc tttaatgctt ctttcatctt 720
actagtaagg cctattaaaa aaaataatac cacttgannng gtgaaggctt ngcaatagga 780
agaagataga atccaggttt angg 804

<210> 8

<211> 720

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (714)

<223> n equals a,t,g, or c

<400> 8

gtgaggacag cctgccagag tctggtctct ggacactatg ggcacacgac tcctcccagc 60
tctgtttctt gtcctcctgg tattgggatt tgccccacgg gctctcctga cacactctcc 120
ccctgcagag gtccagggga cccaacagcc ccagcaagat gagatgccta gcccgacctt 180
cctcaccag gtgaaggaat ctctctccag ttactgggag tcagcaaaga cagccgcccc 240
gaacctgtac gagaagacat acctgccgc tgtagatgag aaactcaggg acttgtagag 300
caaaagcaca gcagccatga gcaactacac aggcattttt actgaccaag ttctttctgt 360
gctgaaggga gaggagtaac agccagaccc cccatcagtg gacaagggga gagtccccta 420
ctcccctgat agcccagggt cagactgagc tcccccttcc cagtagctct tgcatectcc 480
tcccaactct agcctgaatt cttttcaata aaaaatacaa ttcaagttgc ttctcatgga 540
tggcactgct tttctgagga ctcaaagggt caagatggag gggctgactc agtccagcca 600
acatttaatg agcacctact ttatgtatgg agctctaacc catgggtcca tgggaataaa 660
gcagtgaata gtaacaataa ataattgtaa cagcaaaaaa aaaaaaaaaa aaanaaaaaa 720

<210> 9

<211> 540

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (463)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (524)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (535)

<223> n equals a,t,g, or c

<400> 9

```
ccaacagaga tgtatccaga gatccacagt tctggaggct gagaagtcta aaatcaaggc 60
accagcagat tccacatctc gtgaaggctc actctctgct tcacagatgg cactgtcttg 120
ctgtgttctc acatggcaga aggggcaaac aagccccctt gggcctcttt tataaaggca 180
ctaactctat gcctaaaggc agggccctca tgactctatc acctaccaa aggctccact 240
tctttatact attggagggg tagaaggaaac ttcctttcta gaccttgaag gtttaagaat 300
ttgaatctat aaaacaagct gacaatagac agattaacag gagaaaaagc atatacattt 360
tttaatgtgg gccagatggc agaagcttaa ataacacccc aagctacagg gaagtgaggc 420
ctctgatggg ggaggtagtt gacacaggct gtggggaggg gtnaggggga ggaatctgtg 480
gtggagcaaa tttgccttat tacactgata aatggtaatt tacnctaatt aagcntgggt 540
```

<210> 10

<211> 561

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (406)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (450)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (546)

<223> n equals a,t,g, or c

<400> 10

```
agaatgtcac tgagtggcct gaaccaaagc tcatggatac tggagatgaa gaaccagcag 60
gagacccagc gtgtgctctc atccgtactc tgtgacaccc cttagtggac tgttccaacc 120
agttggggccc gggagaagga ctagggaagc ttccttccca tctgcacctc ccagcctgga 180
gaggagcctg tgacttgccc tgttccttgg ctcctcaggc tggggtaagg tgtgggggtg 240
gctccagcct tgaggccagt ctcctgatgc tggattcctt gttttccctg caggcatcaa 300
acagaaaggc cttttgctaa gtagcagcct gatgcactcc gagtcagagc tggacagcga 360
tgatgccatc tttacatggc cagaccgaga gaagggcaaa ctccctngcat ggtcagaatg 420
gctctgtacc caacgggcag acccctctgn aaggccagga gcccgcgga rgagatcctg 480
tagccacctg gtctgtctyc tcagggcarg gccaggcaca attgccgggc cagttcttct 540
aacttnccga gttttgcggg g 561
```

<210> 11

<211> 393

<212> DNA

<213> Homo sapiens

<220>
<221> misc feature
<222> (346)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (381)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (383)
<223> n equals a,t,g, or c

<400> 11
tcgacccacg cgtccgggtg agtgccatct taaaacttac tggagattgg ttttatattt 60
agatttatat aactggttat gtgaatatat ttaaatactg gggaaattgc ttcactgtct 120
tagaaccaag caagattcac ctgtgttttg tgttcatggt catttgcttc ttaaaggcaa 180
gggttgaaga taaataaggt agcaatgtct atagttttgg ccttaactat gccaatctaa 240
ttataattcc ctgtatttaa aatggtttct tttacttatt gaaaggcatt ttagtggtgt 300
ktatgtgtaa tattaaagat tattcaacac ctctcaraaa aaaaanraaa aagggggggc 360
ccccctgggg gtcccaagct nancgtacgc ggg 393

<210> 12
<211> 322
<212> DNA
<213> Homo sapiens

<400> 12
taagaataca aaattagcca agcatgctgg cacatgcctg taatcccagc tactcgggag 60
gctgaggtac gagaatcgct tgaacctggg aggcagagga tgcagtgagc cgggatcacg 120
ccattgcact ccagcctggg ggacaagagt gaatctgtgt ctcacaaaaa aaaaaagaa 180
aaagaaagat gcttaacaaa ggttaccata agccacaaat tcatraccac ttatccttcc 240
agtttcaagt agaatatatt cataacctca ataaagttct ccctgctccc aaaaaaaaaa 300
aaaaaaaaaa aaaaaaaaaa ag 322

<210> 13
<211> 1907
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1834)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1843)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1888)

<223> n equals a,t,g, or c

<400> 13

```
gagaaataat aatatagctt tatagaatth tccatcttgt attaaaaataa tcacatgtac 60
atcattgttaa ctcatgtccat aacataagat tttgtacaac aatttctttt tgtgtgctgg 120
catcattaag gtttagtctg cccagatcac ctattagtac ctaatttata tattctgaat 180
taaaattatc tgttaattta aaaacatttt atctattgtc tttcaaaata gtattaactg 240
aggggttttt tgtgtgtgtt tttctatttt gcttggtctt ttgaacatta ctggactctc 300
gttttagaag gaaaaacctt tcagctctac tctcacatc ttatagcttt gtttgaacat 360
gccaaaaaac caggattagc tgcccatatt caaactcaca ggtttccaga ccgaataacta 420
ccaagaaaat tcgctttaac aacaaagatt cctgatacaa aaggctgcc acaatgtgtc 480
atagtcagaa acccttacac gggacataaa tacctctgtg gagctttaca gtctggaatt 540
gttttacttc agtggatga gccaatgcag aaattcatgt tgataaagca ctttgatttt 600
cctttgccaa gtcctttgaa tgtttttgaa atgctgtgtg tacctgaaca ggaataacct 660
atggctgtgt tagctattag caaaggcact gaatcgaatc aggtagtcca gtttgagaca 720
atcaatttga actctgcac ttcattggtt acagaaattg gtgcaggcag ccagcagtta 780
gattccattc atgtaacaca gttggagaga gataccgttt tagtgtgttt agacaaattt 840
gtgaaaattg taaatctaca aggaaaatta aaatcaagta agaaactggc ctctgagtta 900
agttttgatt ttcgcattga atctgtagta tgccttcaag acagtgtgtt ggctttctgg 960
aaacatggga tgcagggtta agccttcaag tcagatgagg ttaccagga gatttcagat 1020
gaaacaagag ttttccgctt attaggatca gacagggttg tcgttttgga aagtaggcca 1080
acagaaaatc ctactgcaca cagcaatctc tacatcttgg ctggacatga aaatagttac 1140
taagcaacag aaactgatct caaatgacag gaaaatgaat atactccatt gaaaggaaaa 1200
ataaggaaat tcaatacaaa ctgcactatg atttgcttta actattatgg gttatattgc 1260
aatgatctg tactttaggg tagaattcaa tttttctgc agctggaaac agctagtcta 1320
tctcttgcca ctgtgtgtg gttatatcaa gtttgcttaa taaaagctat gagacaaata 1380
gtcctctagt tccaggaaac acagtctttt tttaaaaaaa acaatgtttg taacaagggt 1440
gccatgggat ttttagataa ctctgtatta tcttaagaga ggtaaattha gtgatcattt 1500
tatatcatgt cttattcctt cttaatgaac ataatttgtt aaattctcaa gcaaggtttt 1560
cacttttata ttggccattc tgtatgtttt tgtaaaacag aatatttaat ccttattttt 1620
taatctcttg ctggagtggg gtaatgtatc taacttttag caaaggaggg ttgcagagca 1680
gcttaaatth tttttataat gtataagaat tttgtttatc ttttaagagt agtaaagtac 1740
tttgagtgtt tgggggtcaa cacacacatg caatttgctt aacaaaagta ttttataata 1800
cagttcatat agaataccyt aaaaggagc ytangtttcc acnacagata agtggttaagg 1860
gtcataccgg agataatgat gatagtgnaa tatcctagaa gggggtg 1907
```

<210> 14

<211> 1140

<212> DNA

<213> Homo sapiens

<400> 14

```
agaataaatt ccagggtctc agtctgcctt tggttcgcaa gtttgccac tcgattctgc 60
agtgttggga tgctttgcac aaaaacagaa taattcactg tgaccttaag cccgagaca 120
ttttgttaaa gcagcagggt agaagcggta ttaaagtaat tgattttggc tccagttgtt 180
acgagcatca gcgtgtctac acgtacatcc agtcgcgttt ttaccgggct ccagaagtga 240
```

```

tccttggggc caggtatggc atgcccattg atatgtggag cctgggctgc attttagcag 300
agctcctgac gggttacccc ctcttgccctg gggaagatga aggggaccag ctggcctgta 360
tgattgaact gttgggcatg cctcacagaa actgctggat gcatccaaac gagccaaaaa 420
ttttgtgagc tccaagggtt atccccgtta ctgcaactgtc acgactctct cagatggctc 480
tgtggtccta aacggaggcc gttcccggag gggaaactga gggggccacc ggagagcaga 540
gagtggggka acgcgctgaa ggggtgtgat gatccccctt tccttgactt cttaaaacag 600
tgttttagagt gggatcctgc agtgcgcatg accccaggcc aggccttgcg gcacccttg 660
ctgaggaggc ggttgccaaa gcctcccacc ggggagaaaa cgtcatgaaa aggataactg 720
agagcaccgg tgctatcaca tctatatcca agttacctcc accttctagc tcagcttcca 780
aactgaggac taatttgcg cagatgacag atgccaatgg gaattattcag cagaggacag 840
tgttgccaaa acttgttagc tgagctcacg tcccctgatg ctggtaacct gaaagatacg 900
acattgctga gccttactgg gttgaaaagg agtagctcag acctgttttt atttgctcaa 960
taactctact catttgatc ttttcagcac ttaattttta tgtaagaaag ttgttcattt 1020
tgtttttata aaatacatga ggacaatgct ttaartttta aaaaaaaaaa aaaaaaaaaa 1080
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaagg gcggccgctc gcgatctaga 1140

```

<210> 15

<211> 2008

<212> DNA

<213> Homo sapiens

<400> 15

```

gccaggagac cctgaaacat gaaacaaaac aggccttgat attttttttt ttttaattac 60
tttcccccctt tgcttgattc ttcttccttg gtatcatatt caaaggagga aagaatggat 120
gacacttagg gacagggtcac taagcagaat aggtattagt tggtttttta ttatttttaa 180
atttatatag ctctcatgta ttgaagggtc tactttaaaa actgtgttaa tgtacttgca 240
aatctcctgc tggctcatga aacagcagaa ttccagccaa ggcaaatca gttggatcta 300
gggcttgaaa actgcccag attaaagagc taagatgaaa tagtttgaat aaactggtaa 360
aataacaatt catactatct ctacatattt ttaggatatg aatttttttt tcctgctggg 420
tagattgtcc agtgacttat ggcatgaatt tcttttcatg ctacctcaa tcttcaacta 480
aagatttcta aaaggggtaa gaaatgaggc agtagtttat taaatacata gaagtacaca 540
gctattatat gctgtggctt tgagaagtta acttttgagg aatatgaaac caaaggaaga 600
atttccatgt agataaatta agaattagga aaaacatcta cctaaaagat ggtgtgccct 660
aaaaaactgg aaacatctga aatgctctat ttatgttatt attcctggga gcttatagcc 720
cattattttt actttttttt ttaattcag aagactagaa cagaacttaa attttacaaa 780
cttttgatca taatgctttt gccatactg tctcgtcctt agccccagtg acatgtgctg 840
atcctcctgc ctttagttct aaatggttct gaatagcttt aaaagatgaa taagaaaaaa 900
gatcaagctc ttttgaggct agagggtca tatggagatc taaaaccctc catgttcatt 960
tccagacttg gtcaggcaca tacatagtaa attataatca atcatgggaa ttacataagg 1020
ataatgaggc ccaactgaaa ccaagtttca gtctgcctta cctttaccog tccttgagaa 1080
cagcgggtcca ggagaatcag gcagtctgtg gcctcctgta gcagggcgct cttaacagac 1140
tcagggtgaa ggtttggatc ccttgctaca tcacaaatca gtttcaagag ggccttcagt 1200
aagaccacaa gtctgcagga aactctggaa tcaggaagaa aagctatgtt catactctaa 1260
atctggatat taaaattgga agagacttca aagactgtca agtggaagc tatcatttcc 1320
taaatgaagg aacttaggtc aaagttaaat gaaaaagacc taaaaacaga accaaagaca 1380
gcctctagat ttcttaccct caagtctcct gttagcatac tgcctatata cacagacaca 1440
ccctctgcca cactgctctc ttccctccag aagcttggtt ctactgtaag caacagctcc 1500
ttcatgggct gagaagggga aaccaggatc tccccaccca agggcctata cttttatttt 1560
tcaactctat taactgaact atgcttgagg agagccctga atcctttaac tttaatgcca 1620
tcttaggcct ggaagttaga gtgagggaaa tgagcttggt ttctgggact gtgtcaagtt 1680
gaattttctg tcatttaatt tcagatccta gctcatcatc agtgaagtcc agcctactta 1740

```


cttgtaacta ttaacctttc taaaagtttt catcgccttg gtaaagctcc agaaatggac 1800
aatgctagtg actgcacaat attgtgactg tacttaatgc cactgaactg tacacttta 1860
aatggaaaaat ttatatgtat ttttaccaca ataaacaaaa aaccctaaaa aaactttaat 1920
gaaagggtgga aaataattta acttayaatg tgaaaatata atgtgaaatg tacaataaat 1980
catatttatg gcaaaaaaaaa aaaaaatt 2008

<210> 16

<211> 371

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (350)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (360)

<223> n equals a,t,g, or c

<400> 16

agagagctag agggactagg agataatgtg tatgtagggtt tatgtgatgg gatatcaccc 60
tgaagagttg tgtcttttgt ggccagtgc aaatccagga aatgaatgtt gctgatagg 120
ataaatcttg aggctgaggg cgggtggtac agatgtgtat gggaaacccc aaccctata 180
tattgtaaat agatgggctg ggctaaacat tgttgccgtt tcatacttct accaactcag 240
cttttacaca ataaagctct actgtctctg aaaaaaaaa aaaaaaaaa aaaaaaaaa 300
aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaaaa aaaaaaatn gggggggggn 360
ccccccccc c 371

<210> 17

<211> 763

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (6)

<223> n equals a,t,g, or c

<400> 17

aaaaaaaaa aaaaaaaaa aaacggaaat aaagtttaaa aaatcaatca acatggcctt 60
taattttaac aattttaaca gcaagtgggtg gggggagttc tcagatgagc aactggagct 120
ggaagcactt ctgtggtcaa gcaggcagcc catgggggtg catcttctctg ttgggggatc 180
atccattttc ttcaatgaat agttttaagt ctgtcgaat gctcacacag aggcccgcta 240
ttaaggaggc agacaggcaa cattcaatac gaaggcagga caagctcagc cccgctcctt 300
cattcgggca tgtgtcatta gggatgacat tctctgaagg ctgcccggct tgaatggcca 360
aatccctgca tcatggcttt cttaattcc ctctgctccc aactcacaaa atgaggacct 420
ctcttttaag acgaraaagg cactgttctt caaagggtata catttggaac ttcaataatg 480
aaagcatctc ttgcttgga ggtggaatat aggcaatttt ggatttttaa tgcattggcat 540
ggggcgaggag tgaaatatct tgccagggtt tgtttgccct ataattggag agaaccaggc 600

ctctggatga tacggtatca aacactgctg ctcctttctg ttttcttttg tgggaaaggg 660
aggaggatag aatggagagg aattagtggg agcctggggg aagttcaaaa taaagaaact 720
gtgaaatcct ccacctcaaa gttgggtctg caccaggatt ctg 763

<210> 18

<211> 1926

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (898)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1024)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1083)

<223> n equals a,t,g, or c

<400> 18

aataaacttga ctggaattct agaaaccgct gtcagagcca ccaacgcaca gtttgacagt 60
cctgagatcc tgcgaaggct ggacgtgctg ctgctggagg tctctccagg tgacactgga 120
tgggatgtct tcagcctcga ttatcatgtt gacggaccaa ttgcaactgt gtttactcga 180
gaatgtatga gccactacct aagagtattt aacttcctct ggagggcgac ggatggaata 240
catcctcact gacatacgga agggacacat gtgcaatgca aagctcctga gaaacatgcc 300
agagtcttcc ggggtgctgc accagtgtca cattttggcc tctgagatgg tccatttcat 360
tcacagatg cagtattaca tcacatttga ggtgcttgaa tgttcttggg atgagctttg 420
gaacaaagtc cagcaggccc aggtatttga tcacatcatt gctgcacacg aggtgttctt 480
agacaccatc atctcccgtc gcctgctgga cagtgactcc agggcacttt taaatcaact 540
tagagctgtg tttgatcaaa ttattgaact tcagaatgct caagatgcaa tatacagagc 600
tgctctgga gaattgcaga gacgattaca gtttgaagag aaaaagaaac agcgtgaaat 660
tgaggggcag tggggagtga cggcagcaga ggaagaggag gaaaataaga ggattggaga 720
atttaaagaa tctataccaa aaatgtgctc acagttgcga atattgaccc atttctacca 780
gggtatcgtg cagcagtttt tgggtgttact gacgaccagc tctgacgaga gtcttcggtt 840
tcttagcttc aggtggact tcaacgagca ttacaaagcc agggagccca ggctccgntg 900
tgtctctggg taccaggggg cggcgcatc ccacacgtga arctcgcggt cctcccargg 960
agctgcgggt gatgttcggt gcaactgctag acacgaaatt cccattgacg tcctgcagga 1020
actnatkret gcaggtgtcc tgcccttcgc cccacgagtg cgccatgttt cagcggaggc 1080
cgntgtggga gaagccacgt cgtgtttcam atgtcggagt cgaatgcatt tktaaattccc 1140
taagtcaagt aggtggctg cactgttcac atttgtctct aaaagtcttc atcgctaaaa 1200
gataccataa tttgctgagg cttcttaagc tttctatggt ataatttata tttgtcactt 1260
taaaaaatcc atttctttta gaaaaaatta ggggtgatag atattcatta gttaagatgg 1320
taacgtcatt gctatttttt taacatcctc ttttagaggta atttttgtta acataaccaa 1380
aaattaaatt gaaacaaaat gtcccaacta agaaaatata tagagcattt tatttttttt 1440
tagtgttgta aaatattaac ctctgtgaga tcctttgtat cttaatgcat tacctttaca 1500
catatttatt cttattttct ctcctttcag agtttacatt tttatattta atttactatt 1560

tcagattttt aaaatagtat agaaaaaagt aggagtgata gagaacaaaa atactcttat 1620
acagtgcac ccaaataccg cgaatgcac agctaaagca gcgtgtaaat aggagtgcg 1680
agaaagttaa tggagtattt tttttcaaa gttcctgata agcattggaa agaaatcgac 1740
atggataatg aagatttcct ttttccttgc ctattttttc attgtaaata tttatatact 1800
actgmccaaag atgttgggtt gggggggatt gttttttgta aaaatgtcat tatcaggtca 1860
cataaatctg cctttatggt gcataagtga aaatttagaa aattaaaagc aattatcttt 1920
magaaa 1926

<210> 19

<211> 2301

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (190)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2052)

<223> n equals a,t,g, or c

<400> 19

tcttatcaca tctaagcctt accaggtaga gaaggtagta aatacacttt agaagtaaaa 60
atatgaagta ccgagaggct aaacccactg gcctaagatc tcaccaaagt tcatgaaaac 120
caggactagg acccacggtt cccaaagccc gttcttgctg tgttgtgctg cctccatata 180
cgtcagaagn agcctttcca gaatgattct gggcatatac taagaagagc aggtatggaa 240
agatctattt tcagggaatc ttagaattcc ctacacgagt gggagaaaaga tgcccaaatt 300
ccttacgcat ggtattcatg atggtgccct atctaagtcc aggactgttt tcctacagcg 360
tgctcaaaaa gtgtttaga gggcaggatt ctacattcac agcctgttcc atctacgaga 420
ttttccagat gctacttggt gtagacattc ctaactcatg gtacttagcc accagagatc 480
atgatggaat gagggtgtg cttttctacc tgccattccc tcagaattca tgarggggtg 540
gggacagggg gaccggaatt gtcttagcac cccaatgtta tgacaaaact atgctacttt 600
agaaacgcag tctgtttttc accaattgac atactactga tctgaagtaa ccagtgccat 660
cataagaaat tactgcatta agaaaatcct tgctgtgcc tttgaaaagc tgttcagaaa 720
tcatttacag tgatctttca tctcgtcgc ttagtgtaac attttagtgt gataaatttc 780
aaaattctaa acaaattacc cacttttata ttgaaaatct ctaccagaac tccctcttca 840
tttttwaagg catacatttg cttgttttca agatcaagaa ttctgagcta gctttaagta 900
gcaaactgat ttatatgtgc aattatagga tgcattaaga tgaatgatag cctttacata 960
ttgaaaactt tgcagacgtt ttgttttgaa aatggcattg tatagtaaag gcaaattaat 1020
tttgtaaaat tatgttaaag agtatgttca gacactttct gccatggcca aaaagtatgt 1080
atgaaagtat gtgtgtattt gtttgtaaaa ggatgccaat gttttacctg atatcttagt 1140
gacacttcag ttatctatgc attctttaga tctgtgattc ggtaaacagg cagccatgtt 1200
cacgatgcct tctatgtctt accatatttt taattaacct gttaaatata gcttaaaata 1260
tttttatttt atttattcta tttttactga aatatactgc attattgtgt taatgtatta 1320
tctttcctgg atattatctc ccagtgtatc cagatctaag taatctcagt gaactataca 1380
ttgcctaaaa agtgggtttt taatgatttg tagtcacatt tctattggga tatgtagaag 1440
aaaaggcaaa atgcttaaag ttccttttat tttttaaaag cagctagata gacacagact 1500
tgccacctca tacatctgct ccttgccaac atcaagggga acgactagcc aacatgccta 1560
tggtctaaaaa ctttcctttt cagactaaaag cactgcttgg tgcttcgttt ttctaccctt 1620

```

cacaacatgt gtgatttcat ctaagagata tatacatgta cacatgccct ttgtttccac 1680
ctggatacaa gatcactcat agctaattag gaccattgtt ttttggtcat ctgtcttgtt 1740
gcatgaaggg acattagacc catttccatt aaaataagtt cttggtgata aactgtggca 1800
ctgctacttc tttttaaatc cactttatga tttcaagatg gacacttgta agatgactcg 1860
acacaaggcc attgacctgga agccccagag ctttcctctg tttgtatggc ccgttcatgt 1920
cccaggcatt gcaacacaaa ctcaagattt caccacaaca tgacaagcat tttcctaact 1980
gatattagca caatttaact aataagcccc ttcgctctct agttggccag gcttaacct 2040
atacacatct anactgtgtg ccacacggcc agtagaaagt ttaacttcag cttcagggca 2100
aagataccca ctcacaccgt gtcaacgcaa gcagtagttc ctggcctcca gagcagctta 2160
cttcccctga aagaacgctt tgttttcctt tatgcccttt tctgtgtgac cacttttaca 2220
catttaaagt taatttggtg tgagaataaa tttagctgca taaaaaaaaa aaaaaaaagg 2280
gcggccgctc gcgatctaga a                                     2301

```

<210> 20

<211> 538

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (507)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (514)

<223> n equals a,t,g, or c

<400> 20

```

agaggccgcc aacatgatcc tgggtggatg tgacttctca gccatcatga atgcagtgga 60
ggaaggcaag ggtatttttt acaacatcaa aaactttgtc cgattccagc tgagcacgag 120
catctccgcc ctgagtctca tcaactctgtc caccgtgttc aacctgcccc gccccctcaa 180
cgccatgcag atcctatgga tcaacatcat catggatggg ccaccgggca gaggtgaggc 240
aggggcggctg ggagccctgt gtctctttac ctacctgcgg ggcttcctcc aggggctgct 300
ggctgtgccc aaggctatag ggatgaacaa atacagccac tttccatcag gagttccag 360
aaaactgaag tgtgttgacac tggagtgaga ctgggagtag aaggcagagg agaaagtacc 420
tgggccggca gagctgggtg aggatggaac tttctgcttc ctctggctgg atgctctctc 480
tgggcaaacc tgcattgggtt aattctnatg cttnaatttc aagtcaccca gtcactgg 538

```

<210> 21

<211> 1403

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1370)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1386)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1393)

<223> n equals a,t,g, or c

<400> 21

```
ggcacgagtg gccacatttc cagtacccag tagtcatctg tgccaggggg ttatccaggt 60
acagaacatt cccatcggtg cagaagggtc tatcagctag cactgggttg gacgacactt 120
gccaagacga gctggctaga ggatrgttct ccggacctgg tcccacgtgg ttcccagctg 180
gctggaggcg tgatcctggg tgtggccctg tggtccgcc atgacccgca gaccaccaac 240
ctcctgtatc tggagctggg agacaagccc gcgcccaaca ctttctatgt aggcactctac 300
atcctcatcg ctgtggggcg tgtcatgatg ttctgtggct tcctgggtcg ctacggggcc 360
atccaggaat cccagtgcct gctggggacg ttcttcacct gcctgggtcat cctgtttgcc 420
tgtgagggtg ccgccggcat ctggggcttt gtcaacaagg accagatcgc caaggatgtg 480
aagcagttct atgaccaggc cctacagcag gccgtgggtg atgatgacgc caacaacgcc 540
aaggctgtgg tgaagacctt ccacgagacg cttgactgct gtggctccag cacactgact 600
gctttgacca cctcagtgtt caagaacaat ttgtgtccct cgggcagcaa catcatcagc 660
aacctcttca aggaggactg ccaccagaag atcgatgacc tcttctccgg gaagctgtac 720
ctcatcgcca ttgctgccat cgtggctcgt gtgatcatga tcttcgagat gatcctgagc 780
atggtgctgt gctgtggcat ccggaacagc tccgtgtact gaggccccgc agctctggcc 840
acagggacct ctgcagtgcc ccctaagtga cccggacact tccgaggggg ccatcaccgc 900
ctgtgtatat aacgtttccg gtattactct gctacacgta gcctttttac ttttgggggt 960
ttgtttttgt tctgaacttt cctgttacct tttcagggtc gacgtcacat gtaggtggcg 1020
tgtatgagtg gagacggggc tgggtcttgg ggactggagg gcaggggtcc ttctgccctg 1080
gggtcccagg gtgctctgcc tgtcagcca ggcctctcct gggagccact cgcccagaga 1140
ctcagcttgg ccaacttggg gggctgtgtc caccagccc gccctcctg tgggctgcac 1200
agctcacctt gttccctcct gcccgggttc gagagccgag tctgtgggca ctctctgcct 1260
tcatgcacct gtcttttcta acacgtcgcc ttcaactgta atcacaacat cctactccgt 1320
catttaataa agaaggaaca tcaggcatgc taaaaaaaaa aaaaaaaaaa ksgggggggg 1380
gcccgnatcc canttggccc aat 1403
```

<210> 22

<211> 478

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (474)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (475)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (476)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (478)

<223> n equals a,t,g, or c

<400> 22

```
gagcagaaaag tagatgctta atggcagtggt tctctgaccc agacatgaat cagaatcacc 60
tggaagggtct tgtaataaca aattgctagg ccacaaccct aaagtttctg attcagggta 120
gggcaaggcgc aggcttaaac ttcaggccag gggccacttt aagaattgct atatggccag 180
ggccgggcgc ggtggctcac gcctgtaatc ccagcacttt gggaggccga ggtgggcgga 240
tcacaaggctc aggagatcga gaccatcctg gctaacacgg tgaaaccctg tctgtagtaa 300
aaatacaaaa aaattagcca ggcatgggtg tgggtgcctg tagtcccagc tacttgggag 360
gctgaggcag gagaatgggtg tgaaccagag aggtggagct tgcagtgagc cgagatcgtg 420
ccactgcact ccagcctggg caacagagcg agacttccgt ctcaaaaaaa aaannncn 478
```

<210> 23

<211> 1252

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1227)

<223> n equals a,t,g, or c

<400> 23

```
tttcgagctc tgcaccgagg agctgccctg gacttgagtc ccttgcatcg gagtcccat 60
ccctcccgcc aagccatatt ctgttggatg agcttcagtg cctaccagac agcctttatc 120
tgcttgggct tcctgggtgca gcagatcctc ttcttccctg gaaccacggc cctggccttc 180
ctggtgctca tgctgtgct ccatggcagg aacctcctgc tcttccgttc cctggagtcc 240
tcgtggccct tctggctgac tttggccctg gctgtgatcc tgcagaacat ggcagcccat 300
tgggtcttcc tggagactca tgatggacac ccacagctga ccaaccggcg agtgctctat 360
gcagccacct ttcttctctt cccctcaat gtgctgggtg gtgccatggg ggccacctg 420
cgagtgtctc tctctgccct ctacaacgcc atccaccttg gccagatgga cctcagcctg 480
ctgccaccga gagccgcact ctgcaccccg gctactacac gtaccgaaac ttcttgaaga 540
ttgaagtcag ccagtcgcat ccagccatga cagccttctg ctccctgctc ctgcaagcgc 600
agagcctcct acccaggacc atggcagccc ccaggacag cctcagacca ggggaggaag 660
acgaagggat gcagctgcta cagacaaagg rctccatggc caagggagct aggcccgggg 720
ccagccgcgg cagggctcgc tggggtctgg cctacacgct gctgcacaac ccaacctgc 780
aggtcttccg caagacggcc ctgttgggtg ccaatgggtg ccagccctga gggcagggaa 840
ggtcaaccca cctgcccatc tgtgtgagg catgttctg cctaccatcc tcctccctcc 900
ccggctctcc tcccagcatc acaccagcca tgcagccagc aggtcctccg gatcacggtg 960
gttkggtgga ggtctgtctg cactgggagc ctcargarg ctctgctcca cccacttggc 1020
tatgggagag ccagcagggg ttctggagaa aaaaactggt ggggttagggc ctgggtccag 1080
gagccagttg agccagggca gccacatcca ggcgtctccc taccctggct ctgccatcag 1140
ccttgaaagg gcctcgaata aaccttctct tggaaacct ccaagcccag ctccactcag 1200
ccttggcctt cagccttgtg gaaacancca aggcattcct ccaccctca ag 1252
```

<210> 24
<211> 1074
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (928)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (934)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1028)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1031)
<223> n equals a,t,g, or c

<400> 24
gctggccttt gctctgcgga ctggcgcccc cctggcacag aggtgacatc tcaagggccc 60
aggcagccct ctcttagtgg tgccaagagg cggaggctgc gggctgccct tggccccag 120
cccactcgct cagccctgag gtttccctct gcttcccag ggagcttgaa ggccaagcag 180
tccatggcgg gaatccstgg tagggagagt aatgccccat ctgtgcccac tgtctccctg 240
ctgccggggg cgccctggagg caatgccagc tccaggacag aggctcaggt gcccacggg 300
caaggcagcc cagggggctg tgtctgttca agtcaggctt ccccgcccc tcgcgagca 360
gcgcctccac gggcagcccg gggccccacc ccacgcaactg aagaggccgc ctgggctgcc 420
atggccctga ccttccctgct ggtgctgctc accctggcca cgctctgcac acggytgac 480
agaaacttcc gacgcgggga gagcatytac tgggggcccc cagcggacag ccaggacaca 540
gtggctgctg tgctgaagcg gaggtgctg cagccctcgc gccgggtcaa gcgctcgcgc 600
cggagacccc tcctcccgcc cagcccgagc agcggcccg aaggcgagag ctcgagtgta 660
cggcctggga cctgccactg tggcgtgcgg ctctccccc cgccgcgagg ccgcgacctc 720
tgccacgtgg accgcgcgcg gggcgctccc tggtggcgat ggcgcggcac tggccgagca 780
ctgcgggggc ttctctcctt gttggttgcg gagtgggcgg ccaaggggag aaaaggagcc 840
gcttctgcct cccttgccaa aactccggtt ctaattaaat tatttttagt agaaaaaaaa 900
aaaaaaaaaa aaaaaggcgg ccgtttanag gatncctcga aggggcccga gttaacgcgt 960
gcatgcgaag tcataactct ctccctataa tgatcgatt ataagtaagc actggccgtc 1020
gtttttanaa ngtcgtgaat ggggaaattt gctaactttt ggaacttttt gaaa 1074

<210> 25
<211> 1186
<212> DNA
<213> Homo sapiens

<400> 25

```

tatcagctca agccttacac cayccacctc atcaaggacc tccacttttt ccttcgagtg 60
ctcatccaac tgtaccaccg tatccctcac aagctacaca tcataccact ttgggaccgg 120
gacccagca ccagccttct ggaacagggc cacattgtcc attacctgtc acaggctctc 180
atctccagcc ccaaggacca aacagtattc caacacctac tgcttcaggg ttctgtcctc 240
atcctggctc tgtggccctg ccacatgggg ttcaaggacc tcagcaggca tctccagtgc 300
ctggacagat tccaattcac agagcacagg tgccaccaac atttcaaac aattaccatg 360
ggtcagggtg gcattaaaat ggactccaaa aacatttttt taaatgttct gtaagataaa 420
ctgtatatatt catatgtacc tgtaaggta ctttttaaaag ctgtacatg aacctttgta 480
taaaaaaacac cagtgtctct tcgttgtatt tttctcattt ttgcttttta aaattccttt 540
aaaaaatgtg ctgttaagcc agtattaggt atctttatct tgtaagtga cattccagct 600
gtttttttct ggcagatctg atgctgattt gatgctgtat gatctttttt ttttttttag 660
ttaaattcat ttagtgaatg ttctattatt ttatacatc acattaagta ctcagctaa 720
taatggcact atgaggattt tttttttctt tcctgtcagc agcagttctg tgaatgcac 780
ttaggtataa aaatgcaata cagattttta tattttgggtg tggacatggc tcattttgtt 840
ttaccagtta ttgcaagca aaatgtaatt taatgtatag atgatttcta atgtctcctg 900
acaaactgta aatactgcat ttcttttgcg tatataattg cttacagctt ttctcatttg 960
atatatagca ttgtacatat gacaagtctt ttgcaaaact gtgtgatctt tgtgaaagta 1020
gtacagtata tgacctttaa ttcttttttt attttaaata tactgtcaca ctgaagcact 1080
ggttgggcat ttaattcat gttaataaat cacaattatg tcagttccca cgcgtccgcg 1140
gacgcgtggg cggacgcgtg ggcggacgcg tggcgggacg cgtggg 1186

```

<210> 26

<211> 888

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (670)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (675)

<223> n equals a,t,g, or c

<400> 26

```

gccccaccc gctgtgactg cattctggga tccccagcct gggaatccaa gagctgtcgg 60
ccatttttat tcctccctcc cagacctgac cccttcacg gggctcaaga gacctctctc 120
tccaaatctc catttgccctc ctctggctaa gctggaaaat gcacactctg ccctgggtgt 180
ttccatatta tccgcctgcc ctctcctctg ggtgcctccc gtagccttag taagggtctc 240
gctttcctgg gcccttagag ctgagccatg ctttgccata aagggtgctc cggcttgcaa 300
ccaatgtgtc tgcttggtga tctgtctgtg ggtgtggtgg ggaggagggg gaccaggtgg 360
gtactggcac tctggggtcc ggactttatg tccatggagg ccccaattga ctcagttcaa 420
gggtcactga ggctttgctg atgtaggag agggccagag ggaggctcca cccagcccg 480
gctgagccag ggaacctggg acaaaagtca ggtggctgat tccaggtagt gttttggagc 540
tgggcagtca gtggctgggc ggggacatat gcccagagc caccatgaac tcccaggggc 600
ctccaggcag gggccctcca tcccgcacca agtctttcaa catgatgtcc ccgacgggcg 660
acaactcggc ctacnggctg agattaaggc aggcaagagc ctgaagccga cgcgccagag 720
caaggggctg accacagtgt tttaaggcat cgggcagccg gccttcagg taggcggggc 780
cagcaggagc ctgcgacctg gcttccctgg ccctaggcca ccgggcgctc agccccaccg 840

```


cttctccctg cagccccgatt cgccgctgcc ttctgtgtca cctgcact 888

<210> 27

<211> 789

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (420)

<223> n equals a,t,g, or c

<400> 27

gggtcgaccc acgcgtccgg ttggtcttgt gtctataaga atgatcaagc tgcaaaggat 60
aatccaacta aaagtcttca ggaagaagaa ccatgtccaa ggtttgccca tcagcttgta 120
tacgatgagc tacacaaggt tcattactta ttggtggga atccaggaaa atcttgctct 180
ccaaagatga gattagatga cttctggtca ctgaagttgt gtagacctc aaaagattat 240
ttactgaggc attgcaagta cctcataaga aaacacakgt ttgaagaaaa ggcccaagtg 300
gatcccccta gtgctctgaa atatttaca aatgatcttt atataactgt ggatcattca 360
gaccagaag agacaaaaga gtttcagctc ctggcatcag ctctattcaa atctgggtcn 420
agattttaca gctctgggct tttctgatgt ggatcacacc tatgctcaaa gaactcagct 480
ctttgacacc ttagtaaatt tctttcctga cagcatgact cctcctaaag gcaacctggg 540
agacctcatc acactgtaac tgaagagtca ctggacacag aaatggaaaa caggagtcga 600
ttttccgctcy ttggattgc agctccactg actgacagta aagctgcagt gattgaggac 660
tgcaccagag ttctgaaggg atcttaacca tcacaagttt ttaccctctt ccttcatgcc 720
tgacctcaac ccgctctcc tcactctatt cctaaattag gctaataaag tgaaattggg 780
atactttcc 789

<210> 28

<211> 847

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (799)

<223> n equals a,t,g, or c

<400> 28

ctgtcctgag ctgttcaactg cttggtggct accttctaag agtgagcatc tcacaagcag 60
gtgtcttaag tgaggaaatg gaagctgcag cttcttaaag ctgggaatta gtagagcatt 120
acctctgctg tattctatta gaggtcaagc catcactgaa gccatattca aggagaggga 180
tcgtgcacac tcttcgggaa atacgaccac atgagggctg ccatcctgga aaaaatgcct 240
cttgtggagc gagatggccc tcaggctgat gaggaagcaa aggaagcaa agaagcagcc 300
cagctttcag aagcagcccc agtgcacaca gagccccagg cctcacagct cctggatctg 360
ctagatctcc tggatggggc ttctggggat gtccagcatc ctccccatct ggacctctcc 420
ccaggaggtg ccctgttaca cctgcttgac cttccctgtg twcctccacc ccagctccc 480
atcccagatc tcaaagtgtt tgagcgtgar ggagtacagc tgaatctgtc ttctattcga 540
ccccctgaaa accctgcttt gctgttaatc accatcactg ccaccaactt ctgagagggt 600
gatgtcacc atttcatctg ccaggctgct gtgcccaaga gtctccagct gcagctgcag 660
gccccagtg ggaacacagt tccagctcgg ggtggccttc ctatcaccca gctcttcaga 720

atcctcaatc ctaacaaggc cccctcgcg ctaaagctgc gctcactacg accactttca 780
ccagtcggtg caggagatnt ttgaggtgaa caactgcctg tggaatcgtg gcagtaatgt 840
ctccaat 847

<210> 29

<211> 666

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<400> 29

gganccatgg gagtatgcac atgggatcat aatccattct gtgggttgga aaaagaaaat 60
gttaacctct gctttagagg gtagctacta gctttgttg ggataaaaagt gtaatacatg 120
cacttttgaa ctctgaaaagt ttgccaatct gaaaaggggt gtttctgaag accactatct 180
tttcaaca cttaaaaata agtggttgca gttgtgtatg ggcacgatac tgtattcttt 240
acatttttat ggccctacag ctacttctta tccctgcaag tatataaatt aaaaccaagt 300
cacttttagaa cagctttgaa actagagttt caaaggtaaa aggatctcat gtttctgaat 360
ctgcgtaaag caagatggct gtgatttgac aggtttaatt gctagktttt ataggtggat 420
agaaatgaat agtttgagct ctttaaaatg ttttaaaaaa tgtttgctta ctatctatat 480
atatgacatt attcccaatt agttttatat ctccaagata tatatatgta tataggata 540
tacacatatg tatatatata tagtctatat attctatata agaatatatt ccaataagaa 600
tatattccat acgggaatat attagtcatt gatgtatttt gccggtaaaa ttaaaagata 660
ttttaa 666

<210> 30

<211> 517

<212> DNA

<213> Homo sapiens

<400> 30

cgctcggtt cctgggtctg gctgctgccg cccgcccgtg tccgcccgtg tcgcccggg 60
gcaccaagga gccgttgag ggtccggcg gagggccgct cgtgtggaag tcgtcgacgc 120
cgccgctcgt ccgtctccc gtccgttctc gctcccggc gccatcatgc tggcgtcat 180
ctcccgcctg ctggactgg tccgttcgct cttctggaag gaagagatgg agctgacgct 240
cgtggggctg cagtactcgg gcaagaccac cttcgtcaat gtcacgcgt caggtaatt 300
cagtgaagat atgataacca cagtgggctt caacatgagg aaggtaacta aaggtaacgt 360
cacaataaag atctgggaca taggaggaca accccgattt cgaagcatgt gggagcggta 420
ttgcagagga gtcaatgcta ttgtttacat gatagatgct gcagatcgtg aaaagataga 480
agcttcccga aatgagctaa cataatcttc tagataa 517

<210> 31

<211> 2675

<212> DNA

<213> Homo sapiens

<400> 31

gcgaggtcac gtgacggagc gccggagcgg agggagccgg ggctgggagt tctcctgagg 60

gaagaggagt ggagtagggg ggacgcggcg gcggcggtga caatgagttt tcttgagggc 120
ttttttggtc caatttgtga gatcgatatt gttcttaatg atggggaaac caggaaaaatg 180
gcagaaatga aaactgaaga tggcaaatga gaaaaacact atctcttcta tgacggagaa 240
tccgtttcag gaaaggtaaa cctagccttt aagcaacctg gaaagaggct agaacaccaa 300
ggaattagaa ttgaatttgt aggtcaaatt gaacttttca atgacaagag taatactcat 360
gaatttgtaa acctagtga agaactagcc ttacctggag aactgactca gagcagaagt 420
tatgatattg aattttatgca agttgaaaag ccatatgaat cttacatcgg tgccaatgtc 480
cgcttgaggt attttcttaa agtgacaata gtgagaagac tgacagattt ggtaaaaagag 540
tatgatctta ttgttcacca gcttgccacc tatcctgatg ttaacaactc tattaagatg 600
gaagtgggca ttgaagattg tctacatata gaatttgaat ataataaatc aaagtatcat 660
ttaaaggatg tgattgttg aaaaatttac ttcttattag taagaataaa aatacaacat 720
atggagttac agctgatcaa aaaagagatc acaggaattg gacccagtac cacaacagaa 780
acagaaacaa tcgccaaata tgaaataatg gatggtgcac cagtaaaagg tgaatcaatt 840
ccaataaggc tatttttagc aggatatgac ccaactccaa caatgagaga tgtgaacaaa 900
aaattttcag taaggtaact tttgaattta gtgcttggtg atgaggaaga ccggagtagc 960
ttcaaacagc aggagataat tttatggaga aaagctcctg aaaaactgag gaaacagaga 1020
acaaactttc accagcgatt tgaatctcca gaatcacagg catctgccga acagcctgaa 1080
atgtgaactg aacaggagaa aaaaagaaaa gcmaaaaact cctgtwacc ttgarattaa 1140
gttcagcagg ttaaatgatg ttgcagctgg agggggcgga aaaaggccaa aactccatat 1200
atgttagtct tcctttatct tacagcgay wtttatttta tgatataatg aaatgttcgt 1260
tcatgtatat acatttttaa aagtgtttc tttgaaacac tggaaacttg ttaagctgcc 1320
ttttttttt taacttccta ctttgatgat aagcactcag atatatatca gcgtaaacat 1380
gaaaaatttt catgtgagta ggctgttatt tgtaattttg cttcttttct gcataatgtt 1440
gattataaat cctctctttt caggctaattg attacctctt attctctaca tgcaaaaaat 1500
taaatatttt gtgttcaaat aaaattagaa aacytgagtg gcctcttggtg tcctgcagag 1560
atttaaaaca tggcatctca atatttttga gaactacatt tgtttttaac atatgtgtt 1620
gagaaaagca tatggagtgt ttcaccgcag gcacttctga gtaccattcc atggcttcca 1680
gaattttatc ctcttgagg tcttctgtgc tatgaatatt agatttctt cccaaggga 1740
ttatgtggca ggtcattatg gccttctttt ttttggccat attagtaac agttttgcta 1800
tattccagta gtaccgttgt gtgttttctg caatggggag ttgacttagc ttggcatttt 1860
agatttgta aaactatttt ttccataaat actttgaaac gtatttataat atttcaattt 1920
aaggaatctt tttgccatgt gtatgcaaat attattttct tcatacatc attttctttt 1980
caggggaaaa ttttgggatg ggggactcag gaggacctgt gaagcatgta gttatctaga 2040
tctgggtaat tcatgttta ttaaaactga actttggcta gttaaactca tattgaaact 2100
tcatctagtc tcttaatttt ttaacactaa attcaagtca tttgttttaa gtctctaaaa 2160
aagaagattg cagtcattca ttcatatgca tggggtctga tcgcaaatc actaaatgtg 2220
gagtgtagga accaaaatga aacctgctgt atggaaacta ctttacttta tggttcattg 2280
gtttttgtac caatattttt tatgcacttc agtgcaagtc ttgtcagtta accttacttt 2340
atgagtaagc taaataaccc aaattacatt tctttaaacc tgttttacta ctatggcact 2400
ttgataaaat ggtcaggaac caactttact ggcaaaaggg tccatgtacc accatgtgct 2460
ggagcatctg ttctacatgt ggatatcwat gaatggtaat gttttccttc atgtaagtgc 2520
ctattcagag tttcagaatt ttaaaatgcc aaatattttc atggtcattt gcatgtagta 2580
agccagaaaa tattcaaaga gattttgaaa accaattgta ttaaccagc ctcaaatgtg 2640
gcaaccatga tgtataataa agaatttgaa acaga 2675

<210> 32

<211> 277

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (109)

<223> n equals a,t,g, or c

<400> 32

```
agccgcaccc ccagcacttc ggacatggcc gaggaaggca gagtggccag cgggggcccc 60
ccagggtcgg agacctcgga gtctctcagt gactcactct acgactcgnt gtcctcttgt 120
gggagtcagg gctgaggggc tgcgccacgc cacggccccg ctggagctgg ggaccacaga 180
ctggaccggc tctcttcattg cccagccccg ggagacgggg accccttccc tgaagggacc 240
aaggagggcag gtggataaga aggttgaaaa ggggggtt 277
```

<210> 33

<211> 921

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (839)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (846)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (886)

<223> n equals a,t,g, or c

<400> 33

```
cngtggcttt cctgcaccac tgccccccac caggatgatg gagagtaaga tgattgctgc 60
catcactcc agcagtgcag atgccaccag cagttcaaat tatcattcct ttgtcactgc 120
ttcatccacc tctgtggacg atgcattgcc ttaccactt cctgtcccac aacctaaaga 180
tgcttctcag aaaacagttt actcctcctt tgctaggccc gatgtcacca ctgaaccctt 240
tggtcagat aactgtttgc atttcaatat gactccaaac tgccagtacc gtccccagag 300
tgtacctccc catcacaata aattggagca gcaccaagtg tatggtgcca ggtcagagcc 360
accagcctcc atgggtcttc gttataaac atatgtggcc ccaggaagaa acgcatctgg 420
acaccactcc aagccatgca gccgggtcga gtatgtgtct tctttgagct cctctgtcag 480
gaatacctgt taccocgaag acattccacc gtaccctacc atccggagag tgcagtctct 540
ccatgctccg ccgtcttcca tgattcgctc tgttccatt tcacggacag aagttcccc 600
agatgatgag ccagcctact gcccaagacc tctgtaccaa tataagccat atcagtcctc 660
ccaggcccg ctagattatc atgtcactca gcttcagcct tactttgaga atggccgggt 720
ccactacagg tatagcccat attccagttc ttctagttcc tattacagtc cagatggggc 780
cctgtgtgat gtggatgcct atggacartc cagttgagac cctttcaacg gctttccant 840
```

cgagantttg ttttttacia tcctaggttg caaggaaaga gctttntaca gttatgctgg 900
gtttgggtcc aggtccccgg g 921

<210> 34

<211> 1467

<212> DNA

<213> Homo sapiens

<400> 34

aaaaaaaaa gaaaagtctt gtgttgatgt acagtttctc ctaagaagaa gcgagggtgg 60
tgaatttttg aagcacttct tgaatcggat taacccatgc tcttattgaa ttttttcac 120
tgctctgttt agtttgatat taaagcaaaa ttaagaggtc ttagtttttc ctatagaact 180
tttaatatgt caaaagctat attgtctaaa ttacagtact taagcaaaata ctgagtagtg 240
ttttaaatc agaaatagag cttctattat gaacacatga gaatgatttt tttctcttaa 300
tcattattaa ggaatatatt taatttcacg gtcataataa ggtgataagt aatacctgat 360
tgtttccttt tctgttctag taactcagag gagatacgtg ttttatttgt gatagcaaat 420
tcctaaatga acattaggca agtggatatca ttatcaggcc agctgcagcc tctgccttg 480
acctgcattc ctagaatttc tttgttgctg taattcttga ttaagtgacc ttgactttca 540
ttttgtaatt ttgctaatac tcagcaaat cacttgcatg acgttactgc caaatatgaa 600
ggcagttgaa ttattatgag tgattgtggc agaggtttgt gccatggtga aaactttgat 660
gtttgtctgt gttcattgga tccatctttt taaatgacat taccatgagt ctgttgtaa 720
acctaaatat cttgttttga atttaaaatg ggactctata ttgttgtagt tcaggctctc 780
attgactaag agattgagag aaatctgaca taagaaaata ttgttttcac tgcaggaata 840
aagaggaggt aacagtgaat ccaatatagt tcatattgtt attgtccaat catcaagtta 900
actaagcatt atcagattac gtttatttct catacatatg gatattaact taaggtaaaa 960
aagctggatg tgaaggatct gaaaaggcat taatttatgt actaattcta taaacatgta 1020
ttaataattg cagtattatt aaatacagat ggactcaatg tacctttgaa aagaccacta 1080
atttagaaaa caaagctaag tgcagtcatt acaagaagca aagaaatact taagttagaa 1140
aaaaattaaa atgaaggat ggtctaagtt ttcttcacg tggaacaaat gttaaagaag 1200
cagtgttgcc ttacaatgta tgtgataaaa taataccttt cacaatcaaa attttaatag 1260
taaatataag ataaaattta tattaataaa tgaaaacgta tttgtactga atttagtcac 1320
tagagaacat cgtaacaaaa tacatgaaac aaaagtagcc agaaatgtta gaacaggtgg 1380
aaatgtatac attatttgat ggtttgtttt tttatggaaa taaacaacat acatagaatt 1440
aaatggtgat caaaaacatg gaaaaaa 1467

<210> 35

<211> 2077

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (15)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (423)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (730)
<223> n equals a,t,g, or c

<400> 35
cggacggtgg gncgncgaca caatggggcca yggagttccc gttcgatgtg gacgcgctgt 60
tcccggagcg gatcacggtg ctggaccagc acctggaggc cccagcccg ccgacccgga 120
accacaacgc cggcccggtg tgatctacag cagcaaatta tgaccattat agatgaactg 180
ggcaaggctt ctgccaaggc ccagaatctt tccgctccta tccactagtgc atcaaggatg 240
cagagtaacc gccatgttgt ttatattctc aaagacagtt cagcccgacc ggctggaaaa 300
ggagccatta ttggtttcat caaagttaga tacaagaagc tctttgtact ggatgatcgt 360
gaggctcata atgaggtaga accactttgc atcctggact ttacatcca tgagtctgtg 420
cangccatgg ccatgggcca gaactcttcc agtatatgtt gcagaaggag cgagtggaaac 480
cgcaccaact ggcaattgac cgaccctcac agaagctgct gaaattcctg aataagcact 540
acaatctgga gaccacagtc ccacaggtga acaactttgt gatctttgaa ggcttctttg 600
cccatcaaca tcctccagca aggaagctgc cacccaagag agcagaggga gacatcaagc 660
catactctc tagtgaccga gaatttctga aggtagctgt ggagcctcct tggcccctaa 720
acagggcccn tcgccgcgc acacctccag cscacccacc ccccgctcc agcagcctgg 780
gaaactcacc agaacgaggt cccctccgcc cctttgtgcc agagcaggag ctgctgcgtt 840
ccttgcgct ctgcccccca caccctaccg ccgccttct gttggctgct gaccctgggg 900
gcagcccagc tcaacgctgt cgcaccagct cccttccccg ctctgaggag agtcgatact 960
aacagctacc ctctccctgc cctgggagac ctgggggtggg cagggaaacc ctccctgaga 1020
acctcagacc cactcttcca ttgcatcctg taggaccag tggaacctga cagagcccat 1080
aggattccct cttctacttt cttagacagc agggatgtca gggctctaaa ctgcctaaca 1140
ctttgtagct tttcttaaca caaaagcacc ccttctctcc taacttgggc tctgaatact 1200
ttcccaacag gaagtctgat ctgttgccag acttcttggt tagatggctc atacatttat 1260
ctagagaagc acactcttgc ttgctgtcaa acttttagamc accatggaag gtctaagggc 1320
atcctgtgcc agggaaactt tttaaggaat tttatctatg ggataaacc catattccct 1380
ctagtgtcta ctggtggctc taatactgct ttgtgctgcc tgccacactt gccctttgag 1440
cctgcgaatg cgcgctagt agcaagctct gcttcagagc agtctagtta ggtagaacag 1500
ggacttacca gcttcccaaa gggatctact caccattgcc aaactcttca tttccacatt 1560
ttgtgtaggt gtcagggaac cccaaactgg tgttgctttg gggctcttaa aggagattgg 1620
ctgacaccac catttcccc agatccagat tctctgaggg aggttgttc ttgagagtag 1680
atccagagtg tcaaggatct gttagatcct ggaatccctt cttgcatcca tccctccctg 1740
gtagctaggt ccgatatata tctgtcttgg tgagattgtc gagatgagat gggggaccac 1800
tcttctctg tcttctctc ctcctttcct ccatagcaag gacgacctc cctgctccat 1860
gccagagta tagctagatc ccttccctc cctaccctct gaatgtgtgc tagatcaggt 1920
gccccactgt gtttctgaa atccttgga gccggtctc ccatctccc ctactcactc 1980
ttcccttttc ttctctcagt gttgtctgaa taaagtgtga aatcttttgt gttttctaaa 2040
ttgacatttt caatgaaaaa aagaatcaca aaaaaaa 2077

<210> 36
<211> 384
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (366)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (370)
<223> n equals a,t,g, or c

<400> 36
ccagcaggag aatcctcccc tgccccttgg ctcaaaggcc ctggagccca cctcccagaa 60
gccccgtgtg ggggggggcc acgggggaga tcccaagctc agtccccaca aagttcaggg 120
ccggtcggag gcagggggcag gtccgggtcc aaagcaagga caccacagct cttccgactc 180
cagcagcagc tccagcgatt cggacacgga tgtraagtcc cagctgtgtg gctccaagca 240
gcacgagagc atcccgggca aggccaagaa gcccaaagtg aagaagaagg agaagggcaa 300
gaaggagaag ggcaagaaga aggaggctcc ccaactgaagg gcctggacaa ggctcattaa 360
acttctctctn tgccaaaaaa aaaa 384

<210> 37
<211> 468
<212> DNA
<213> Homo sapiens

<400> 37
gcgatgctgg gtgtggcagg cctcctgagc cggtctggagg aggacaggct gctgctgcta 60
ccgaaggagg atgkctgtcg ctgggccttg gctgtaggca gctgggctta ctgccggggc 120
ctgcatacac agcgccctcca gtgggagtga cagttggata cagccaggca gggtttctgc 180
cctgccgaac actttccctc ccacctgcct gctcctggcg ccttctccct aggggtagac 240
tcttctgcct actgaagtgg gtttgctgca cattgactgg tcaggggcag agtctgggtg 300
ctgtcctttg gccacgtgtg gggacttgtc tagaccagaa tgaaaggaca ggggtcccaga 360
cacgtttggg ggtcctgatt ctgggctgga cacggttgtg gatccagaga agaggcctag 420
tctccaataa atcttaggaa ttttgaggaa aaaaaaaaaa aaaaaaaaaa 468

<210> 38
<211> 1095
<212> DNA
<213> Homo sapiens

<400> 38
ggcagcagga taatgagcat aagcgttcac tgaccaagac tccagccaga aagtctgcac 60
atgtgaccgt gtctgggggc acccaaaaag gcgaggctgt gcttgggaca cacaaattaa 120
agaccatcac ggggaattct gctgctgtta ttacccatt caagttgaca actgaggcaa 180
cgcagactcc agtctccaat aagaaaccag tgtttgatct taaagcaagt ttgtctcgtc 240
ccctcaacta tgaaccacac aaaggaaagc taaaaccatg ggggcaatct aaagaaaata 300
attatctaaa tcaacatgtc aacagaatta acttctacaa gaaaacttac aaacaacccc 360
atctccagac aaaggaagag caacggaaga aacgcgagca agaacgaaa gagaagaaag 420
caaaggtttt ggggaatgcga aggggcctca ttttggtgta agattaataa ttttttaaca 480
tcttgtaaat attcctgtat tctcaacttt tttccttttg taaatttttt ttttttgctg 540
tcatccccac tttagtcacg agatcttttt ctgctaactg ttcatagtct gtgtagtgtc 600
catgggttct tcatgtgcta tgatctctga aaagacgtta tcaccttaa gctcaaattc 660

tttgggatgg tttttactta agtccattaa caattcaggt ttctaacgag acccatccta 720
aaattctgtt tctagatttt taatgtcaag ttcccaagtt cccctgctg gttctaatat 780
taacagaact gcagtcttct gctagccaat agcatttacc tgatggcagc tagttatgca 840
agcttcagga gaatttgaac aataacaaga atagggttaag ctgggataga aaggccacct 900
cttcactctc tatagaatat agtaaccttt atgaaacggg gccatatagt ttggttatga 960
catcaatatt ttacctaggt gaaattgttt aggcattatgt accttcgttc aaatatcctc 1020
atgtaattgc catctgtcac tcactatatt cacaaaaata aaactctaca actcatctaa 1080
catgcttact taaaa 1095

<210> 39

<211> 1757

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (596)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (647)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (648)

<223> n equals a,t,g, or c

<400> 39

gccagccacc attttatctc tctaaagtct ggtcccagta ttaaacctat tctttagtaa 60
actcatatta ctgttctaaa ttgaagaaat tatttattac tctgtacttc tagactcaaa 120
attctttatc aaagatagtc tcaaagaggt agtacaagtc ctgtttaact gcactttttc 180
acattcacag tgcttctctt gatattcttc cttacatcat tatacactgt tgatatcatt 240
ttactcttct tctctcttca catttcttaa attttggttc ttttctgta catgtgtttt 300
agcggggccc ttttctttga actttgtcta attagcctgt acatttttgt ttcttttaag 360
gtagaacaga tctttttttg tttctccttt taagtctact ggttttaaaa gaggtaaatg 420
tatccataga ccacagtgcc ttgtcttttc ctctgccagc acatggagca cgggattaga 480
tgcacaaacc tatttaggga actatttttg tagatgtttg agtttataca gaaattgcag 540
ctggtatttt attttgctgt acatttactc aacttgtcca ttagtattta actatntcca 600
gagtttggtt aggagtaaga attgacccat tcgttagttt accatanntt ttcctgggat 660
aaaaaggagc cagaaataag cttatttgct aaataattaa ttatgtaagc ccacctaggt 720
cctgcataag atccccctca catacttcac aatataatag tgtgtgtgtg tgtgtgtgtg 780
tgtgtgtgtg tgtgtgtgtg tgtgtatktg gctaaaaaat tatactgccca aaattactga 840
ttataaatac ttgactacac tgattgatgg gacaaaatga ttaaagtatt ttcagggatc 900
ttattccata tgcaccacc aaagatttct acagtgttat aaagtatata aatattccaa 960
atttctgtgg ttaaatattt ttttcttttt tttccttttt tagaataaca cagtctgtgc 1020
tttccaaaaa tgcttgaact tttatgttgt taagaaatat ataagtatat cttacattaa 1080
gcatgagtct aatttgattt aattgggatg gactaaattt tcatttgatt atcaggaaaa 1140
ttaaggagtt atatatttaa aagcaatttt ctgtgttttc tcttttgtaa gttgactcat 1200
ttgtgaagca attaggcaaa ttttgagaag atcattgtta ttgtggtttg cagtatatat 1260


```

ttcttagtaa atatcactta agattaaatt ttccagaaaag aaaattatag cttttttccc 1320
aaaatatttt taagatttaa tctttttgta gtatgctaca gatttaatta tattaactct 1380
tttttaagac attgaccatg acttaacatt ttgccttcta acacctttta aatctatgta 1440
ctttaatagt taagagaaaa taagtttgca gatttttaat aatctgtttg taaaaggcta 1500
tctctaagcc tagtatgtgg gtaattttac aggtgtgttt ttgataact ttaataataa 1560
ataaactcat tttatttggt gcaattcgcg ttcttttttt tatgccagag tacatatgtt 1620
ggattccatg aattggtatt acttattatt atgtgttgat taaatatatg cacacactta 1680
ggattacaga tcacagagca aattatgaaa atcataaaca ttctggtatg gtcattcata 1740
ggattatgaa aaagaaa 1757

```

<210> 40

<211> 1945

<212> DNA

<213> Homo sapiens

<400> 40

```

accgggagct ggtgacggat ggcggggccc ccagcccctg gcgctgcaac tgggaacagt 60
tggtgaaccc gcgaccaagc gaggcggacc ctgaagcgga ccccgaggaa gccactgctg 120
ccagggtgat tgacaggttt gatgaagggg aagatgggga aggtgatttc ctatgtagtg 180
gtagcattag aaaactggca tcagcctccc tcttgacac ggacaaaagg tattgcgcca 240
aaaccacctc tagaaaagca tggaatgaag accattggga gcagactctg ccaggwtcgt 300
ctgatgagga aatatctgat gaggaagggt ctggagatga agattcagag ggactgggtc 360
tggaggaata tgatgaggac gacctgggtg ctgctgagga acaggagtgt ggtgatcaca 420
gggagagcaa gaagagcaga agccactctg caaaaacacc gggcttcagt gtccagagta 480
tcagtgactt tgagaaatth accaaggga tggatgacct tgggagcagt gaggaggagg 540
aagacgaaga gagtggcatg gaagaagggg atgacgcgga agactcccaa ggcgagagtg 600
aggaagacag ggctggagat agaaacagtg aggatgatgg tgtggtgatg accttctcta 660
gtgtcaaatg ttctgaggaa gtggagaaaag gaagagccgt gaagaaccag atagcactgt 720
gggaccagct cttggaaggga aggatcaaac tacaataaagc tctgttgacc accaaccagc 780
ttctcaacc agatgttttc ccattgttca aggacaaaag tggcccagaa ttttccagt 840
ccctgaaaaa tagtcacaag gcacttaaaag cattgttgag gtcattggta ggtcttcagg 900
aagagttgct tttccagtac ccagacacta gatatctagt agatgggaca aagccaatg 960
cgggaagtga ggagatttct agtgaagatg atgagctggt agaagagaag aagcagcaac 1020
gargaagggt ccctgcaaaag aggaagctgg agatggagga ctatcccagc ttcattggca 1080
agcgctttgc cgactttaca gtctacagga accgcacact tcagaaatgg cacgataaga 1140
ccaaactggc ttctggaaaa ctggggaagg gttttggtgc ctttgaacgc tcaatcttga 1200
ctcagatcga ccataattctg atggacaaaag agagattact tcgaaggaca cagaccaagc 1260
gctcygtcta tcgagttctt ggcaaacctg agccagcagc tcagcctgtc ccagagagtt 1320
tgccagggga accggagatc ctctctcaag cccctgctaa tgctcatctg aaggacttgg 1380
atgaagaaat ctttgatgat gatgactttt accaccagct ccttcgagaa ctcatagaac 1440
ggaagaccag ctctctggat cccaacgatc aggtggccat ggggaaggcag tggcttgcaa 1500
tccagaagtt acgaagcaaa atccacaaa aagtagatag gaaagccagc aaaggcagga 1560
aacttcggtt tcatgtcctt agcaagctac tgagtttcat ggcacctatt gaccatacta 1620
caatgaatga tgatgccagg acagaactgt accgctctct ttttggccag ctccaccctc 1680
ccgacgaagg ccacggggat tgacatcgcc cactccgac acccagtggt cgccttggt 1740
ggtgcggtg ctggtccaga tggaggaaac cagtgacttt atggggctga gctagtaggg 1800
aagcccctgg aaagatgctg cgttccgaac ctgtgcctaa tacacgcaag ggcgctgtcc 1860
cgcccaacc cgcttttaaa cgccacaaat aaagagcatt gttaccgcca agtacgagc 1920
ggccgcgaat taccggaccg gtaac 1945

```

<210> 41

<211> 588

<212> DNA

<213> Homo sapiens

<400> 41

```
aggcgtatac caccatgact gaaaacaaaa gacttttttt tgagactccc tctcaaaaac 60
aaaacaaaaac aaaaaaatta gacaaatgct acattaatgt ttgggtgggc agattctact 120
ttgaatctga agtttgacaga tatgcctata gatttttgga gtttaccact ttcttattct 180
gtatcattaa tgtaatatTT taaattacta tatatgttac catttttctg gatttagtaa 240
gaaatttgca gttttgggtt gatgtaacaa gggttttaat gtaatttatg ttagattttg 300
catttttttc attactgtta tattttaacc tgactgactg atctaattgt attagtattg 360
tgaataatca tgtgaaatgt tttgagacag agtactatat ttgtgaatat aattttatgg 420
tttttttcac ttagaacctt tctgtgtgga aaactaagaa aattgctttc tgctgtataa 480
tctggcattc attgtagatt aaagcttatt tttctgtgaa taaaacgtat tcaataaaaat 540
actattctttt aaaattawaw mawaaaaaaa aaaaaaaaaa aaaaaaaaaa 588
```

<210> 42

<211> 1568

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (104)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (112)

<223> n equals a,t,g, or c

<400> 42

```
cattagattt cactttttta aaaaacccag tggacattgc tataaataag atttatttgg 60
ctacaaataa cctgggatgt tgcttattat gattgatgcc tgcnggtttg tncccaagct 120
gagtgaattt gaacctcgtc ctccctactc attttgatga ctgaggtctg tttataagaa 180
aagggaagttt ggagaagaaa accgagatta gaaaatatca tgttttggtt ggagataaga 240
accagggatg gcaagtacca gtgtgtacaa atgtatttca cgaggtttga aggaacgcat 300
aatcaagagg gaaaacaatt tgtccttcatt tggacgtatt atttggattt gggtagcaaa 360
caaaatggaa tgtggtctgt taggagcatt ctgtttgttc ttttgcctct gatgtgatga 420
atcattgccca catgctagat ggactcttca tatccagggt ttgtccctca gggctgagca 480
ctgtattaaa gagtttttgt tgagtcattt aaccttagtg tccacatcca gatcagctgt 540
aaaatgggga agacgtgtgc tgatttgga tgaatgcaaa atatcactat cattttccta 600
attacagagg agcaaagggtt atcttcagcc ctttcagttc tatgctcaca tattcaaata 660
tcaaatgtaa tttagctgaa gttatttaaa atcaagctct ttcaatatct gttcaaagaa 720
aaagaacaca ctttgaaaat tctgcaaagc tgtctcccag tctttaaaat gtctggaagc 780
actctccttc tttacaatac caacatcact ggcccagaat ctccctgtg ctaggttgta 840
aatataaata aattacttgt tttgtaaaat ttgttaaaga atattttggt agaaataact 900
caaacatatt ctttgggtta tatttatata tatgtgaaat aaatatacta tcaaaagggt 960
atattttata caaaaagtaa attgctacct ttgtatgct aatatgcaaa gttttgtata 1020
atatgatggt ttatttttag ctctacactt aaaccatagg tgggtgagtg ggaacttttg 1080
aaaactatca agaggcttgt tagacaaatt tatattctga aacctcaata agaaagcatt 1140
```

```
ccaggtttca atccttggtt tttgtcctgc tcccaaattc ttttttaaac ccatagttct 1200
tgtgtcttat ttgattcttc tgcgtgacac attgtattgg tccttggtgc atgtagtcta 1260
ctgtgtgttt tccgatttta taaggcagca tttctccata caaaaaraaa aaaaatgatg 1320
tacatataaa cgcttttrtt gtatggctcc tccatgttac tgtatatatc tgccagcact 1380
tcccagttac actcctgtga gtcagcttat tttacccta acataaatag tatgttttgt 1440
agtagttatc aaatttaaga gataaagcaa tcagaatgtt tggattttct tctatcttaa 1500
tgtgaatttc ataattaatg tctatttatt cagctattca ttaaaatata ggattctttg 1560
gggaaaaa 1568
```

<210> 43

<211> 1060

<212> DNA

<213> Homo sapiens

<400> 43

```
gcttgtcatg agaaggtggt aaatatccaa aaagaccccg gtgaatctct cggcatgacc 60
gtcgcagggg gagcatcaca tagaraatgg gatttgccta tctatgtcat cagtgttgag 120
cccggaggag tcataagcag agatggaaga ataaaaacag gtgacatttt gttgaatgtg 180
gatggggtcg aactgacaga ggtcagcccg agtgaggcag tggcattatt gaaaagaaca 240
tcacctcga tagtactcaa agctttgga gtcaaagagt atgagcccca ggaagmctgc 300
agcagcccag cagccctgga ctccaaccac aacatggccc caccagtgga ctggtcccca 360
tcctgggtca tgtggctgga attaccacgg tgcttgtata actgtaaaaga tattgtatta 420
cgaagaaaca cagctggaag tctgggcttc tgcattgtag gaggttatga agaatacaat 480
ggaacaaac cttttttcat caaatccatt gttgaaggaa caccagcata caatgatgga 540
agaattagat gtggtgatat tcttcttgct gtcaatggta gaagtacatc aggaatgata 600
catgcttgct tggcaagact gctgaaagaa cttaaaggaa gaattactct aactattgtt 660
tcttggcctg gcactttttt atagaatcaa tgatgggtca gaggaaaaca gaaaaatcac 720
aaataggcta agaagttgaa acactatatt tatctgtcga gtttttatat ttaaagaaag 780
aatacattgt aaaaatgtca ggaaaagtat gatcatctaa tgaaagccag ttacacctca 840
gaaaatatga ttccaaaaaa attaaaacta ctagtttttt ttcagtgtgg aggatttctc 900
attactctac aacattgttt atattttttc tattcaataa aaagccctaa aacaacaaaa 960
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa gggcggccgc tctagaggat 1020
ccaagcttac gtacgcgtgc atgcgacgtc atagctcttc 1060
```

<210> 44

<211> 1344

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (144)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (163)

<223> n equals a,t,g, or c

<400> 44

```
cccacgcgtc cggggccacc agggcctcct ggccccctg ggccccagc cctgttggg 60
```

```

ccaccccatg cccggatctc ccagcatgga gaccatttgc tgtccaacac ctactgag 120
accaacaacc actggcccca gggnaaccac tgggcctcca ggncytccag ggcccatggg 180
tccccctggg cctcctggcc ccacaggtgt ccctgggagt cctggtcaca taggaccccc 240
aggcccccact ggacccaaaag gaatctcttg ccaccagga gagaaggcg agagaggact 300
gcgtggggag cctggccccc aaggtctctg tggggcagcg gggggaactg gccctaaggg 360
agaccctggg gagaagagcc actgggtctc tagcttacag agcttcctgc agcagcaggc 420
tcagctggag ctcttgcca gamgggtcam cctcctgga gccatcatct ggccagaacc 480
agagctgggg tctggggcg gccctgccc caccagcacc cccagcctcc ttcggggcaa 540
gaggggcgga catgcaacca actaccggat cgtggccccc aggagccggg acgagagagg 600
ctgagggttg tggcgccccc tgaggcagac caggccaggc tccccctcct acctggactc 660
ggccagctgc ctccagggac cggcctgcca tatttattaa tgcctcagg gtcccttctg 720
ccatctaggc cttaggggta agcaggtctc agtctggca ccatgcacat gtctgaggct 780
gagcaagggc tgagaggaga ggcttggcc tcagtttccc tctgtgaagt ggggggaggc 840
aggccttcaa ggaggatag aggtacaagg cttcgtctca tctgtgtct gagcatccag 900
gccccaaagg actgaggag tcaggagctg gggctcggca catgcagaga tgacagggca 960
gggggcagtc ttcctcccc tccccacca aacctcggg agccctcctg tgccctccc 1020
tccttgttgt ccagtgtctg gytccccacc ccgaggtcag gctgccaat cctctgactg 1080
gatcaccggg ggcttcttg ctcagttctt ccctctgagc cccagggccc tcccgcactc 1140
caggttgggg atggggacat ggagaggaag gggccgcta ctctgcaaa tgctgtgac 1200
agatgccagg aggtagatgt gtgctggcca ataaaggccc ctacctgatt ccccgcaaaa 1260
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa gcgccgctc tagaggatcc aagcttacgt 1320
acgcgtgcaa cgcggtcat agct 1344

```

<210> 45

<211> 892

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (890)

<223> n equals a,t,g, or c

<400> 45

```

ttgagaagtt ggatgaatat atatatagac acttcttttg tcacactttt tccccccat 60
atggaccag tcgacctgat aaaaagcaac gtatggtaaa tattgaaaac tccaggcatc 120
gaaaacaaga gcagaagcac cttcagccac agccttataa aagggaagg aaatggcata 180
aatatggctg cactaatgga agacaaatgg caaatcttga aatagaattg gggcaattac 240
cttttgatcc tcaatactga ttcacaattg agttaaatta gacaactgta agagaaaaat 300
ttatgctttg tataatgttt ggtattgaaa ctaatgaaat taccaagatg acaatgtctt 360
ttcttttggt tctaagtatc agtttgataa ctttatatta ttcctcagaa gcattagtta 420
aaagtctact aacctgcatt ttcctgtagt ttagcttcgt tgaatttttt ttgacactgg 480
aatgttcaa ctgtagtttt attaggaag ccaggcatgc aacagatttt gtgcatgaaa 540
tgagacttcc tttcagtgtg agagcttaaa gcaagctcag tcatacatga caaagtgtaa 600
ttaacactga tgtttgtgtt aaatttgag cagagcttga gaaaagtaca ttgttctgga 660
atttcatcat taacatttta taatcttaca ctacttctt gtctttttgt gggttcaaga 720
gccctctgac ttgtgaagaa ttgctgccc tcttaagagc ttgctgactt gttttctgtg 780
gaaatttttt gcacatctga atatcgtgga agaaacaata aaactacacc atgaggaaaa 840
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaccccg ggggggccc ga 892

```

<210> 46

<211> 496
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (496)
<223> n equals a,t,g, or c

<400> 46
aattcggcag agtcggagtg tggactttct cctagttgca gtcaggcttc atacgtatt 60
gtcctgcccg taagttcccg ttttgtgtgt ggtagagca gccagcgggt acagaatgga 120
ttttggaaga gggagtcacc actggacctc caaggaagcc acgtgcagac atctacaacc 180
ttcgatctcc tgacgagttt attgttgcc aaaaccaggc tttgattgaa ccaggatgaa 240
tgcgggtgtt ggaagtagaa tatatatata catataaaat tgaaactggc gatggaatat 300
gagaggagcc ctctggaaag aaaaggacag accctgtgct ttcataaaag tgaagatctg 360
gctgaaccag ttccacaagg ttactgtata catagcctga gtttaaaagg ctgtgcccac 420
ttcaagaatg tcattgktag actttgaaat ttctaactgc ctacctgcat aaagaaaata 480
aaatctttta aatcan 496

<210> 47
<211> 1229
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (764)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1165)
<223> n equals a,t,g, or c

<400> 47
attcggaaacg agagctgata tggaagggtt atttctttct ggccaatact ttttggtatt 60
tctaaatatt gcaatcttga tttttactat taaatttggt aattgtcagt tctggctttt 120
ttgcataaag agttggtcca ttaacttgcc aatttgaagc ttctaactag atattcccta 180
ctgaaagttt tggatttggt tttagtttgt ggagcagtct tagctgggga caggtaattg 240
acaacggcag agatactttc ttttcctagg attctaagtc tgtaatccac atcctcaatg 300
tattcacagg actttaaaaa tctctccaaa tgaggragg aaatatcctg gtgctttcta 360
atgggttact aaaagttgtg tttagaacaa cagattttta taggcatctt cctttgttat 420
gtgtcattag ccttttgccc gtttacctta gggctctttg aaggagaaat ggatgggaga 480
aaacctgtca cttggcgaaa gtaaaaggga taattaactg gctcagagct tatgtgcaga 540
gttccaagcc ccaaagttaa tctagaacca ctcgataaca ccaataaaaa tatttatttc 600
acatctgtta tatatctgga aaatgktcta agcatcttac acatatttct cattaaatcc 660
acaggtgacc attgtgaggt agawattttg ktctaawttt ccagatgagg aagctgagac 720
cctaaaaggt taggtgacag gttatacaac ttggagtgtg ggangaggag agaggaacct 780
gaacagggca agttggggat ctgacttttg tttgggtaga tgtaagcaca ttgtattttt 840
ggcttagatg ctttattcat catggctgaa ggaataacca ttactcact caccgaaaat 900

```

tgtttacaat aatctagatg aatttgctgt ctttggacat ctgtcttttg actggacccc 960
agtatatagt ctgtggaagc tcacttaagg agaragctcc tttttgtttg gtttagagaaa 1020
ttttctgtcc taaaagtaga aatagcccct tctaggtaag gatggagcat ttgatcatac 1080
tggtttcatt atattcctct aacagggttg aaccgattgt ttttgagtac ttgtttcaaa 1140
cttctgagta ttttccttct ggaanatagc tcagtgtttt aaaatttaca tgaacttaaa 1200
aggttaattt ttttttaaaa gaatggtta 1229

```

<210> 48

<211> 1411

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1410)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1411)

<223> n equals a,t,g, or c

<400> 48

```

catcctggct ccttgagtgg gagtratata acctctatcc taactgtcct tgtccctcct 60
ctccccacag ctggattatc accgaggcct cttgggtggac cgccctctg aaactaagac 120
agaggagcag ggaataccac ggccctgca cccccaccc ccaccccg tccagccacc 180
acagcacccc cgggcagagc agcgggagca ggaacgggca gtgcgggaac agtgggcaga 240
acgggaacgg gaaatggagc ggcgggagcg gactcgatca gagcgtgaat gggatcggga 300
caaagtgcga gaagggcccc gttcccgatc aaggtcccg gaccgccgcc gcaaggaacg 360
tgcgaagtct aaagaaaaga agagtgagaa gaaagagaaa gccaggagg aaccacctgc 420
caagctgctg gatgaccttt tccgaaagac caaggcagct ccctgcatct attggctccc 480
actgactgac agccagatcg ttcagaaaga ggcagagcgg gccgaacggg ccaaggagcg 540
ggagaagcgg cgaaaggagc aagaagaaga agagcaaaag gagcgggaga aggaagccga 600
gcgggaacgg aaccgacagc tggagcgaga gaaacgtcgg gagcacagtc gggagaggga 660
caggagagaga gagagagaaa gggagcggga caggggggac cgagatcggg atagggaaaag 720
ggaccgagaa cgaggcaggg aaagggatcg cagggaacac aagcgcaca gcagaagccg 780
gagtcggagc acacctgtgc gggaccgggg tgggcgccgc tagctgggaa aacactagag 840
ctgcaggtag cagccactcg gccccagggg gttatggcca cagagggata ggcacagtct 900
ccaccacctt ggagccaagg gtctttcaca tcacctatcc ctacatacat accaaatgga 960
aaagtggcca tccttttccc cccaaacaca ccccttaac ctatctcttg ggacttagcc 1020
cgaccctccc tctcatttcc cattaagtct gagaggcaag agctaggtta ggcaaggagg 1080
tggttggcca gagatgggga acagccaggt gccccagtc tctgattttt cctccatcct 1140
gcttaccacc tccttggtga cttacagcct tctcttggga acagccgggg ccaggactgg 1200
gtcacctatg agctgaatca gcctctcctc ctgagtccca gggccctgc agttcccagt 1260
ctcttctgtc ctgcagccct tgcctctttc ccacaggttc cactttatat ccaccttttc 1320
cttttgttca atttttattt ttattttttt tattattaaa tgatgtgggc tatggaaaaa 1380
aaaataaaaa tctgacttag ttttaaaaan n 1411

```

<210> 49

<211> 1685

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (344)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1606)

<223> n equals a,t,g, or c

<400> 49

```
cgctnttccc cccacacccc gtgtggccag ggatccccgc atggcccac ttagaaactc 60
aactatattg tggatgctaa acacttcact tcaggcaatc ccaaggcatt tgctccaggg 120
tatccgatga gattacagct gttaagcttg ctttccattt cataacttgc tgtgcagcta 180
gttaccaccc ccatgctgaa gagtaaagca aagtgccgtg gttcggcagt ggaatccacc 240
cccagcactc tgctcgact ggagcgttca agtccggtta tgtgagaaca gactaggact 300
ctcttgctgc ctctaattgc atttcactgt caccctcccc agtnttctga tgggtgtgcat 360
gtgaggagaa gatgagggtta ggactgagaa gtgcagaagt tggaacagtg gtaaggctgt 420
tttaaaataa gatgttttgt tttaataata tgctcctggc acaaagctag gagtaaagt 480
gactccaaag ggagttcagt taatctctga aatgcacaaa acctagctat tttctccctc 540
tcacacagct ctgagctctg tccattgcta cccaattct ctggggacat aaaaccaggc 600
tggaaggga ccaggaaagt tgaaatagt acatatcatc cactagtccc aagggctaag 660
gaatagttag tttattctgg aaggaaactg gaagcttagt ctaattagt cctggggatg 720
acctatgcaa tcacaccgct tatgaccatc cttagagagg ccctgagcac cagcttgatc 780
ttagggattt ccaaagtaac ctgctttttg cctggatagg gttaaaatag acctttcttg 840
cctatccttg ccttaacctt tctgcctgag gttggcctga gattgtgagt caacgacttt 900
gctatctttt cctcagtgtt gaactttcat taagaaataa agtcctagct tcttacagag 960
aggggtccaa atggtgaatg ctcacctcgc ctgggattca aggaattagc tcagagrttg 1020
gccctagct tttctgcctt ttagggggac agcaaaagg gaaaatttgc tgcagaaaaa 1080
tccaaaagat tgctgtagct ctcacaggga agtggtaaag atcagctaaa cctggggttg 1140
ggtgctttct gccagtgagg tcttggcata agtagattaa tctgctctt ttaagaaaag 1200
gcaacttatt caggcagtct ggaaaggggg ttctcagaaa actcagtttc tttattcctt 1260
cttttctccc aactactgtt actggttata gaggtctttg gactctaaag accaatgttt 1320
ggccactaac tggactaata tgtatctttc tgtgatttca tcatagaggt ctgttttgtg 1380
agggtttggt gtgcagaaaa ctttgattaa atcttaatgg gaggttggtt gacctggatt 1440
atctacagtg agcagactta aatggaacag aagtttatgt gtccaaatga tggaatcatt 1500
aaacctgagt gacttgacct gtgtggttcc ttaatagtat ctatatatct agacaaaaat 1560
agattgtgaa rgtaaagtgt gaatgaaaag gatggaaata atgttntcat atgttaatcc 1620
atgagcttga atccaggag gaatacctcg gtgctttaac caccttagtt ataacacatt 1680
tctta
```

1685

<210> 50

<211> 660

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (515)

<223> n equals a,t,g, or c

<400> 50

```
cggcacgcgt gggcctactt tcacgcttcc tccctccccc ctctccctt atcccttcgc 60
tttcgctctt ttccgtcgag gccgaccctt gagttgtgag tctggggtct ggttggtgaa 120
aaagagccct tgaagctgga agacgggaga ggacaaaagc atgtcttccc ttcctgggtg 180
cattggtttg gatgcagcaa cagctacagt ggagtctgaa gagattgcag agctgcaaca 240
ggcagtgggt gaggaactgg gtatctctat ggaggaactt cggcatttca tcgatgagga 300
actggagaag atggattgtg tacagcaacg caagaagcag ctagcagagt tagagacatg 360
ggtaatacag aaagaatctg aggtggctca cgttgaccaa ctctttgatg atgcatccag 420
ggcagtgact aattgtgagt ctttggtgaa ggacttctac tcgaagctgg gactacaata 480
ccgggacagt agctctgagg acgaatcttc ccggnctaca gaaataattg rgattcctga 540
tgaagatgat gatgtcctca gtattgattc aggtgatgct gggagcagaa ctccaaaaga 600
ccagaagctc cgtgaagcta tggctgcctt aagaaagtca gctcaagatg ttcagaagtt 660
```

<210> 51

<211> 1572

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1555)

<223> n equals a,t,g, or c

<400> 51

```
tnagtgaaag aatgtgccat attacatatt gcaacctaatt ttgttaaaac taactccagc 60
actaaagctg aaatgccaca aacactaaaa gtataaatat gtctgatttt tgaaacacat 120
aagctttgct ctttaggcag gaatgatctt ttcaaatcat tagcacaata tttaaatatc 180
taaaaattta agagatccat actttctgta gctttacaat taatttaagt actaaaaaga 240
caaggatttc ttttaagaaa ttatagcat ttactgtgtt atttaaatgc taagccaaag 300
tatctgcact taggtatacc tctttatgcc aataatgatt ttaatgaagg ctcttttcag 360
atgtaacctt atgaaggaaa tatctgcttt gtgtatatgc cagttagaat actggtttct 420
aaagtctgtc aaattgtatt tcagtggcac aaaaaccagt tttgaggctc tagacttata 480
attctttgaa taaaactgat aacttatatt tataattgga gtggagacct acctccataa 540
ttagataaac tctttttgga ttataatcag aattttgcct tttttcttct caaattatta 600
catatgtatg tattatatat ccacatatat agttttccct gattaaatgg atattaaaa 660
aattgcgggt gcttcaggac tttttgcttc tatatttaag tatattgttt ttatagcaag 720
aacatattct gaatgtttta taaatcttta ataatttata ttaggtaat atttttgtat 780
cacaatgcat ttttttttct ctcttttcc tccaaactat accactgtat ttaccacttc 840
taagagtgac tgacgacggg ccagatgacc cttgaagtag tcattatgta gcaataaatg 900
```



```

aagcctgaaa cagggttttt tacttccact ttaatcctta gaaatttctt ggcaacttcg 960
catattttca ttgacaccgg tgtataagta taaattttaa tgaactaatt acttttgcg 1020
attttaaaatt ctttatatgg tagttatttt ttataacagg atattaacat aagttaaadc 1080
ctatgtattt gaaatttgta cagagctttc ctctttactt caaacagcaa aaaagtgggg 1140
ggcatattgt agtcctgtca tttaagttat gtwaaaaatt taatcattat tttgatgctt 1200
taaacattct catgtgtaat atatgttttt gtatcaaaaa cactcatata tttcaagaaa 1260
aagaaattat gttaaatagc cctgttttaa gaaaaatatt tatgaagcat ctcaacttga 1320
agatcaagtc aaagttataa ctcaggatct gargtctcaa gctaggagag actgagaatt 1380
ttaatcagtt tgggcatata rtttgactg aatcacatct gtagtactta gccaaagaca 1440
atttgarga raatatcagc cttctggaar tagtacttc ctgaacaatg taaagtgtcg 1500
caratattca ataaaatggc aacctgttaa aaaaaaaaaa aaaaaaaaaa acccnaaggg 1560
ggggggggcc cg 1572

```

<210> 52

<211> 635

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (632)

<223> n equals a,t,g, or c

<400> 52

```

gctgctccag ctgttcgaag gtgatccaga cgcaagatgg ctgtcctctc taaggaatat 60
ggttttgtgc ttctaactgg tgctgccagc ttataaatgg tggcccacct agccatcaat 120
gtttccaagg cccgcaagaa gtacaaagtg gagtatccta tcatgtacag cacggaccct 180
gaaaatgggc acatcttcaa ctgcattcag cgagcccacc agaacacgtt ggaagtgtat 240
cctcccttct tattttttct agctgttgga ggtgtttacc acccgcgat agcttctggc 300
ctgggcttgg cctggattgt tggacgagtt ctttatgctt atggctatta cacgggagaa 360
cccagcaagc gtagtcgagg agccctgggg tccatcgccc tcctgggctt ggtgggcaca 420
actgtgtgct ctgctttcca gcattcttgg tgggttaaaa gtggcttggg cagtggaccc 480
aaatgctgcc attaaagaat tataggggtt taaaaactct cattcatttt aaatgactta 540
cctttatttc cakttacatt ttttttctaa atataataaa aacttacctg gcacagccct 600
catacctaaa aaaaaaaaaa aaaaaaaaaa tnggg 635

```

<210> 53

<211> 1367

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (106)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (958)

<223> n equals a,t,g, or c

<400> 53

```

ggccacggtg ttatctcttt gatcaagttg ttttacgtgt cacagtagag tctgaccagt 60
taccataacc aaacattttg tgcgtgtgct attggtgctg cgatantctc ctttgggtgg 120
aacacagtga agatcgacat gagtgcagcc cggagagatc ctcttccaat tgttccattt 180
ggattagctg catttgctac cactttgttt gccttgggat tagctttagg aacaaccata 240
gctgttgga tggtgttttt tatccagatg aaaataattc tcagaaacaa aacttctatt 300
gagtcattga ttgaagagaa ggctaaagat cgaattcagt attatcaact agatgaagtc 360
tttgtttttc catatgatat gggagtaga tggaggaaact ttaaacagggt atttacgtgg 420
tcagggggtcc ctgaaggaga tggacttgag tggccagtaa gagaaggctg tcaccaatac 480
agcttaacaa tagaacagtt gaaacaaaaa gcagataaga gagtcagaag tgttcgctat 540
aaagtaatag aagattatag tgggtgcctgc tgccctctga ataaaggaat caaaccttc 600
ttcacaaagtc cctgcaccga agagcctcga atacagctgc aaaaagggga attcatttta 660
gccacaagag gtttacgata ctggttatat ggagacaaaa ttcttgatga ttccctttata 720
gaagggtgtt caagaataag ggggttggttc cctagaaaaat gtgtggaaaa gtgtccctgt 780
gatgctgaaa cagatcaagc cccagagggg gagaagaaaa atagatagct gctgttaaaa 840
caaaattatc ctttaagtct gcttaattac ttgaaaattg tacatattac taaagaatta 900
tgcaatgagc ctactctggt taagatgttc ttttcctcaa aggtgcccta gtgccatnga 960
tttaaatatt tttattacca ttttgaaatg gagaagccat tctgcatatg cctttgaatt 1020
cctgccctc tttaccacct cttcctcccc ctcaaaggaa aaacatttca tccaagtaag 1080
ttaacggcat tttctgtagg attttcctta tgcaactgcac actctggacc tcacctgcag 1140
atacagttcc ccccttgcca ggagcatctg catgtggtac ttctcttttc cctcagttga 1200
tatttcttat atgatattct agatactata gaactcaatt tgtcagattc agtataacct 1260
cagattttgt tacctgtctt ttaaaaatgc agattttgtc aaatcaaata aagatcaatg 1320
gatgttggtg ataawmaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaa 1367

```

<210> 54

<211> 378

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (14)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (29)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (363)

<223> n equals a,t,g, or c

<400> 54

```

ggcagtggtg gggngtgaga accctgggnc ctgctgaaaa gcctcctctg taggaggtca 60
cccagcagga caragaagca ggaggaggac agggccacar aggaagccaa gaacggtgaa 120
aaggccaggc ggwggtcara ggagggtgac ggccagcacc cggcccaaga ggaggccccg 180
gaatcgcccc agacctctgg cccagagcag aaaataggtg tgggagcccc agggaggaaa 240
agccarytgg agaggaagca gagwtggaag aggtacagag ggtgaagggg gagaggggtgc 300

```

aaaatgaaga ggtgggacct gaacatgaca gccagaaaac aaagaagctt gaggagggag 360
ctncagtga ggcgaccc 378

<210> 55

<211> 1058

<212> DNA

<213> Homo sapiens

<400> 55

tcgggtatga ggctgggact aagccaaggg attcaggtgt ggtgccggtg ggaactgagg 60
aagcgcccaa gcttttttgt tctgaactcc cactgcgttg tggattcctg aggatgggat 120
gactgtatct tgattaccg gagttttcaa gatggcagca tctatgcatg gtcmgcccag 180
tccttctcta gaagatgcaa aactcagaag accaatgggc atagaaatca tagaaaaaa 240
ttttgactat cttagaaaag aaatgacaca aaatatatat caaatggcga catttggaac 300
aacagctggg ttctctggaa tattctcaaa ctctctgttc agacgtgct tcaagggtta 360
acatgatgct ttgaagacat atgcatcatt ggctacactt ccatttttgt ctactgttgt 420
tactgacaag ctttttgtta ttgatgcttt gtattcagat aatataagca aggaaaactg 480
tgttttcaga agctcactga ttggcatagt ttgtggtgtt ttctatccca gttctttggc 540
ttttactaaa aatggacgcc tggcaaccaa gtatcatacc gttccactgc caccaaaagg 600
aagggtttta atccattgga tgacgctttg tcaaacacaa atgaaattaa tggcgattcc 660
tctagtcttt cagattatgt ttggaatatt aaatggtcta taccattatg cagtatttga 720
agagacactt gagaaaaacta tacatgaaga gtaacaaaaa aaatgaatgg ttgctaactt 780
agcaaaatga agtttctata aagaggactc aggcattgct gaaagagtta aaagtaactg 840
tgaacaaata atttgttctg tgccttttgc ctggtatata gcaaatactc aaaaagtatt 900
caataattca atcaataaat ataagtttca tcttacacgt aagatacagg tcttatctcc 960
tgatggtgtg tccattttgc ctggtatata acagataata aatatccagt gtcaataaat 1020
gtaacaataa aagtttcatc ttctctcttt gtatgtgg 1058

<210> 56

<211> 682

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (667)

<223> n equals a,t,g, or c

<400> 56

gggnccggaa catattccct tactcaaaag attgcatgac tgaatttgct taaggaaaaa 60
aaaaattgta tcaagtccat taatacaatt atacattaat tatattacat taatacaata 120
tatggtttgt gaattcagag acattaccag tttgcctoct tctctcaata gaacttgtat 180
tttcattttc ttggttaagc agttgtctcc taatattatc ccatatgcta cctagtttgc 240
tggtcccaag cagtttactg tacttcaacta gatttggtac ctgctctccc ctggacttct 300
ttttcaatat tctagccttt cctagatgta aatctttacc tccttggttag tgaaattaga 360
tataagccat gatttgagga ggaagaaat ctggaatact taatttcatt taattatcta 420

tgctgatgaa tgcctgtatc attgttaata aaggagaatt gaaaatactc atttctactt 480
tctgccctca aatttctgtt tctatctcaa ctaggcaaga atcagcaggg tgcattgaygc 540
cattttaagc tgcttcacat cagactgaaa tcctaattac agttcataag tgaaacagac 600
taattcmatg ggcaatacct ttkgtawagg tccygtgctt aaaggaggca agtataaatt 660
ttcccantaa ggaatccccg gt 682

<210> 57

<211> 644

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (27)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (458)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (619)

<223> n equals a,t,g, or c

<400> 57

ggagncctggg cctgtgcggc ggccgcngta gcgctttgga aggcgcacgg ggcgaagatg 60
gcggcgagacg acaggaggcg ctgaggaggt tcgtggcggt gacgggcgcc gaggaggacc 120
gggcccgcct ctttctcgag tcggccggct gggacttgca gatcgcgcta gagcttttta 180
tgaggacgga ggggatgaag acattgtgac ctttcgcag gcaacccccca gttcagtgtc 240
cagaggcaca gccccagtg ataataagatg gacatccttc agagacctca ttcattgacca 300
agatgaagat gaggaggaaag aggaaggcca gaggtgagtc ttctagaggg ggtcaggggg 360
acagttcaca gggaaagtcca gggtaattgtg taaatcacct agaacaggac ctggtaaaac 420
atgtttggtt tattgttagc cttactact gtgggctngt ctgtgtgggt tatatcttag 480
gaagtctctt ctactccttg tagcttagaa gtgacccttg ttccgttact taatgtattt 540
attgaggaat attaggggag ggaaaccaag gaaaagatct agcattccca ctttttggtta 600
ttgactaaaa aggatttgna aatcagtttg taaaagaagg tgct 644

<210> 58

<211> 766

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (760)

<223> n equals a,t,g, or c

<400> 58

```
gggtcgaccc acgcgtccgg aatgttttgg tgaataaatc tgttcttcag caaccctacc 60
tgcttctcca aactgcctaa agagatccag tactgatgac gctgttcttc catctttact 120
ccctggaaac taaccacgtt gtcttctttc cttcaccacc acccaggagc tcagagatct 180
aagctgcttt ccatcttttc tcccagcccc aggacactga ctctgtacag gatggggccg 240
tcctcttgcc tccttctcat cctaataccc cttctccagc tgatcaaccy ggggagtact 300
cagtgttcct tagactccgt tatggataag aagatcaagg atgttctcaa cagtctagag 360
tacagtccct ctctataag caagaagctc tcgtgtgcta gtgtcaaaag ccaaggcaga 420
ccgtcctcct gccctgctgg gatggctgtc actggctgtg cttgtggcta tggctgtggt 480
tcgtgggatg ttcagctgga aaccacctgc cactgccagt gcagtgtggt ggactggacc 540
actgcccgct gctgccacct gacctgacag ggaggaggct gagaactcag ttttgtgacc 600
atgacagtaa tgaaaccagg gtcccaacca agaaatctaa ctcaaactgc ccacttcatt 660
tgttccattc ctgattctwg ggtaataaag acaaactttg tacctcaaaa aaaaaaaaaa 720
aaaactcgag ggggggcccc gaamcaattc gggctatagn agagcg 766
```

<210> 59

<211> 2361

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1174)

<223> n equals a,t,g, or c

<400> 59

```
gttctagatc gcgagcggcc gccctttttt tttttttttt tttggaaagc aaggatcaca 60
cttccccctc cctgttccct aatccccctt ctaaaaaggg gggaaaatcc ggatggattt 120
tagggattgg tctggtgtca gctgtgtttt attgcacacc taaatcctga ttataggctt 180
ttcattttct cgcaaagcct ttattttggc agttaagcca aatgtgtttt ccagaaagt 240
agttattttc tcctctttct ttcttttctt tcctcccttt tcccgctctg accccaaacg 300
ttattgtcca aacatgactg gacagcagct tttgtttctt gaccctgtaa tatgacagtc 360
tgctaataat gacagaaggt gcagtttttg gggtatagtc gtgattttcg ctaatcaatc 420
atattagcag gaaaaaaaaa gacttgtttc tgttgtaact gagtcttaag aaaaagtgcc 480
catagttagt tgacaatttc caaaggcttt agtaccacct gtatttcaaa atgggggacc 540
caaactcccg gaagaaacaa gctctgaaca gactacgtgc tcagcttaga aagaaaaaag 600
aatctctagc tgaccagttt gacttcaaga tgtatattgc ctttgtattc aaggagaaga 660
agaaaaagtc agcacttttt gaagtgtctg aggttatacc agtcatgaca aataattatg 720
aagaaaaatat cctgaaaggt gtgcgagatt ccagctattc cttggaaagt tccctagagc 780
ttttacagaa ggatgtggtg cagctccatg ctctcgata tcagtctatg agaagggatg 840
taattggctg tactcaggag atggatttca ttctttggcc tcggaatgat attgaaaaaa 900
tcgtctgtct cctgttttct aggtggaag aatctgatga gccttttagg cctgttcagg 960
ccaaatttga gtttcatcat ggtgactatg aaaaacagtt tctgcatgta ctgagccgca 1020
aggacaagac tggaatcggt gtcaacaatc ctaaccagtc agtgtttctc ttcattgaca 1080
gacagcactt gcagactcca aaaaacaaag ctacaatctt caagttagtc agcatctgcc 1140
tctacctgcc acaggaacag ctcacccact gggncagttg gcaccataga ggatcacctc 1200
cgctcttata tgccagagta gagtactgac cagcaaatg gagaagatca gagaatgcag 1260
cagcagtttt ttttcttgtt ttcttaccac tttattcttt cagagtttaa agaaaaatgga 1320
```

ctcatgcaca gaacactatg cattttgaaa cttgttcac cttggattttt ttaaactcatt 1380
tttatctcag aacttaaaca aaaattagat gtcgtgcacg gactgtgtga aagaagatgc 1440
tttgcataatt tgctgcactg catcagatc ttactaaaaa tgtgaaatga aaggactatt 1500
gtacactgaa atgcttaaat gtatctgaaa gcacaagggtg atactcattt ttatgggtctt 1560
cccatttgtg ctgggtttttg cctctttgac atctgtcac agtattttaga ggggtgagaag 1620
tgaatgtaac aggtataaat aacattttta aaaacaataa ctttgctata atcacagttg 1680
ttccagagca ctgtcagata cattctaattg accagaactg gtttaaaaaa agaaaataca 1740
accatgggaa rgaaatctta aatgaaaaac gcatctcatt gtaggcattt ttgcctcata 1800
ttttactggg ccatgtttgt ttctgtgtac tcatgtattt tttttttcca gatctctttc 1860
cccaagttgc tattgtaaga gtattctgct gcgtgtggat gcagttatac acattaaagc 1920
agatctggag tctgaagtag ctataaagca gctataaaac agaaatacat gcatagctgc 1980
agaaacccatg ataggtagag gacttttctt ttgggtttgt tttgtttgt tttgtttgt 2040
ttttgggttt acagagaaga gatttttatt acaaagaaaa aaattccagt gaattgtgca 2100
gaaatgctgg tttttacacc atcctaaaga aaaactttac aagggtgtt tggagtagaa 2160
aaaagggttat aaagttggaa tcttaaattg taaaattaac cattgagtgt caaagttcta 2220
aaagcagaac tcattttgtg caatgracat aagggaagac tactgtatag gtttttkttt 2280
tctcctttta aatgaaraaa arcttgctta aggggtgcat acttttattg gagtaaatct 2340
gaatgatcct actcccttgg a 2361

<210> 60

<211> 1472

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (129)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (130)

<223> n equals a,t,g, or c

<400> 60

aattcggcac gagcccagat ggcgctggag gaccaggccg ccacactgga gtataagacc 60
atcaaggaac atctcagcag caagagtcac aaccatgggg tgaaccttgt ggagaacctg 120
gacagcctnn ccccaaagt tccacagcgg gaggcctccc tgggtccccc gggagcctcc 180
ctgtctcaga ccggtctaag caagcggctg gaaatgcacc actcctcttc ctacgggggtt 240
gactataaga ggagctaccc cacgaactcg ctcacgagaa gccaccaggc accactctca 300
aaagaaacaa cactaactcc tccaattcct ctcacctctc cagaaaccag agctttggca 360
ggggagacaa cccgccgccc gccccgcaga ggggtggactc catccagggtg cacagctccc 420
agccatctgg ccaggccgtg actgtctcga ggcagcccag cctcaacgcc tacaactcac 480
tgacaaggctc ggggtggaag cgtacgccct cgctaaagcc ggacgtaccc cccaaacct 540
cctttgtctc cctttccaca tccatgaagc ccaatgatgc gtgtacataa tcccaggggg 600
aggggggtcag gtgtcgaacc agcaggcaag gcgagggtgc cgctcagctc agcaagggtc 660
tcaactgcct cgagtaccca ccagaccaag aargcctgcg gcagagccga ggacgctggg 720
tcctcctctc tgggacacag gggtaactcac gaaaactggg ccgcgtggtt tggtagaagg 780
ttgcaacggc ggggactcac cttcattctc ttctttcact tccccccaca ccctacaaca 840
ggtcggaccc acaaaagact tcagttatca tcacaaacat gagccaaaag cacataccta 900
ccccatcccc ccccccmca cacacacaca catgcacaca acacatacac acacacgcac 960

agaggtgaac agaaactgaa acattttgtc cacaacttca cgggacgtgg ccagactggg 1020
tttgcgttcc aacctgcaaa acacaaatac atttttttaa atcaagaaaa tttaaaaaga 1080
caaaaaaaaa agaattcatt gataattcta actcagactt taacaatggc agaagtttac 1140
tatgcgcaaa tactgtgaaa tgcccgccag tgttacagct ttctgttgca gcagataaat 1200
gccatgttgg gcaactatgt catagatttc tgctcctcct ctcttttaat gaaataacgt 1260
gaccgttaac gcaagtaact ctttatttat tgttcacctt ttttttcctt aaggaaagga 1320
ctcttccaaa tatcatccta tgaacagctc ttcagaaagc ccattgaaag ttaaaactatt 1380
taacgtgaaa tccattaact ggaataattg agtttcttta tttttacaat aaattcactg 1440
agtaaataaa aaaaaaaaaa aaaaaaaaaa aa 1472

<210> 61

<211> 1672

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (884)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1583)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1645)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1663)

<223> n equals a,t,g, or c

<400> 61

ccagcctcca ggcacccggg atccagcgcc gccgctcata acacccgcga ccccgagct 60
aagcgagct cccgacgcaa tggacccggc gctggcagcc cagatgagcg aggctgtggc 120
cgagaagatg ctccagtacc ggcgggacac agcaggctgg aagatttgcc gggaaggcaa 180
tggagtttca gtttcctgga ggccatctgt ggagtttcca gggaacctgt accgaggaga 240
aggcattgta tatgggacac tagaggaggt gtgggactgt gtgaagccag ctgttgagg 300
cctacgagtg aagtgggatg agaatgtgac cggttttgaa attatccaaa gcatcactga 360
caccctgtgt gtaagcagaa cctccactcc ctccgctgcc atgaagctca tttctcccag 420
agattttgtg gacttggtgc tagtcaagag atatgaggat gggaccatca gttccaacgc 480
caccatgtg gagcatccgt tatgtcccc gaagccaggt tttgtgagag gatttaacca 540
tccttgtggt tgcttctgtg aacctcttcc aggggaaccc accaagacca acctggtcac 600
attcttccat accgacctca gcggttacct cccacagaac gtggtggact cttcttccc 660
ccgcagcatg acccggtttt atgccaacct tcagaaagca gtgaagcaat tccatgagta 720
atgctatcgt tacttcttgg caaagaactc ccgtgactca tcgaggagct ccagctgttg 780
ggacaccaag gagcctggga gcacgcagag gcctgtgttc actctttgga acaagctgat 840
ggactgcgca tctctgagaa tgccaaccag aggcggcagc ccanccttc ctgcctcctg 900

ccccactcag gggtggcgtg tgatgagcca ttcatgtgtt ccaaactcca tctgcctgtt 960
acccaaacac gcctctcctg gcagggtaga cccaggcctc taaccatctg acagagactc 1020
ggcctggaca ccatgcgatg cactctggca ccaaggcttt atgtgcccac cactctcaga 1080
gaccacgttt cctgactgt catagagaat catcatcgcc actgaaaacc aggcctgtt 1140
gccttttaag catgtaccgc tccctcagtc ctgtgctgca gcccccaaa tatatttttc 1200
tgatatagac cttgtatatg gctttaatgc cgaaaaatat ttatttttcc ttaaaaaagg 1260
tgtcaacttg gaaataatgg tttaaaaaca ggataagcat taaggaaaaa cactttcaat 1320
gtgtcttcca ttgatgaat ttgtkttkct ctctttatcc ccgcaagtgg agtttcatgt 1380
cctcggtgaa accagacagt gtgaatctgt tccagcccaa atctgcagca ttagggatga 1440
gttctcrgaa gtgattctga actgagcacg cactcatgtc tgcattggga actctgggga 1500
gaagagcctt ccttttcttt cccttgggcc atttgctttt ccttgctgctc ttactgaggg 1560
cggaggcagg gaggggtctct gtnctttcca gggcctggg cagggccatc ctggccattc 1620
agggaaagat gggaagagtt agggnctccg ttttaggcag cntgggtgg ga 1672

<210> 62

<211> 1540

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1265)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1468)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1507)

<223> n equals a,t,g, or c

<400> 62

gcggacgggtg ggtcgaccca cgcgtccgct actaacaact taccacagtgc cggagactgc 60
tttctgaaaa ggccactcac gtgaacacta gggatgaaga tgagtrtacc cctcttcac 120
gagcagccta cagtggacac ttagatattg ttcaggagct cattgcacag ggggccgatg 180
ttcatgcagt gactgtggat ggctggacgc ccctgcacag tgcttgaag tggaataata 240
ccagagtggc ttctttctta ctgcagcatg atgcagatat caatgcccaa acaaaaggcc 300
tcttgacccc cttgcatctt gctgctggga acagagacag caaggatacc ctagaactcc 360
tcctgatgaa ccgttacgtc aaaccagggc tgaaaaacaa cttggaagaa actgcatttg 420
atattgccag gaggacaagt atctatcact acctcttga aattgtggaa ggctgtacaa 480
attcttcacc tcagtcctaa caattctagt aatttctcta agtttctaaa taccagtgcc 540
tcctgtgtgt gagatgtatt ccataatca aagttgacgt caaacatctt actacaaaaa 600
ttcagtgcac ttattataa cattcttcca agtgaaattgc ctgactttra tgcataaatg 660
tatttgaaag taatttgcac atatctttaa ttattctgt ggagtttgtg atttttttt 720
cagaaataat tttaattgtg gtatacttaa aaacttgaca cgggttgtac agaaactgg 780
atttttggtg ctgatacaag agaaatgtat ttttaaatat ccacatcct ggatctttgt 840
tggttattta gtatattgac atatatttt ataagggtgag gtaactcaga acttaattta 900
aaagtcttaa atattctgat acaattcagc tgtcttctct accttaccat agccagtgtc 960

tttcatttta aaccagagca agtaacatat tagtgacttg aatcttcata agttaagta 1020
aaaaacagca aaaaacctag atctttgtct tttagaacac agaccatttt caggaaagca 1080
gttagctaag tgtttaattc atgaatattg tatactgcat cccctaccac aatttacaca 1140
atcctgtgga tagtcctacc tcaccctggg caacctacat gatccttaag ctaatggcga 1200
atcacgatga ccttgtagac atgcacacaa ctataccttt gtccaacaga tcataatata 1260
tctgnctatc caactggttt tacctgccta atcctactga tttgggcact gcttgatar 1320
tctctcaagt tcacagaaa tgttgatttt ctaaggctct catttttaca gagtatacag 1380
gcaaaagtgc aggggaaaag gaattagtct aagagtaagg ggatgattat tatrttgagg 1440
staaaaccac aaagtggctc aggccttnaa aaaaaaacac tgtggataat gacaaaaagc 1500
ataagtnaaa atatttgaga aaaataaagt acaagaaatg 1540

<210> 63

<211> 1044

<212> DNA

<213> Homo sapiens

<400> 63

aacaccttca tccatgaaga tatctggaac attcgtagta tctgcagcac caccaatata 60
caatgcaaga acggcaagat gaactgccat gaggggtgtag tgaaggtcac agattgcagg 120
gacacaggaa gttccagggc acccaactgc agatatcggg ccatagcgag cactagacgt 180
gttgctcattg cctgtgaggg taaccacag gtgcctgtgc actttgacgg ttagatgcca 240
ccatgtaggg attatcgga gtggttgacc ttacacttac tccttaaata gcagtgaata 300
atgcatttga gctgtcccag gctctgtctc ctacagctcat ttctactct tttctctat 360
ataactcatt ctattaaata cattgcacca aagagatatg gagacataaa cctgtaata 420
atgaggctgg gcttttctgt aataagcttc cttttataat actggtcagc ttagctctct 480
cagatcctat cctgtggaat ttagttatta tgtgtattta ttagtattt caaacatttc 540
aaaatgcttt catctatgtt tatcacattt taataccaca gcacttataa tgatgtcact 600
acatatagaa gctcaaagt aagggtattg ctgaagactg taaagttaat ggaagaattg 660
agacaaaaat ccagtgtagc tggccactta tccagggtt tttctacttc atcacaagga 720
atgttttgaa agtgtctgct ttttttatcc ttaaaattca cctgtcaggg aggcattaaa 780
aatttggaat tgatgccag caaatgtga gctctgtatt ttttggcatt cttatgtttg 840
ggtttaataa gattaagaaa atgatactgg gaattttctt tttcctgaaa ctttgaatca 900
ccctagtaag tcaaagtact aaaaaatgta ctagatcatt aagacttatg tgctcttact 960
gattgaaaga ttttttatgt tttccttgta ataaaggacc taaaccgaag gtacctgaaa 1020
aaaaaaaaa aaaaaaaact cgag 1044

<210> 64

<211> 851

<212> DNA

<213> Homo sapiens

<400> 64

cacgagagaa ggtggttatt tatacaaaaca tggacatact cactcccaag ggctgatgag 60
atgctgaatt ttctttggg gcattcatta attgtcccag ctgcagcgac tggagcaagt 120
ctggaagctg cctgtgctaa gaccacccag ctgtccctgg gttctcatcc tagggccttc 180
tttgcttcca ggtcagggga cctgcttcaa tgagaaagca actgaattga ggctaggaga 240
ggtagggaga gctgagtct gacttcacct gtgcagaact ctctgcccc atgttacctg 300
gactggaaca gactgtgaat atagcagaag gttccaagaa ctctgggtgc tgacctagaa 360
gaggcacagt tctctctact ggaaagaaaa cgatgtagcc gattgcacaa gggtgccaa 420
ggaagaccca ggtggccca tcaaaggaac ctgggggagg atgcaggagg ctgaaggga 480
gcacctggca tttctctcac tgtgctctta ccgcatcagc aacccccaac ttttgggcct 540

actctgcccc ccatgcggtga ataccctgct tggatgctgt gcttttccgg tttgtctcta 600
agcccccttc tccagggcat gttgggttcc ctggcctctc agtgtcctaa ctggagccca 660
gagtgccttg ttctgagcca ggagacggct gagcactggc cctccacacc taagcgtcct 720
ttacattaac ttattggtct tgtataacac ctggtgccat tgccaagtgg ctgtgtcctc 780
agctacagag ctggaattgt gtggggttta gtgctaaata cttcaataaa gtctgttttt 840
tgtgattggc t 851

<210> 65

<211> 2793

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2793)

<223> n equals a,t,g, or c

<400> 65

ccacgcgtcc ggattacatg tagttattga gaatcctttc gaattcagtg gcttaatcat 60
gaatgtctaa atattgttga cattaggatg atacatgtaa attaaagtta catttgttta 120
gcatagacaa gcttaacatt gtagatgttt ctcttcaaaa atcatcttaa acatttgcac 180
ttggaattgt gttaaataga atgtgtgaaa cactgtatta gtaaaacttca tcacctttct 240
acttccttat agtttgaact tttcagtttt tgtagttccc aaacagttgc tcaattttaga 300
gcaaattaat ttaacacctg ccaaaaaaag gctgctgttg gcttatcagt tgtctttaaa 360
ttcaaatgct catgtgactt ttatcacatc aaaaaatatt tcattaatga ttcaccttta 420
gctctgaaaa ttaccgcgtt tagtaattat agtgggctta taaaaacatg caactctttt 480
tgatagttat ttgagaattt tgggtgaaaa tatttagctg agggcagtat agaacttata 540
aaccaatata ttgatatttt taaaacattt ttacatataa gtaaaactgcc atctttgagc 600
ataactacat ttaaaaaata agctgcatat ttttaaatca agtgtttaac aagaatttat 660
atTTTTtatt ttttaaaatt aaaaataatt tatatttcct ctgttgcatg aggattctca 720
tctgtgctta taatggtag agattttatt tgtgtggaat gaagtgaggc ttgtagtcac 780
ggttctagtg tttcagtttg ccaagtctgt ttactgcagt gaaattcatc aaatgtttca 840
gtgtggtttt ctgtagccta tcatttactg gctatttttt tatgtacacc tttaggattt 900
tctgcctact ctatccagt gtccaaatga tatcctacat tttacaaatg ccctttcagt 960
ttctattttc tttttccatt aaattgccct catgtcctaa tgtgcagttt gtaagtgtgt 1020
gtgtgtgtgt ctgtgtgtgt gtgaatttga ttttcaagag tgctagactt ccaatttgag 1080
agattaaata atttaattca ggcaaacatt tttcattgga atttcacagt tcattgtaat 1140
gaaaatgtta atcctggatg acctttgaca tacagtaatg aatcctggat attaatgaat 1200
ttgttagtag catcttgatg tgtgttttaa tgagttattt tcaaagtgtg gcattaaacc 1260
aaagtggca tactggaagt gtttatatca agttccattt ggctactgat ggacaaaaaa 1320
tagaaatgcc ttcctatgga gattattttt cctttaaaaa attaaaaagg ttaattattt 1380
tgaaaaaaa aaatcgaccc acgcgtccgg attacatgta gttattgaga atcctttcga 1440
attcagtggc ttaatcatga atgtctaaat attgttgaca ttaggatgat acatgtaaat 1500
taagttaca tttgttttag atagacaagc ttaacattgt agatgtttct cttaaaaaat 1560
catcttaaac atttgcattt ggaattgtgt taaatagaat gtgtgaaaca ctgtattagt 1620
aaacttcac acccttctac ttccttatag tttgaacttt tcagtttttg tagttcccaa 1680
acagtgtctc aatttagagc aaattaattt aacacctgcc aaaaaaggc tgctgttggc 1740
ttatcagttg tctttaaatt caaatgctca tgtgactttt atcacatcaa aaaatatttc 1800
attaatgatt caccttttag tctgaaaatt accgcgttta gtaattatag tgggcttata 1860
aaaacatgca actctttttg atagttattt gagaattttg gtgaaaaata tttagctgag 1920
ggcagtatag aacttataaa ccaatatatt gatattttta aaacattttt acatataagt 1980

```

aaactgccat ctttgagcat aactacattt aaaaataaag ctgcatattt ttaaatcaag 2040
tgtttaacaa gaatttatat tttttatttt ttaaaattaa aaataattta tatttctctt 2100
gttgcatgag gattctcatc tgtgcttata atgggtagag attttatttg tgtggaatga 2160
agtggaggct gtagtcatgg ttctagtgtt tcagtttgcc aagtctgttt actgcagtga 2220
aattcatcaa atgtttcagt gtgstyttct gtagyctatc atttactggc tattttttta 2280
tgtacacctt taggattttc tgcctactct atccagttgt ccaaatagata tcctacattt 2340
tacaaatgcc ctttcagttt ctattttctt tttccattaa attgccctca tgcctaatg 2400
tgcagtttgt aagtgtgtgt gtgtgtgtct gtgtgtgtgt gaatttgatt ttcaagagt 2460
ctagacttcc aatttgagag attaaataat ttaattcagg caaacatttt tcattggaat 2520
ttcacagttc attgtaatga aaatgttaat cctggatgac ctttgacata cagtaatgaa 2580
tcttgatat taatgaattt gttagtagca tcttgatgtg tgttttaatg agttattttc 2640
aaagttgtgc attaaaccaa agttggcata ctggaagtgt ttatatcaag ttccatttgg 2700
ctactgatgg acaaaaaata gaaatgcctt cctatggaga gtatttttcc tttaaaaaat 2760
taaaaagggt aattattttg aaaaaaaaaa acn 2793

```

<210> 66

<211> 303

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (108)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (278)

<223> n equals a,t,g, or c

<400> 66

```

agattaagca tacttaacca ctccattatt gtagattcac tttcaacctt aaaaattaat 60
accagtttgc ataaaccaat atctgaaaag aacaggaaat gttaatgnca agcaacagct 120
attaatactg atgtgaatgg atgcatttgt tttgcagtgg tgactggcct aggcagggtt 180
gggatctgtg aaagaattga ttcattttca aaattattcc ataaagttaa aaagttacac 240
tttaaaggca acaggtcata cagttcttta aaatctgnat ccaactgtag ctttatttaa 300
aag 303

```

<210> 67

<211> 1410

<212> DNA

<213> Homo sapiens

<400> 67

```

ccacgcgtcc ggtccctagg agataagagt atcttgacac gcagggtgcag gtttcccagc 60
agtcaggga agagtcggat gtttgtgcc tctgatcctg atgtctggag agatagccat 120
gtgtgagcct gaatttgga atgacaaggc cagggagccg agcgtgggtg gcagggtggc 180
agtgtcctgg tacgaacggt ttgtgcagcc atgtctggtc gaactgctgg gctctgctct 240
cttcattctc atcgggtgcc tgcgggtcat tgagaatggg acggacactg ggctgctgca 300
gccggccctg gccacgggc tggctttggg gctcgtgatt gccacgctgg ggaatatcag 360
tgggtggacac ttcaacctg cgggtgtccct ggcagccatg ctgatcggag gcctcaacct 420

```

ggatgatgctc ctcccgactt ggggtctcaca gctgctcggg gggatgctcg gggctgcctt 480
ggccaaggcg gtgagtcctg aggagagggt ctggaatgca tctggggcgg cctttgtgac 540
agtccaggag caggggcagg tggcaggggc gttgggtggca gagatcatcc tgacgacgct 600
gctggccctg gctgtatgca tgggtgccat caatgagaag acaaagggcc ctctggcccc 660
gttctccatc ggctttgccg tcaccgtgga tatcctggct gggggccctg tgtctggagg 720
ctgcatgaat cccgcccgtg cttttggacc tgcgggtggt gcccaaccact ggaacttcca 780
ctggatctac tggctggggc cactcctggc tggcctgctt gttggactgc tcattagggtg 840
cttcattgga gatgggaaga cccgcctcat cctgaaggct cagtgaagca gagctcgtgg 900
gattcctgct gctccagggt tcctcagctc acctgtccca gactgaggac aggggagttc 960
ctgcatttcc tgccagggca gaggccaga ggagcgaccc cctgcttcca ctgcttgggc 1020
ctgctttctc agatagactg actgctgagg aggtcttagg ttcttggaa tcctttgtgc 1080
tcacagaga cccagcctg gggaacacgc tgcccgact gccagagag cagtgcaaac 1140
accacaacac gagcgtgttt cttgagagga atgtcccca gttggacaag gaggctgttt 1200
ctgcacatca gctcatttcc cgcaccccat ttcttgcttg attgctttgt tgggggcctg 1260
gccacttct tgcttctcaa gctgacaatt ctcactttgc aataaatagt ccagtgtttc 1320
cttccaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1380
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1410

<210> 68

<211> 1024

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<400> 68

angggangga agggaaggga agggaagggg ggggtcacgcg ggggcgcgcg cgcgcaccgg 60
gagccgctcg raggcgagtg gaactggatc ggggttgctg ccagcggcgt gagcttcggc 120
cgccatttta caacagctcc actcgcgccg gacacaggga gcagcgagca cgcgtttccc 180
gcaacccgat accatcggac aggttttctc cgcctcagcc caacggggag atctctggaa 240
acatggctac agaacatgtt aatggaaatg gtactgaaga gcccatggat actacttctg 300
cagttatcca ttcagaaaat tttcagacat tgcttgatgc tggtttacca cagaaagttg 360
ctgaaaaact agatgaaatt tacgttgacg ggctagtgtg acatagtgat ttagatgaaa 420
gagctattga agcttttaaa gaattcaatg aagacggtgc attggcagtt cttcaacagt 480
ttaagacag tgatctctct catgttcaga acaaaagtgc ctttttatgt ggagtcata 540
agacttacag gcagagagaa aaacaaggga ccaaagtagc agattctagt aaaggaccag 600
atgaggcaaa aattaaggca ctcttggaag gaacaggctc acacttgatg tgaccactgg 660
acagagggaag tatggaggac cacctccaga ttccgtttat tcaggtcagc agccttctgt 720
tggcactgag atatttggtg gaaagatccc aagagatcta ttgaggatg aacttgttcc 780
attatttgag aaagcataag gttctcagag tgatagagct ggaaggatca taaaggatca 840
ctagtctaat ggttctcaaa ctcgaaggaa cattaggggc atcctggaga gtttactggg 900
ccccgtctca gaggttctga tgctgtaaat caggcgggac ccctgaattt tcatttctaa 960
gaagctcaca ggtgctacta atgttgaaag gaacctatga tctagttcaa cttctgggat 1020

tctt

1024

<210> 69

<211> 1848

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1761)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1844)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1847)

<223> n equals a,t,g, or c

<400> 69

agactcggat ggtagtccgc tgtccaacag ccarccttcc ttcccagtgg agatcttgcc 60
cttcctctac ttgggctgtg ccaaagactc caccaacttg gacgtgttg aggaattcgg 120
catcaagtac atcttgaacg tcaccccca tttgccgaat ctctttgaga acgcaggaga 180
gtttaaatac aagcaaatcc ccattctcga tcaactggagc caaaacctgt cccagttttt 240
ccctgaggcc atttctttca tagatgaagc ccggggcaag aactgtggtg tcttggtaca 300
ttgcttggtc ggcattagcc gctcagtcac tgtgactgtg gcttacctta tgcagaagct 360
caatctgtcg atgaacgatg cctatgacat tgtcaaatg aaaaaatcca acatatcccc 420
taacttcaac ttcattgggtc agctgctgga cttcgagagg acgctgggac tcagcagccc 480
atgtgacaac agggttccag cacagcagct gtattttacc accccttcca accagaatgt 540
ataccaggtg gactctctgc aatctacgtg aaagacccca caccctcctt tgctggaatg 600
tgtctggccc ttcagcagtt tctcttgga gcatcagctg ggctgcttcc tttgtgtgtg 660
gccccagggtg tcaaatgac accagctgtc tgtactagac aaggttacca agtgcggaat 720
tggttaatac taacagagag atttgctcca ttctcttttg aataacagga catgctgtat 780
agatacaggc agtaggtttg ctctgtaccc atgtgtacag cctacccatg cagggactgg 840
gattcgagga cttccaggcg catagggtag aaccaaatga tagggtagga gcatgtgttc 900
tttagggcct tgtaaggctg tttccttttg catctggaac tgactatata attgtcttca 960
atgaagacta attcaatttt gcatatagag gagccaaaga gagatttcag ctctgtatatt 1020
gtggtatcag tttggaaaaa aaaatctgat actccatttg attattgtaa atatttgatc 1080
ttgaatcact tgacagtgtt tgtttgaatt gtgtttgttt tttcctttga tgggcttaaa 1140
agaaattatc caaagggaga aagagcagta tgccacttct taaaacagaa caaaacaaaa 1200
aaagaaaatt gtgctctttt ctaatccaaa gggatatatt gcagcatgct tgactttacc 1260
aattctgatg acatctttac ggacactatt atcactaaga ccttggttatg gcgaagtctt 1320
tagtcttttt catgtatttt cctcatgatt ttttctcttt atgtagtttg actatgcctt 1380
acctttgtaa atatttttgc ttgtgttgct gcaaagggga taatctggga aagacaccaa 1440
atcatgggct cactttaaaa aaagaaagaa taaaaaaacc ttcagctgtg ctaaacagta 1500
tattacctct gtataaaatt cttcaggagg tgtcacctca aatgcaatac tttgggttg 1560
tttctttcct ttaaaaaaat ttgtataaaa ctggaagtgt gtgtgtgtga gcatgggtac 1620
ccatttgata agagaaatgc atttgattgt gaagaaggga gagttaaat ctccattatg 1680

```
ttcgtggtgt aaagtttwwa gctggaattt attataagaa tgtaaacct taaattatta 1740
ataaataact attttggcta naaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1800
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaggg gggncnc 1848
```

<210> 70

<211> 2682

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (647)

<223> n equals a,t,g, or c

<400> 70

```
gaatacccca ggatttatgt ataaaaacct gcagtgtctg gttattgatg aagctgatcg 60
tatctttgat gtggggtttg aagaggaatt aaagcaaatt attaaacttt tgccaacacg 120
tagacagact atgtctcttt ctgccacca aactcgaaaa gttgaagacc tggcaaggat 180
ttctctgaaa aaggagccat tgtatgttgg cgttgatgat gataaagcga atgcaacagt 240
ggatggtctt gaacagaaga accgaaagaa gaagcttatg gtcttctttt catcttgtat 300
gtctgtgaaa taccactatg agttgctgaa ctacattgat ttgccgtctt tggccattca 360
tggaagcaa aagcaaaata agcgtacaac cacattcttc cagttctgca atgcagattc 420
gggaacacta ttgtgtacgg atgtggcagc gagaggacta gacattcctg aagtcgactg 480
gattgttcag tatgacctc cggatgacct taaggaatat attcatcgtg tgggtagaac 540
agccagaggc ytaaatggga gagggcatgc cttgctcatt ttgcgcccag aagaattggg 600
tttcttcgt tacttgaaac aatccaaggt tccattaagt gaatttngac ttttcctggg 660
ctaaaatttc tgacattcag tctcagcttg agaaattgat tgaagaagt tactttcttc 720
ataagtcagc ccaggaagca tataagtcac acatacagc ctatgattcc cattctctga 780
aacagatctt taatgttaat aacctaaatt tgctcaggt tgctctgtca ttgggtttca 840
agggtgcctc cttcgttgat ctgaacgtca acagtaatga aggcaagcag aaaaagcgag 900
gaggtggtgg tggatttggc taccagaaaa ccaagaaagt tgagaaatcc aaaatcttta 960
aacacattag caagaaatca tctgacagca ggcagttctc tcaactgaaca catgccttcc 1020
tttcatcttg aataactttg tcttaaaatg aatttttttt ccccttgatt taacaggatt 1080
ttttagact ttagaatttg gacttaccta acaagagtat aaattgactt gggttgcaag 1140
cactgagcac tgttacttct atcacgtctc tcttttatth ctgggatata aaacaggctt 1200
taagtttctt ggttgcccaa gggcagagca aggaatatct ggtgtttctt gtgatgataa 1260
tattttaatt ttaaataatc ctccctcata caagtgtatg ttaccatttt aatataattc 1320
ttttgtacc tttccttctt gttttgtgaa gatttttgtg gcatggattg ctgtgtctac 1380
tgctgtaaaa ggtgacctag tgtactgggc agctggtggc ggtgcagaaa agagtctcag 1440
gttatttttt gtttttagtt atttcttgga ccttgacagt atctaagac tcctcctgaa 1500
aatgctgcag tataaaagag caaagagctt tgggaaatac ctaagaagca ccttaagatt 1560
aggtggtcat tgcttttata gattcttgat ttttaagcaa caggcctttc tcaggtgttg 1620
catttttttg agcaaaaact atgggttgta atttgaataa agtgtcacta agcagttata 1680
acgtttgatg gctggggggg aggaagagga tggaaattgag atgtttgagc ctcatttaca 1740
tcaatagagg tgtaatgtac tgcatttctt catttggtta cataacaaag actttcatac 1800
aaagaacgat gatgctctc attaagattt gtttaattca aggtgggttg gatttggtta 1860
gcctttgcac tctgtagagt acttagaaga caaggccaac ttacttgag ttagagccaa 1920
gctgtcagac ggtgcccagc acacattaat gtttagcttct tcttgagaaa aaaatacctc 1980
ttccaggccc tgaaacaaaa aatacatttg ctgtgaagat tgaagaatgaa caaagttaga 2040
aaaaaaaaa gcaaatcag tgatttagtc agatgagttt ttcgttgtag gagcacttga 2100
tttctagtgt gttttgtaca gtatataact acaagatagt acattttgta gcagttcaaa 2160
```

gccaaagtgtg ctacgcatcat tttgctgttg tgccagttaa tcataggatc ccattaaata 2220
agtggtgctaa catcgaatat agagaaaact ggtaaagaac attccagtag gaaaagaaaa 2280
gaacaatctt ccattttctgg gcttgggccac catcaccctg gtcggacctg tcctggactt 2340
ccaaccttga ctgctgagct cctggcttag cttcttgggt tcctaattcc tgggtgttaa 2400
taattctctc cacgatcatg tttttctgat ttttttttc agaaataatg ttttttaaaa 2460
gacaaaaaca aagggaagaa tatttaatta ctgagcagaa gtaaatactg ttggtatttt 2520
gtacataatc taatttttat atgcatgtty atgcttttta atttttttat caaaaattaa 2580
gtcatctacc tactacttgt aaccagcttg tttcataaca tggtattttc ctgtgtcatt 2640
aaataattac ttcaatgttg aaaaaaaaaa aaaaaaaaaa aa 2682

<210> 71

<211> 412

<212> DNA

<213> Homo sapiens

<400> 71

ggcagagcca acagacaccc tccactctga ggtctcacct tcgcctttgc tgaagtctcc 60
ccgcagccct ctccaccag aggtctccct ataccgagac ccaccatssw wccatcctga 120
ggaccgcccc aacctcggga gccccccact cagtaggtct gaaggcctcc atttgtaccg 180
aaacaccccg ctacgctga cagcctccta ggytccctga ggtaccttc caccagacc 240
ctccttcccc accccataag ccctgagact cccgcctttr acctgacgat ctccccctt 300
ccgccttca ggttctctct aggcgctcag aggcgctctt ggggggttgc ctcgagtccc 360
cccacccctc cccaccacc accgctcccg cggcaagcca gcccgtcag aa 412

<210> 72

<211> 1361

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (46)

<223> n equals a,t,g, or c

<400> 72

taagaacttt tgtccttccc ttcttgaggt ccttttttga ttgtgncttt agtatcaagt 60
tcagcagttg tttcaagagc agctgagata atwaattcaa ccaaagggcg ccttggttct 120
ttawggatc tgawactcct ctaggacttg cgagggccgt gtcctgagta gtgtgtgmcc 180
cttggtctac gttgcctctg ttttctccta gttgccagac ttagaagmct tgatgaaaag 240
ggagaaccag aagattctga ctcccttggg gagectggac accccaggga aagcaacagt 300
acaggtggtc attctggccg accctgacgg acatgaaatt tgctttgtcg gggatgaagc 360
atttcagagaa ctttctaaga tggatccaga gggaagcaaa ttgttggatg atgcaatggc 420
agcagataaa agtgacgagt ggtttgccaa acacaataaa cccaaagctt cagggttaacg 480
gaagacatga tgcagagcaa gcctctgtga ttcctgcccc gcacctgtga ggctgacgt 540
gtcagttccc aataaatgct cttctgattt gtttcccgta caggcaagga ggcttgggta 600
gtgcagattt gtgtatttca atctttgaaa gctctgatgt aatttagaaa tgaaatccaa 660
tcatgagtc aggtagagaa cgcctgctgt aatctacact gttgctggga ctgcgcattc 720
tgtatataac tgtgttgat gagtgacaga tgattgtcca gactaggaca gcggcatgaa 780
catgactttg gttgggattg cggatagtta gggttacctc tgaatcgtgt agcttttatg 840
agagcagctg tgcaagtga tccacattaa tgccttgctg tggtgccatt cccagcgcct 900
gacgatacgc tcttctattg tcttattctg gcaggttttg acgttttaaa ttttttaaa 960

```

aaattttatt ccttggaacca aaagggtttgg ttaaccacccc ccctcttact tgctttcaca 1020
ttttgagtgt ccagaggaaa cagaaaggaa tgagtgtgtg acgttgctgc acgcctgact 1080
ctgtgcgagc ttctttctgt gtatatattt tgttttattt ttttccgtgt atatttttaa 1140
tcccgcagaga acatcatgtg agatttcttt aaaatggatt aaacgatttc ttcagcctga 1200
aaaaaaaggt tttgaaaatg ttttcttgta gttttgtttg gttctaaaca acaaataggt 1260
tttaatcact cgaaatggaa ttatattgtg tattcattga ataaattttt tttgaaagta 1320
agcttctgaa atcaaaaaaa aaaaaaaaaa aaaaaaaaaa a
1361

```

<210> 73

<211> 928

<212> DNA

<213> Homo sapiens

<400> 73

```

cagcaatggc agcatctgcc ttgatatcct gcggtctcag tggctctccag cgttgactgt 60
gtcaaaagtt ctcttgcca tctgtctgct gctctgcgac cccaaccccg atgacccct 120
ggtgccagag atagcacaca cctacaaggc cgacagagag aagtacaaca gactagcaag 180
agagtggaca caaaaatatg ctatgtaagt gccttggagg ttttacctga gacactgtcc 240
aagagaagct ggcagagagg tcttccctta aaactttggg ctggttggtg agccattcaa 300
agagcatcat ctgttcttca aacaaatgtt ggtcaccac tctctccagc tgcagcatgt 360
tggtgccatt ttcagcaatt acggccttga cagtgccacc tctttgatgc caaatcagca 420
accattggtt ttatgatctg cagtcttcct ggtgacctg gaatctctct ctctgccgcc 480
tcagtttgct tgctggtctc ttggggggcc aggcctgcac gtctctccta cccggcctca 540
aatggtgctg ctgcccata tgggtaccaca ccagggcctc agcctggccc ctcaccacat 600
accctttgcc ttttagaact cagtgccatc ctgggtgccc agggcagarc aggccttgtt 660
cgcacctcat ctgctgcaga accacatcct gaggagtctc agcttatcct ggagggaatt 720
gggaacagtg tcaactgggaa gtgaaggcct agccctgtgg ctccaccag tctcctcctg 780
cagtgccacg tgggtgaattt tctcgcttc acaccaagaa agcagcaaag tggaaaattt 840
tcaaggatac aaaggcacat aacamcccca taagragatg attaaaggtt tttagaagca 900
agagcaaaat tttgaaaacc tctaggag
928

```

<210> 74

<211> 1186

<212> DNA

<213> Homo sapiens

<400> 74

```

gcggacgcgt gggcagcccg ggcggctgcc cttgggtgct cccttccttg cccgacaccc 60
agaccgacct tgaccgcca cctggcagga gcaggacagg acggccggac gggccatgg 120
ccgagctccc ggggcccttt ctctgcgggg ccctgctagg ctctctgtgc ctgagtgggc 180
tggccgtgga ggtgaaggta cccacagagc cgctgagcac gccctgggg aagacagccg 240
agctgacctg cacctacagc acgtcggtgg gagacagctt cgccctggag tggagctttg 300
tgcagcctgg gaaacccatc tctgagtcct atccaatcct gtacttcacc aatggccatc 360
tgtatccaac tggttctaag tcaaagcggg tcagcctgct tcagaacccc cccacagtgg 420
gggtggccac actgaaactg actgacgtcc acccctcaga tactggaacc tacctctgcc 480
aagtcaaaa cccaccagat ttctacacca atgggttggg gctaataaac ctactctgct 540
tggttcccc cagtaatccc ttatgcagtc agagtggaca aacctctgtg ggaggctcta 600
ctgactgag atgcagctct tccgaggggg ctccataagc agtgtacaac tgggtgcgtc 660
ttggaacttt tcctacacct tctcctggca gcatggttca agatgaggtg tctggccagc 720
tcattctcac caacctctc ctgacctcct cgggcaccta ccgctgtgtg gccaccaacc 780
agatgggcag tgcactctgt gagctgacct tstctgtgac cgaaccctcc caaggccgag 840

```


tggccggagc tctgattggg gtgctcctgg gcgtgctggt gctgtcagtt gctgcgttct 900
gcctggtcag gttccagaaa gagaggggga agaagcccaa ggagacatat gggggtagtg 960
accttcggga ggatgccatc gctcctggga tctctgagca cacttgatg agggctgatt 1020
ctagcaagg gttcctggaa agaccctcgt ctgccagcac cgtgacgacc accaagtcca 1080
agctccctat ggtcgtgtga cttctcccga tccctgaggg cggtgagggg gaatatcaat 1140
aattaaagtc tgtgggtacc aaaaaaaaaa aaaaaaaagt cgacgc 1186

<210> 75

<211> 933

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (791)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (794)

<223> n equals a,t,g, or c

<400> 75

gtctggatca tattgtagat aaagtaaaag aatgtgtgga tcatttaagt agagatgagg 60
atgaagagaa actggtagcc tcactatggg gagcagagag atgtttacga gttttagaaa 120
gtgtaactgt gcataatccc gaaaatcaaa gctacttgat agcatataaa gattcccaac 180
ttattgtttc atcagctaaa gcattacagc attgtgaaga actgattcag cagtacaacc 240
gtgctgagga cagcatatgc ttagctgaca gtaagcctct gcctcaccag aatgtaacta 300
accatgtagg caaagcagtg gaggactgca tgagggccat catcggggtg ttgcttaatt 360
taactaatga taatgagtgg ggcagcacca aaacaggaga gcaggacggt ctcataggca 420
cagcgctgaa ctgtgtgctt cagggtccaa agtacctacc tcaggagcag agatttgata 480
ttcgagtgtt gggcttaggt ctgctgataa atctagtggg gtatagtgtt cggaatcggc 540
actgtcttgt caacatggaa acatcgtgct cttttgatcc ttccatctgt agtgagagaag 600
gggatgatag tttaaggata ggtggacaag ttcattgctgt ccagcttttag tgcagctatt 660
ccttgagcga gageggggcag cccagctagc agaaagtaaa acagatgagt tgatcaaaga 720
tgctcccacc actcagcatg ataagagtgg agagtggcaa gaaacaagtg gagaaatata 780
gtgggtgtca ntgnaaaaga ctgatggtac agaagagaaa cataagaagg aggaggagga 840
tgaagaactt gacctccaat aaaggtatat ttactttgaa tgggtgtattt aaattgaaga 900
tagtggtttg attttgtttt ttccctccaa acc 933

<210> 76

<211> 1964

<212> DNA

<213> Homo sapiens

<400> 76

ggtttggcag ggcggggcgc ctgcgaaga tgggtggcgc cgtggcgtgt ggctcccgtc 60
gtctggccaa gtctcagcgc agcgcaccgg ccggcgtctc gttggcctgg agccacacc 120
caccgggtcc ctgacccgc gccccgcgc cccggttccc ggcattgcctc gcgcccgtaa 180
gggcaacacg ctccggaagg gtggtcagcg ccgtggagga ggtgcccga gcagtgccca 240
agctgactcg ggttccagtg acgatgaggc agccagttag gccgcagca ccgccagtga 300

```
atgccccagc cttctcagca ccactgcaga ggacagcctt ggggggggatg tcgtggatga 360
gcaggggccag caggaagacc ttgaggaaaa gctgaaggag tatgtggact gtctcacaga 420
caagagtgcc aagaccgggc aggtgctctt gagagcctgc gcctggccct agcgtcccg 480
ctcaaccccg acttcttgct ggagcgccgc ctcacgctag ccgatgccct ggaaaagtgc 540
ctcaagaaaag ggaagggcga ggaacaagcc ctggctgctg ctgtgctagg cctgctctgc 600
gtgcagctgg gccctggacc taagggtgag gagctgtttc acagcctgca gcctctgctg 660
gtctctgtgc tcagtgcag cactgctagc cctgctgccc ggctccactg tgcttctgcc 720
cttgccctgg gctgctacgt ggctgcccgt gacatccagg acctggctctc ttgccttgcc 780
tgcttagaaa gtgttttcag ccggttctat ggctggggg gcagctccac aagtccctgtg 840
gttcttgcca gcctgcacgg cctgctctct gctgccctgc aggcctgggc attgctgctc 900
acctctgcc ctagcaccca aatcagccac atccttgaca ggagctgcc ccggtgccc 960
cagctcttgt ccagtgaag tgtgaacctg cggatcgctg ccggtgaaac cattgactg 1020
ctctttgagc ttgccggga ccttgaggag gagtttgttt acgaggacat ggaggccctc 1080
tgcagtgtcc tgcgactct ggccactgac agtaacaagt accgtgccaa ggctgatcgt 1140
cggcgccagc gctctacttt ccgcgccgtg ctgactccg tggagggcg tgaatgcgaa 1200
gaagagatag tgcgcttcgg ctttgaggtg ctctacatgg acagctgggc tcggcaccgg 1260
atctacgtg ccttcaagga agtgcctggg tcgggcatgc accaccacct ccagaacaat 1320
gagctactcc gtgacatctt tggcctgggc cctgtgctgt tgctggatgc cactgccctg 1380
aaggcctgca aggttccacg ctttgagaag cacctgtaca atgctgctgc cttcaaagcc 1440
cggaccaagg ctggaagccg tgtgcgggac aagcgggcag acatcctgtg aagcaggacc 1500
tgctgaagag gagactttct atgcccttgg tccgtatttt taacagaaga cagtgcaca 1560
actggtctcc accagtattt gtcactttat tttttttaat gacaaaacca aaaacagaca 1620
tggggtgggt agctgggggc ccggacactt gggaccctga ccccttgct cctgactca 1680
gccctgtggc cccttcctgt cctgtctcag gccaggctaa atatgtgcct tcctcagggc 1740
tgtggggcag gcactagggg gcctttccct tcctttcctt tctcaggcct tgctccccc 1800
ggatgaccca ctcttagggg ggtggtggca tctggacaaa tgccaccaca gcaggtgggg 1860
tggaagagct acctggaatg gatttgtgtg ctgattttta aggattatta cagataatta 1920
aacagaacgg tcagccttca aaaaaaaaaa aaaaaaaaaa aaaa 1964
```

<210> 77

<211> 1802

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1680)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1747)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1757)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1800)

<223> n equals a,t,g, or c

<400> 77

acgcgtccgc ttccatctgc tctggaatta aatatgcatt tcaggtgatt ggagagctac 60
attcccaact cgatggatcc gaagtactgc tgctgactga tggggaggat aacactgcaa 120
gttcttgtat tgatgaagtg aaacaaagtg gggccattgt tcattttatt gctttgggaa 180
gagctgctga tgaagcagta atagagatga gcaagataac aggaggaagt catttttatg 240
tttcagatga agctcagaac aatggcctca ttgatgcttt tggggctytt acatcaggaa 300
atactgatct ctccsagaag tcccttcagc tcgaaagtaa gggattaaca ctgaatagta 360
atgcctggat gaacgacact gtcataattg atagtacagt gggaaaggac acgttctttc 420
tcatcacatg gaacagtctg cctcccagta tttctctytg ggatcccagt ggaacaataa 480
tggaaaattt cacagtggat gcaacttcca aaatggccta tctyagtatt ccaggaactg 540
saaagtgagg cacttgggca tacaatctty aagccaaagc gamcccagaa acmttaacta 600
ttacagtwac ttctcgagca kcaaaattct tctgtgcctc caatcacagt gaatgctaaa 660
atgaataagg acgtaaacag tttccccagc ccaatgattg tttacgcaga aattctacaa 720
ggatatgtac ctgttcttgg agccaatgtg actgctttca ttgaatcaca gaatggacat 780
acagaagttt tggaaactttt ggataatggt gcaggcgctg attctttcaa gaatgatgga 840
gtctactcca ggtattttac agcatataca gaaaatggca gatatagctt aaaagttcgg 900
gctcatggag gagcaaacac tgccaggcta aaattacggc ctccactgaa tagagccgcg 960
tacataccag gctgggtagt gaacggggaa attgaagcaa acccgccaag acctgaaatt 1020
gatgaggata ctacagaccac cttggaggat ttcagccgaa cagcatcccg aggtkcattt 1080
gtggtatcac aagtcccaag ccttccttgc ctgaccaata cccaccaagt caaatcacag 1140
accttgatgc cacagttcat gaggataaga ttattcttac atggacagca ccaggagata 1200
atthttgatgt tggaaaagtt caacgktata tyataagaat aagtgcagat attcttgatc 1260
taagagacag ttttgatgat gctcttcaag taaatactac tgatctgtca ccaaaggagg 1320
ccaactccaa ggaaagcttt gcattttaaac cagaaaatat ctcagaagaa aatgcaacc 1380
acatatttat tgccattaaa agtatagata aaagcaattt gacatcaaaa gtatccaaca 1440
ttgcacaagt aactttgytt atccctcaag caaatcctga tgacattgat cctactccta 1500
ctcctactcc tactcctgat aaaagtcata attctggagt taatatttct acgctggat 1560
tgctctgat tgggtctgtt gkaattgkta actttatttt aagtaccacc atttgaaact 1620
taacgaagaa aaaaatcttc aagtagacct agaagagagt tttaaaaaac aaaacaatgn 1680
aagtaaagga tatttctgaa tcttaaaatt catcccatgt gtgatcataa actcataaaa 1740
ataattntaa gatgtcngga aaaggatact ttgattaaaa taaaaacact catggatatn 1800
ta 1802

<210> 78

<211> 995

<212> DNA

<213> Homo sapiens

<400> 78

cggggtgcct ggggcagacg agggcggctt ctccgcggac agctaggag agtgtcctgg 60
gtgtcagcca gaacatgtct ttcaacctgc aatcatcaaa gaaactgttc attttcttag 120
gaaaatcact gtttagtctt ctggaggcta tgatttttgc cttactccca aagccacgga 180
agaacgttgc tggtgaaata gtcctcatca caggtgctgg aagtggactc ggaaggctct 240
tagccttgca gtttgcccg ctgggatctg ttcttgttct ctgggatatc aataaggagg 300
ggaatgagga aacatgtaag atggctcggg aagctggagc cacaagagtg cacgcctata 360
cctgcgattg cagccaaaag gaaggagtgt atagagtasc cgaccagggt aaaaaagaag 420
tcggcgatgt ttccatccta atcaacaatg ccggaatcgt aacaggcaaa aagttccttg 480
actgtccaga tgagcttatg gaaaagtcac ttgatgtgaa tttcaaagca catttatgga 540

```

cttataaagc ctttctacct gctatgattg ctaatgacca tggacatttg gtttgcattt 600
caagttcagc tggattaagt ggagtaaatg ggctggcaga ttactgtgca agtaaatttg 660
cagcctttgg gtttgctgaa tctgtatttg tagaaacatt tgtccaaaaa caaaagggga 720
tcaaaaccac gattgtgtgc ccctttttta taaaaactgg aatgktgaa ggttgacta 780
caggctgtcc ttctctgttg ccaattctgg aacccaaaata tgcagtgaa aaaatagtag 840
aagctattct acaagaaaaa atgtacttgt atatgccaaa gttgttatac ttcatgatgt 900
ttcttaaaag gtaattacat cagcttctat tacttcccta acatgccagt ctacagtta 960
ctccaaatcc cmccaggaaa aaccactttt aaaaa 995

```

<210> 79

<211> 1215

<212> DNA

<213> Homo sapiens

<400> 79

```

gcaggaccgt cattgacgcc atgagcgcgc tgctgcggct gctgcgcacg ggtgccccag 60
ccgctgcgtg cctgcggttg gggaccagtg cagggaccgg gtcgcgcctg gctatggccc 120
tgtaccacac tgaggagcgc ggccagccct gctcgcagaa ttaccgcctc ttctttaaga 180
atgtaactgg tcactacatt tcccccttc atgatattcc tctgaagggtg aactctaaag 240
aggaaaatgg cattcctatg aagaaagcac gaaatgatga atatgagaat ctgtttaata 300
tgattgtaga aatacctcgg tggacaaatg ctaaaatgga gattgccacc aaggagccaa 360
tgaatcccat taaacaatat gtaaaggatg gaaagctacg ctatgtggcg aatatcttcc 420
cttacaaggg ttatatatgg aattatggta ccctccctca gacttgggaa gatcccatg 480
aaaaagataa gagcacgaac tgctttggag ataattgatcc tattgatgtt tgcgaaatag 540
gctcaaagat tctttcttgt ggagaagtta ttcatgtgaa gatccttgga attttggtc 600
ttattgatga aggtgaaaca gattggaaat taattgctat caatgcgaat gatcctgaag 660
cctcaaagtt tcatgatatt gatgatgta agaagttcaa accgggttac ctggaagcta 720
ctcttaattg gtttagatta tataaggtac cagatggaaa accagaaaac cagtttgctt 780
ttaatggaga attcaaaaac aaggcttttg ctcttgaaat tattaaatcc actcatcaat 840
gttggaagc attgcttatg aagaagtgt atggaggagc tataaattgc acaaactgac 900
agatatctga tagcccttcc cgttcacac aagagggaagc aagatcatta gttgaatcgg 960
tatcatcttc accaaataaa gaaagtaatg aagaagagca agtgtggcac ttccttgga 1020
agtgattgaa acatctgaaa ttctgctgtc aagattccca tctctaagga ctccaagtgc 1080
tagagacaag ggggtctatg agcatttact gacttcctgt taaaacttca ttttttcaa 1140
ctttttgagc tatgcaatat ataaataaac agtaagaatt ttaaattaaa aaaaaaaaaa 1200
agggcgcccg ctcta 1215

```

<210> 80

<211> 2660

<212> DNA

<213> Homo sapiens

<400> 80

```

gcagacttga ttagcctgcc taccactgta gagggacttc agaagagtgt agcttccatt 60
ggcaatactt taaacagcgt ccactctgct gtggaagcac tacagaaaac tgtggatgaa 120
cacaagaaaa cgatggaatt actgcagagt gatatgaatc agcacttctt gaaggagact 180
cctggaagca accagatcat tccgtcacct tcagccacat cagaacttga caataaaacc 240
cacagtgaga atttgaaaca ggatatcctg taccttcaca actctttaga ggaggtaaac 300
agtgccttag tggggtacca gagacagaat gatcttaaac tcgagggaat gaacgagaca 360
gtcagtaatc ttaccagag agtcaacctg atagaaagcg atgtggtgac tatgagcaag 420
gtagaaaaga aagcaaactc gtccttcagc atgatgggtg atagatctgc cactctgaaa 480

```

agacagtctt tggwtcaagt caccaacaga acagatacag taaaaatcca aagcataaag 540
aaagaagata gttcaaattc tcaggatcc aagctaagag agaaactcca gctgatcagt 600
gctcttacia acaaacctga gagcaacmgg cctccagaga ccgccgatga agagcaagta 660
gagagtctca catcaaagcc atcagcattg ccaaaatttt cacagtctct tggagacca 720
gttgagaaag ctgcccact aagacctatc tccctaccag gagtttctag cactgaagat 780
cttcaggatt tattccgcaa gactggccag gacgtggatg ggaagctgac ctaccaggaa 840
atctggacct ccctagggtc tgctatgcca gaaccagaga gcttgagagc atttgattcc 900
gatggagatg gaagatactc attcctggag ctaagggtag ctttaggtat ctacttccat 960
caggcatatt ttagaaatgg actgccta atctattttac ctaacaacaa aacaacctt 1020
acttacctat cagtctctta gtcctccaaa ctactgtagc agatactttg ccacctttta 1080
acttggttga agaagctata taaaagttat ttttttaaag aagaagacca ttttacttat 1140
gatgttcaga aatctatgat ttcctacaac cagtaagatc ttacatttta aaattgccag 1200
aaaaaaaaatt aaagccctct tttttctct ttcctttttt tgaggggagg agaccttatc 1260
ttttaagct gggaaatgta tatagagaga gaataagcca cttttatatt tcaactaaat 1320
ttgccttaaa ttagctgcac tttatagaga ctcaaaaaat gtcttttctt taaaagatag 1380
gccttttctg tttgtaaata tttaaatgaa agaaagcatt gtgcatattg tgtggaaagt 1440
aggaagaatg gttttgaaca ggatatgaac aaatgactta ttaaaaattg ctgatctggg 1500
gtaggtggca gctgaaacta catccatgtc tccataaggy atccctcaaa ggcccaggcg 1560
ctgccagggg gtttgcctg gtagctggag gaaccgattt caggagtag aactggaga 1620
caatactgac tccaggcatg gctcatggaa gtaggattct ggttctttgt tcctattccc 1680
tcagctaata ccaacctggg aatcagagaa gtcttgggga tttttctcat ttttagtact 1740
atttcagggt ttatgagcat aaaaagttat ccattgggga gctccatttt ccctgctgag 1800
tgagctagat tgcttcccc acccaccac ttaagtctgt cttaaagccg tagctggctc 1860
ccaccaccag taccatctcc atttgaatgg cagggtctaa tccccccagc cattatctca 1920
cactgaccac ccagagcttt agaagagagc tgtgcttcta attttgacct agaaaacct 1980
acccttgag attttacct gaagctaacc aagagcctaa tatgtttctc tgggggatga 2040
ctaaagccaa aaaggctgtg agatgaaaca tgtgaaataa tattcagttt ccttaccatt 2100
accagctcag aagtagctag aggttttcta ccaaaggat gccaaagtat agcaggcgag 2160
gcctggagct agggccttca catggtggtg gcaagttttt caaatctaat acaatcaagt 2220
acaatacttc ctttaaatgc ttctgtggac ctggcatgaa agatccctag attgaaagga 2280
ataatacttc catgtctcct gtatgttgag tctagaattg ctgtgttgtt cttagaagca 2340
gtctttgggc aacaacttga aaggggaaaa aaaaactaca aaaacttaac tttggtatag 2400
gccaagtcag ggagaaagta gagaaagctg tcatgccaca gacttcttta gtggagatca 2460
tttctttttt aactttgttc aggttgccct tcaccatgga tacagtccgg tacccttaaa 2520
catttaaggg ctgttttttt tttctttaca ttagtctcag cttggtatta accaaactta 2580
aatttttttt ccagaagtat taaaatttag ttaaagcaaa aaaaaaaaaa aaaaaaaaaa 2640
aaaaaaaaaa gggcgccgc 2660

<210> 81

<211> 1790

<212> DNA

<213> Homo sapiens

<400> 81

ccttgcccta tctgaccaca ctatttaatg tttttatctt aaatttaaaa aactgaaatc 60
tgaaactgta tgatagtcag taggcctaaa actccccctt gactttgaga atttgaaggg 120
attcccttcc atcctgaagt ctactttctt gtcattttaag ggtgagagaa gggaagcttc 180
tgtgcctttc tagaaattaa caaagaacct gttgggtcct agaccagaaa tccagcacgt 240
tatcctagca tgaaactccc ttttaaaaag tgaacacca aagagatgtt acaaatgaag 300
atatagcccc ttcagcatct ctggagacca ataccacaat aaggaataat aacaacctag 360
gaataaaggc aggagttttt acataggatg cacgatatcc aaagggactt tataaacaca 420

agccattgag tcttaagacc tgacacatgg caaggrrtagg acatgtgaga gaaaatggat 480
ggacggttga agaaaagagg ggaaaagacc atgcagttga actaacttct ttttaatttc 540
aaacaggtat tttttgcacg cgtgcgatct ggaggaagta gccaagtgtt tttcatgacc 600
ctcaacagaa attccatgat gaactggtaa cagaagagca cttggcactt atcttcatgg 660
cgttatttct aattttaaag aacataactc atgtggactt atgccagtct agaggcagaa 720
tcagaaggct tgggtgaaca tatcgcttcc cctttttcct ctccctccgc ccctcccagt 780
acagtccatc tttcaatggt gcagcctggt tgagaaggag agaaaaaggt ggcaggaatt 840
tccaggagat cccaagaat gctgccttgt ctgtggacaa agatggacca tgtgcccttc 900
ggaattaggg atagaaacaa atattgtgtg ctcttaacga ttaagctgtg ttatggtggg 960
ttttcaggtt tttacctttt ttctttaccc ctttactctg caagaatggg gaaagaatgc 1020
atactgcgaa aatgagtctt ttaaattctg tctgcctact agttttaagt atatggtatg 1080
ttgtaaaatt tccaatgatg agagacagca caataaatgt accttatctc cttaggctga 1140
aggccataac tacatagtgg agtaatttaa gaactctctt gccttcacca acccaaaagg 1200
ttgctttttg atagcaactg gctaataaat tttttaaaga gaagaaaaat actagttttc 1260
ccctcttttg ggaaatagat tttaaatggc taaactacta gccttaaaac tactagtcta 1320
ataaaaacaa ctaccacttt tgtgaatctg acaggccaca tttttatatg gccctttaca 1380
gaatggagtg tgtgaacag gataactaac ccattgagtt gagctggcct agcgatggag 1440
ggacactcta acacaacttt ccctcagcta ttatgcaaca gatcaggga aaagatggga 1500
tgacagatgg ggtcagacag aaagagcttc tgggaaacaa gcttacatag tcttttttaa 1560
aatgcacaaa gcctcccagc taagagggtca cttggttttg gcttcattag gactgagact 1620
ttgttgagtt ctttctggga cttggagagt ggatgatatt caggctctga acattcccag 1680
cgctctcccg agggtgccac tttctcaaga tgaaaactgt gactgaaaaa attaataata 1740
aatgtttctg agctgcaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1790

<210> 82

<211> 1350

<212> DNA

<213> Homo sapiens

<400> 82

agtgggaggg ccatttcttg ttctctctcc cgctctcgga agctttcgtc tcgtgggtgc 60
gaaaggtaac cgaagcggct caggaaggca gctgtcactg agccctcgga acagagcgag 120
agtatcgtaa gtaaccaggc tcagccggtt tctcaggccg ctctagtcaa ataaaccata 180
aagatcagac tcgggcttct tcaacttctt ctctccgtgg tttcgccatt agcttccggt 240
tccggggagg ggccgagttt tcttcgaaga tttggggctc cgcgatacag ttaggatggc 300
tgtagtacct ctgctgttgt tgggggggtt gtggagcgct gtgggagcgt ccagcctggg 360
tgtcgttact tgcggctccg tgggtgaagct actcaatacg cgccacaacg tccgactgca 420
ctcacacgac gtgcgctatg ggtcaggtag tgggcagcag tcagtgcagc gtgtaacctc 480
tgtggatgac agcaacagtt actggaggat acggggggaag agtgccacag tgtgtgagag 540
gggaaccccc atcaagtgtg gccagcccat ccggctgaca catgtcaaca ctggccgaaa 600
cctccatagt caccacttca cttcacctct ttctggaaac caggaagtga gtgctttttg 660
tgaggaaggt gaagggtgatt atctggatga ctggacagtg ctctgtaatg gacctactg 720
ggtgagagat ggtgaggtgc ggttcaaaca ctcttccact gaggtactgc tgtctgtcac 780
aggagaacaa tatggtcgac ctatcagtgg gcaaaaagag gtgcatggca tggcccagcc 840
aagtcagaac aactactgga aagccatgga aggcactctc atgaagccca gtgagttggt 900
gaaggcagaa gccaccatg cagagctgtg aatctagagg ctctgagcca ctgttaacgc 960
acaatgttca cagacatctg ttgctgcctc accttgggat ccctgccaca agttccttgg 1020
gcagtggcca tgtcaccatt gagatgaaga tatacaacag aaaatagtgg ctgtgttttg 1080
aagcttcagc cctgcacatt tgaactagtc actctcccag acttgcgtgg gtcagttctt 1140
tctgagtaga ggacttgctg gtaaaggggc agatgctttt tattagtact gataaaacaa 1200
actgagggaa acatccctct tagctgggaa acttttactc ttcaggagct tggcatcatg 1260

gactgttaat gtatgtgatt ttccccctat tttctctctc caaaatgata aaaacaataa 1320
ttttaaaaaa aaaaaaaaaa aaactcgagg 1350

<210> 83

<211> 1746

<212> DNA

<213> Homo sapiens

<400> 83

ggcatcagta aggtctgtat ttaaatgtgg atgtagacat cataattacc aagacaagaa 60
attgttttga gaaattctct gatgtttttc ttcttcaggt ttcacgtgcc acgatcatgg 120
tgccacggta ctgcagtatg cacccaaaca gcaactccta atctcggggg gtaggaaaag 180
acacgtctgc atttttgaca tcargcaaag gcagtcatt cacacgttcc aggcccatga 240
ctcagctatt aaggctctgg ccttggatcc ctatgaggaa tattttacca caggttcagc 300
agaaggtaac ataaagggtt ggagattgac aggccatggc ctaattcatt catttaaaag 360
tgaacatgct aagcagtgca tttttcgaaa cattggggct ggagtcagtc agattgacat 420
catccagggc aatcggtctt tctcctgtgg tgcagatggc acgctgaaaa ccagggtttt 480
gccccatgct tttaacatcc ctaacagaat tcttgacatt ctataaagat tgggggtttt 540
tttttatata catttcagtt aaaaggcaca ctacagtcac cactaggcaa ttctgctttc 600
taagcagttg tattgaaaac agagaatctc tgtgtagaat ttgaatatga cccaagctga 660
gtattatcta aacaggttgg tggaatgaat gcgcagtgtac cttattatgc tgacatacta 720
aaaaaaaaataa aacctagtat tgtatgaagg atagctattc ttacagcat ttagcaaac 780
tgattcagaa aacatttgag attagcaaat tagtaacttg aaataatgaa aaggacgttt 840
ataccaaatt aaggaagaaa atgttgctga tttgggtttt tcttcctgtt cttaccactg 900
actgaagcat gcctgcagtc tctcctctgt tggaatgaag gataatcata aggtgtttgt 960
taggagcgct agaccacctg gaaaactttc ttagctgtgg agcagtgcgc agtgaccagt 1020
tctctgctgt gagaggccgt ttccattctt tctgctgaa ttttttcct gttagtgttt 1080
atactgagct agtactgtaa cttgcaaatt agtgcaaatt taaatgcaat gttttactca 1140
caatttgac attcacattt ttgggactgc tagtttttct atttaaatat ttgccttcat 1200
gttaggaatg tactatgtga acatgacata ttgtagtta accaaacaca cttcttagt 1260
ccagtttagt actttttctt ttcgtgtatt caagggttaa cacccaaaca ttttaaggata 1320
tgttgaaact acaccaatag agcatttcat atcataatta aaatgaatgt taggcttctt 1380
gtggccagtt aatagttgat gagattgggt acattattta ttgccacagc ctattgtata 1440
aactatgcag agttaaatat ttgcttgtaa aatattagcc aatgttgtca ttattttgat 1500
gtatttcctt ggttatgacc aaaaatatgt tgagatactg aaactaatgt ctgtgtgttt 1560
aatgttttac cagcaaattg tcttatcatg ttaatgagaa tgttcaatgc ctgtgtgtta 1620
aatagtaaat acaatggcat aaaagtaact ttctctgaag atgtgatgtt caggctgtga 1680
aatatatatg taaaagaaaa ataatgtta tttgttagaa aaaaaaaaaa aaaaaaaaaa 1740
ctcgta 1746

<210> 84

<211> 1491

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (176)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (711)

<223> n equals a,t,g, or c

<400> 84

```
cccacgcgtc cgaatcaaca aagaatctga agttttacaag atgcttcagg agaaacagga 60
ggtgaatgaa cccctgaaac agtctacctc ttccctgatt ttgcaggaaa tcctggagtc 120
tgagataaaa ggggatctca acaaccctca ggattcagaa gtgttaaaac tcctgnaccc 180
traagtgrmt gcatcaattg gaaatgctca gaagggtgcc atgtgtgaca aatgtggtcc 240
tggcattgta ggcattgttg taaagctgcg gggaccatca tccccctccct gaatgttacg 300
tgtgcactga ttgtgacamt cacccgaaaag agaagggtct tttttttttt tttccatcaa 360
atctactgtg aagagcatgc ccaggagcaa gtcacaccac ttgagggtac gatgtgatca 420
tcattgtccc caagtgatcc agcagacctg cactgaccac cgttctccag ccagcctctg 480
ctgcagctag ttctctaaat gttctggcct tgtctcttga aagttctgta cttattcccc 540
caccceccaa cccgcttacc ttggttttat cctgtctcat tgaaccttga gtcccttgct 600
ttgggttaact gactcacact gactgtgtgg tgcccgccca cttttacaat gaatggcaaa 660
ctgtttttgt tcagtgtccc cttgccggca acactgtgta ccttctctcc ntcagtctct 720
ctctgtgca aatggacatc agccacattt gaaccaaatac aaatataata tgtctgacat 780
tgatttcatt tttgtctcat ctatagactc ccagaaaaaa raaawaaaaa aatgcaaaag 840
aaaatttaaa tgaacaaaca gacccttctg gaacctactg tgacatgaga tgtgtccaca 900
aggcgaggtt tcaggcacac actgaaggag catctgtctg acagccctcc ccgcccccaa 960
gtccttacca acctgtgccca ctgcataccc tgtaaaaagt gtgtgggtgt tggtaatgga 1020
ggagttttga agaataagac attrggrgaa aaaatcgacy cctatgatgt aataataaga 1080
atgaataatg gtcctgtttt aggacatgaa gaagaagttg ggagaaggac aaccttccga 1140
cttttttata cagaatctgt tttttccaga tccctattca caatgaccct aatcgcacag 1200
tgattcctca ctgcttttaa gccacatgat ttaagggtggc tgttggatt gttgatgggt 1260
gacaaaataa acactaatgg tttttggaag aaaccagcct taaacctgat ttataaacct 1320
tatcaaatcc gaattattga tcctttcatt atcagaacag cagcttatga actgcttcat 1380
tttccaaaag tgtttcccaa aaatcagaaa cctaaacacc caacaacagg aattattgcc 1440
atttttacat aacctctatg attttaataa aaaaaattat gtaaaaaaaa a 1491
```

<210> 85

<211> 968

<212> DNA

<213> Homo sapiens

<400> 85

```
ctgcagttaa gtgaaaagaa aatccagcct cccctccaa aaaaaaaaaa aaaaaaaaaa 60
atttaatttt taaaaattag tggtatggca ataagacact tcagaggcta tcttaacctc 120
tgaataccca tcttctagtt taaagacaga gacatcccat ctggaaaatg ttaacttgtg 180
ttgccatctc gttgccggag taagtagaca taagacagag tttaagaagt aaaaatatag 240
aaaaattttg atgggtcaca tgagataaat attagaatat tactattcca atgattaaat 300
gaggatcttg aaataaatc tgaagtcttc caatttttac atttattgga ggggtccctg 360
agtctctgca acttttttat ttaagtctct tgctcttatt ttgtgcataa atgttaaacc 420
tccaaaaaat gaaatgttag ctttctttct tttacttttt attaaattta atagaaaata 480
tgacctgagt agttaaaaag tattttgcat tatttgagc aagatgtctc tagcactgct 540
caaagggcaa attttaaaac ttcagtctgg gtgaaagatt tgctagtttt acagaaagat 600
ttgctatctt aaactcaagc tgggttttct gttctcatgt aagtactggt gatgctgtct 660
tatgaattct tccaagggtc tggttggtgaa ataaacatta catgagagct ttcctgtcat 720
ctacactata tggtgtctgg agtggtgaac aaatttattt tagtttctaa gttgtaattc 780
atcctcatat ggtctatacg attttgaatg tgtgccacta catactgaga tgataatgct 840
```


gtacaatttt aagtggtagc agtttctgta tgcagtaggc tgaaatattt tgatgaactg 900
cttaattttt ggatttttatt ttttaagttg tataatttat tttcttgcaa aataaaaaatg 960
taatataa 968

<210> 86

<211> 3068

<212> DNA

<213> Homo sapiens

<400> 86

cgacagaagg ggaggttggt tttataagag gcgtcattgg cgcccgagct gtgaccgccg 60
ccactggggc agccagcaca atcgggcgga ggtggcgtg ccccttcaga cctgaaagat 120
gtctgaaaat tccagtgcaca gtgattcatc ttgtggttg actgtcatca gtcagtggg 180
gtcagatata gaaatgttga attctgtgac cccactgac agctgtgagc ccgccccaga 240
atgttcatct ttagagcaag aggagcttca agcattgcag atagagcaag gagaaagcag 300
ccaaaatggc acagtgccta tggaagaaac tgcttatcca gctttggagg aaaccagctc 360
aacaattgag gcagaggaaac aaaagatacc cgaagacagt atctatatg gaactgccag 420
tgatgattct gatattgcta cccttgagcc acctaagtta gaagaaattg gaaatcaaga 480
agttgtcatt gttgaagaag cacagagttc agaagacttt aacatgggct cttcctctag 540
cagccagtat actttctgtc agccagaaac tgtattttca tctcagccta gtgacgatga 600
atcaagtagt gatgaaacca gtaatcagcc cagtcctgcc tttagacgac gccgtgctag 660
gaagaagacc gtttctgctt cagaatctga agaccggcta gttgctgaac aagaaactga 720
accttctaag gagttgagta aacgtcagtt cagtagtggt ctcaataagt gtgttatact 780
tgctttgggtg attgcaatca gcatgggatt tggccatttc tatggcaca ttcagattca 840
gaagcgtcaa cagttagtca gaaagataca tgaagatgaa ttgaatgata tgaaggatta 900
tctttccag tgtaacagg aacaagaatc ttttatagat tataagtcag tgaaagaaaa 960
tcttgcaagg tggtggacac ttactgaagc agagaagatg tcctttgaaa ctcagaaaaac 1020
gaaccttgct acagaaaatc agtatttaag agtatccytg gagaaggaag aaaaagcctt 1080
atcctcatta caggaagagt taaacaaact aagagaacag attagaatat tggaagataa 1140
agggacaagt actgaattag ttaaagaaaa tcagaaactt aagcagcatt tggaagagga 1200
aaagcagaaa aaacacagct ttcttagtca aaggagact ctgttgacag aagcaaagat 1260
gctaaagaga gaactggaga gagaacgact agtaactacg gctttaagg gggaaactcca 1320
gcagttaagt ggtagtcagt tacatggcaa gtcagattct cccaatgtat atactgaaa 1380
aaaggaaata gcaatcttac gggaaagact cactgagctg gaacggagc taaccttcga 1440
acagcagcgt tctgatttgt gggaaagatt gtatgttgag gcaaaagatc aaaatggaaa 1500
acaaggaaca gatgaaaaa agaaaggggg cagaggaagc cacagggcta aaaaataagtc 1560
aaaggaaaca tttttgggtt cagttaagga aacatttgat gccatgaaga attctaccaa 1620
ggagtttgta aggcatacata aagagaaaat taagcaggct aaagaagctg tgaaggaaaa 1680
tctgaaaaaa ttctcagatt cagttaaatc cactttcaga cactttaag ataccaccaa 1740
gaatatcttt gatgaaaagg gtaataaaa atttggtgct acaaaagaag cagctgaaaa 1800
accaagaaca gtttttagtg actatttaca tccacagtat aaggcaccta cagaaaacca 1860
tcataataga ggccctacta tgcmaaatga tggaaggaaa gaaaagccag ttcaacttaa 1920
agaattcaga aaaaatacaa attcaaagaa atgcagtcct gggcatgatt gtagagaaaa 1980
ttctcattct ttcagaaagg cttgttcttg tgtatttgat tgtgtcaac aagagtccat 2040
gagccttttt aacacagtgg tgaatcctat aaggatggat gaatttagac agataattca 2100
aaggtagatg ttaaaagaac tggatacttt ttgtcactgg aacgaacttg atcagttcat 2160
caataagttt ttcctaaacg gtgtctttat acatgatcag aagctcttca ctgactttgt 2220
taatgatgtt aaagattatc ttagaaacat gaaggaatat gaagtagata atgatggagt 2280
atgtgagaag ttggatgaat atatatatag acacttcttt ggtcacactt tttcccctcc 2340
atatggacc aggtcggttt acataaaacc gtgtcattac agtagtttgt aacatttgta 2400
gattggatag catttttatg atttgatgag tttcttgtaa ggttaccgtt tctaagagtt 2460

```
gtgctttatg gccactgaga gaattcagaa taaattgaaa gatggagtct aaaaattatt 2520
agctgttaca aatggaacat ttcattataa cgtgatcact ttgacttgag caaatggttt 2580
aattttttatc ttaaaaatca gttaagaata tataaaatcc tactttggcc aagtttgttt 2640
cttttcatta tagtttatat gaaaagatca ccttaagtga aattattttc ctttaatctt 2700
ttatgtatgt attcactttt ggaagctagg aatgagcaac acaaatttta ctctgaagtc 2760
agaagagctc atatataata attctaattgt cccacctatt ttcacttgtc cattccatgt 2820
accagcttag ttatgatact tagtcacata attatctttg ataaaggtag aggcacaaag 2880
aggcaacta agcaagtcaa attctaattgt gtgtacttca taataatgtt ttatccattt 2940
tcacttttat attctgtaac atgaaactta cctaactctc aaatgttagc ttcatTTTTT 3000
acctttgaaa tacttaatct ttctgaataa atataatgtg tctataaaat aaaaaaaaaa 3060
aaaaaaaaa 3068
```

<210> 87

<211> 2230

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2227)

<223> n equals a,t,g, or c

<400> 87

```
ggcacagccg tgctccgggt aagatggcag cggacggaca gtgctcgctc cccgcttcat 60
ggcgccgggt gaccctcacc cacgtcgaat atcctgcagg tgatctctcy ggccacctcc 120
ttgcctacct gagcctcagc cctgtatttg tcatcgtcgg ttctgtgacc ctcatcatat 180
ttaagcggga gtgcacacga tctccttcct tgggggcctg gcactgaacg aggggggtcaa 240
ctggctgatc aaaaacgtca tccaggagcc acggccctgt ggaggccccc acacagcagt 300
gggcaccaag tacgggatgc cctccagcca tccccagttt atgtggttct tctccgtcta 360
ttccttcctt ttcctgtatt taagaatgca ccaaacaac aacgccaggt tcctggactt 420
gctgtggagg cacgtgctct ccctgggact cctcgtctgt gccttccatg tctcctacag 480
cagggtctac ctgctgtacc acacctggag ccagggtgct tatggaggca tcgctggagg 540
cctcatggcc atcgcttgt tcatcttcac ccaggaggct ctcaccccgc tgttccccag 600
gatagcagcc tggcctgtct ccgagttctt cctaattccga gacacaagcc tcattcccaa 660
cgtactctgg tttgagtaca cggtaacccg ggcagaagcc aggmaccagc aacgcaagct 720
ggggacgaaa ctgcagtgc cagtgggtgt ggctgggtcc agcctccaga tctggcccgc 780
acgatgcctt gcaggatgga caggatgaca gacaggagc aagcagagac ctctacagac 840
ccaagtcacc aagtggagcc ttttttttct ttattttaat tttaatgaac aaggtggacc 900
aaagggtgta accagcccct caaccaggac cctggggggc ctgctgcctg ggggccgtgg 960
ccagagaccc tcgctgtgct gctgccagcc cctggctggg gcagaaagtg ccctcggcat 1020
gggcacctgg gtgtgggggt gaggaggagg gctcgcgcct ctggtctcgg ggccaggaaat 1080
tccagtggtg gtgagaagta cacactatgt attttttggt tttktcagag gcaggcagga 1140
ttttggagct ggaagaatct gctctccggt ggctgcctg tgaacagagg gctcccggct 1200
agcttcccag gcccttcgcc ctatgccagc aggcagactg cctctccctg ggccgggggt 1260
gcctgggtgc caggaggagg ggagcatacc ccacaccctc cctgccaccg ttgccgttcc 1320
agaacctcgg tcagtgtttc cctgtctggg ggcaggggcc agagcgagca cgcgtctggc 1380
ggctgctgtc gttgtgttct acccgtact gaccacaac cacaagggtt ttctctgggt 1440
ccctgtccct aagacaataa tcgctttctg acaaaggagc ctgcacattt gggtgagcag 1500
acccaagctg ttacagctc tttcttgtcc tgccatccag tagcagttag tcttcatccc 1560
cacgtgaaca aaatgggaag gagccgtgag gagaggagtg aggcaacagg caccggaagt 1620
ccctcgtcct tccctctgtg tgctctgaat atgtccttgt ccttcctgac ccactctcta 1680
```

```
ccagctggga acctgcttgg ggtccccctc aaacctgtgt ctgggggtgtg ggctcacaga 1740
tccctatcag cctgggttcgt gggagggtc ttcctaaagg gacccccatc tctaagtcac 1800
tctgaaaagg agttgtggag aggagacgcc tccagactct cagaagtttt gaggactgaa 1860
ctgggtcact cgggatctgt gttcgaatcc tccccacccc tttctttgtg gagtttccta 1920
acctgctgct gaagcacaat gttttggtgc tttcttttct catttggtta aggcagtgtc 1980
caaaagccat tccagatgcc aagaccaggg gcttatttct agggaaggta ggtcggtttc 2040
catgtttccc tcccgttatt tttatttttt actttttgcc tgagacaagc cgagtatgag 2100
gtgggttgat ttaagaaaaa tcaatgaaat tgtttactac tgttttaaaa taaaaccgta 2160
aactctgaaa aaaaaaaaaa aaaaaaaaaa aaaaactcga gggggggccc ggtacccaat 2220
cgcatcncat 2230
```

<210> 88

<211> 1163

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (159)

<223> n equals a,t,g, or c

<400> 88

```
actgccccaa gctcaaggag atcaatttcc gtgggaacaa gctgaggggac aagcgcctgg 60
agaagatggg cagcggctgc cagaccagat ccatacctgga gtacctgcgc gtcggaggcc 120
gtggtggcgg gaaagggcaa gggccgtgcg agggctcgna gaaggaagag agccggagaa 180
gaggaggggag aggaagcaga ggcgggaagg tggtgatggg gargagcagg acgtgggaga 240
tgccggcccg ctgctgctca gggtcctgca cgtctctgaa aaccccgta ctttgacagt 300
cagagtgagc cccgagggtcc gggatgtgcg gccctacatt gtggggggccg tgggtgcgagg 360
catggacctg cagccaggga atgcactcaa gcgttctctc acctcgaga ccaagctcca 420
cgaagatctc tgtgagaaga ggacggctgc cacccttgcc acccagagc tccgtgccgt 480
caaaggggcc ctgctgtact gcgcccggcc cccacaggac ctcaagattg tccccttggg 540
gcggaaaagaa gccaaaggcca aggagctggg gcggcagctg cagctggagg ccgaggagca 600
gaggaagcag aagaagcggc agagtgtgtc gggcctgcac agatacctc acttgctgga 660
tggaatatgaa aattaccgt gtcttgtgga tgcagacggg gatgtgattt cttcccacc 720
aataaccaac agtgagaaga caaaggttta gaaaacgact tctgatttgt ttttggaagt 780
aacaagtgcc accagtctgc agatttgcaa ggatgtcatg gatgccctca ttctgaaaat 840
ggcagaaatg aaaaagtaca cttagaaaa taaagaggaa ggatcactct cagatactga 900
agccgatgca gtctctggac aacttccaga tcccacaacg aatcccagtg ctggaaagga 960
cgggcccctcc cttctggtgg tggagcaggt ccgggtgggt gatctggaag ggagcctgaa 1020
ggtggtgtac ccgtccaagg ccgacctggc cactgcccct cccacgtga ctgtcgtgcs 1080
ctgacscag ggcgcctgt ccgcgtttgt ttggccggtt ttgcggagggt ttctatgcgg 1140
caatgctgaa ttatccgtta gat 1163
```

<210> 89

<211> 1939

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (20)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (33)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1609)

<223> n equals a,t,g, or c

<400> 89

```
agcaacctat agatcatgan aggcaacggt nanctgacag taccgggtcgg aattcccggg 60
tcgaccacacg cgtccrgcgg taaccgccac agctgccagc gacaggatgg agagcgactc 120
agactcagac aagagtagcg acaacagtgg cctgaagagg aagacgcctg cgctaaagat 180
gtcgggtctcg aaacgagccc gaaaggcctc cagcgacctg gatcaggcca gcgtgtcccc 240
atccgaagag gagaactcgg aaagctcatc tgagtcggag aagaccagcg accaggactt 300
cacacctgag aagaaagcag cggtcggggc gccacggagg ggccctctgg ggggacggaa 360
aaaaaagaag gcgcgcgtcag cctccgactc cgactccaag gccgattcgg acggggccaa 420
gcctgagccg gtggccatgg cgcggtcggc gtcctcctcc tcctcttctc cctcctcctc 480
cgactccgat gtgtctgtga agaagcctcc gaggggcagg aagccagcgg agaagcctct 540
cccgaagccg cgagggcgga aaccgaagcc tgaacggcct ccgtccagct ccagcagtga 600
cagtgcagac gacgaggtgg accgcatcag tgagtggagg cggcgggacg aggcgcggag 660
gcgcgagctg gagggccggc ggcggcgaga gcaggaggag gagctgcggc gcctgcggga 720
gcaggagaag gaggagaagg agcggaggcg cgagcggggc gaccgcgggg aggctgagcg 780
gggcagcggc ggcagcagcg gggacgagct cagggaggag gatgagcccg tcaagaagcg 840
gggcagcgaag ggcggggggc ggggtccccg gtcctcctct gactccgagc ccgaggccga 900
gctggagaga gagggcaaga aatcagcgaa gaagccgagc tcctcaagca cagagcccgc 960
caggaaacct ggccagaagg agaagagagt gcggcccagc gagaagcaac aagccaagcc 1020
cgtgaagggt gagcggaccc ggaagcggtc cgagggtctc tcgatggaca ggaaggtaga 1080
gaagaagaaa gagccctccg tggaggagaa gctgcagaag ctgcacagtg agatcaagtt 1140
tgccctaaag gtcgacagcc cggacgtgaa gaggtgcctg aatgccctag aggagctggg 1200
aaccctgcag gtgacctctc agatcctcca gaagaacaca gacgtggtgg ccacctgaa 1260
gaagattcgc cgttacaaag cgaacaagga cgtaatggag aaggcagcag aagtctatac 1320
ccggctcaag tcgcgggtcc tcggcccaaa gatcgaggcg gtgcagaaag tgaacaaggc 1380
tgggatggag aaggagaagg ccgaggagaa gctggccggg gargagctgg ccggggagga 1440
rgccccccag gagaaaggcg gargacaagc ccagcaccga tctctcagcc ccagtgaatg 1500
gcgaggccac atcacagaag ggggagagcg cagaggacaa ggagcacgag gagggtcggg 1560
actcggagga ggggccaagg tgtggctcct ctgaagacct gcacgacanc gtacgggagg 1620
gtcccagacct ggacaggcct gggagcgacc ggagggagcg cgagagggca cggggggact 1680
cggaggccct ggacgaggag agctgagccg cgggcagcca ggcccagccc ccgcccgagc 1740
tcaggctgcc cctctccttc cccggctcgc aggagagcag agcagagaac tgtggggaac 1800
gctgtgctgt ttgtatttgt tcccttgggt ttttttttcc tgccctaattt ctgtgatttc 1860
caaccaacat gaaatgacta taaayggttt ttttaatgaaa aaaaaaaaaa aaaagggcgg 1920
ccgctctaga ggatccctc 1939
```

<210> 90
<211> 2032
<212> DNA
<213> Homo sapiens

<400> 90
ggcacaggcg gccgcccgcg cgccaagtgc ctcagccctt ggctcctgcc cagtgttttag 60
ggtgttgccg gagaaaaggg gaagagtcac cgcctgtcgg ggctaggata tgatgggtga 120
gaggtgtcaa accaaattct ctcggtttgg aaacggagaa aatctaaaaa tgaggatgtg 180
aggaaagagt ccgctctcaa ggcgcgttgt ggtctatccg agccccgtcc cctgggctcc 240
ctcgggctgg ggtgaggcgg gcagctacgc gtggggtagg accatcttac gctgggaccc 300
cgccaaggag ccccggaag taggtgaaag ggcaggggcg tggctctcgg ggcgccaccc 360
acgctcttga aatctgggtg attgcgagcg gccgctcagc gtccccaca ccacagaccc 420
gcgcgcgcga cgaccagca gccgccatgg ctctgctccg aggtgtgttt gtagttgctg 480
ctaagcgaac gccctttgga gcttacggag gccttctgaa agacttcact gctactgact 540
tgtctgaatt tgctgccaa gctgccttgt ctgctggcaa agtctcacct gaaacagtgt 600
acagtgtgat tatgggcaat gtcctgcaga gttcttcaga tgctatata ttggcaaggc 660
atgttggttt gcgtgtggga atcccaaagg agaccocagc tctcacgatt aataggctct 720
gtggttcttg ttttcagtcc attgtgaatg gatgtcagga aatttgtgtt aaagaagctg 780
aagttgtttt atgtggagga accgaaaagca tgagccaagc tccctactgt gtcagaaatg 840
tgcgtttttg aaccaagctt ggatcagata tcaagctgga agattcttta tgggtatcat 900
taacagatca gcatgtccag ctccccatgg caatgactgc agagaatctt gctgtaaaac 960
acaaaataag cagagaagaa tgtgacaaat atgccctgca gtcacagcag agatggaaag 1020
ctgctaatag tgctggctac tttaatgatg aaatggcacc aattgaagtg aagacaaaga 1080
aaggaanaaa gacaatgcag gtacagcagc atgctcggcc ccaaaccacc ctggaacagt 1140
tacagaaact tctccagta ttcaagaaag atggaactgt tactgcaggg aatgcatcgg 1200
gtgtagctga tgggtctgga gctgttatca tagctagtga agatgctgtt aagaaacata 1260
acttcacacc actggcaaga attgtgggct actttgtatc tggatgtgat ccctctatca 1320
tgggtatttg tctgtccct gctatcagtg gggcactgaa gaaagcagga ctgagtctta 1380
aggacatgga tttggtagag gtgaatgaag cttttgctcc ccagtacttg gctgttgaga 1440
ggagtttgga tcttgacata agtaaaacca atgtgaatgg aggagccatt gctttgggtc 1500
accactggg aggatctgga tcaagaatta ctgcacacct ggttcacgaa ttaaggcgctc 1560
gaggtggaaa atatgccgtt ggatcagctt gcattggagg tggccaaggt attgctgtca 1620
tcattcagag cacagcctga agagaccagt gagctcactg tgacccatcc ttactctact 1680
tggccaggcc acagtataac aagtgcctt cagagcagct gccacaactg gccatgccct 1740
gccattgaaa cagtgattaa gtttgatcaa gccatggtga cacaaaaatg cattgatcat 1800
gaataggagc ccatgctaga agtacattct ctcagatttg aaccagtga atatgatgta 1860
tttctgagct aaaactcaac tatagaagac attaaaagaa atcgtattct tgccaagtaa 1920
ccaccacttc tgccttagat aatatgatta taaggaaatc aaataaatgt tgccttaact 1980
tcaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aagggggagg gg 2032

<210> 91
<211> 1788
<212> DNA
<213> Homo sapiens

<400> 91
ttcccaccgg cttccccaga ggtggaagaa acccgaracg ttccgaagtc aacgcaagca 60
aaggggagtg cgggtcgggg aggaatatc ttttgaaaac gtaatatgtg ccttggggct 120
ctccagccct ttgggacttc caatgggatc ttagaagcag ccgaagcagc gtgagggcgg 180

```

cagccagggc cagccacgat ttgaacgctc tgccttgag ctcttctgga ccgaggagcc 240
caaagcccta ccctcaccat tcaccagggt acagttctta tccgcgtgaa tacacatggc 300
tctgttacga aaaattaatc aggtgctgct gttccttctg atcgtgaccc tctgtgtgat 360
tctgtataag aaagtccata aggggactgt gcccaagaat gacgcagatg atgaatccga 420
gactcctgaa gaactggaag aagagattcc tgtggtgatt tgtgctgcag cagggaggat 480
gggtgccact atggctgcca tcaatagcat ctacagcaac actgacgcca acatcttgtt 540
ctatgtagtg ggactccgga atactctgac tcgaatacga aaatggattg aacattccaa 600
actgagagaa ataaacttta aaatcgaggga attcaaccgg atggctctca aagggaagat 660
cagaccagac tcatcgaggc ctgaattgct ccagcctctg aactttgttc gatcttatct 720
ccctctactt atccaccaac acgaagaaag tcacttattt ggacgatgat gtaattgtac 780
aagtgatgat ccaagaactg tatgacacca ccttggccct gggccacgag gcggctttct 840
cagatgactg cgatttgccc tctgctcagg acataaacag actcgaggga cttcagaaca 900
catatatggg ctatctggac taccggaaga aggccatcaa ggaccttggc atcagcccca 960
gcacctgctc tttcaatcct ggtgtgattg ttgccaacat gacagaatgg aagcaccagc 1020
gcatcaccaa gcaattggag aaatggatgc aaaagaatgt ggaggaaaac ctctatagca 1080
gctccctggg aggggggtg gccacctccc cartgctgat tgtgtttcat gggaaatatt 1140
ccacaattaa cccctgtgg cacataaggc acctgggctg gaatccagat gccagatatt 1200
cggagcattt tctgcaggaa gctaaattac tccactggaa tggaagacat aaaccttggg 1260
acttccctag tgttcacaac gacttatggg aaagctggtt tgctcctgac cctgcaggga 1320
tatttaaaact caatcaccat agctgatata actctacct taaaatattc cctgtataga 1380
aatgtggaat tgtccctttg tagccaacta taacattgtt ctttatgaat attacctttg 1440
atacatatga tccacaatat aaaaaccaa aactactgtg tgcaaattat accttgagc 1500
atataggcat tgattaaact ctttaagtac atgtgataac tatggaaatc aagattatgt 1560
gactgaaaaa cataaaggaa gagacccatc tagataacag caatcaacct gcttaattct 1620
gaatgacaat tatatccaca aatttttaaa acttctacat gtatttttca catgaagatc 1680
tccttaacag gttgccaaac ttttctttta taaaactatt acatttaaaa tatggacgct 1740
tgaaaaataa aatattcatc atttttaaaa aaaaaaaaa aaaaaaaa 1788

```

<210> 92

<211> 495

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (402)

<223> n equals a,t,g, or c

<400> 92

```

gggtcgaccc acgcgtccga ccacattcct cctctaaaga agccccctggg agcacagctc 60
atccaccatgg actggacctg gaggttcctc tttgtggtgg cagcagctac aggtgtccag 120
tcccagggtgc agctgggtgca gtctggggct gaggtgaaga agcctgggtc ctcggtgaag 180
gtctcctgca aggtctctgg aggcaccttc agcagctatg ctatcagctg ggtgcgacag 240
gcccctggac aagggtctga gtggatggga gggatcatcc ctatctttgg tacagcaaac 300
tacgcacaga agttccaggg cagagtcacg attaccgagg acgaatccac gagcacagcc 360
tacatggagc tgagcagcct gagatctgag gacacggcca tntattactg tgcgaraagk 420
ccctmagcgg gtatctatc ccaactacta ccacgggatg gacgtctggg gccaaaggag 480
cacggtcacc gtctc 495

```

<210> 93

<211> 1377

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1367)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1371)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1376)
<223> n equals a,t,g, or c

<400> 93
ggacccacgc gtccgagact tttgaaggaa gaaatcaatt atgatgataa actacaggta 60
agagttgggt gttctaggtc ctgttcttca ggctttttat attcagaaat atgatacatg 120
tataacagct taggcagtc ttcctagtt ggtcacatta gggtgacaac caaagattaa 180
caccaacatg tatttaattg tatgatatt atgtgtccctt ttggcttaac actttcaaac 240
atttttctta tatttggtt tcctagttcc ctattatttt tttaaactctg gaagttgtca 300
ttttttaatt aatttctgtc gaactgaagt gattagaaat atcaacatct ctttcagggt 360
aaaaatattt tgtatcatgc agtcaaagaa atggtgagag ccttgaagat acacgaggat 420
gaagtagasg atatggaaga aaattaagtg tgatccagtt tgatattttt aggttggtga 480
actgggatta cttaaccttg aatgatgata tgtatgcaca ctgacttta gcttcataaa 540
accatcagtg ccaagaaatt ctctttgtag taattacttg ttactgacac cacagcagta 600
tagcatatgt cacagctcct gtgattcaat gttataaaac aagcagaatt ttaaaagcag 660
cactatatag ctgttttgta ttatagtgt tatgatgtt gtgaaaatgc cagattttaa 720
atgatgtatt tatttttgggt aaaaaataaa aaattctatg ctatattgtt gatcaagtgt 780
aaatgtgacc ttgtacagtt tactaaaatt actgatattt ttcactacat tgagacagtt 840
actgtgagaa taggacacaa acaccagcta ttgcctgcat ctgggaaatt gctgaatcgc 900
acagcagtc tgtcataatc agaaaattac tgccaaataa ttgtaaaatt tgtaaagtat 960
aaagtatata aagtagatac taaatacaga cacttcaata ttttgttgaa gctattgact 1020
gtacaattaa acattttcaa aaggtgtaatt ttatttaaaa ttgtctcatt ttggtaaaa 1080
ttatgtgaac ttttaaagct aaatattaaa cttaatatgc tatgtaaata tatacatata 1140
tacatttaat gatgtatttt tttaaaacat tggcttgctt ttgttaaagt gcaagtgtta 1200
catatggctt tgtacattaa agttgaaagg ggttttacct tttccattaa aaggacttta 1260
tcaaaaaaaaa aaaaaaaaaa aaamamamaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1320
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaggggnggc nttttna 1377

<210> 94
<211> 2819
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (82)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (83)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (84)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2816)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2817)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2818)

<223> n equals a,t,g, or c

<400> 94

```
ttttgtattt ttttttagta gagacagggt ttcactgtgt cagccaggat ggtcttgatc 60
tctggcccaa gcttttaaaa cnnnaaaacg gtggccaggc acggcagcac acacttgtaa 120
ttctagcact tcaggaggcc gagatagaag gatcactcga gcccaggaat tcgagaccag 180
ccctgacgat ataaataggg aaagagagag agagagagag agagcgcgtg cacgcgaggg 240
agaatgcctg cccctctcca gcctggccaa cagaaggaga tcctgtctca aagaaagtga 300
aatgcttca gctagaatta taggataatc ctgkgtgaga taatttacat agygactgga 360
agagttatta aatattgctt tcttgytgct aagttctgta gttgggtag atatctaag 420
cgtaactatt gtgaaagtca agagattcta atttttccag aaaataatgt tgagtttcca 480
tgaagaacag gaagtactgc cagaaacttt ccttgcta atcccttctc tgataaagat 540
ggacattcac aaaaaagtaa ctgacccaag tgtggccaaa agcatgatgg cttgcctcct 600
gtcttcactg aaggctaag gctcccgggg agctttctgt gaagtgagac cagatgataa 660
aagaattctg gaattttaca gcaagttagg atgttttgaa attgcaaaaa tggaaggatt 720
tccaaaggat gtggttatac ttggtcggag cctgtgacat ttgttgacac tgtgaactgt 780
ccaaaagtct cttaactgca ccttgtaaat ggtagttgag gtcttcatac agttcagcct 840
ctagaatggg aacaaatcag ccaattggat tcgaaacaaa gaagactatg taaaactcac 900
ccatcacact ttgagactac tcaactggtg gaagaatata gtattgcagc aaatcctgta 960
tgaaagagag atgtgggctt cctttttgag tcttgtgtta ggtgctgaga ccttttacat 1020
gggcttatac agggagagag tcttcaataa atgtagtcag cactattttc tgcattccag 1080
gtggttgctg tcttcacctg agagtaatca agataacatc tgtcatcttc cttggtttat 1140
tgagtgaat gctctcagt cttaggggac atggcagaga tgaaagaaag aaagagtggg 1200
tttcagaagt gtcagggtgg agtgattcca agtgggatgg ttgtggcatt agtttaagct 1260
gaataaataa tttcaatttg gggcagttat tctgcttttt gtaaagccgt ggccaattgt 1320
ctcctgtaat gactgttggt tcaggcatgt tgtactttgt agggacaaat gtgcatttgt 1380
```



```
ttgtggcaaa agcctacaat tgacaaactt gtaaatttct ttgtatataa actagctgta 1440
acctgactat cctttgtggt tactgttttt gtaaatttct ttcctctata aatgaaaggg 1500
tgttggttca gaatggcact ttgaataatg taaaccagtg aaaagtggat tttctttact 1560
tttgtccttg gggttggggg ttgtttttgt ctttttgaag ttttattatt tttaaagtgc 1620
ctcccaccta ggcgtaggcc atgaccattt ggggtacgag agcctaattt tgtaggactt 1680
aatctgttga aaagtgcagt tacttctgga aattaacctc aatattagggt cagcatgtga 1740
aatgttggat ttgacatgtc aggtaggggt cagggactga ttggtcccat ttgccctcag 1800
gtcagttggt taatctcaag acctgttact actgatttta ttaaatacaga gtctttaatt 1860
cttgcatggt tgtatctaatt ttctgaacga atgagcacac ttttaaccagt tatttacagt 1920
tacctttttc ctttaaccgg attgtgaaag cttcatgtat ttttaatttag attctgtggt 1980
tttaaggggt ctgagcatga agctggcaga tagtcggcag gactcatttt ttcacatggt 2040
ctggctgatt tctccataga ttgataacag tattttgtta tcttgcttct ctgtagtttt 2100
gcatcagctg ttttaactttg agctgagtga ggggagaggg gtaaagagaa agaaacttaa 2160
gttttctttc acagaactcc accattgtgg gctttgagag agccctaaag cattgtacct 2220
agtgttacct agtgacttcc aaccaaagcc tttgagtatg cactaaatag gtgagaagaa 2280
aggagagaag gtttttaggt tagaaacctt taaccgatag aaggatatgg tatgttgtaa 2340
agctggaacc aagtttgcat ttttgagggc ttgagatgaa ggggaagactc ttaccagata 2400
gtaagacagc tgagttttcc tcagttttct cgtcttaaca ctagtggaca attctagcat 2460
tttgtttgga ggatttcaga gtttaacctca tggaattcag gatttttttag caagtttgct 2520
tttggtttta tcttggtctt tagtaatcat gttggctggt ctggtcacag gtgactgtga 2580
aacagatgcc ctggtcttgc tttcatcact ctaggatcat gaagtgcatt gctatttcct 2640
ggttatgaat attaaggttg gaattacatt tttattgatt gtttgatca gagctcagtt 2700
cctgtagaaa acgaactgta aaagaccatg caagaggcaa aataaaactt gaagtgaatg 2760
cttaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaannna 2819
```

<210> 95

<211> 705

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (488)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (682)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (684)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (687)

<223> n equals a,t,g, or c

<400> 95

gcggcctggc taactcctgc caggcagtgc ctttcccga gcgtgccctc gccgctggtg 60
cggctcctac ctggtctgga aagagctggg aggttcaca gagaaggctg tggttccct 120
gggctctac actgggcagc tggccctgaa ctgggcatgg ccccccattc tctttggtgc 180
ccgacaaatg ggctgggcct tgggtgatct cctgctggtc agtggggcgg cggcagcact 240
accgtggcct ggtaccaggc gagcccgctg gccggccgcc tgctctacc cttacctggc 300
tggctggcct tcgcgaccac actcaactac tgcgtatggc gggacaacca tggctggcgt 360
gggggacggc ggctgccaga gtgagtggc ggcccaccag ggactgcagc tgcaccagca 420
ggtgccatca cgcttgatgt gtggtggcgc tcacgcttc atgaccactg ggcctgctag 480
tctgtcangc ccttgggcca ggggtcagca gagcttcaga ggtggcccca cctgagcccc 540
caccggggag cagtgtcctg tgctttctgc atgcttagag catgttcttg gaacatggaa 600
ttttataagc tgaataaagt ttttgacttc ctttaaaaaa aaaaaaaaaa aagggggggc 660
cgtaaccaat tggccctaaa gngngtngtt taaaattaat ggccg 705

<210> 96

<211> 3472

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (55)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (69)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3457)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3466)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3470)

<223> n equals a,t,g, or c

<400> 96

attaaagcca gaaccagatg atctcattga cgaagacctc aactttgtgc agganaatcc 60
cttatctcng aaaaaaccta cagtgcactc tacatatggt tcttctcgcc cttctattga 120
aatttatcga ccacctgcaa gtgaaatgc agatagtggc gttcatttaa acaggttgca 180
atttcaacag cagcagaata gtattcatgc tgccaagcag cttgatatgc agagtagttg 240
ggtatatgaa acaggacggt tgtgtgaacc agaggtgctt aacagcttag aagaaacgta 300
tagtccgttc tttagaaaca actcggagaa aatgagtatg gaggatgaaa actttcggaa 360
gagaaagtgt cctgtggtgaa gttcagttgt taaagtaaaa aaattcaatc atgatggaga 420

```

agaggaggaa gaagatgatg attacgggtc tcgaacagga agcatctcca gcagtgtgtc 480
tgtgcctgca aagcctgaaa ggagaccttc tcttccacct tctaaacaag ctaacaagaa 540
tctgatattg aaggctatat ctgaagctca agaatccgta acaaaaaaa ctaactactc 600
tacagttcca cagaacaga cacttccagt tgctcccaga actcgaactt ctcaagaaga 660
attgctagca gaagtgggtc agggacaaa taggaccccc agaataagtc cccccattaa 720
agaagaggaa acaaaaggag attctgtaga aaaaaatcaa ggaactcaac agaggcaatt 780
attatccccg ctgcaaatcg acccagtaat ggcagaaact ctgcagatga gtcaagatta 840
ctatgacatg gaatccatgg tccatgcaga cacaagatca tttattctga agaagccaaa 900
gctgtctgag gaagtagtag tggcaccaaa ccaagagtcg gggatgaaga ctgcagattc 960
ccttcgggta ctttcaggac accttatgca gacacgagat cttgtacaac cagataaacc 1020
tgcaagtccc aagtttatag tgacgctgga tgggtgtccc agccccccag gatacatgtc 1080
agatcaagag gaggacatgt gctttgaagg aatgaaaccc gtaaaccaaa ctgcagcctc 1140
aaacaaggga ctgagaggtc tcctccaccc acagcagttg cacttgctga gcaggcagct 1200
tgaggacca aatggttagt tttctaaccg tgagatgagt gaactgagtg tggcacagaa 1260
accagaaaaa cttttggagc gctgcaagta ctggcctgct tgtaaaaatg gggatgagtg 1320
tgccatccat caccctatct caccctgcaa agccttcccc aattgtaaat ttgctgaaaa 1380
atgtttgttt gttcacccaa attgtaaata tgatgcaaag tgtactaaac cagattgtcc 1440
cttcactcat gtgagtagaa gaattccagt actgtctcca aaaccagttg caccaccagc 1500
accaccttcc agtagtcagc tctgcccgtta ctccctgct tgtaagaaga tggaatgtcc 1560
cttctatcat ccaaaacatt gtaggtttaa cactcaatgt acaagaccgg actgcacatt 1620
ctaccatccc accattaatg tcccaccacg acatgccttg aaatggattc gacctcaaac 1680
cagcgaatag caccagtcct tgccctggcag aagatcatgc agtttggaag ttttcatgta 1740
ctgatgaaag atactctaca gaacttgta aatctttgaa acttggaata tattgctttc 1800
ataatatgaa gttttattgc ctatctatct gaagtgtcta atttttcaag tttgtaagtt 1860
tattatgtgg ttttaacatt ggggtgtttt gttttgtttt tactatgaaa agacagctta 1920
aggaagagct aaattctggt aaaatatttg gggcatgttt gtgcactgct gttgtgagga 1980
tcagcatatg aaattgacat catgggttagt catggtagtg cagcttaggg ggctacacgg 2040
ttgctgtgtg agtgagagag tgcaagtggg cagttgtcat tattctaaaa attgtactac 2100
tttacttttt cccaaagatt atataatgtt cataatccac catgaaaaca gcattggcca 2160
aaggtagtga ggctgcttaa aatattcaat tctgcttttt aatttttaag tgaatttagt 2220
ttgaaaagca tgattataca ggcctctcag gctgagtgtc actttggtaa agttcccagt 2280
tttctgcctt tctgtgacag gatgaatgag gtgggtatgg acagtggagg cagctggaat 2340
ggcaagtgca gaaaatagga acagttctat acagtgtctc catttactaa taacataatg 2400
ccttctaaat aatttttttg ggaaactaca ttatcacaaa attatacaaa tttttttaca 2460
agtatttaca tactgtatct gaaaacagac tttaaagtca caagattata aatgtacata 2520
tgtatttcca cattctgaaa aataacattc tcagaatcca cagaaaaat acttagttac 2580
tactgaagat aatttttgaa atgtaaaaat tagattttaa tagtatattt taaatgacag 2640
aactataatt acagagatca gatcagatag gtaaaactgca agatagatag gatgaaactt 2700
ttggcctact gtattactta cagagttttt ttgtgtgtgg tttttaaaac tgtaaggca 2760
agaagtgtca aatgcttttag agttaataa cagatcactg atttcaaaga cttgggtgtat 2820
agtgttaaaa attaaagctt aaaaggtggt tagaaaagtg gattaatgca aaaggggtaa 2880
taaagactgc aacattctca ggaccaaatt aaactgctaa aaaaaaaaaa aaagttcatt 2940
gacttgctta gtogtatact caaatgatga taaacctaca tgtgcaaagg ctcacgttta 3000
agattgtcaa gccagcagtc tactgttgtg ttgccattgc ttttccattg ggagaagaaa 3060
gaattaacca gtcattaaac catttggtaa gttgcacttt gctgtgtgta tcccacagga 3120
aaggcttgaa acacgagaag cagcaaagac agagcacaca agtgcataag gctgtgtgtc 3180
tcggcttggg tgaaatgaca gttcctcttc attctaaagg tttactccat tgaatttaag 3240
gcatttgctt attccagtgt tgagatgctt tgcactctct cagaagaaat ttattttaaa 3300
ttgtttaaat atctggaaat acttttagct atcatttata aagatagttt tgttctcagt 3360
ttcactataa attatagaac aaatgggaaa caagggttta atttagttca gccattttac 3420
aaggaaataa taaaatacta aaatctggak tgttcmntta ccttttcccc cc 3472

```

<210> 97

<211> 1216

<212> DNA

<213> Homo sapiens

<400> 97

```
ggagcgtgac ttcattctcaa aaaaaaaaaa aaaaaaatta ttctgtacca tcacarcctt 60
tcacaacgat ggcaagcctt atgtcttggg agcctgtttt gctaggcaaa gttacaagt 120
acctaattggg agctcaaagt tgtgtgtgtc tctctgtgtg tttgtgtgtg tgtgtgcaact 180
caagacctct aacagcctcg aagcctgggg tggcatcccg gccttgccat tagcatgcct 240
catgcatcat cagatgacaa ggacaaccct catgacgaag caacatgaat tagggggcct 300
cttggccttg gtccaaaatt gtcaatcaga aatgaacata aaggactcca gagcagtggg 360
actgtctgtc aaaagactct gtatatcttt tgtggatgag ttttgtgaga gaacagagag 420
accattgtac ctggcacaag ggctsttcat gaaaagggag acttactggg aggtgcaaga 480
cagtggcatt tctcctctcc tcttgctgct cagcacagcc ctggattgca gccccgaggc 540
tgagaccaga caaagcccgg gaggcagaaa gatgctccaa gaaccaacac tatcaatgtc 600
tttgcaaatc ctcacaggat tcctgtgggt ccagctttgg aactgggaaa ccttctctcg 660
gatccgcact cattccactg atgccagctg cccctgaagg atgccagtac tgtggtgtgt 720
gagtctcagc agccgcccac acgctcctaa ctctgctgca tggcagatgc ctaggtggaa 780
atagcaaaaa caaggcccag gctggggcca ggccagagg ggaaggccct ggattctcac 840
tcatgtgaga tcttgaatct ctttctttgt tctgtttgtt tagttagtat catctggtaa 900
aatagttaaa aaacaacaaa aaactctgta tctgtttcta gcatgtgctg cattgactct 960
attaatcaca tttcaaattc accctacatt cctctcctct tcactagcct ctctgaagg 1020
gtcctggcca gccctggaga agcactgggt tctgcagcac ccctcagttc ctgtgcctca 1080
gccacaggc cactgtgata atggtctgtt tagcacttct gtatttattg taagaatgat 1140
tataatgaag atacacactg taactacaag aaattataaa tgtttttcac atcaaaaaaa 1200
aaaaaaaaaa aaaaag                                     1216
```

<210> 98

<211> 1186

<212> DNA

<213> Homo sapiens

<400> 98

```
ggcacgagaa ataatcacct ggagtttggt aaaccatatg gattctcagg ctctctctct 60
gaagattctg attcagtagg tctgggagtg gcgccttggg ttttgatcaa aattgtagag 120
cattttaagg tgagtacctg agggagaact taaagacatc ttagttgggg agtagtcct 180
ttgaatttta cagctagata taatcttcag tcagataaaa tttatgggag ctggtgtctt 240
atgcctgact cttagtaatt tcataccggt ttgaagtacg tgtgcccag cctaaagcct 300
tgactttcag aatgttgtct tttgattctt ctgtcttgat ttgattaggg gtgaaattta 360
gaagtcttag taatgtaact tgaagatggt aaacaaaaat ctcaagtaaa atgaaaagca 420
aatatgggct actgaattaa gaaactggca ttctagtatt aaatcctcac ttcaggagct 480
tttaaaaaata ctgagacccc ccataacca gagattcaga ttcaaagact gaggatagga 540
ccttagcatt gtagctatct aaagtttcta atgtgcaccc agggttggga atcaccaatg 600
tgggtgtgaa aatgcctaca aagggtttta gtgccttaga agtcctaaga agcccaatct 660
gtatcaaagc agatccattt tgcaaggatc tttcttttag aactttctca gttctcttag 720
taagaacttt agaagtaatc ttgataataa gcacagacag cctaacagca gaggcaactt 780
aaataactcc tgagcagttg gcactagaac agaatacttg gaatgacacc aaagttaacc 840
aagtccagca tatgtccaaa gagttaagt tttcatttac tgtagcattc tgggtgagaa 900
attggttgct gaaatcttaa gacagtgtgc tcaaccttgg ctgcacattg gaatcacctg 960
```

tagggtttta aagcatccaa atggtaatta acaggcagca aaacttcaga actagttctg 1020
catctactgt gcaaagatca tgattaactg tcaagacact ggtagaacag aacaagcaaa 1080
agattaagag ttcaaaagta aatgcaacca wtttaacatg tagtgttatt aaaaaattac 1140
aaaggcctag accagcctgg gcaacagaga ccatgcttaa aaaaaa 1186

<210> 99

<211> 1120

<212> DNA

<213> Homo sapiens

<400> 99

aattcggcac gaggtgacca ggagtcgacg tgtgcagaag tcctggtaat ctggtccttg 60
ttcccgctctg gataccagct tccttcagca gcgcaggcgg tggccctga ggcccgctga 120
aggagtcaaa cttgcgggaa ttttgcagtt tatctgcagg gctgttggtt ccagcaagac 180
ccaaagctag aaaaggagga ggaagaaact gacccgatca gtgccagaag tcattgtatt 240
caaagaagaa taagcaagaa agaaaagaag gaaggaagag aggtagacag atacaagatg 300
aaatcctgtc aaaaaatgga aggaaaacca gaaaatgaga gtgaaccaa gcattgaggaa 360
gagccaaagc ctgaggaaaa gccagaagag gaggagaagc tagaggagga ggccaaagca 420
aaaggaactt ttagagaaaag gctgattcaa tctctccagg agtttaaaga agatatacac 480
aacaggcatt taagcaatga agatatgttt agagaagtgg atgaaataga tgagataagg 540
agagtcagaa acaaacttat agtgatgcgt tggaaaggta atcgaaacca tccttaccct 600
tatttaattgt agtttacctt gatttttatac tgatattaac aataccatat agcttgcttt 660
ttattagcat ttcctgatat tcctttgtcc atattttctac ttataacctg ttgctattaa 720
tggtttttaga tgtatctctt gttatctgca tctcattggt tattgtattt tgaaccaatc 780
tacaagtctc tgtcttttaa taaaagaact ttacacattt gtaaaaaaga ggttcttggt 840
aagatataaa atggaaaaag gctaagtaat atgtgaatat catatttttg aaaggtaaaa 900
agtacatttg tatattacat atatggacat aacttgtaga ggatgaaaga aagtacagcc 960
tctcgggtggt gggattatga atgatttttc tccttttgct tgtttgattt ttctatattc 1020
ctaaaattaa cacacattat tattgctaga ataataaaag ttttataaaa aagaagcaaa 1080
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa agggggggga 1120

<210> 100

<211> 1225

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (286)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (288)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1213)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1225)
<223> n equals a,t,g, or c

<400> 100
cttgctctgt caccagggt gggagtgcag tggcatgac tctgcaacct ctacctccca 60
ggttcaagca attcttgtgc ctcagcctcc caagtaactg ggactacagg tgcacgctac 120
cacacctggc tgattttttt tatgttttag tagagacagg gtttcaacca tgttgcccag 180
gttggtctca aactcctgag ctcaggcaat ccaccgcct tggcctccca aagtgcagg 240
attacagggt tgagccacca caccagcta ttttttcttt cggttnnnaa ttttaaagtt 300
gggggggggtc tcaatttgtt taccctgggt ggtctcgaac tcccggactt aagcgatcct 360
ctggctccaa gccactacc agtctcaggt ttctttacta aaagatcact accttttttt 420
ctcttatctg ctgccatgtg agatgtggct ttcaccttc gccatgattg tgaggccttc 480
ccagccacgt rgaactgtaa gtccaataaa cctcttttgt aaattaaaaa aaaaaaatca 540
ctatttaaga tactaggatg gattgtgact gttgaggagt acttacatat cctacatttg 600
actacattat ttccaaacca agtattccat ccaaaggaac atactgctat catagagacc 660
aaggagggac tgtttaaggt tgccaagggt aagcgagctg agaggctttg tcctcgtgcc 720
agtaactctg aaatctctct taattcctgc tgtccaggca gcagaatgcc atgggttccc 780
caagtaggta gctgcttttag cakttaaagc ccaaagtctt gttctgttga tcagaggtct 840
ctgaatttct gaagtgggtg ttcgtttctg gtgactgagt taatccttta caatccctct 900
tgtaaagtgt gctaatagaa agaatccacc tttcaaagct gcagaaccag accgtgccct 960
aaattgacca acgtarctga tgtgcctcag gaagtctctt gccagctgtc cctgtgaaga 1020
ccccctctc cccccagct gctgccttgc acactgaagc atctcagact gtgcaaaagcc 1080
rtgtagtcat caagacagta aatcccaggg cttgggttaag tgctgtgtga taacttgttt 1140
ggatgagact taacttaaaa ccacttacia taaacttggg aaactaccgt caaaaaaaaa 1200
aaaaaaaaat ttnggggggg cccgn 1225

<210> 101
<211> 1213
<212> DNA
<213> Homo sapiens

<400> 101
ggtagataaa ttgactagct gataagtaaa attaaaagag gaattttctc tttcaggtag 60
gattattgac tgtgctttta ctgtcacttt taatcccaaa tatgatacgt tattaagaagc 120
tgtaaaagat gctactaaca ctggaataaa gtgtgctgga attgatgttc gtctgtgtga 180
tgttggtgag gccatccaag aagttaggta gtcctatgaa gttgaaatag atgggaagac 240
atatcaagtg aaaccaatcc gtaatctaaa tggacattca attgggcaat atagaatata 300
tgctggaaaa acagtgccga ttgtgaaagg aggggaggca acaagaatgg aggaaggaga 360
agtatatgca attgaaacct ttggtagtac aggaaaagggt gttgttcatg atgatattga 420
atgttcacat tacatgaaaa attttgatgt tggacatgtg ccaataaggc ttccaagaac 480
aaaacacttg ttaaatgtca tcaatgaaaa ctttggaacc cttgccttct gccgcagatg 540
gctggatcgc ttgggagaaa gtaaatactt gatggctctg aagaatctgt gtgacttggg 600
cattgtagat ccatatccac cattatgtga cattaaagga tcatatacag cgcaatttga 660
acataccatc ctgttgcgtc caacatgtaa agaagttgtc agcagaggag atgactatta 720

aacttagtcc aaagccacct caacaccttt attttctgag ctttgttgga aaacatgata 780
ccagaattaa tttgccacat gttgtctgtt ttaacagtgg acccatgtaa tacttttattc 840
catgtttaaa aaagaaggaa tttggacaaa ggcaaaccgt ctaatgtaat taaccaacga 900
aaaagctttc cggactttta aatgctaact gtttttcccc ttctgtgcta ggaaaatgct 960
ataaagctca aattagtttag gaatgactta tacgttttgt tttgaatacc taagagatac 1020
tttttgata tttatattgc catattctta cttgaatgct ttgaatgact acatccagtt 1080
ctgcacctat accctctggt gttgcttttt aaccttcctg gaatccattt tctaaaaaat 1140
aaagacattt tcagatctga gaaaaaaaaa aaaaaaaaaa aaaaaaacyc gggggggggc 1200
ggtmccaat tcg 1213

<210> 102

<211> 1564

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1509)

<223> n equals a,t,g, or c

<400> 102

ctgatgcagt aatatattca gtttatctat gcattgctgg ggtcttgata taagatggaa 60
acagtgtttc ttatctaaag ttttgattat cagcaagttg aatggacact ttagttattt 120
aaataaagat ttttgaccaa aatccagaaa atgggtatgag agaaaaataa attccagcta 180
gtttaataga tttttaaatg tcaatctgat ttttagctctt agagagtgtc aactagctkg 240
taatgtttac ttttaattcc ttgttaaaat gaggcaataa tttgcaagat tttgttatat 300
aagtgtaaaat tcttcataac tttttagatc taattcaata ttttctgact acttctgcc 360
tgtataaac catttttgtg catgcttgga tgtgacattc ataaattgta ccactatgac 420
tttatccatg taaaatagct atttattgaa ttttctttta aaagctggat ttactgggtt 480
tttttttct ctcagtttaa acatcttaac agaraawttg ttactgttta ttaaaagaat 540
tgcatttgga agcaagaaaa caaccggatt tgcaatcatg tgaagaaaca ataatgcctt 600
tattatcaag tgtaagcac aattcttata acaaactgtg ataaattctt ttgtttttt 660
tttcttttg cccattattt tcttatgaac aaaccaaaaa ttcattggtg gcagttgcag 720
tgttggtgga tatatctttt atgtacaggg aatttgaaaa ggacagtgga ttcatttaga 780
agtgtaaact gtgctgtgat tatagcaata catttgtag ttgtacttca tctttttcat 840
gctagctttt taaatgttta gttttcctct tgtcatggtc agctgctgaa ttacttgaa 900
ggatgtagaa tactgtttaa aaaatactaa aatttgatac attagatcaa agaattgtgc 960
aatcatttcc tttttaattt ttaaaatgtt gaggtcata aatatttgag aacatcagat 1020
ctaataagac atagtgtatc tatttaatta accaaagtct ctagtgaata tttcaacttt 1080
gaatgtaaac taacaaataa acctgaccac caaggagatt gtttgcccag agtttcaaag 1140
cacattgtct acaaatggaa attgaaataa tttataaaat attgacgtta ctatgttttt 1200
taaaaagtgc ctaatttttt cactaaatgg aggaactat tagttttatt gttaaataatg 1260
gtagatatta atattcctct tagatgacca gtgattccaa ttgtccagat ttgaaataag 1320
taccctgtga gtatgagata aattagtgtc aatcagaaca agtttcagta tcagatgttc 1380
aagaggaagt tgctattgca ttgattttta tttttgtaca taaacactga tttttttgag 1440
cattattttg tatttggtgt actttaatac ctggtgtaca gttccagaaa taaaaatctg 1500
ggaatcttna aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa gggcgccgc 1560
tcta 1564

<210> 103

<211> 1457

<212> DNA

<213> Homo sapiens

<400> 103

```
aattcggcac gagccgttca agggctctta gaggtggcca aagactcaat cccccaagt 60
cactggaaaa agacccaggt ggtcctaaag gcaacagcag gactacgctt actgccagaa 120
cacaaagcca aggctctgct ctttgaggta aaggagatct tcaggaagtc acctttcctg 180
gtaccaaagg gcagtgttag catcatggat ggatccgacg aaggcatatt agcttggtt 240
actgtgaatt ttctgacagg tcagctgcat ggccacagac aggagactkt ggggaccttg 300
gacctagggg gagcctycac ccaaatcacg ttctgcccc agtttgagaa aactctggaa 360
caaaactccta kgggctacct cacttccttt gagatgttta acagcactta taakctctat 420
acacatagtt acttgggatt tggattgaaa gctgcaagac tagcaaccct gggagccctg 480
gagacagaag ggactgatgg gcacactttc cggagtgcct ktttaccgag atgkttggaa 540
gcagagtggg tctttggggg tgtgaaatac cagtatgggt scaaccaaga aggggaggtg 600
ggctttgagc cctgctatgc cgaagtgtgt aggggtgggtc gaggaaaact tcaccagcca 660
gaggaggtcc agagaggttc cttctatgct ttctcttact attatgaccg agctgttgac 720
acagacatga ttgattatga aaaggggggt attttaaaag ttgaagattt tgaaagaaaa 780
gccagggaag tgtgtgataa cttggaaaac ttcacctcag gcagtccttt cctgtgcatg 840
gatctcagct acatcacagc cctgttaaag gatggctttg gctttgcaga cagcacagtc 900
ttacagctca caaagaaaag gaacaacata gagacgggct gggccttggg ggccaccttt 960
cacctgttgc agtctctggg catctcccat tgaggccacg tacttccttg gagacctgca 1020
tttgccaaca cctttttaag gggaggagag agcacttagt ttctgaacta gtctggggac 1080
atcctggact tgagcctaga gatttaggtt taattaattt tacacatcta atgtgaactg 1140
ctgcctaacc actcaagagt acacagctgg caccagagca tcacagagag ccctgtgagc 1200
caaaaagtat agttttggaa cttaaccttg gagtggagag ccaggggacag gtccctggaa 1260
accaaagaaa aatcgcattt caaccctttg agtgcctcat tccactgaat atttaaattt 1320
tcctcttaaa tggtaaactg acttattgca atcccaagac ccatcaatat cagtattttt 1380
ttctcccta tacagtggcc tgcccacct tatctgcacc cacctccctt gaaaaagaaa 1440
aaaaaaaaa ctcgtag 1457
```

<210> 104

<211> 785

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (748)

<223> n equals a,t,g, or c

<400> 104

```
gtgctgatga aaatatggag gaaatgaaac tctcatgggt tttgcaggga atctaagtca 60
gtgctgtgtt gtgaatgtag gtgtaccctt tgaattcata tgttgaatcc taaccccaa 120
agcaatggca ttaagaggtg gggccttttg ggctgggata caggagttag ccactgcgcc 180
tggctgatcc cagcactttt caaatgatgc cgctcaaagc cgtgacttgg cctactttga 240
acagcaaaact tgttgctgct gttgtcaacc tgaaggcctc tcaaatgcca gcttcaagca 300
gggtgtgaat tggccagtgt cagatctcag gagtctgtg ttgagagtgt ggctttcagc 360
tgcggggagc tgcacttggg ggggaaagcc aggcaggtca ccctcacagc cagataatgt 420
ggagggtcaga acccaaggaa gggagtgaga cctccactcc cagtggggga cctggccacc 480
catccttggg gacctgagaa agcgtacttc accttggggg gaaggctggg tggggccaga 540
gggaccagtg ccctcctcag tgcttagggg cagagccacc tgcagcaatg gtatctgcat 600
```


attagcccct ctccaccttc tttctcccg c tgaatcattt ccctcaaagc ccaagagctg 660
tcactgcttc tttctccctg ggaagaatgc gtggactctg cctggtgata gactgaagcc 720
agaacagtgc cacaccctcg ccttaatncc ttgctaggtg tctcagattt atgagacttc 780
ttagt 785

<210> 105

<211> 921

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (40)

<223> n equals a,t,g, or c

<400> 105

aggggaaggac cctcccctag agtctctgga ggggaactggn atacaattgc agagtgcact 60
aaacagttgc ccagaaaga catgtcttgt tttaaagccc agaacctgaa attattatag 120
atthttattcg gtaataagga actttgcatg tgtaattact taaggatatg aagatgagat 180
tgtcctggat tattaagcac cctaaatgcc atgtacaggt gtccttccaa gagaacagaa 240
gaggagacac agacacagag caggaggaca cgtggagaca gaggcagact ggagtgatgc 300
ggccacaagc ccagggacac ctggagcccc caggagctgg gagaggcagg aaggatcctc 360
ccctagagcc tccaggggga actggaggat gcgtaagaga ccagaaactt ccacagaagg 420
aggaaaatta acctcctgct tctctagact gttccaaaagc tgaaccctag aaagcaaagc 480
tgatacagaa gcattccaggc tgcaggagta caggctcgaa gtgctgagcg tgggccttg 540
gtgtgtctca tgggggaaaa aaaactgtga aaaacctcag agtagcatct tcacagtaac 600
gcacggacga tccctaaact gccttgtaaa caaaaatgag agcttgagtc agaggaagcc 660
gagacaatat ccttcctcga caacgtgcga gaacctgac gtccccagc aaaggaagac 720
gttgcaagca ggcaaaatgc gtcgattttt tttttttgtc agtatgatga tttttgcagc 780
cacttggtcta tggagagcag ccgacacccc ctcttacagc cgtggatgtt tcctggaagc 840
tgactcagtc tgttcactgg ttgagctttg agtgaaaaga taacacaggt ctattgactc 900
acacacatgt ttttaagatgg a 921

<210> 106

<211> 592

<212> DNA

<213> Homo sapiens

<400> 106

gtgatctgca tgtggcaggg ctgcgcagtg gagcgccag tgggcaggat gacgagccag 60
acctctctgc ccagtcctcc cgggccagc cggccgacga tgtctactgt tgtggagctg 120
aacgtcgggg gtgagttcca caccaccacc ctgggtaccc tgaggaagtt tccgggctca 180
aagctggcag agatgttctc tagcttagcc aaggcctcca cggacgcgga gggccgcttc 240
ttcatcgacc gccccagcac ctatttcaga cccatcctgg actacctgcg cactgggcaa 300
gtgccacac agcacatccc tgaagtgtac cgtgaggctc agttctacga aatcaagcct 360
ttggtcaagc tgctggagga catgccacag atctttggtg agcaggtgtc tcggaagcag 420
tttttgctgc agtgccgggc tacagcgaga acctgggagc tcatggtgcg cctggcacgt 480
gcagaagcca taacagcacg gaaktccagg tgyttgtgtg cctggtgaaa cttgagggag 540
caggatgcat tattattcag aggtcctgtg tttttcttgc aggataagaa gg 592

<210> 107

<211> 2248

<212> DNA

<213> Homo sapiens

<400> 107

```
tctgggagta cgacatggcg cgcgagctgc gggcgctgct gctgtggggc cgccgcctgc 60
ggcctttgct gcgggcgcgc gcgctggcgc ccgtgccggg aggaaaacca attctgtgtc 120
ctcggaggac cacagcccag ttgggcccga ggcgaaaccc agcctggagc ttgcaggcag 180
gacgactgtt cagcacgcag accgccgagg acaaggagga acccctgcac tcgattatca 240
gcagcacaga gagcgtgcag ggttccactt ccaaacaatga gttccaggcc gagacaaaga 300
agctttttgga cattgttgcc cgggccctgt actcagaaaa agaggtgttt atacgggagc 360
tgatctccaa tgccagcgat gccttgaaa aactgcgtca caaactgggt tctgacggcc 420
aagcactgcc agaaatggag attcacttgc agaccaatgc cgagaaaggc accatcacca 480
tccaggatac tggatatcgg atgacacagg aagagctggg gtccaacctg gggacgattg 540
ccagatcggg gtcaaaggcc ttcctggatg ctctgcagaa ccaggctgag gccagcagca 600
agatcatcgg ccagtttggg gtgggtttct actcagcttt catggtggct gacagagtgg 660
aggtctattc ccgctcggca gccccgggga gcctgggtta ccagtggctt tcagatggtt 720
ctggagtgtt tgaatcggc gaagcttcgg gagttagaac cgggacaaaa atcatcatcc 780
acctgaaatc cgactgcaag gagttttcca gcgaggcccg ggtgcgagat gtggtaacga 840
agtacagcaa ctctgcagc tccccctgt acttgaatgg aaggcggtg aacaccttgc 900
aggccatctg gatgatggac cccaaggatg tcsgtgagt gcaacatgag gagttctacc 960
gtacgtcgc gcaggctcac gacaagcccc gctacaccct gactataag acggacgcac 1020
cgctcaacat ccgcagcatc ttctacgtgc ccgacatgaa accgtccatg tttgatgtga 1080
gccgggagct gggctccagc gtttgcaactg tacagccgca aagtcctcat ccagaccaag 1140
gccacggaca tcctgcccga gtgctgcgc ttcattccag gtgtggtgga cagtgaggac 1200
attccccga acctcagccg ggagctgctg caggagagcg cactcatcag gaaactccgg 1260
gacgttttac agcagaggct gatcaaattc ttcattgacc agagtaaaaa agatgctgag 1320
aagtatgcaa agttttttga agattacggc ctgttcattg gggaggggcat tgtgaccgcc 1380
accgagcagg aggtcaagga ggacatagca aagctgctgc gctacgagtc ctcggcgctg 1440
ccctccgggc agtcaaccag cctctcagaa tacgccagcc gcatgcgggc cggcaccgcg 1500
aacatctact acctgtgcgc cccaaccgt cacttggcag agcactcacc ctactatgag 1560
gcatgaaga agaaaagacac agaggttctc ttctgctttg agcagtttga tgagctcacc 1620
ctgctgcacc ttcgtgagtt tgacaagaag aagctgatct ctgtggagac ggacatagtc 1680
gtggatcact acaaggagga gaagtttgag gacaggtccc cagccgccga gtgcctatca 1740
gagaaggaga cggaggagct catggcctgg atgagaaatg tgctggggtc gcgtgtcacc 1800
aacgtgaagg tgacctccg actggacacc caccctgcc a tggtcaccgt gctggagatg 1860
ggggctgccc gccacttcct gcgcatgcag cagctggcca agaccagga ggagcgcgca 1920
cagctcctgc agcccacgct ggagatcaac ccaggcacg cgtcatcaa gaagctgaat 1980
cagctgcgcg caagcgagcc tggcctggct cagctgctgg tggatcagat atacgagaac 2040
gccatgattg ctgctggact tgttgacgac ctagggcca tgggtgggccc cttgaatgag 2100
ctgctgtgca aggccttga gcgacactga cagccagggg gccagaagga ctgacaccac 2160
agatgacagc cccacctcct tgagctttat ttacctaaat ttaaaggat tctttaaccc 2220
gaaaaaaaaa aaaaaaaaaa cctctgcc
```

2248

<210> 108

<211> 785

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (769)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (771)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (785)
<223> n equals a,t,g, or c

<400> 108
cccagcccat aaagatctgt gaccggcagc cccagacctg cctgccttcc tgactttctgt 60
tccagagcaa aggtcattca gccgcttgaa tcagcctttt cccccacccc ggtccccaac 120
tttgtttacc cgataaggaa ggtcagcatt caaagtcaag aagcgccatt tatcttcccg 180
tgcgctctac aaatagtctc gtgagaaaga tggccgggaa ctcgatcctg ctggctgctg 240
tctctattct ctcggcctgt cagcaaagt attttgcttt gcaagttgga aaggcaagat 300
taaaatacaa agttacgccc ccagcagtca ctgggtcacc agagtttgag agagtatttc 360
gggcacaaca aaactgtgtg gagttttatc ctatattcat aattacattg tggatggctg 420
ggtggtattt caaccaagtt tttgctactt gtctgggtct ggtgtacata tatggccgtc 480
acctatactt ctggggatat tcagaagctg ctaaaaaacg gatcaccggg ttccgactga 540
gtctggggat tttggccttg ttgacctcc taggtgccct gggaattgca aacagctttc 600
tggatgaata tctggacctc aatattgcc aaaaactgag gcggcaattc taactttttc 660
tcttcccttt aatgcttgca gaagctgttc ccaccatgaa ggtaatatgg tatcatttgt 720
taaatataaaa taaagtcttt aaaaaaaaaa aaaaaaaaaa gggggggccnc nctaaaaaat 780
ccaan 785

<210> 109
<211> 611
<212> DNA
<213> Homo sapiens

<400> 109
ggcacagtgg tggaaggtgg gtgtgtagtc gtggtacttt acgcctcggg gtttagggag 60
gagcctaagg taaggagtca gaaacgggga gtaaccgagc tgcggctttt atataaggtc 120
agtggtaggt aaggaagggg ccttaacctc tgctggtgac cagaagcctg catttctgca 180
ttctgcttaa ttccctttcc ttagatttga aagaagccaa cactaaacca caaatataca 240
acaaggccat tttctcaaac gagagtcagc cttaacgaa atgaccatgg ttgacacaga 300
gatgccattc tggcccacca actttgggat cagctccgtg gatctctccg taatggaaga 360
ccactcccat tcctttgata tcaagccctt cactactgtt gacttctcca gcatttctac 420
tccacattac gaagacattc cattcacaag aacagatcca gtggttgag attacaagta 480
tgacctgaaa cttcaagagt accaaagtgc awtcaaagtg gagcctgcat ctccacctta 540
ttattctgag aagactcagy tctacaataa gcctcatgaa gagccttcca actccctcat 600
ggcaattgaa t 611

<210> 110
<211> 664
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (72)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (614)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (616)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (633)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (649)
<223> n equals a,t,g, or c

<400> 110
aattcggcac gagtttggtt tttncctcta tgtaggtatc tggtaaagta atgaatttat 60
cacaaatttt tncaggccc tttctggtta ctgtggcttt atggcagcca rcctttatgc 120
tcgttccata tttggtgaag atgcacttgc aaatgtcagc attgagaagc caattcacca 180
gggaccagat gctgctgtta ccggccatat aagaattcgt gcaaagagcc agggaatggc 240
cttaagtctt ggagataaaa tcaacttgct acagaagaaa actagtatat aaaaataaac 300
aaaaagtcct tgaagcttta cagttaattt aggtatgggc ttactggact ccaacatctt 360
ttgtactctt tcatgcttat atagaatctg agttcatgct gaatactttt cagccaataa 420
tttatagcct ttcccttaaa tcaagattga gtttaaaatt atagtttgct ttttgtctta 480
acagttctga atgctgtcct caaagtatat aatgtttcat gtaccaagac ccttttcaca 540
gtcaataaac agatctattc ataaaaaaaa aaaaaaaaaa ggcggcccgc tctagaggac 600
cccaagctta cgtncncgtg cattgcgacg tcntagctct tttataggng tcaccctaaa 660
tttc 664

<210> 111
<211> 4065
<212> DNA
<213> Homo sapiens

<400> 111

tcccgcgcgc gcccgctgcg tccgcccagc gccagtcgcg gtcccgaccg gcccgcggca 60
gcctgcgcgc gccatggcca cctccccgca gaagtcgcct tctgtcccca agtctccac 120
tcccaagtcg ccccggtccc gcaagaaaga tgattccttc ttggggaaac tcggaggagac 180
cctggcccg aggaagaaag ccaaggaggt gtccgagctg caggaggagg gaatgaacgc 240
catcaacctg cccctcagcc caattccctt tgagctggac cccgaggaca cgatgctgga 300
ggagaatgag gtgcgaacaa tgggtgatcc aaactcacgc agacsccaa gcttcaagaa 360
ctgatgaagg tattaattga ctggattaat gatgtgttg ttggagaaag aatcattgtg 420
aaagacctag ctgaagattt gtatgatgga caagtcctgc agaagctttt cgagaaactg 480
gagagtgaga agctaaatgt ggctgaggtc acccagtcag agattgctca gaagcaaaaa 540
ctgcagactg tcttgagaa gatcaatgaa accctgaaac ttcctccag gagcatcaag 600
tggaatgtgg attctgttca tgccaagagc ctggtggcca tcttacacct gctcgttgc 660
ctgtctcagt atttccgtgc accaattcga ctcccagacc atgtttccat ccaagtgggt 720
gtggtccaga aacgagaagg aatcctccag tctcggcaaa tccaagagga aataactgg 780
aacacagagg ctctttccg gaggcataa cgtgatgcct ttgacacctt gttcgacct 840
gccccagaca agctgaatgt ggtgaaaaag aactcatca ctttcgtgaa caagcacctg 900
aataaactga acctggaggt cacagaactg gaaaccaggt ttgcagatgg ggtgtacctg 960
gtgctgctca tggggctcct ggagggctac tttgtgccc tgacagctt ctctctgacc 1020
ccggacagct ttgaacagaa ggtcttgaat gtctccttg ccttgagct catgcaagat 1080
ggaggggttg aaaagccaaa accgcggcca gaagacatag tcaactgtga cctgaaatct 1140
acactacgag tgtgtacaa cctcttcacc aagtaccgta acgtggagtg aggggctgcc 1200
ctgggcccac cactgcccga gagttcttgc tgttgccgta ctggaccctc ctccgaactg 1260
ccttaccctg cttattcctg tctcttgac tgtgctctc cacaagtcca gctgcaacct 1320
agagatagt gaaactgaaa ttaggaagga aatcatcaat aactcagtgg gctgacctat 1380
ccctcccagg cgctggggac caacctagca atgaagggtt ggaagggtgt tcccttccc 1440
gtgccaggtc cagatttccc tccatgattt gggaaaccagc ttaggcaaaa gagtccccac 1500
aagatgaaaa taaagatcct agttaccatt caaaggatgc taactgtgtg tcaggcccca 1560
cactaagtgc tctgctctga tatactcaag gccattaatc ttcaggactc ccattgacgt 1620
aggtgtttca tcccccttt acagatgagg aaactaaggc ttggagggtt aatgacttgc 1680
cagaagtgtg aattttttct ctctttgaac ataacctctc cttctccct aaaggtaacc 1740
actattctga gtccaatcat caaggttttg ctttctttt tagctaagta tgcattcctc 1800
aatagtagac agtacaacat gtttataaca agccaattac attatgttct ttgcatgttc 1860
taaagtgtg tatgtgtgtg cacatctgag cacgtgcaca tgtacacctg agccaaaaac 1920
acgagaacct actgatctca ccactggggc aagctagggtc agagcttagt gattcacact 1980
gaaattggca aattggattt aacccaatta atagtgtgtg tgtggcagga gtcattgtcc 2040
tcacatcctt tgtacaaatg aaaattactc ttaattcctt cagatttata ataactctgt 2100
actttggttt cagggtgaca tttgggaagg attttgttta gaattaatgg agtggcacat 2160
tttgagcct ttttgcttga ttgcatgtaa tggaaatgcc ctatattttc ctgcaaaaata 2220
agtactaaat tcattatcgt taagcaaatg tacaatatgc tcaggcaccg cagagagctg 2280
ggcacgggcc catgtgagca tcactttgga agtagggctc ttcaacaggg acccttgaac 2340
tttaaagaaa ggaacttctt tttgccttct aattgatcat ttagactatt ctggctaagt 2400
ctgcccacat gtaattaccg gctaattcaa gcgaggaaaa atgtaagtca tttagacca 2460
agccaagcag tttctttgcg tgggttactc aagggcttgt ggttacttgt atctcctcta 2520
tgtgaacttg actttgaaag acagagctct agtgtgccag cctgctaagt cctgtaagaa 2580
tagggaagg cggagggggg tgggcagtga ctaggggacg agaagcatgg ggaaaaatatt 2640
tgcaactctaa acatacagag atagaggtgg ggcttgtgtg acttaacact tgtagccatc 2700
aactgactga gacctgggc taaataatca attgtgctga tattacatct cgttatggaa 2760
tgttctctaa tatgccaggt agacaccagc ccaagtacc tctccagaa gtctgtgact 2820
acctgtcac tactttagc ccattccaca aagccatct ctggtttgag aattcatttt 2880
gatctgtatc tacaccacc aaagttaggc ctctataat gtccaaaaca ttcctttcag 2940
cctttttatt tcttactgta ctgtctctta ctgtactgtc tatctgcagt aattgaggac 3000
ccataaaatt tagataacta catgtctttg ctcttagaat tgtcactcag cataatgagc 3060

```

atttaacata caaaggcaat gtactgtttt gtgttgatct atgtaaaaga atacaattct 3120
tttttacata attagtgaat ttttattttt tattaggaaa cactaaatag tgtaatattt 3180
cttttgcttt taaaaaaatt cctggtagca aatcaagata aataattgct tcattttctt 3240
gagcaatact gaagcaggat gaagtaagag gaatgcattc atttaaacad gctttgcttt 3300
atgaattttg tctctttttt ggtctctttt tcttatattc aagttacaaa tgtacaagta 3360
tccttactaa gagtgctcct tttgtatttt acatatatac agtatgaaaa tacattggaa 3420
cactaggaaa gtttttaaat aacagttcta atttatcaga aaattgtgtt ttgggattga 3480
gttctttgtc tcagcccaga atcccaggtc ctgggcctgg ttttctaatt ctgtcatctc 3540
agttcgatat tttactttag aayctggaat ctctactta atatatgacc atgactttga 3600
aaggcaaaag aggaatcaag aataaataaa acaaacttaa tcttcattct taaaaaaaag 3660
aaaaaagaaa ccaaaatgag aatcaacact tcataggctc actgggtttt cttttctttt 3720
ctgttaattt aaactcagtt atttttaatg cttaatacat acatggtgca aaatttaaaa 3780
agcgcaaaata ggtatctagt ggaaaacctc agcctccctc tctctccggc acccattacc 3840
tctccctgga ggcaactgtt ttgatccatt tcttacacac actgccagag atactctagg 3900
catgtaaagc acaaacatac atataaaatc tgcgggcttc aaaaaatata agtaggatgt 3960
catctatact gtcatacact ttgtttttta tcacttactt aatgttatat cttggatatt 4020
gtattaccct ggggtattaaa aagaactcct ttcacatttt aaaat 4065

```

<210> 112

<211> 1492

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1487)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1491)

<223> n equals a,t,g, or c

<400> 112

```

atctctgnct cgtatccgcc ttgcctccac aagtgtgtggg attacagggtg tgagccacca 60
caccggcct atattgtttt gaaagcatac tctatatata gttaygggca gaggcacagg 120
catcctcagc agctgattca ggagatgatg gtaaaagctag ctaactatga attaaacatt 180
cacatatcca gtctacctgg tccagtaata atacaagcaa atcttgtatt tcaggaacaa 240
atcaagggtt tcttaatttt ttggcttata tacaatgaag taaaaacttg ataaacatgg 300
tttcaaattg aggaggagag tcttggtatg atgttttaat atgtatacct tataattctg 360
cctctagcca aatgctatgt ttgcaaaatg tggcatctgt tagtttttat tgtctgtgtc 420
ttctttgttt actatacctt gggtaatttt gtgttaccaa aaaaaaaaaa aaaagggaagt 480
gtaatgtcag acacacaaga aaagcaaatc agtgttgtta gcttaaagta caatttcaaa 540
ggtcattacc aacagcaggg ttttttttat actttaaaaa cattatgcta catatcattg 600
ccattttcat attttggggt tttgtacttc ttatacaatg gaatcaatgg aatgtcatc 660
cagccactga attgccatta ttatatctaa aaagtttcta agatgacagt tatcactatt 720

```

```
ttgtttttatc tccatgctga catttgaaag aagggtactag tatccctcta gccagattgc 780
ttagtttttc gttggtaatc aaacaacagt tgtactaaag gaaagtaaag ctaggacctc 840
aatcagaatc atagttgcct gcatatatgg taacaagggtc gtgtgcattt gctttcacag 900
tgatgagtga gaggatgaga agaaattatt tgacattttt ctgtggttga atagaagaca 960
cctttctttt gtcttttaggt ttagggaggag atactaagat actggatgtt tatcctatct 1020
tagtttggtt ggagtaataa gagagaagaa gaggggtggac tttggctttt cagtgttttt 1080
tcccctaaaag agtgatattg ctgacgtttc tatcaatttt acacataata tgtggctatg 1140
aaaccatata tctcacttaa gtaacaaagt aatcactttg tctatcacta agtaatagac 1200
aaaaatcatt gtctattatt taaagccaac aaaacagtgt aacagtttta agttcaataa 1260
tgtaagtat tgtatagaaa tatattggag gcaaagttca gttgatgaca attgtgtata 1320
tgttactgat gctgtaaaatt atttttaata aagaaaattg tattatcaaa aaaaaaaaaa 1380
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1440
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaggg gggcccnttt na 1492
```

<210> 113

<211> 1482

<212> DNA

<213> Homo sapiens

<400> 113

```
atTTTTTTTT tTTTTTTTT tTTTtaagct caaatTTTT ttaaccttgg aacaaatgca 60
ataattcctt acagtcatat agcattttac aattctgaaa gaattttcac atataccaat 120
tgaaccttca caacagccct gtgaggtagg cagggtagggt gttactttcc tactgctagt 180
gaggagaaca aggctaagag gtgaagtaag gtcacacagt tggtaaatga tggagctagg 240
actagaacct aggccttctg acttgctgcc tccctgagaa cattcaagta ataatatcct 300
atgctatata atgaatatTT atctcctatg atgttgatca ggtctatttt tatacatttt 360
acctgttttt atattacctt tccctaatac tgaaaaacac taaaactgat acgagacatt 420
tgagacattt tctctctgtt actggacttt cttagatata tcaaacacta tcctaagatg 480
aaacttctgt tttgaggatc cttggctata aattctcaat tatgatacga acatttattt 540
tacaattctt acaaaagtat gttgaaacta agaaaaacaa aacagggtgtg tttatatctt 600
caaaagcttt gccatctttc ccaatatgac tttaatatgt gtataaatgg aaaacaggga 660
aagggtgttca gcaacattac agtattattg ggaaacctat gcagagtgcata cataaatgca 720
agttacatca cattaatTTg ttttctttgt tggccttata gaggatgaac cattagtgtc 780
tttcagggtt gatcctgtct tatatgtgac atcctcagca acaagctaaa atgatctgaa 840
ttataccaat agcgggggtc aagcagcagt gatcaaaaca taatatgtat gtggtaaaaa 900
ggaacacttg ttcttcagtg gttttgccta ggtccagtgt aattacttat caaaacatgt 960
tttaagctac agcatctctg ttaaatttag aaatgtgaaa catggaagag ttatcaactt 1020
ggaatacaat actaaaactg aatataactg acccaaaact acctgtcaaa agcacagtaa 1080
agcttaacag gcataaaaaa taaggggtaa aacccaactg ctctgacata agagtgaaaa 1140
acagcaaatg caaacatca ggttaaagtg aatcacaaaa cattaggcat gtagagcatg 1200
ttttaaaaaat aggaaaatga cactgagctg tattaccagg aataatattt ttaaaattcc 1260
aggtatggcc ttctacattt aatagcaatg gagagaaaca agcctatggg ggaggtgcag 1320
aaaatgaatg ggatctagaa agggagtTga gtttggaagt taaggaagtc aaagaaacaa 1380
atacatccaa aatggattta gaatctctcc tctaaatcca tcctttccaa aaaaatgtttc 1440
ctatattgag agatttggtg aaaaaaaaaa aaaaaaagtc ga 1482
```

<210> 114

<211> 3731

<212> DNA

<213> Homo sapiens

<220>
 <221> misc feature
 <222> (652)
 <223> n equals a,t,g, or c

<220>
 <221> misc feature
 <222> (3730)
 <223> n equals a,t,g, or c

<400> 114
 aaaaacttag tttagggtga cacgtataga aggtacgccg tgcaggtagc ggtccggaat 60
 tcccgggtcg acccacgcgt cgggagctcc atccccgcct ccccgccag ccgacccggc 120
 tccttcacct tcccggggga cagcgactcc ctccagcggc agacaccccg ccacgcggmc 180
 cctggcaagg atactgaccg catgagcagc tgctcctcgg ccagcgagca gtccgtgcag 240
 tccaccaga gcaacgggag tgaaagcagc agcagtagca acatctccac catgttggtg 300
 acacacgatt acacggcagt raaggaggat gagatcaacg tctaccaagg agaggctcgtt 360
 caaattctcg ccagcaacca gcagaacatg tttctggtgt tccgagccgc cactgaccag 420
 tgccccgcag ctgagggtcg gattccaggc ttgtcctcgg gccacaccag tgcagtcata 480
 gtggagaacc cggacgggac tctcaagaag tcaacatctt ggacacacgc actccgttta 540
 aggaaaaaat ctgagaaaaa agataaagac ggcaaaaggg aaggcaagtt agagaacggg 600
 tatcgggaagt cacgggaagg actcagcaac aagggtatctg tgaaagcttc tncaatccca 660
 actacattta tgaacgttcc ccccagaatt cgtcattcca ttgagttagg tcacgtgtga 720
 gacaggggag accgttggtc ttagatgtcg agtctgtggc cgcccaaag cctcaattac 780
 ctggaagggc cctgaacaca acaccttgaa caacgatggt cactacagca tctcctacag 840
 tgacctgga gaggccacgc tgaagattgt gggcgtgacc acggaagatg acggcatcta 900
 cacgtgyatc gctgtcaatg acatgggttc agcctcatcg tcggccagcc tgagggtcct 960
 aggtccaggg atggtggga tcatggtgac ctggaaagac aactttgact cctctacag 1020
 tgaagtggct gagcttgga ggggcagatt ctctgtcgtt aagaaatgtg atcagaaagg 1080
 aaccaagcga gcagtggcca ctaagtgtgt gaacaagaag ttgatgaagc gcgaccagg 1140
 caccatgag cttggcatcc tgcagagcct ccagcaccct ctgcttgctg gcctcctcga 1200
 cacctttgag acccccacca gctacatcct ggtcttagaa atggctgacc agggtcgcct 1260
 cctggactgc gtggtgcgat ggggaagcct cactgaaggg aagatcaggg cgcacctggg 1320
 ggaggttctg gaagctgtcc ggtacctgca caactgcagg atagcacacc tggacctaaa 1380
 gcctgagaat atcctggtgg atgagagttt agccaagcca accatcaaac tggctgactt 1440
 tggagatgct gttcagctca acacgacct ctacatccac cagttactgg ggaaccctga 1500
 attcgcagcc cctgaaatca tcctcgggaa ccctgtctcc ctgacctcgg atacgtggag 1560
 tggtggagtg ctcacatacg tacttcttag tggcgtgtcc cccttcctgg atgacagtgt 1620
 ggaagagacc tgcctgaaca tttgcccgtt agactttagc ttcccagatg actactttaa 1680
 aggagttagc cagaaggcca aggagtctgt gtgcttctcc tgcaggagga ccccgccaag 1740
 cgtccctcgg ctgcgtggc cctccaggag cagtggctgc aggcggcaa cggcagaaag 1800
 cacgggcgtc ctgcacacgt ccagactgac ttccctcatt gagcggcgca aacaccagaa 1860
 tgatgttcga cctatccgta gcattaaaaa ctttctgcag agcaggcttc tgccatagat 1920
 ttgacctatc cagaagttct ttctcattct ctttcacctg ccaatcagct gttaatctga 1980
 attttcaaga gaaaacaagc aaacataact gatcagctgc cggatgttc atcgtgtgaa 2040
 attgcattcc aagttagctg tgctcagcag tgcttgaca cagagctgca agctgcgtg 2100
 gggtaggaga ccgtcacta cactctkccc aaggcagagg tcgcattgct gtatcacagt 2160
 attttattca ggtttctgca aaaaaataaa aagataactt ttttaacaa acatgaatag 2220
 aattttgcaa atttaacgtt ttcaagattt attcaaggaa acaaaatgcc tatgttcaac 2280
 cactggtgtt aatgaacaaa gatactgtgc gtctctgggg aagacgcacc taggtggcgg 2340
 ccactcccat ggccttgtct aggactcaga gaccactcgg ctctgagctt ccaggcgcct 2400


```

cgtctgtgtg catctcacgc ccgacgtggc ttctgaaacg tgcattcaac ctcaaacttt 2460
tgcataaaat agaataatc gttttgctct gatgaaatgt aggccttact tgtatataag 2520
actgttcctg ccttcgggtc gtcatthttc cacctgcctc ccctaccac cccccacca 2580
ccacctgggg ctctctctgg ggggtccgag gtcttcccat cacatgaaga catcaggttg 2640
ggctctgccc cactgcccc ctccctgttc ctgccccaa ggcgcaatca gattgtggag 2700
cagtacacag tcagatgaaa atactgtaaa tgcactcatt gggggttttt tgggtttact 2760
tcataatcatg tgcaatgttg tggctttaac attttatgca actatthtat aagacctctg 2820
ttgtacctgt aataaatata tagaaaaagc acatacttcg tatggtgagc tttatggtt 2880
tgtgtgtgtg ttgggggttg cgggtgggtg ggtagggtcg tagccctgtg ccacgggttc 2940
aaagagactt ttcgtgaaat ttgttggtt tgaggactgt aaaagtgtt tcatactctg 3000
aatataaaac tggataatag ggtaatgtt taaaatttat tatgctatta ttcagaatgc 3060
caaagtatta ttttttttcc caaaatcagt ctggacattt actactttt agactttttg 3120
acgttgaact ttctgtataa aaattggctg ggttttgagc ttttggttag aaataaaagc 3180
cgattaagca ctggccgccc cgcggctggg acccaatgcc cgagtcactg tggcagcatt 3240
cgactgggtg tggggagtcc tttcaactca aggaggtcg gtttctgggc accctcgaag 3300
ttttctggat gtctttttat ctttctctg tgaactgcac taaaaagag accagccgc 3360
ttccaagcc agccagacac ctgggtcttg agccataaac tggcgtagt aagctttgca 3420
gttccagtg tattttattt attctttgt tgggttttg tttgttctt cttgtaaaat 3480
tgtacagaaa ctttttaaaa gaaattggat tcgaaactgg atgtgtattc gtaacctcat 3540
aatttttatt tgtgtattgt ttcttttatt atttcagtta aatttttaca ttattttgca 3600
tgtatatttc tttgtacaga gaccttacct gtttacacag tatgatgtga tgtaaatatt 3660
ttattttgcc atcagttatt ttaaaaaatt aaacatattt gcctgaaaaa aaaaaaaaaa 3720
ggggggcgcn c 3731

```

<210> 115

<211> 1315

<212> DNA

<213> Homo sapiens

<400> 115

```

ggcggcagcg ccggaagga ggccaagagc gcggggcgcg aggcaagatg gcggcaacca 60
agaggaaacg gcgtggagggc tttgcagttc aggcgaagaa gccaaaaaga aacgaaatag 120
atgaggagcc gccagctaag cggcacgcca cagcagagga ggtggaggaa gaagagaggg 180
accggatccc agggcccggtt tgcaaggga agtggaaaaa taaggaaacg attctcatct 240
tttcttccag aggaataaat tttagaacaa gacatttaat gcaggacttg agaattgtga 300
tgcctcattc taaagcagat actaaaatgg atcgtaagga taagctattt gtgattaacg 360
aggtttgtga aatgaagaac tgtaataaat gcatttattt tgaagctaag aaaaaacagg 420
atctctatat gtggctttca aattcacctc acggaccatc tgctaaattc cttgttcaaa 480
atattcatac cctcgctgaa ctgaagatga ctggaaactg tttgaaaggt tctcggcccc 540
ttttgtcttt tgacctgct tttgatgaat taccacatta tgctttgkta aaagaactct 600
taattcagat ctttagtaca ccacggatc atcccaaaag ccaaccattt tgggaccacg 660
tgtttacttt caccattttg gataatagga tatggttctg gaactttcag atcatagaag 720
aagatgctgc tctttagtaa ataggacct gttttgtctt aaatctcata aagattttcc 780
agggaagttt tggaggacca acttttatatg aaaatcctca ctaccagtca ccaaactatg 840
atcggcgctg cataagatcc atcacagctg caaaatacag agagaaacag caagtgaag 900
atgtgcaaaa actgagaaag aaagagccga agactcttct tccacatgat cccactgcag 960
atgtttttgt aacaccagct gaggagaaac caatagaaat acagtgggtg aaaccagagc 1020
caaaagttga tttgaaagca agaaagaaac ggatttaca aaggcaaga aaaatgaaac 1080
agaggatgga cagtgggaaa acaaaataag tcaatggaaa cctgatttgt ttttcagtta 1140
ctttatattt attttgtatt caatgtgtaa atacttttat tatctaatac tatcttacgt 1200
ctaattagtg tagcatthac aagaaagaaa aattaagatc ttaaaatcag tgattatctt 1260

```

tttctaaata aaatatcacc agaattcatc agttaaaaaa aaaaaaaaaa aaaaa 1315

<210> 116

<211> 1320

<212> DNA

<213> Homo sapiens

<400> 116

ggctggcttc tgcgtgggtgc agctgcgcac gtgtttcagc cggcagcgct ttaagatttc 60
cggggatgga atccgaaatg gaaacgcaga gcgccrgggc agaggagggc ttaccacagg 120
tcacccgcaa ggggtggccga cgggcgaaga aacgacaggc tgaacagctg tccgcagcag 180
gagagggcgc ggatgcgggc cgcacggaca cagaggaggc caggccggcg aagaggcccg 240
tcttcccacc cctctgtggg gacgggctcc tgagtgggaa agaagaaaca aggaaaattc 300
cagtcccgac taacagatac acaccattga aagaaaactg gatgaagata tttactccta 360
ttgtggaaca tttgggactt cagatacgct ttaacttgaa atcaaggaat gtagaaatca 420
ggacttgtaa rgaaaccaag gatgttagtg ctctgacaaa agcagctgat tttgtgaaag 480
cttttattct cggctttcag gtggaggatg cacttgccct catcagggtg gatgacctct 540
tcctagagtc ttttgaaatt acagatgtta aaccctaaa gggagaccat ctatccaggg 600
caataggaag aatcgtggc aaaggaggaa aaaccaaatt caccatagag aatgtgacac 660
ggacaaggat agttttggct gatgtgaaag ttcacatcct tggctccttc caaaatatca 720
agatggcaag aactgccatt tgcaacctaa tcttgggaaa tcctccttcc aaggtttatg 780
gcaatatctg agctgtggct agcagatcag cagatcgatt ctgatttcaa gtcagagact 840
ttttatcttg cctttggact ctggtgaaaa atactttaca gtggtcggtc acaagaaacc 900
atctgaacaa tttcagtcac ttgaagcctc cgtcccttct tccattctca gccagaagca 960
taaacagaaa agaaagattt aagaggattc aactcaaca ggttttagga taatttaaat 1020
atcaaaaatt gattgttata cttacacatt aggtataatt tatcatttat ctgaaatcac 1080
atgtagcaga ttgcatagtc tgtaatcctc tcagagggaa acttcttggt taaacagctc 1140
tatatggatt tatactttta tatttataaa tttataactt catacaaatt tataaacatt 1200
tctttataaa ttgtaattta atagattatc tcagaaaaac ctctctgaat gatgaccctt 1260
ccttaatact ggggtgatgtg tgaatatttg tttgttgcca gacagggctc cactttgtca 1320

<210> 117

<211> 2025

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1339)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1916)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1944)

<223> n equals a,t,g, or c

<400> 117
ttatttttaca gcctcaaata tttctcatta tcttgtcact tagttcttca tgtttctcct 60
tctgactttt aataatggta ataggaaaac aaaacccaaa gcttttcaga acttcagtgt 120
gagggttcct attttracaa gttaacttgt aaatactcag gttttacgat gtataattta 180
cctaatagac caaactaact catggagata ttttgaacta ttatttaggt acaaacttta 240
taaagaatgt tagtatgtca taaaatataa cattacagct tatttaaaac caaatatagt 300
tgaacatatt taaaatacat tttcacagaa tggatgaatt agttgtttct tcagagttac 360
ttatgaacag ttgaatgctt taaaatgttc tgtctgtagg taacatctaa aacacaagtg 420
ggtttattta aattttttaa atttgaaatt ttttatttgc aaaaaattgt tttatgcttt 480
attatatcgc aaatgagtgt cagatttttg agtaccatg atcatgcttc catttttttt 540
agtttttaa acaccaaacca atatttttcc tttaaatttt aatcttataa tatagaaatc 600
ttatgtaaat gaaattttgt catgtttcaa ataaagagaa ctgaagtaga aaatagaaat 660
gccagtaa acacataatgt ttaatttaca acttacatta ggggtttggg ggaatgctaa 720
ttatatattg agaataatac ttagaactct tcaaaatggg ctcttctaag gaggtcacta 780
ctgaacaaaa ttgttccctc ttctgttaaa tagaataggt ttaaatagact agtcaaatga 840
attattttct tcttgttaaa taaattaaat cttactttct tttaatgacc aaccttaggt 900
aaaacaaaaa tattgtaatc ctagaataa tctccagct tctcacctg aaaatctatt 960
gaagtgtacc ctggtcatcc taataatggg atgaggggag tttccagcag atttcagggt 1020
gttcttaaa gttttgttgg tcattttctc aatagtacat gaaatcaaga tgcttatgag 1080
catggaaatg tatttaaaatg ttttgcctgt gtcctcctca gtcagaatag aaaagtaact 1140
gaaatactct tacctttctg tccttgataa aatagtaaag aaaaccaa acacccaggc 1200
ctgatgggaa aaatgattcc tttattctag caacttactt tctgttggtg tgggaaatgt 1260
tattaatttc tattactaaa gtccatatca caaaatgata ttttaataa accttgggg 1320
aaatcatgaa ttttttttnc tacgtgtgag tataaaagac aaaagttgaa cagcatggaa 1380
tctcattgcc aaattattag tgaatgtata gtccaggtat tctttgagac acacagtatc 1440
attaatttcc gaattgtatt tcagtgttat tttttgttg tgaccactaa gcttctgtct 1500
taatacaaa ctgttaccct ctacagaatt taagtctgaa gatgtaaaga gagaacaggc 1560
cttgtgtaac agaagatact cttttttatg ctccttactg tgatcacaga aaaattaaaa 1620
atccaagtgc tctctagatt tgttgataaa ctttttatgc ttgcatttaa acttgaaatg 1680
tatgagcaga atgagacaat cagttaaact agaaatgaga agtattataa tgtaaaggcc 1740
ttgttttgc gtagcaataa aatgaccaag tgcaatgact tgatttaata aaatcatatt 1800
ttaaagtg tgstatgaaw atttttggct ataaaatttt accctgactt gstttcaata 1860
actgttacgt aatgcagtt gatgtgttaa cctaacattc caaaaaaaa attganaggg 1920
ggaatctcaa aatagtatat actnactaa cttgtttaca ggtgctgtat ttaaagcat 1980
gcttctctct caaaaagaaa aattaaagga ttttattgcc aaacc 2025

<210> 118

<211> 1295

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1286)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1292)

<223> n equals a,t,g, or c

<400> 118

```
ggcacgaggt gcagttacac gttttcctcc aaggagcctc ggacgttgtc acgggttttg 60
ggtcggggac agagcgggtga ccatggccag gctggcggtg tctcctgtgc ccagccactg 120
gatggtggcg ttgctgctgc tgctctcagc agctgagcca gtaccagcag ccagatcgga 180
ggaccggtac cggaaatccca aaggtagtgc ttgttcgagg atctggcaga gcccacgttt 240
catagccagg aaacggggct tcacggtgaa aatgcactgc tacatgaaca gcgcctccgg 300
caatgtgagc tggctctgga agcaggagat ggacgagaat cccagcagc tgaagctgga 360
aaagggccgc atggaagagt ccagaacga atctctcgcc accctacca tccaaggcat 420
ccggtttgag gacaatggca tctacttctg ccagcagaag tgcaacaaca cctcggaggt 480
ctaccagggc tgccgcacag agctgcgagt catgggattc agcaccttg cwagctgaa 540
gcagaggaac acgctgaagg atggtatcat catgatccag acgctgctga tcatcctctt 600
catcatcgtg cctatcttcc tgctgctgga caaggatgac agcaaggctg gcatggagga 660
agatcacacc twcgagggcc tggacattga ccagacagcc acctatgagg acatagtac 720
gctgcggaca ggggaagtga agtggtctgt aggtgagcac ccaggccagg agtgagagcc 780
aggtcgcccc atgacctggg tgcaggctcc ctggcctcag tgactgcttc ggagctgcct 840
ggctcatggc ccaacccctt tccyggaccc cccagctggc ctctgaagct ggcccaccag 900
agctgccatt tgtctccagc ccctggctcc cagctcttgc caaaggccct ggagtagaag 960
gacaacaggg cagcaacttg gagggagtcc tctggggatg gacgggaccc agccttctgg 1020
gggtgctatg aggtgatccg tccccacaca tgggatgggg gaggcagaga ctggtccaga 1080
gcccgcgaat ggactcggag cygagggcct cccagcagag cttgggaagg gccatggacc 1140
caactgggcc ccagaagagc cacaggaaca tcatctctct cccgcaacca ctcccacccc 1200
agggagccct ggcctccagt gccttcccc gtggaataaa cgggtgtgtcc tgagaaacca 1260
maaaaaaaaa aaaaaaaaaa aaaaangggg gnccc 1295
```

<210> 119

<211> 1257

<212> DNA

<213> Homo sapiens

<400> 119

```
tggacagacc tcagcaacac agcaatgtgc actttgtagt aaagaaatac caacttgagt 60
ttttaaaaaa cctagttagt cctacattcc cttttcttta aaaaaacttt attgttcaca 120
ttattttaat gactatgaga taatgtatat gacagcactt tgagaaaata tcaactgtaa 180
tataactata aggttgtagt attgtctgtt taaaagataa gacagttgat tcaatgtgga 240
tggaccctgt ggggtacctg aaaatgtaga tacgtaagaa tcacctctgt catttatcac 300
atttagaaat atgaaactgc ttaacaggta tgagcaggta tagcaagtgt ttgctaaagt 360
tgtagttcag agctgaatta cttcaggaaa ctagggacca actttttggt ttaattccga 420
tctttaaaaa gtaagaatgt gtactcactc cagaacacag aagctcttcc aaggaccttg 480
actcaagaag gatgaggctc tcttactctt ctccatttat ccactatatg cttggccatt 540
tatcctaaat gtggtgggaa cagacttggt atctgttgat gttgacagtg tcttttttaa 600
cctatgtcct gcatagtttt gttaggttta cagggggagg tggatggcca taaaaccagt 660
gcactttggg aattactttt ctaggatttc ctaccagtta taaatgacat tgacatttgt 720
catctttttt ttctttttct aaaaagaata gctgaattta attcacctat tataaaatac 780
tcaaaagtaa attgcmittg tggccacttc tgaattatag ctacatttca ttatgacccc 840
ttctgctcwc ttccattttg ctactgatgt cattcttggt atcaggctgt gccctacca 900
ggagttcata ttgggttgac agggttatct atatttttgt tcttgatttt tgagttttat 960
catcttgcat taaattgttc cacctggatt tggggttcat ctctgtgccc taaggatctg 1020
ctatgaccaa tcctcttttt gtaggtgggt ctctggctta agtattgata ggcttcagcg 1080
gtttgtgtct ctgtcttagc ttgtatcaag ccagtagtag ctcacttcct ttgtaaatto 1140
ctgcttcagt ctgggatccg taggagtatg tgagaacttc tgaacgtct ccaactctta 1200
acagtcaaga tatctatatc atttgawag agttctggkt ttccaactac taagacc 1257
```

<210> 120
<211> 397
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (378)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (395)
<223> n equals a,t,g, or c

<400> 120
ccccaggaaa ataccaaacc taggtttaaa aatagggag ggaagggaga gaagaaaggg 60
aagataaaga aagcccaacc tccttccaaa atgtcatgag aatcttgagc acatatgggc 120
cttgcatga ccacatgacc tgcagagccc ctgttataga actcattttt atattttcct 180
tagtataaca gttaataataa tatgtcattt ttgttaatag tgtctttttg tcattttact 240
ttttaaaaga ttttattgaa atatacatag aggaaagtgc atctatcata agtgtgccaa 300
attgatggaa ttctaaaatc tttantggta cctgttttagc acatagattg gacacggaa 360
ataactaaca accagaantc tccgtgtact ccctncc 397

<210> 121
<211> 876
<212> DNA
<213> Homo sapiens

<400> 121
cccacgcgtc cggaaggtgc ttactcgtcg tctccatgag ccggttcctg ggcgtcttag 60
agccaaggcg cgaggctcgg agtgagaggt agagctggag gggaccctaa gcgcctccg 120
cccgggacgt gagccgctgc gccaccggg ctagaccggg cgccatcatg ctgcttctgc 180
caagcgcgcg ggacggccgg ggcaccgcca tcaccacgc tctgacctct gcctctacac 240
tctgtcaagt tgaacctgtg ggaagatggt ttgaagcttt tgtaagagg agaaacagaa 300
atgcttctgc ctcttttcag gaactggagg ataagaaaga gttatccgag gaatcagaag 360
atgaagaatt gcagttggaa gagtttccca tgctgaaaac acttgatccc aaagactgga 420
agaaccaaga tcattatgca gttcttggaac ttggccatgt gagatacaag gctacacaga 480
gacagatcaa agcagctcat aaagcaatgg ttttaaaaca tcaccagac aaacggaaag 540
cagctgggtga accaataaaa gaaggagata atgactactt cacttgcata actaaagctt 600
atgaaatggt atctgatcca gtgaaaagac gagcatttaa cagtgtagat cctacttttg 660
ataactcagt tccttctaaa agtgaagcaa aggataattt cttcgaagtg tttacccag 720
tgtttgaaag gaattccaga tgggtcaaata aaaaaaatgt tcctaaactt ggtgatatga 780
attcatcatt tgaagatgta gatataattt attcttctg gtataatttt gattcttgga 840
gagaattttc ttatttagat gaagaagaaa aaaaaa 876

<210> 122
<211> 1278
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (107)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (128)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (149)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1228)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1231)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1262)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1269)
<223> n equals a,t,g, or c

<400> 122
ccaggaaggc ctttggaatg caaacatttt tgggtcccat cgaaatgttc gaaagactaa 60
ggaagataaa aagagttgac gggcgccgac ctgaaggact gcgtcanaca acagcctgag 120
cagcaatnca gcctcccag cgtgcaganc tgccggcgcc tgcgtgagag gagggtcgcc 180
agctggggccg tgtcctttga gcgctgctg caggaccccg tcggtgtccg ctacttctct 240
gatttttctaa ggaaagaatt cagtgaagaa aacattttat tctggcaggc ctgtgaatat 300
tttaatcatg ttcctgcaca tgacaaaaag gagctttcct acaggggcccg ggagattttc 360
agtaagtttc tctgcagcaa agccaccacc ccggtcaaca tcgacagcca ggcccagcta 420
gcagacgacg tcctccgcgc acctcaccca gacatgttca aggagcagca gctscagatc 480
ttcaatctca tgaagtttga tagctacact cgctttctga agtccccgct gtaccaggga 540

atgcacacctg ggcggaagtg gagggccgtg cactcccga ctcgcagcag gtccccagca 600
gcccggcttc caagcacagc ctccggttcag accactccag tgtgtccacg ccaaaaaagt 660
taagtggaaa atcaaaatcc ggccgatccc ttgaatgaag agctggggga tgaggacagc 720
gagaagaagc ggaaaggcgc gtttttctcg tggctcgcga ccaggagcac cgggaggtcc 780
cagaaaaaga gggagcacgc ggaccacgca gacgacgccc tgcatgcaa tgaggacctg 840
tgtcgcgcag agtcgcaggg ctctgtgtcc tctgcgggga gcctggacct gtccgaggcc 900
tgcaggactt tggcaccgga gaaggacaag gccaccaagc actgtgcat tcatctccc 960
gatgggacat cctgcgtggt ggctgtcaag gcgggcttct ccatcaaaga catcctgtcc 1020
ggactctgtg agcggcatgg catcaacggg gcggccgcgc acctcttctt ggtgggcggg 1080
gacaagcttc tgggtgcttg accaagacag tagcatcttg ggagtcaagg gacctgcgc 1140
ttaggaaaag cggcaccttt gttttcgggt tggatctttg tttccgattt aaccgggtca 1200
tggggattca aggccaagcc caccaagncc nttacggagg tgctgcggcc ctggtggcca 1260
gntacggcgt ggactcat 1278

<210> 123

<211> 3115

<212> DNA

<213> Homo sapiens

<400> 123

ggcacgagcg cgggcagcga gcgcactcgg cggacgcaag ggccggcggg agcacacgga 60
gcactgcagg cgccgggttg ggacagcgctc ttccgtgctg ctggatagtc gtgttttcgg 120
ggatcgagga tactcaccag aaaccgaaaa tgccgaaacc aatcaatgct cgagttacca 180
ccatggatgc agagctggag tttgcaatcc agccaaatac aactggaaaa cagctttttg 240
atcaggtggt aaagactatc ggccctccggg aagtgtggta ctttgccctc cactatgttg 300
ataataaagg atttcttacc tggctgaagc tggataagaa ggtgtctgcc caggaggtca 360
ggaaggagaa tccccctcag ttcaagttcc gggccaagtt ctaccctgaa gatgtggctg 420
aggagctcat ccaggacatc acccagaaac ttttcttctt ccaagtgaag gaaggaatcc 480
ttagcgatga gatctactgc cccctgaga ctgcccgtgct cttgggggtcc tacgctgtgc 540
aggccaagtt tggggactac aacaaagaag tgcacaagtc tgggtacctc agctctgagc 600
ggctgatccc tcaaagagtg atggaccagc acaaacttac cagggaccag tgggaggacc 660
ggatccaggt gtggcatgcg gaacaccgtg ggatgctcaa agataatgct atgttggaat 720
acctgaagat tgctcaggac ctggaaatgt atggaatcaa ctatttcgag ataaaaaaca 780
agaaaggaa agaccttttg cttggagtgg atgcccttgg actgaatatt tatgagaaag 840
atgataagtt aaccccaaag attggctttc cttggagtga aatcaggaac atctctttca 900
atgacaaaaa gtttgcatt aaaccatcg acaagaaggc acctgacttt gtgttttatg 960
ccccacgtct gagaatcaac aagcgatcc tgcagctctg catgggcaac catgagttgt 1020
atatgcgccg caggaaagcct gacaccatcg aggtgcagca gatgaaggcc caggcccggg 1080
aggagaagca tcagaagcag ctggagcggc aacagctgga aacagagaag aaaaggagag 1140
aaaccgtgga gagagagaaa gagcagatga tgcgcgagaa ggaggagttg atgctgcggc 1200
tgcaggacta tgaggagaag acaaagaagg cagagagaga gctctcggag cagattcaga 1260
gggccctgca gctggaggag gagaggaagc gggcacagga ggaggccgag cgcctagagg 1320
ctgaccgtat ggctgcactg cgggctaagg aggagctgga gagacaggcg gtggatcaga 1380
taaaagacca ggagcagctg gctgcggagc ttgcagaata cactgccaag attgccctcc 1440
tggaaaggcg gcggaggcgc aaggaggatg aagttgaaga gtggcagcac agggccaaaag 1500
aagcccagga tgacctggtg aagaccaagg aggagctgca cctggtgatg acagcaccac 1560
cgccccacc acccccgtg tacgagccgg tgagctacca tgtccaggag agcttgacag 1620
atgaggcgcg agagcccacg ggctacagcg cggagctgtc tagtgagggc atccgggatg 1680
accgcaatga ggagaagcgc atcactgagg cagagaagaa cgagcgtgtg cagcggcagc 1740
tgctgacgct gagcagcgag ctgtccagc cccgagatga gaataagagg acccacaatg 1800
acatcatcca caacgagaac atgaggcaag gccgggacaa gtacaagacg ctgcggcaga 1860

```

tccggcaggc caacaccaag cagcgcatcg acgagttcga ggccctgtaa cagccaggcc 1920
aggaccaagg gcagaggggt gctcatagcg ggcgctgccg gccccgccac gcttgtcttt 1980
agtgtccaa gtctaggaac tccctcagat cccagttcct ttagaaaagca gttaccaaac 2040
agaaacattc tgggctggga accagggagg cgccttggtt tgttttcccc agttgtaata 2100
gtgccaaagca ggcctgattc tcgcgattat tctcgaatca cctcctgtgt tggtctggga 2160
gcaggactga ttgaattacg gaaaatgcct gtaaagtctg agtaagaaac ttcattgttg 2220
cctgtgtgat acaagagtca gcatcattaa aggaaacgtg gcaggacttc catctgtgcc 2280
atactgttgc tgtattcgaa atgagctcaa attgattttt taatttctat gaaggatcca 2340
tctttgtata ttacatgct tagaggggtg aaaattattt tggaaattga gtctgaagca 2400
ctctcgaca cacagtgtt cctcctccc gtcactccac gcagctggca gagagcacag 2460
tgatcaccag cgtgagtggg ggaggaggac acttggatat ttttttagtt tttttttttt 2520
tggtcttaaca gttttagaat acattgtact tatacacctt attaatgac agctatatac 2580
tatttatata caagtgataa tacagatttg taacattagt tttaaaaagg gaaagttttg 2640
ttctgtatat tttgttacct ttacagaat aaaagaatta catatgaaaa accctctaaa 2700
ccatggcact tgatgtgatg tggcaggagg gcagtgggtg agctggacct gcctgtgca 2760
gtcacgtgta aacaggatta ttattagtgt tttatgcatg taatggacta tgcacacttt 2820
taattttgtc agattcacac atgccactat gagctttcag actccagctg tgaagagact 2880
ctgtttgctt gtgtttgttt ggcagtctct ctctgccatg gccttggcag gctgtggaa 2940
ggcagcttgt ggaggccgtt ggttccgcc actcattcct tctcgtgcac tgctttctcc 3000
ttcacagcta agatgccatg tgcaggtgga ttccatgccg cagacatgaa ataaaagctt 3060
tgcaaaggca aaaaaaaaaa aaaaaraaaa maaaaaaaaa maaaraaaaa aaaaa 3115

```

<210> 124

<211> 379

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (340)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (344)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (366)

<223> n equals a,t,g, or c

<400> 124

```

atgaatcggg ttattgaaga agatggaacc agtgcttaat ttgtcttgtc taagggtggaa 60
atcttgtgct gtttaaaaag cagattttat tctttgcctt ttgcatgact gatagctgta 120
actcacagtt aacatgcttt cagtcaagta cagatttgtt ccaactggaaa ggtaaatgat 180
tgctttttta tattgcatca aacttggaac atcaaggcat ccaaaacact aagaattcta 240
tcacacaaaa aataattcgt ctttctaggt tatgaagaga taattatttg kctggtaagc 300
atttttataa acccacycat tttatattta graaaatccn aaanggggtg gggacgccct 360
tgtagngacc ttccaatac 379

```


<210> 125
<211> 1267
<212> DNA
<213> Homo sapiens

<400> 125
ggcacagttag aggctggtga ctgtggccgg ctgggagtag gcggcagtga gtttccttgg 60
gagggcagcg cgcttggcgc ttctcccttc ccccgatct gcctccagtc tcggacttgg 120
ttgttgcgcg ctccggctcc ggctgagctg ggagagttgg aggaggtggc ggcgggcaga 180
ggtgatgtct gggagccctt ccttgacagc ccgggcccag aagagtcctt gcaggaagca 240
tcaccacaggc tggcagatca tggtagcagc agcgggggtg gctgggaagt gaaacggagc 300
cagcggctga ggagggggccc cagcagcccc cgaaggccct atcaggacat ggagtatgaa 360
agacgtggtg gtcgtggtga caggactggc cgctatggag ccactgaccg ctgcgaggat 420
gatggtgggg agaaccgcag ccgagaccac gactaccggg acatggacta ccgttcatat 480
cctcgcgagt atggcagcca ggagggcaag catgactatg acgactcatc tgaggagcag 540
agtgcggagg attcctacga ggcctccccg ggctccgaga ctacgcgtag gcggcgccgg 600
cggcacaggc acagccccac cggcccgcga ggcttcccc gagacggcga ctatcgggac 660
caggactatc ggaccgagca agggaggagg aggaggagga rgaggatgag gaggaggagg 720
agaaggccag taacatcgct atgctgagga tgctgccaca ggcaccact gaggatgaca 780
tccgtggcca gytgcagtcg cacggcgtgc aagcacggga rgttcgggtg atgcggaaca 840
aatcttcagg tcagagccgg ggcttcgcct tcgtcgagtt tagtcacttg caggacgctc 900
acgatggatg gaagccaatc agcactccct caacatcctg ggccagaagg tgctgatgca 960
ctacagtgac cccaagccca agatcaatga ggactggctg tgcaataagt gtggcgctcca 1020
gaacttcaaa cgccgagaga agtgcttcaa atgtggcgtg cccaagtcag aggcagagca 1080
gaagctgccc ctccgmacga ggytgatca gcagacactg ccactgggtg kccgggagct 1140
gagccagggc ctgcttyccc tgccgcagcc ctaccaggcc cagggagtcc tggcctccca 1200
agccctgtca cagggtctcg agccaagctc agagaacgcc aatgacacca tcattttgcg 1260
caacctg 1267

<210> 126
<211> 841
<212> DNA
<213> Homo sapiens

<400> 126
gcgttacsat cgtccgtgcg caccgcccgg cgtccagggt agtctcccat ctgcagagac 60
gcggacgcgc cggcccgcag ttggcctgcg gagcgcggtg gacggtttgg cgcccaccag 120
gcgatcaata ctttgattt ttaatttcta gatttggcaa ttcttcgctg aagtcatcat 180
gagctttttc caactcctga tgaaggaa ggaactcatt cccttggttg tttcatgac 240
tgtggcgggc ggtggagcct catctttcgc tgtgtattct ctttggaata ccgatgtgat 300
ccttgatcga aaaaaaatac cagaaccttg ggaactgtg gaccctactg tacctcaaaa 360
gcttataaca atcaaccaac aatggaaacc cattgaagag ttgcaaaatg tccaaagggt 420
gaccaaatac cgagccctcg cctctttctt ctgaagagta ctctataaat ctagtggaaa 480
catttctgca caaactagat tctggacacc agtgtgcgga aatgcttctg ctacattttt 540
agggtttgtc tacatttttt gggctctgga taaggaatta aaggagtgc gcaataactg 600
cactgtctaa aagtttgtgc ttattttctt gtaatttga atattgcata ttgaaatttt 660
tgtttatgat ctatgaatgt ttttcttaaa atttataaag ctttgtaaat tagattttct 720
ttaataaaat gccatttgtg caagaaaaaa aaaaaaaa aaagggcggc cgctctagag 780
gatccctcga ggggccaag cttacgcgtg catgcgacgt catagctctg tccctacgaa 840
g 841

<210> 127
<211> 1172
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (119)
<223> n equals a,t,g, or c

<400> 127
tactatttat agcttttagat agggcctccc ttcccctctt ctttctttgt tctctttcat 60
taaaccctt cccagtttt tttttatact ttaaaccctg ctccctcatgg ccttgccnt 120
ttctgaagct gcttcctctt ataaaatagc ttttgccgaa acatagtttt tttttagcag 180
atcccaaaat ataatgaagg ggatgggtggg atatttgtgt ctgtgttctt ataatatatt 240
attattcttc cttggttcta gaaaaataga taaatatatt tttttcagga aatagtgtgg 300
tgttccagc ttgatgttg tgggtggtg agtgagtga ttttcatgtg gctgggtggg 360
ttttgcctt tttctcttg cctgttctg gtgccttctg atggggctgg aatagttgag 420
gtggatggt ctacccttgc tgcctctgt ttgggacca gctggtgtt tttggtttgc 480
tttcttcagg ctctagggct gtgctatcca atacagtaac cacatgcggc tgtttaaagt 540
taagccaatt aaaatcacat aagattaaaa attccttctt cagttgact aaccacgtt 600
ctagaggcgt cactgtatgt agttcatggc tactgtactg acagcgagag catgtccatc 660
tgttggacag cactattcta gagaactaaa ctggcttaac gagtccagc ctgagctgtg 720
ctgggacgac ccttgtctcc ctgggtagga ggggggggaa tgggggagg ctgatgaggc 780
cccagctggg gcctgtgtc tgggaccctc cctctcctga gaggggaggc ctggtggctt 840
agcctgggca ggtcgtgtc cctcctgacc ccagtggctg cggtagaggg aaccaccctc 900
ccttgctgca ccagtggcca ttagctcccg tcaccactgc aaccagggg cccagctggc 960
tgggtcctc tctgccccca gtgccttcc ccttgggctg tgttgagtg agcacctcct 1020
ctgtaggcac ctctcacact gttgtctgtt actgattttt tttgataaaa agataataaa 1080
acctggtact ttctaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aaaaaggcg gccgctcgcg atctagaact ag 1172

<210> 128
<211> 891
<212> DNA
<213> Homo sapiens

<400> 128
accacgcgt ccgtgagatt ctctcccagg ccacaagaca tttcctgctc ggaaccttgt 60
ttactaat tccactgctt taaggccctg cactgaaaat gcaagctcag gcgccggtgg 120
tcgttgtag ccaacctgga gtcgggtccc gtccggcccc ccagaactcc aactggcaga 180
caggcatgtg tgactgttgc agcgactgag gactctgtct ctgtggcaca tttgtttcc 240
cgtgccttgg gtgtcaagt gacagtgata tgaatgaatg ctgtctgtgt ggaacaagcg 300
tcgcaatgag gactctctac aggaccgat atggcatccc tggatctatt tgtgatgact 360
atatggcaac tctttgctgt cctcattgta ctcttggcca aatcaagaga gatataca 420
gaaggagagc catgcgtact ttctaaaaac tgatggtgaa aagctcttac cgaagcaaca 480
aaattcagca gacacctctt cagcttgagt tcttcacat cttttgcaac tgaatatga 540
tggatatgct taagtacaac tgatggcatg aaaaaaatca aatttttgat ttattataaa 600
tgaatgtgt ccctgaactt agctaaatgg tgcaacttag tttctccttg ctttcatatt 660
atcgaatttc ctggcttata aactttttaa attacatttg aaatataaac caaatgaaat 720
attttactga taagattctt catgcttctt tgctctcctt aaaatgtctt tttcactagt 780

tagttccaag ggtcagtctc ataattttgt tcttatactt tgatttcctt tttctttttt 840
ttttttttta atagagttgg gggatcyttc yttctttttt tttttttttt t 891

<210> 129

<211> 2461

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (14)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1164)

<223> n equals a,t,g, or c

<400> 129

tttcaattct tggntttctg ttcccttcta ccaaaaccca gcataggact cagtgtacac 60
ttactttaaa acaaaaactc acatttctta gctgcagggtg atggcctcag tcattttaat 120
acagcctgga atgagtttgc ttattaccca gtcactttcg tagtgaatgt tcaaacccca 180
aagcaaatgt ttgcatctcc ttgtccataa aggagaaaagc caggttatag gagaaagaga 240
gagaaaggcg catgtctgtt tgcacagaga gaggcaattt tgtctacctt tcgagaatca 300
gttataaaca gaagggcctc ttaggatttt gagctctcct gacaatgaag gaaaagctct 360
cttgagtata caagttccac actcattacc tttcagtggt gacccatcac ccactacaat 420
ttgttgaagg aaggagagtgt cagagatatg ctttaaagtt gtactttagg ctgaaattct 480
ctctgtattt ggacctgcag atgttgtgta cagaaccgat gcatggcagg gtcaggaaagc 540
taggagagtgt aaggcgcttg tggagacagc tctgtagcaa atgaaacacg gaagcctccg 600
ggaatgtggt tgtgtcacca gcagcaggca ttccctgtgc ctccccaccc ccagtctcca 660
catccccagc agcctcttcc agaagcatgt caggggagacg gacagggtgt ctccttctcg 720
tgcaccacat ccaagtccac ccaccgtgga cgcagtgtga ctaaagtctg gccttgaaga 780
gaaaccctca gccagcttgt ctgttgtcca ggtccctcag tgtctctggt ttccctgccc 840
tgcataattt aagcattagt gccaaataca tctgtcataat tcctctccct ggagactgcg 900
aaatgtccag actttttcag actagtggag ggaaggaatg ttacataact aagtggaggc 960
ctggaactgt caggatttga cagggctgga ccagagaccc ttccgcctct gcctagtgtg 1020
tctgtcaggc aggcagcagc catcaatcca agaattgagcc atggcgga gacactktgtg 1080
gagaaaggaa cccagccgag gtctgagttt cagacagaaa ctggggagtt gggacatctt 1140
ctctgtgccca ggcttcattg acancatcgg tccttcattg asctgagccc aagccctcag 1200
tggaacctgt caagagcatc agcagcatgg agctgaagac cgagccctt gatgacttcc 1260
tgttcccagc atcatccagg cccagtggct ctgagacagc ccgctccgtg ccagacatgg 1320
acctatcttg gtccttctat gcagcagact gggagcctct gcacagtggc tccctgggga 1380
tggggcccct ggcacagagc tggagcccct gtgcactccg gtggtcacct gtactcccag 1440
ctgcactgct tacacgtctt ccttcgtctt cacctacccc gaggtgact ccttccccag 1500
ctgtgcagct gcccaccgca agggcagcag cagcaatgag ccttcctctg actcgtctag 1560
ctcaccacag ctgctggccc tgtgaggggg cagggaaagg gaggcagccg gcaccacaa 1620
gtgccactgc ccgagctggt gcattacaga gaggagaaac acatcttccc tagaggggtc 1680
ctgtagacct atggaggacc ttatctgtgc gtgaaacaca ccaggctgtg ggcctcaagg 1740
acttgaaagc atccatgtgt ggactcaagt ccttacctct tccggagatg tagcaaaacg 1800
catggagtgt gtattgttcc cagtgcact tcagagagct ggtagttagt agcatgttg 1860
gccaggcctg ggtctgtgtc tcttttctct ttctccttag tcttctcata gcattaacta 1920

atctattggg ttcattattg gaattaacct ggtgctggat attttcaa at tgtatctagt 1980
gcagctgatt ttaacaataa ctactgtgtt cctggcaata gtgtgttctg attagaaatg 2040
accaatatta tactaagaaa agatacgact ttattttctg gtagatagaa ataaatagct 2100
atatccatgt actgtagttt ttcttcaaca tcaatgttca ttgtaatgtt actgatcatg 2160
cattgttgag gtggtctgaa tgttctgaca ttaacagttt tccatgaaaa cgttttattg 2220
tgttttta at ttatttatta agatggattc tcagatattt atatttttat ttattttttt 2280
tctaccttga ggtcttttga catgtggaaa gtgaatttga atgaaaaatt taagcattgt 2340
ttgcttattg ttccaagaca ttgtcaataa aagcatttaa gttgaatgcg aaaaaaaaaa 2400
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaagggcggc cgctcgcgat ctagaactag 2460
t 2461

<210> 130

<211> 2197

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1381)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2194)

<223> n equals a,t,g, or c

<400> 130

gacgagatct gtgtcagaac gtgcgtgtga gcggatacaa aacccgagag aggcgtgagc 60
agcgtgtgtt ttgcgagcgg kagcgaggrg cgccggctgr ggtgtgtgct cctgagctct 120
tcagaaacca ggctgctttc aggaacattg ctgtggattc ccagggccta ttccactaga 180
agcaagatgg ctgaactcaa tactcatgtg aatgtcaagg aaaagatcta tgcagttaga 240
tcagttgttc ccaacaaaag caataatgaa atagtccttg tgctccaaca gtttgatttt 300
aatgtggata aagccgtgca agcctttgtg gatggcagtg caattcaagt tctaaaagaa 360
tggaatatga caggaaaaaa gaagaacaat aaaagaaaaa gaagcaagtc caagcagcat 420
caaggcaaca aagatgctaa agacaagggt gagaggcctg aggcagggcc cctgcagccg 480
cagccaccac agattcaaaa cgccccatg aatggctgcg agaaggacag ctctgccaca 540
gattctgcta acgaaaaacc agcccttatc cctcgtgaga aaaagatctc gatacttgag 600
gaaccttcaa aggcacttcg tggggtcaca ggcccaaata ttgagaaatc agtgaaggat 660
ttgcaacgct gcaccgtttc tctaactaga tatcgcgta tgattaagga agaagtggat 720
agttccgtga agaagatcaa agctgccttt gctgaattac acaactgcat cattgacaaa 780
gaagtttcat taatggcaga aatggataaa gttaaagaag aagccatgga aatcctgact 840
gctcgtcaga agaaagcaga agaactaaag agactcactg accttgccag tcagatggca 900
gagatgcagc tggccgaact cagggcagaa attaagcact ttgtcagcga gcgtaaatat 960
gacgaggagc tggggaaaagc tgcccgggtt tctgtgaca tcgaacagct gaaggcccaa 1020
atcatgctct gcggagaaat tacacatcca aagaacaact attcctcaag aactccctgc 1080
agctccctgc tgctctgct gaatgcgcac gcagcaacct ctgggaaaca gagtaacttt 1140
tcccgaat catccactca caataagccc tctgaaggca aagcggcaaa ccccaaatg 1200
gtgagcagtc tccccagcac cgccgacccc tctcaccaga ccatgccggc caacaagcag 1260
aatggatctt ctaacaaaag acggagattt aatccacagt atcataacaa caggctaaat 1320
gggcctgcca agtcgcaggg cagtgggaat gaagccgagc cactgggaaa gggaacagc 1380
ngccacgaac acagaagaca gccgcacaac ggcttccggc ccaaaaacaa aggcggtgca 1440

```
aaaatcaaga ggcttccttg gggatgaaga ccccgaggc cccggcccat tctgaaaagc 1500
cccggcgaag gcagcacgct gcagacacct cggaggccag gcccttccgg ggtagtgctg 1560
gtagggtttc acagtgcagt ctctgcccc aagagaataga agtttccaca gatgcagcag 1620
ttctctcagt cccggctgtg acgttggttg cctgagctag gaggaaaaag agcagttttc 1680
actcagtttt gggtccctgc ccgagggtgct gacccaattc gctgccaaaa gagtgtcaat 1740
cagaatatac aaatcccgta tggttgtgtc atcctctctt aatcattttt actaattcta 1800
ataatcagct ctagcttgct tcataatttt catggctttg cttgatctgt tgatgctttc 1860
tctcatcaag actttgcagc attttagcca ggcagtattt actcattatt aggaaaatca 1920
agatgtggct gaagatcaga ggctcagtta gcaacctgtg ttgtagcagt gatgtcagtc 1980
cattgattgt ctttagagag ttaatgttac aaaaaagaat tcttaataat cagacaaaca 2040
tgatctgctg aggacacatg cgcttttgta gaatttaaca tctgggtgtt ttctgaaaaa 2100
atatatatac atatattgct ttatttgaaa caaattaaaa tatgctgcat ttgaaaaaaa 2160
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaanaaa 2197
```

<210> 131

<211> 464

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (397)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (405)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (420)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (449)

<223> n equals a,t,g, or c

<400> 131

```
attcatctcg ttcattgcta caccocgtta gatcctcggc ggtgccaggc ttcgtcgaat 60
ggaaacccaa atgggtataa gcgttccact ttggcagcgt tgctaaagca accaccacca 120
gcgttggaac ggttggttac taccaattta tggaccgcct attaatggc atggttacag 180
ccaatacgat agtgagaaaa ccaaaacgca gcctagttag agtagaaagt gtaacaccat 240
tgcccacaac tggttgctgc ctactgtcgc taaggcggct taggcaaaat ttgctgcagc 300
gtacgcggag agttgtttac caaagatgcc ttacaaccct gcgctgaaaag ggggctgctg 360
cttatgccaa ttatgtttta aggttggtt ctgggtanttt tgcnaaatgg ttttgatttn 420
aggtttacca tttgttttcc ccaaattanc cctttgcgcg cccg 464
```

<210> 132

<211> 1950

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1941)
<223> n equals a,t,g, or c

<400> 132
ggggtggctt gccaccacaca gcaattgtga ctttgaaagg ccattccacc gttctatcaa 60
tccagctgct tcaggatgat gggcaacgtg gtaagaccag tgaattccac gaacatgagc 120
ccactgctgc acttctatgg ccgtaaatta acttgagaaa gtggaatcga attccgatgt 180
tgaattttcc ttctggcccc attcatgtgg cagggtggga ttcaggattg tcctcccggt 240
ctggcagcag ggccccagcg gcaccatgtc tgccctcgga gtcaccgtgg ccctgctggt 300
gtgggcggcc ttctcctgc tgggtgtccat gtggaggcag gtgcacagca gctggaatct 360
gccccagcg cctttccgc ttcccatcat cggaacctc ttccagttgg aattgaagaa 420
tattcccaag tccttcaccc ggttggtcca gcgcttcggg ccggtgttca cgctgtacgt 480
gggctcgcag cgcatggtgg tgatgcacgg ctacaaggcg gtgaaggag cgctgctgga 540
ctacaaggac gagtctcgg gcagaggcga cctcccgcg ttccatgcgc acagggacag 600
gggaatcatt ttaataatg gacctacctg gaaggacatc cggcggtttt ccctgaccac 660
cctccggaac tatgggatgg ggaacaggg caatgagagc cggatccaga gggaggccca 720
cttctctctg gaagcactca ggaagacca aggccagcct ttcgaccca cttcctcat 780
cggctgcgcg ccctgcaacg tcatagccga catcctcttc cgcaagcatt ttgactacaa 840
tgatgagaag ttcttaaggc tgatgtattt gtttaatgag aacttccacc tactcagcac 900
tccttggtc cagctttaca ataattttcc cagctttcta cactacttgc ctggaagcca 960
cagaaaagtc ataaaaaatg tggctgaagt aaaagagtat gtgtctgaaa ggtgaagga 1020
gcaccatcaa tctctggacc ccaactgtcc ccgggacctc accgactgcc tgctcgtgga 1080
aatggagaag gaaaagcaca gtgcagagcg cttgtacaca atggacggtg taaccgtgac 1140
tgtggccgac ctgttctttg cggggacaga gaccaccagc acaactctga gatatgggct 1200
cctgattctc atgaaatacc ctgagatcga agagaagctc catgaagaaa ttgacagggt 1260
gattgggcca agccgaatcc ctgccatcaa ggaataggca gagatgccct acatggatgc 1320
tgtggtgcat gagattcagc ggttcacac cctcgtgccc tccaacctgc cccatgaagc 1380
aaccgagac accattttca gaggatacct catccccaag ggcacagtcg tagtgccaac 1440
tctggactct gttttgtatg acaaccaaga atttctgat ccagaaaagt ttaagccaga 1500
acacttcctg aatgaaaatg gaaagttaa gtacagtgc tatttcaagc cattttccac 1560
aggaacacga gtgtgtgctg gagaaggcct ggctcgcag gagttgtttc tttgtgtgtg 1620
tgccattttg cagcatttta atttgaagcc tctcgttgac ccaaaggata tcgacctcag 1680
ccctatacat attgggtttg gctgtatccc accacgttac aaactctgtg tcattccccg 1740
ctcatgagtg tgtggaggac accctgaacc ccccgcttcc aaacaagatt tcgaattgtt 1800
tgaggtcagg attttctcaa ctgattcctt tctttgcata tgagtatttg aaaataaata 1860
ttttccaga atataaataa atcatcacat gattatttta actawaaaaa aaaaaaaaaa 1920
aaaaaaaaa aaaaaaaaaa naaaaaaaaa 1950

<210> 133
<211> 2093
<212> DNA
<213> Homo sapiens

<400> 133
cacgcgttcg cccacgcgts sgctgggttt ctgaactgct gggtttctgc ttgctcctct 60
ggagatgcag cgtctgttga ctccagtga ggcattctg caactgacaa gagcgggtgca 120

```

ggaaacctcc ctcacacctg ctgcctgct cccagtagcc caccaaaggt tttctacagc 180
ctctgctgtc cccctggcca aaacagatac ttggccaaag gacgtgggca tcctggccct 240
ggaggtctac tcccagccc aatatgtgga ccaaactgac ctggagaagt ataacaatgt 300
ggaagcagga aagtatacag tgggcttggg ccagaccctg atgggcttct gctcagtcca 360
agaggacatc aactccctgt gcctgacggg ggtgcaacgg ctgatggagc gcatacagct 420
cccatgggac tctgtgggca ggctggaagt aggcactgag accatcattg acaagtccaa 480
agctgtcaaa acagtgtctc tggaactctt ccaggattca ggcaatactg atattgaggg 540
catagatacc accaatgcct gctacgggtg tactgcctcc ctcttcaatg ctgccaaactg 600
gatggagtc agttcctggg atggtcgtta tgccatggtg gtctgtggag acattgccgt 660
ctatcccagt ggtaatgctc gtcccacagg tggggccgga gctgtggcta tgctgattgg 720
gccaagggc tctggccctg gagcgagggc tgaggggaac ccatatggag aatgtgtatg 780
acttctacaa accaaatttg gcctcggagt acccaatagt ggatgggaag ctttccatcc 840
agtgtacttt gcgggccttg gatcgatgtt acacatcata ccgtaaaaaa atccagaatc 900
agtggaagca agctggcagc gatcgaccct tcacccttga cgatttacag tacatgatct 960
ttcatacacc cttttgcaag atggtccaga agtctctggc tcgcctgatg ttcaatgact 1020
tcctgtcagc cagcagtgc acacaaacca gcttatataa ggggctggag gctttcgggg 1080
ggctaaagct ggaagacacc tacaccaaca aggacctgga taaagcactt ctaaaggcct 1140
ctcaggacat gttcgacaag aaaaccaagg cttcccttta cctctccact cacaatggga 1200
acatgtacac ctcatccctg tacgggtgcc tggcctcgt tctgtccac cactctgcc 1260
aagaactggc tggctccagg attggtgcct tctcttatgg ctctgggtta gcagcaagtt 1320
tcttttcatt tcgagtatcc caggatgctg ctccaggctc tcccctggac aagttggtgt 1380
ccagcacatc agacctgcca aaacgcctag cctcccga aa gtgtgtgtct cctgaggagt 1440
tcacagaaat aatgaaccaa agagagcaat tctaccataa ggtgaatttc tccccacctg 1500
gtgacacaaa cagccttttc ccaggctactt ggtacctgga gcgagtggac gagcagcatc 1560
gccgaaagta tgcccggcgt cccgtctaaa ggtgttctgc agatccatgg aaagcttct 1620
gggaaacgta tgctagcaga gcttctcccc gtgaatcata tttttaagat cccactctta 1680
gctggtaaat gaatttgaat cgacatagta gcccataag catcagccct gtagagtga 1740
gagccatctc tagcgggccc ttcattctc tccatgctgc aatcactgtc ctgggcttat 1800
ggtgctatgg actaggggtc ctttgtgaaa gagcaagatg gagcaatgga gagaagacct 1860
cttctgaaat cactggactc cagaaatgtg catgcagatc agctgttgcc ttcaagatcc 1920
agataaactt tcctgtcatg tgtagaact ttattattat taatattgtt aaacttctgt 1980
gctgttctg tgaatctcca aattttgtac cttgttctaa gctaataat agcaattaaa 2040
aagagagaaa gagggaaaaa aaaaaaactt tggggggggc ccggacccca att 2093

```

<210> 134

<211> 729

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (646)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (665)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (725)

<223> n equals a,t,g, or c

<400> 134

```
gctcgtgccg aattcggcac gagtcggcac gaggtccaag ggggtgtgtg ttcacgggaa 60
tgctgagtac cagcccgggt ctccagttaa ttccctccaag tgccaggact gcgtgtgcac 120
ggacaagggtg gacaacaaca ccctgctcaa cgtcatcgcc tgcacccacg tgccctgcaa 180
cacctcctgc agccctggct tcgaactcat ggaggccccc ggggagtgct gtaagaagtg 240
tgaacagacg cactgtatca tcaaacggcc cgacaaccag cacgtcatcc tgaagcccg 300
ggacttcaag agcgaccga agaacaactg cacattcttc agctgcgtga agatccacaa 360
ccagctcatc tcgtccgtct ccaacatcac ctgccccaac tttgatgcca gcatttgcac 420
cccgggctcc atcacattca tgcccaatgg atgctgcaag acctgcaccc ctcgcaatga 480
gaccagggtg ccctgctcca ccgtcccggt caccacggag gtttcgtacg ccggctgcac 540
caagaccgtc ctcatgaatc attgctccgg gtccctgcggg acatttgtca tgtamtcggc 600
caaggccagg ccctggacca cagcttgctc ctgctgcaaa gaaganaaaa ccagccagcg 660
tgagntggtc ctgactgccc aatggcggtc gctaacacac acctacacca catcgagact 720
gccantgcc                                     729
```

<210> 135

<211> 1189

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1160)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1175)

<223> n equals a,t,g, or c

<400> 135

```
gtaanacnga cagtacngta cggaattccc gggtcgaacc cacgcgtcct ggggtgagag 60
ggtgacgcat gtgccctctg gcagctctgct gctgtgtcca gagtccgact ccagctgggc 120
```


tgtaactggg cttggccccc gccttaggcc ccgccagcag gcgaagcagg gagatgtcag 180
actgctacac ggagctggag aaggcagtc tttgcctggg ggaaaacttc tacaatatg 240
tgtctaagta cagcctgggc aagaacaaga tcagcaagag cagcttccgc gagatgtcc 300
agaaagagct gaaccacatg ctgtcggaca cagggaaccg gaaggtgag gataagctca 360
tccagaacct ggatgccaat catgatgggc gcatcagctt cgatgagtag tggacctga 420
taggcggcat caccggcccc atcgccaaac tcatccatga gcaggagcag cagagcagca 480
gctagagacc cctttggcca caccttccag gcactggcct gatgccccgc cctgggtgctc 540
tccccaggct ccctcctcag cctcctgccc acccagggcc ctttactctc ttctccctcc 600
agaccttctc ctgaccttg ctgaactggg gtccctttgt gagtgtctca gtctagaggt 660
acctccctcc ctgggggggc tcagctcctg gagtcgcagg cccttggggc ccctctgtga 720
gatctcaatg ctgtctgggg accctaagag ttttctcacc tgttcagtct catctaacct 780
tccaatgtct gatgttctg ccaaattcct gcctgattct gggctccgtcc tgacctcaa 840
aggtcagctt ggtgcttgag gtctccctgc tcttgggtggc agtggttagca gcaacagcag 900
cagcagcagc agcagcagca gcagagacct ctccactttc ccttagcccc tctgctgggt 960
agagaggcac tttcaggac ttcctccag ctgcctcttc atctgggaat gagctaagca 1020
aggctgagcc tccctctgtt gcttgaaata atgatgatat aaaggctgga tttggagttt 1080
gtatccctg gtctctggg atgctcatta aaaccttccc actccttgga aaaaaaaaaa 1140
aaaaaargcg gsgggtttan aggatccaac ttacntaccc gtgcattgc 1189

<210> 136

<211> 1466

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1291)

<223> n equals a,t,g, or c

<400> 136

cngcttagtt cttccatttc acccttgca cagctgttg gccacggccc tgccctctgg 60
tctgccgtgc atctccctgt gccctcttgc tgccctgctg tctgctgagg taagggtgga 120
ggagggttac agccacatcc caccctctcg tccaatccca taatccatga tgaaagggtga 180
ggtcacgtgg cctccaggcc tgacttccca acctacagca ttgacgccaa cttggctgtg 240
aagggaagag aaaggatctg gccttgtggt cactggcatc tgagccctgc tgatggctgg 300
ggctctcggg catgcttggg agtgcagggg gctcgggctg cctggcctgg ctgcacagaa 360
ggcaagtgtc ggggctcatg gtgctctgag ctggcctgga ccctgtcagg atgggcccc 420
cctcagaacc aaactcactg tccccactgt ggcatgaggg cagtggagca coatgtttga 480
ggggaaggg cagagcgttt gtgtgttctg gggagggaag gaaaagggtg tggaggcctt 540
aattatggac tgttgggaaa aggggtttgt ccagaaggac aagccggaca aatgagcgac 600
ttctgtgctt ccagaggaa agcaggggagc agcagcttgg ctgactgctc agagtctgtt 660
ctgacgccct gggggttctc gtccaacccc agcaggggag cagcgggacc agccccacat 720
tccactgtg tctactgctt gaacctatct atttgtatt ttttgaaca gagttatgtc 780
ctaactattt ttatagattt gtttaattaa tagctgtc ttttcaagtt ctttttttat 840
tcatatttat gttcatggtt gattgtacct tcccaagccc gccagtgagg atgggaggag 900
gaggagaagg ggggccttgg gcgctgcagt cacatctgtc cagagaaatt ctttttggga 960

ctggaggcag aaaagcggcc agaaggcagc agccctggct cctttccttt ggcagggttg 1020
ggaagggcctt gccccagcc ttaggatttc agggtttgac tgggggcgtg gagagagagg 1080
gaggaacctc aataaccttg aaggtggaat ccagttattt cctgcgctgc gaggggttct 1140
ttatttcact cttttctgaa tgtcaaggca gtgaggtgcc tctcactgtg aatttgtggt 1200
ggcggggggc tggaggagag ggtggggggc tggctccgtc cctcccagcc ttctgctgcc 1260
cttgcttaac aatgccggcc aactggcgac ntcacggttg cacttccatt ccaccagaat 1320
gacctgatga ggaaatcttc aatasgacgc aaagatcaat gcaaaaattg ttctatctga 1380
acataaact ggagctcgtc caaaaagcac actacgacag aactggacgt tcgcagcgtc 1440
cattcaaagc agcctcctca tcaagt 1466

<210> 137

<211> 140

<212> DNA

<213> Homo sapiens

<400> 137

ggttcaccta cggaaacctt gttacgactt ttacttccto tagatagtca agttcgaccg 60
tcttctcagc gctccgccak ggccgtgggc cgaccccggc ggggccgac cgarggcctc 120
actaaaccat ccaatcggtg 140

<210> 138

<211> 4142

<212> DNA

<213> Homo sapiens

<400> 138

gaggggagcg agcggcgctt tgggggaggg gtcgcgtagg cgccctcacct gaccctgcgg 60
ccgtgcgggtt gctgctccgg ggcaggtctc cttccaggcc agggggcccg aatcatgtac 120
ataaagcagg tgattatcca gggttttcga agttacagag atcaaacaat ttagatccc 180
ttcagttcaa aacataatgt gattgtgggc agaaatggat ctggaaaaag taactttttt 240
tatgcaattc agtttgttct cagtgtatgag tttagtcac ttcgtccaga acagcggttg 300
gctttattgc atgaaggtag tggctcctcgt gttatttctg cttttgtgga gattattttt 360
gataattcag acaaccgggtt accaatcgat aaagaggag tttcacttcg aagagtatt 420
ggtgccaaaa argatcagta tttcttagac aagaagatgg tcacgaaaaa tgatgtgatg 480
aacctccttg aaagcgcttg tttttctcga agcaatcctt atttatattgt taaacaagga 540
aagatcaacc agatggcaac agcaccagat tctcagagat taaagctatt aagagaagta 600
gctggtacta gagtgtatga cgaacgaaag gaagaaagca tctccttaat gaaagaaca 660
gagggcaaac gggaaaaaat caatgagttg ttaaaataca ttgaagagak attacatact 720
ctmgagggaag aaaagggaaga actagctcag tatcagaagt gggataaaat gagacgagcc 780
ctggaatata ccattttacaa tcaggaactt aacgagactc gtgccaaact tgatgagctt 840
tctgctaagc gagagactag tggagaaaaa tccagacaat taagagatgc tcagcaggat 900
gcaagagata aaatggagga tatcgaacgc caagttagag aattgaaaac aaaaatttca 960
gctatgaaag aagaaaaaga acagcttagt gctgaaagac aagagcagat taagcagagg 1020
actaagttgg agcttaaagc caaggattta caagatgaac tagcaggcaa tagtgaacaa 1080
aggaaacgtt tattaagaaga gaggcagaag ctgcttgaaa aaatagaaga aaagcagaaa 1140
gaactggcag aaacagaacc caaattcaac agtgtgaaag agaaagaaga acgaggaatt 1200
gctagatttg ctcaagctac ccaggaaaga acggtccttt atgcaaagca gggtcgagga 1260
agccagttta catcaaaaga agaaagggat aagtggatta aaaaggaact caagtcttta 1320
gatcaggcta ttaatgacaa gaaaagacag attgtgcta tacataagga ttggaagac 1380
actgaagcaa ataaagagaa aaatctggag cagtataata aactggacca ggatcttaat 1440
gaagtcaaa ctcgagtaga agaactggac agaaaatatt acgaagtaaa aaataagaaa 1500

gatgaactmc aaagtgaag aaactacttg tggagagaag agaatgcaga acagcaagca 1560
cttgctgcta aaagagaaga tcttgaaaag aagcaacaac ttcttagagc agcaacagga 1620
aaggccattt taaatggaat agacagcata aacaaagtgc tagaccactt ccgtcgaaaa 1680
ggaataaacc agcatgttca aaatggctat catggtattg taatgaataa ctttgaatgt 1740
gaaccagctt tctacacatg cgtggaagtc actgctggaa acaggttatt ttatcacatt 1800
gttgattcag atgaagtcag cacgaagatt ttaatggagt ttaataaaat gaatcttcct 1860
ggagagggtta cttttctgcc tcttaacaag ttagatgtca gggatacagc ctatcctgaa 1920
accaatgatg ctattcctat gatcagcaaa ctgaggtaca atcccagatt tgacaaagct 1980
ttcaaacatg tgtttggaag gactcttatt tgctcgtagca tggaaagttc aaccagctg 2040
gcccgctgctt tcactatgga ctgtattact ttggaagggt accaagtcag ccacggtggg 2100
gctctaactg ggggttatta tgacacaagg aagtctcgac ttgaattgca aaaagatggt 2160
agaaaagcag aagaagaact aggtgaactt gaagcaaagc tcaatgaaaa cctgcgcaga 2220
aatattgaaa ggattaataa tgaaattgat cagttgatga accaaatgca acagatcgag 2280
acccagcaaa ggaaatttaa agcatctaga gatagcatat tatcagaaat gaagatgcta 2340
aaagagaaga ggcagcagtc agagaaaacc ttcattgccta agcaacgtag cttacagagt 2400
ttggaggcaa gcttgcatgc tatggagtct accagagagt cattgaaagc agaactggga 2460
actgatttgc ttttcaact gagtttgga gatcagaaga gagtagatgc actgaatgat 2520
gagattcgtc aacttcagca ggaaaacaga cagttgctaa atgaaagaat taaattagaa 2580
ggtattatta ctcgagtaga gacttatctc aatgagaatc tgagaaaacg cttggacca 2640
gtagaacagg aacttaatga gctgagagag acagaagggg gtactgttct cacagccaca 2700
acatcagaac ttgaagccat caataaaaga gtaaaagaca ctatggcacg atcagaagat 2760
ttggacaatt ccattgataa aacagaagct ggaattaagg agcttcagaa gagtatggag 2820
cgctggaaaa atatggaaaa agaactatg gatgctataa atcatgatac taaagaactg 2880
gaaaagatga caaatcggca aggcattgcta ttgaagaaga aagaagagtg tatgaagaaa 2940
attcgagaac ttggatcact tccccaggaa gcatgtgaaa agtaccagac actgagcctc 3000
aaacagttgt ttcgaaaact tgagcagtcg acacagaat taaagaagta cagccatggt 3060
aacaaaaagg ctttggrtca gtttgtaaat ttctccgagc agaaagaaaa gttaataaag 3120
cgtcaagaag agttagatak gggttacaaa tcgatcatgg aactgatgaa tgtacttgaa 3180
cttcggaaat atgaagctat tcagttaact ttcaaacagg tatctaagaa cttcagtga 3240
gtattccaga agttagtacc tgggtggcaa gctacttttg tgatgaagaa aggagatgtg 3300
gagggcagtc agtctcaaga tgaaggagaa gggagtgggt agagtgagag ggggtctggc 3360
tcacaaagca gtgtcccatc agttgaccag tttactggag ttggaattag ggtgtcattt 3420
acaggaaaac aaggtgaaat gagagaaatg caacagcttt caggtggaca gaaatccttg 3480
gtagcccttg ctctgatttt tgccattcag aaatgtgacc cggctccatt ttacttggtt 3540
gatgaaattg accaggctct ggatgctcag cacagaaagg ctgtgtcaga tatgattatg 3600
gaacttgctg tacatgctca gtttattaca actactttta ggcctgaact gcttgagtca 3660
gctgacaaat tctatggtgt aaagttcaga aataagggtta gtcattatga tgtgatcaca 3720
gcagagatgg ccaaagactt tgtagaagat gataccacac atgggttaatt ggaataact 3780
acactagggt ttgggagatg tatatagtaa tatgattctc ataccagga actgtaaatt 3840
taaacctaaa tatttgcca atagttttca gacttaaagc atcatagtcc ttttatattt 3900
gtctttgtat tttataagat actctgtaat gtcattgttg tactgatagt ttaagaattt 3960
aatttcctgt acaacttttt gtaaatgtt ctgctcctat tttaaatgtt ttgaaacatg 4020
ctaaatattc tttcctaatt attttatcac ttatactacc tttttatag cttcaattaa 4080
ataatcgggt ttatgactaa aaaaaaaaaa aaaaacysg gsgggggggc cggtaaccaa 4140
tt 4142

<210> 139

<211> 1747

<212> DNA

<213> Homo sapiens

<220>
<221> misc feature
<222> (1659)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1704)
<223> n equals a,t,g, or c

<400> 139
aagactccat ttcaaaaaaa aaaaaaaaaa aaaatccact catataaaaag gtgagctcag 60
ctcactgggc cattttctcag tggcttctcc atcctcattt gcaaacctca gagggataag 120
gcagttgaac ctgatgagca agaattataa cagcaaggaa acattaatgc ttagaattct 180
gagatccagc acaactcagt ctgtgggagc tcagctcgct gccagggat aggtatgacc 240
tatgtctgcc ttaggctgct gggagatgcc attctccagt ttcagaagca ggcagggcaa 300
aggtcaagac tgtggtattg gggctctttg gctctgaagg atcctggaac cactgatttt 360
ggtttattcc ctccagggtc taaagagaac aagagggtgct agctcttacc aaaacagatg 420
gtagagagag ttgctggcta tttaaaaagc tctttcatct ktttaattcac ytcttctttt 480
cacctcttta accactcctc aggaacagaa cacttctagg actgggggtc ttttagctcc 540
ataagcaagt gagcagatgg gacaagttag tcttttctcc ctagaacaa aggggatgcc 600
cagtggtttc cctttgcttc ccaacctaaa atttcaagtt taataaaata gcaattagca 660
gaagtgacca aattgggaga taattatcag tcatgaggaa agacacagat ttcggtcata 720
aagaatgtaa gggctataag tagaaacttt ctataacct aatgatgta tagaattatt 780
tttgagcagg agcagaaaaga ttaaataatga tcacttcata cttctaaatc agaaatagga 840
agattaaaac cacagaacag tttgtgattt ctattgctgt agctagggtat cttactctgt 900
ccactcttgt tcaagtatct aactcttctg gaaaccaa ataggctttaga agagattatc 960
ctatattcct atcagtataa tactaaaatg taacttttta atcatctggt ttttaaaaga 1020
taaacagttt agcccatctc tccagagagc aaacatagga atatgactca ggagcctcct 1080
agggcttctc atcagccctc acaccgctt cccctccaa cccacagcct ttgcttccag 1140
gtggcaggat tactactttg cctcttcagc agcatctact ctaggcatat tgatcatttt 1200
agacactggg agaagagaac ctcaaactag gaggaaga cagagcctcc acttagtttt 1260
gggaggggat ggcagacagt caaggagatg agcgtcctaa ggcagtgttg gatagggtca 1320
gatgcaccac ccatggagag gtttgtcaac acaaagacat ggaagggttag aggtttgtca 1380
acaaaaagac atggaagggt aggtttgtca acacaaagac atggaagatt agaggtttgt 1440
caacacaaag acacaggaag aatgggctgc agaagattta gatgttttcc atttgggcac 1500
attttactta gctggagaac taggttttaa acagcctggg taggaaaatt agaascaagc 1560
tggatgcagt gggctccatg cctgttaatc ccaacacttt tgggaggttc caggcaggaa 1620
agatcacttg ggcctaagaa gtccaacctg ccagcgganc ttaagattca caccacttgc 1680
actcccacc tgggggttaa ttanaaccaa aacccttttc tccgaaaaaa aaaaaaaaaa 1740
aaaaaaa 1747

<210> 140
<211> 1240
<212> DNA
<213> Homo sapiens

<400> 140
gcggacgcgc ctggaggcgc ggcggcaggg ctggggcggc gggcgggcgc cggatcatgga 60
acgccaaagag gagtctctgt ccgcgcggcc gccctggag accgaggggc tgcgttctc 120
gcacaccacg gtgggctccc tgctggccac ctatggctgg tacatcgtct tcagctgcat 180

ccttctctac gtggtctttc agaagctttc cgcccggcta agagccttga ggcagaggca 240
gctggaccga gctgcggctg ctgtggaacc tgatgttgtt gttaaaccgac aagaagcttt 300
agcagctgct cgactgaaaa tgcaagaaga actaaatgcg caagttgaaa agcataagga 360
aaaactgaaa caacttgaag aagaaaaaag gagacagaag attgaaatgt gggacagcat 420
gcaagaagga aaaagttaca aaggaaatgc aaagaagccc caggaggaag acagtccttg 480
gccttccact tcactctgtcc tgaaacggaa atcggacaga aagcctttgc ggggaggagg 540
ttataacccg ttgtctgggtg aaggaggcgg agcttgctcc tggagacctg gacgcagagg 600
cccgtcatct ggcggatgag gctaagaatc ttgttagtgt cacttttgac attagcaaga 660
tgaaccctta accctcgatt caattgcctt acgcacgctt ttcacagtga ctagccaagg 720
ggagggtggg ttgatttctg ttcctaacta cacctgcata tgcagggct ccagtcagca 780
aaaggtatag atgttgccctc taggcattgag gtcattggct acattctact tggagacagt 840
gattgcattc attgatttca tggttaattg ctagtggta ggtaaaggcc tctagatgat 900
tagcaatctt gataaaagag gcctagtaat gttcttttga ggttagaaat ccttgctgct 960
aggacagtct ctgtgacagg ttgcgttgaa tgatgtcttc cttatcaatg gtgagccac 1020
cagtgaggat tactgatgtg gacagttgat ggggtttgtt tctgtatatt ttttttatg 1080
tacagaactt tgtaaaaaacg aaactattta aaaaacaaga ataacatttt tagcatcttt 1140
attcaaggag atttatggac ttcaatttgt ctatcaaaca ttaaataagct ttttattaca 1200
acctccaaaa aaaaaaaaaa aaaaaaaaaa aaawaaacaa 1240

<210> 141

<211> 671

<212> DNA

<213> Homo sapiens

<400> 141

gtatcctgca ggtcccggtc cggaattccc gggctctacc acgcgtccgc gtgcgcaayg 60
tgccctggga atttggcgac gtaattcccg actatgtgct gggccagagc acctgtgccc 120
tgttcctcag cctccgctac cacaacctgc acccagacta catccatggg cggctgcaga 180
gcctggggaa gaacttcgcc ttgcgggtcc tgcttgctca ggtggatgtg aaagatcccc 240
agcaggccct caaggagctg gctaagatgt gtatcctggc cgactgcaca ttgatcctcg 300
cctggagccc cgaggaagct gggcgggtacc tggagacctg caaggcctat gagcagaaac 360
cagcggacct cctgatggag aagctagagc aggacttcgt ctcccgggtg actgaatgtc 420
tgaccaccgt gaagtcagtc aacaaaacgg acagtcagac cctcctgacc acatttggtg 480
ctctggaaca gtcctcgcgc gcatcaagag aagatctggc cttatgcccc ggctggggcc 540
ctcagaaagc ccggaggctg tttgatgtcc tgcacgagcc cttcttgaaa gtaccctgat 600
gacccagct gccaaaggaaa ccccagtggt aataataaat cgtcctccca ggccaggcaa 660
aaaaaaaaa a 671

<210> 142

<211> 3265

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (3234)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3256)

<223> n equals a,t,g, or c

<400> 142

```
ggcaccgtgg acattgagag cctgacgggc taccgcacct accgctgtgc ccacccctg 60
gccacactct tcaagatcct ggcgctcttc tacatcagcc tagtcatctt ctacggcctc 120
atctgcatgt atacactgtg gtggatgcta cggcgctccc tcaagaagta ctcgtttgag 180
tcgatccgtg aggagagcag ctacagcrac atccccgacg tcaagaacga cttcgccctc 240
atgctgcacc tcattgacca atacgacccg ctctactcca agcgcttcgc cgtcttctcg 300
tcggargtga gtgagaacaa gctgcggcag ctgaacctca acaacgagtg gacgctggac 360
aagctccggc agcggtctac caagaacgcg caggacaagc tggagctgca cctgttcatg 420
ctcagtggca tccctgacac tgtgtttgac ctggtggagc tggaggtcct caagctggag 480
ctgatecccc acgtgaccat cccgcccagc attgcccagc tcacgggcct caaggagctg 540
tggctctacc acacagcggc caagattgaa gcgcccgcgc tggccttctc gcgcgagaac 600
ctgcggggcg tgacatcaa gtacaccgac atcaaggaga tcccgctgtg gatctatagc 660
ctgaagacac tggaggagct gcacctgacg ggcaacctga gcgcggagaa caaccgctac 720
atcgtcatcg acgggctgcg ggagctcaaa cgctcaagg tgctgcggct caagagcaac 780
ctaagcaagc tgccacaggt ggacacagat gtgggcgtgc acctgcagaa gctgtccatc 840
aacaatgagg gcaccaagct catcgtcttc aacagcctca agaagatggc gaaacctgac 900
tgagctggag ctgatccgct gtgacctgga gcgcacccc cactccatct tcagcctcca 960
caacctgcag gagattgacc tcaaggacaa caacctcaag accatcgagg agatcatcag 1020
cttccagcac ctgcaccgcc tcacctgcct taagctgttg tacaaccaca tcgcctacat 1080
ccccatccag atcggcaacc tcaccaacct ggagcgcttc tacctgaacc gcaacaagat 1140
cgagaagatc cccaccagc tcttctactg ccgcaagctg cgctacctgg acctcagcca 1200
caacaacctg accttctctc ctgcccagat cggcctcctg cagaacctcc agaacctagc 1260
catcacggcc aaccggatcg agacgctccc tccggagytc ttccagtgcg ggaagctgcg 1320
ggccctgcac ctgggcaaca acgtgctgca gtcactgccc tccagggttg gcgagctgac 1380
caacctgacg cagatcgagc tgcggggcaa ccggtctggag tgcttgcctg tggagctggg 1440
cgagtgccta ctgctcaagc gcaggtcttg tgggtggagg ggacctgttc aacacactgc 1500
caccgagagt gaaggagcgg ctgtggaggg ctgacaagga gcaggcctga gcgaggcccg 1560
cccagcacag caagcagcag gaccgctgcc cagtcctcag gcccggaggg caggcctagc 1620
ttctcccaga actcccgac agccaggaca gcctcgtggc tgggcaggag cctggggccg 1680
cttggtgagtc aggcagagc gagaggacag tatctgtggg gctggccctt tttctccctc 1740
tgagactcac gtccccagc gcaagtgtt gtggaggaga gcaagtctca agagcgcagt 1800
atgtggataa tcagggtctc ctccctggag gccagctctg cccagggggc tgagctgcca 1860
ccagaggtcc tgggacctc actttagttc ttggtattta tttttctcca tctccacct 1920
ccttcatcca gataacttat acattcccaa gaaagtctag cccagatgga aggtgttcag 1980
ggaaaggtgg gctgcctttt ccccttgtcc ttatttagcg atgccgccg gcatttaaca 2040
cccacctgga cttcagcaga gtggtccggg gcgaaccagc catgggacgg taccacagca 2100
gtgccgggct gggctctgcg gtgcggtcca cgggagagca ggcctccagc tggaaaggcc 2160
aggcctggag cttgcctctt cagtatttgt ggcagtttta gttttttgtt ttttttttt 2220
taatcaaaaa acaatttttt taaaaaaaaa aagctttgaa aatggatggt ttgggtatta 2280
aaaagaaaaa aaaaacttaa aaaaaaaaaa aactaacgg ccagttagtt ggagtctcag 2340
ggcagggtgg cagtttccct tgagcaaaac agccagacgt tgaactgtgt ttcctttccc 2400
tgggcgcagg gtgcagggtg tcttccggat ctggtgtgac cttggtccag gagtctatt 2460
tgttctctgg gagggaggtt tttttgtttg ttttttgggt ttttttgggt tcttgttttc 2520
tttctcctcc atgtgtcttg gcaggcactc atttctgttg ctgtcggcca gagggaatgt 2580
tctggagctg ccaaggaggg aggagactcg ggttggttaa tccccgatg aacggtgctc 2640
cattcgaccc tcccctctc gtgcctgccc tgctctcca cgcacagtgt taaggagcca 2700
agaggagcca cttcgcgccag actttgtttc ccaccgcct gcggcatggg tgtgtccagt 2760
gccaccgctg gcctccgctg cttccatcag ccytgtcgcc acctggtcct tcatgaagag 2820
cagacactta gaggtggtc gggaaatggg aggtcgcccc tgggagggca ggcgttggtt 2880
```

ccaagccggt tcccgtccct ggcgcctgga gtgcacacag cccagtcggc acctggtggc 2940
tggaagccac cctgcttttag atcactcggg tccccacctt agaaggggtcc cgccttaga 3000
tcaatcacgt ggacactaag gcacgtttta gagtctcttg tcttaatgat tatgtccatc 3060
cgtctgtccg tccatttgtg ttttctgcgt cgtgtcattg gatataatcc tcagaaataa 3120
tgcacactag cctctgacaa ccatgaagca aaaatccggt acatgtgggt ctgaacttgt 3180
agactcggtc acagtatcaa ataaaatcta taacagaaaa aaaaaaaaaa gggngccggtc 3240
taaagatcaa cttctncctt gatca 3265

<210> 143

<211> 765

<212> DNA

<213> Homo sapiens

<400> 143

gccacgcgt ccgccacgc gtccgcggac gcgtgggggg tgacattgag ctcaccagcg 60
ccaccgtccc cggcgaagtt ctgcgtggt cggcggagta gcaagtggcc atggggagcc 120
tcagcgggtct gcgcctggca gcaggaagct gttttaggtt atgtgaaaga gatgtttcct 180
catctctaag gcttaccaga agctctgatt tgaagagaat aaatggattt tgcacaaaac 240
cacaggaaaag tcccggagct ccatcccgca cttacaacag agtgccttta cacaaacctt 300
cggattggca gaaaaagatc ctcatatggt caggctcgctt caaaaaggaa gatgaaatcc 360
cagagactgt ctggttgag atgcttgatg ctgcaaagaa caagatgcga gtgaagatca 420
gctatctaatt gattgccctg acggtggtag gatgcattct catggttatt gagggcaaga 480
aggctgccc aagacacgag actttaacaa gcttgaactt agaaaagaaa gctcgtctga 540
aagaggaagc agctatgaag gccaaaacag agtagcagag gtatccgtgt tggttgatt 600
ttgaaaatcc aggaattatg ttataacgtg cctgtattaa aaaggatgtg gtatgaggat 660
ccatttcata aagtatgatt tgcccaaacc tgtaccattt ccgtatttct gctgtagaag 720
tagaataaaa ttttcttaaa taaaaaaaaa aaaaaaaaac tcgag 765

<210> 144

<211> 1694

<212> DNA

<213> Homo sapiens

<400> 144

gccacgcgt ccgggttgct ctattattat tattattatt atttgccttt ttttaatcac 60
catttatttg tgtaattctt ctccattgat tggctgtcat tcttataata tctgaactac 120
cccataaaat aattctctga cttaaaggca tttagctttt gcttatcgtt tttacatcct 180
ctattcaact aagacactgt cttagagatt attttttcca gatggatcgt tggcctaaat 240
tttcacactt ctcccctggt catccttttt cctcttccct gcttcctggg aataaaagga 300
acttttttaa aaaaattaat tagtccacag gwctmattat ctttctttat gattaatcta 360
tgactttttg gtacaagaac aatggaaaaa gtgaattaag gtaatgaaca aaacctttca 420
cccacttaaa cattttccag ttttgagatt cctcttcgtg tttgtggtgt cttccccttg 480
ttaccccttc tgcccttttt ctctgactat ggtaatgttg tctttaggct catatcagtc 540
tcccagagac attctgcagt cattatcacc ttttgggtg gattttattt tgttttattt 600
tgtttttttt aaaaaataa ctttttaaca ttggtgcata tttgcttggg atagagcttg 660
tgtaatttac caatcgtatt gattgtaagt gattgtgccc tgagaggta tatttaacaa 720
gacaaaaata atcttggtta ataaaggagc ccatgagatt tgagtcagggt tgtaagtga 780
atcacttaca cttttggata gaatttatac tcctgctctt ataaatcagt ggtagactta 840
ccatttttta aagttttctt gcattttttt gtttttttat tgccacagct ccctattctt 900
tcttgctgc ctccaccccc ctgttcagga aaaaaaaaaa ttgagcctta aagtgcagc 960
tgatttttta attgctgaat tttgtgaaat tttacttttt ccaagtgttt ccaacttta 1020

```
aaagagaagt gaagacaaat aggttggaat ggtgaagaca aatggrttgg ratttcacag 1080
gctgtgaata attccttagg atctggcaaa ccgtgaagtc ttatttgaag accttatctc 1140
ctgagagttc ttttggagta ggaaaaagaa ccctatttga aatagaccgt ttttctcttg 1200
ttttaaatct gtttaatat tctgattttt aagcagcttt caaaacaagt gtggtggaaa 1260
aaaagaaaata gtagtaggaa gatgtttagg gcagcagaac tctgggtcta aataagtaca 1320
tgttcccact tgttgccgat ttttgagagt actagggcca tctttctcaa ttttgtatta 1380
tttgtgtgca tgtttatatc aaagatgccc attttgtaa aatgctattt cctttattac 1440
cttggaact gactcagcct catgttgctc ctaattagtg tttaaggctc ccatgagttg 1500
cagataaaat gatttatttt aacaagtaga aggaggtgat tcaccttttg gattgtaaat 1560
atatgaaaat gtctacaagg tctttatctg ctttctgtca gcatttatat taaatgataa 1620
attaatgagg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1680
aaaaaagggc ggcc 1694
```

<210> 145

<211> 823

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (182)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (731)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (743)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (749)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (755)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (817)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (821)

<223> n equals a,t,g, or c

<400> 145

```
tgtcacagac tacttcatca gtcgcttadc tggagctggg accaagggtt ctgccaggcc 60
ttgggatcag ctcttggggg tcagagcagc cttcccacat cctctggcac tgctgaactt 120
ttgcagcagc tctttcctcc tctyttggat gcccttcgag agcccagggt acgacggatt 180
tnctgccagc ctgcagatcc tgcgcctgtc gccctgcagg tctctgtacc cttcagacca 240
ccttgctctg gttcctgggc agagctcagc agtacttggc agcatgggac ccagcttcct 300
tcctgctcct gatccaaaag gacttacctc ctctgttgca tgaggcagaa gctttgtata 360
gcctggcctc agaggaaagc ttagctctgg aagtggagca gcagctgggc ctggagatcc 420
agaagctgac tgcacagatc cagctcctgc ctgaagagtc actaagtgtc ttttctcaag 480
aatgtcataa acaagccatg caagggttca agctctacat gccacggggt cggtagctgc 540
ggcttcgtct ctgtcctgaa cctcccagtg ctcctagtga gtatgctggt ttagtggtcc 600
gcaccgtact ggagcctgtg ttgcaaggat tgcaagggtt gcacctcaag cccaggcccc 660
tgcccttggt caggctctga cggscatcgt ggggtgcctgg cttgaccaca ttcttaccce 720
tgggattcgg nttcaacctg canggagcnc tkcancttaa acaagacttt ggaatggtca 780
ggagtttgtg gaaaagagca tggacctgtc cctgatntcg nca 823
```

<210> 146

<211> 1134

<212> DNA

<213> Homo sapiens

<400> 146

```
caaaagcata gagaaattat aaaattcaag aacagatggt agaatggaaa ctgatctaga 60
ggttataata aaggataata gtcttgtgct gacaccatca cacatcaaag cctacatggt 120
gatgactctt caaggattag aatatttaca tcaacattgg atcctacata gggatctgaa 180
accaaacaac ttgttgctag atgaaaatgg agttctaaaa ctggcagatt ttggcctggc 240
caaattcttt gggagcccca atagagctta tacacatcag gttgtaacca ggtggtatcg 300
ggccccgag ttactatttg gagctaggat gtatggtgta ggtgtggaca tgtgggctgt 360
tggtgtgata ttagcagagt tacttctaag ggttcctttt ttgccaggag attcagacct 420
tgatcagcta acaagaatat ttgaaacttt gggcacacca actgaggaac agtggccgga 480
catgtgtagt cttccagatt atgtgacatt taagagtttc cctggaatac ctttgcacat 540
catcttcagt gcagcaggag acgacttact agatctcata caaggcttat tcttatttaa 600
tccatgtgct cgaattacgg ccacacaggc actgaaaatg aagtatttca gtaatgcgcc 660
agggccaaca cctggatgtc agctgccaaag accaaaactgt ccagtggaaa ccttaaagga 720
gcaatcaaat ccagcttttg caataaaaag gaaaagaaca gaggccttag aacaaggagg 780
attgcccaag aaactaattt tttaaagaga acactggaca acattttact actgagggaa 840
atagccaaaa aggcaataaa tggaaaaata gtaaacatta agtaaatgct gtagaagtga 900
gtttgtaaat attctacaca tgtaaaatat gtaaaactat gggttatttt tattaatgt 960
attttaaaat aaaaatttaa ttctggtttt tctgattaga gtgcmaaagt gagaaaagt 1020
caatactctt gaatgtagaa ttgaaaatgc attagggaaa acttaataaa aattattacc 1080
agttatttgg aagatctgac ccatatagta tcacaaatct gtagtagcat gggt 1134
```

<210> 147

<211> 1486

<212> DNA

<213> Homo sapiens

<400> 147

```
gcggacgcgt ggggggcggg gtgtcgtttc ctttcgctga tgcaagagcc tagtgcggtg 60
```

gtgggagagg tatcggcagg ggcagcgctg ccgcccgggc ctggggctga cccgtctgac 120
ttcccgtccg tgccgagccc actcgagccg cagccatgtc tggggacgag atgatttttg 180
atcctactat gagcaagaag aaaaagaaga agaagaagcc ttttatgtta gatgaggaag 240
gggataccca aacagaggaa acccagcctt cagaaacaaa agaagtggag ccagagccaa 300
ctgaggacaa ggatttggaa gctgatgaag aggacactag gaaaaaagat gcttctgatg 360
atctagatga cttgaacttc tttaatcaaa agaaaaagaa gaaaaaaact aaaaagatat 420
ttgatattga tgaagctgaa gaaggtgtaa aggatcctta gattgaaagt gatgttcaag 480
aaccaactga accagaggat gaccttgaca ttatgcttgg caataaaaag aagaaaaaga 540
agaatgttaa gttcccagat gaggatgaaa tactagagaa agatgaagct ctagaagatg 600
aagacaacaa aaaagatgat ggtatctcat tcagtaatca gacaggccct gcttgggcag 660
gctcagaaag agactacaca tacgaggagc tgctgaatcg agtggtcaac atcatgaggg 720
aaaagaatcc agatatggtt gctggggaga aaaggaaatt tgtcatgaaa cctccacaag 780
tcgtccgagt aggaaccaag aaaacttctt ttgtcaactt tacagatatc tgtaactat 840
tacatcgtca gcccacacat ctccttgcac ttttgttggc tgaattgggt acaagtgggt 900
ctatagatgg taataaccaa cttgtaatca aaggaagatt ccaacagaaa cagatagaaa 960
atgtcttgag aagatatatc aaggaatatg tcacttgta ccatgccga tcaccggaca 1020
caatcctgca gaaggacaca cgactctatt tcctacagtg cgaaacttgt cattctagat 1080
gttctgttgc cagtatcaaa accggcttcc aggctgtcac gggcaagcga gcacagctcc 1140
gtgccaagc taactaattt gctaactact gattttgcaa agcttgttgt ggagatgtgg 1200
ctggacaggt ttgccatcag agtgatata ccgttgtatt aaaaacaaga taaaaaagct 1260
gccaagattt ttggcgagtg gttggtctga agtccttgca agacgctgat gctcaagctg 1320
ttgacatact cattgcctac tttaacacct gtcagagaaa cgtgatattg ggtaaggagg 1380
tgctttttta aaatcggtca tagacttctg taaaatgcaa gataaattaa agttattata 1440
acagtgataa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaa 1486

<210> 148

<211> 153

<212> DNA

<213> Homo sapiens

<400> 148

cgctgtgca ggcgaagatc gaggtcgtc cggcagagga ctccttcaca ggcttcgtcc 60
ggaccctgta ctttgctgac acctacctga aggagtgcc aggtgagcg gccaggcctc 120
cagaacatgg agctggcgcc tgtgcagcgc aag 153

<210> 149

<211> 882

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (14)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (25)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (870)
<223> n equals a,t,g, or c

<400> 149
acgaaggcca aagntgggtgt cacnncgggtg cgccgtctag actagtggat cccggctcag 60
gattcgcacg tgccctcctgg tctcagtatg gcgctgtcct gggttcttac agtcctgagc 120
ctcctacctc tgctggaagc ccagatccca ttgtgtgcc aacctagtacc ggtgcccac 180
accaacgcca ccctggaccr gatcactggc aagtggtttt atatcgcatc ggcctttcga 240
aacgaggagt acaataagtc ggttcaggag atccaagcaa ctttctttta cttcaccccc 300
aacaagacag aggacacgat ctttctcaga gagtaccaga cccgacagga ccagtgcac 360
tataacacca cctacctgaa tgtccagcgg gaaaatggga ccctctccag atacgtggga 420
ggccaagagc atttgcgtca cttgctgac ctcagggaca ccaagacct catgcttgct 480
tttgacgtga acgatgagaa gaactggggg ctgtctgtct atgctgacaa gccagagacg 540
accaaggagc aactgggaga gttctacgaa gctctcgact gcttgcgcat tccaagtca 600
gatgtcgtgt acaccgattg gaaaaaggat aagtgtgagc cactggagaa gcagcacgag 660
aaggagagga aacaggagga gggggaatcc tagcaggaca cagccttga tcaggacaga 720
gacttggggc catcctgccc ctccaacccg acatgtgtac ctcagctttt tccctcactt 780
gcatcaataa agcttctgtg tttggaacag ctaaaaaaaa aaaaaaaaaa aaaaaaaaaa 840
actcgagggg gggcccgkaa cccaatcgcn tgatattatt ag 882

<210> 150
<211> 1508
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (155)
<223> n equals a,t,g, or c

<400> 150
ctaaggaggg gtgaaccggc ccaggtcgga aacggagcag gtcaaaactc ccgtgctgag 60
gtgggaggat cgcttgagcc caggagtctt gggctgtagt gcgctatgcc gatcgggtgt 120
ccgcantcta gccttggaac cagtgaaga ctgtntcaaa aacagcaaca garagcagga 180
cgtgagactt ctacctgctc actcagaatc atttctgcac caacctggc cacgtttgtg 240
gagctcagta ccaaagccaa gatgcccatt gtgggcctgg gcacttgga gtctcctctc 300
ggcaaagtga aagaagcagt gaagtgggc attgatgcag gatatcggca cattgactgt 360
gcctatgtct atcagaatga acatgaagtg ggggaagcca tccaagagaa gatccaagag 420
aaggctgtga agcgggagga cctgttcac gtcagcaagt tgtggccac tttctttgag 480
agacccttg tgaggaaaagc ctttgagaag accctcaagg acctgaagct gagctatctg 540
gacgtctatc ttattcactg gccacagga ttcaagtctg gggatgacct tttcccaaa 600
gatgataaag gtaatgccat cggtggaata gcaacgttct tggatgcctg ggaggccatg 660
gaggagctgg tggatgaggg gctggtgaaa gcccttgggg tctccaattt cagccacttc 720
cagatcgaga agctcttgaa caaacctgga ctgaaatata aaccagtgc taaccaggtt 780
gagtgacc cctacctcac gcaggagaaa ctgatccagt actgccactc caaggccatc 840

accgttacgg cctacagccc cctgggctct ccggatagac cttgggccaa gccagaagac 900
ccttcctgc tggaggatcc caagattaag gagattgctg caaagcaca aaaaaccgca 960
gcccagggttc tgatccggtt ccatatccag aggaatgtga ttgtcatccc caagtctgtg 1020
acaccagcac gcattgttga gaacattcag gtctttgact ttaaattgag tgatgaggag 1080
atggcaacca tactcagctt caacagaaac tggagggcct gtaacgtgtt gcaatcstct 1140
catttggaag actatccctt cgatgcagaa tattgaggtt gaatctcctg gtgagattat 1200
acaggagatt ctctttcttc gctgaagtgt gactacctcc actcatgtcc cattttagcc 1260
aagcttattt aagatcacag tgaacttagt cctgttatag acgagaatcg aggtgctgtt 1320
ttagacattt atttctgtat gttcaactag gatcagaata tcacagaaaa gcatggcttg 1380
aataaggaaa tgacaatttt ttccacttat ctgacagaa caaatgttta ttaagcatca 1440
gaaactctgc caacactgag gatgtaaaga tcaataaaaa aaataataat cataaaaaaa 1500
aaaaaaaaa 1508

<210> 151

<211> 1232

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1213)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1217)

<223> n equals a,t,g, or c

<400> 151

naactgngct agcatgccca gccacagag agcctccact agagtgatgc taagtggaaa 60
tgtgagggtgc agctgccaca gagggccccc accagggaaa tgtctagtgt ctagtggatc 120
caggccacag gagagagtgc cttgtggagc gctgggagca ggacctgacc accaccagga 180
ctccagcctg ggtgacaggg tgaacgccat ctcaaaaaat aaaaattaaa aaataaaaaa 240
agaacctgga tctcaattta atttttcata ttcttgcaat gaaatggact tgaggaagct 300
aagatcatag ctagaataac agataattcc acagcacatc tctagcaaat ttagcctatt 360
cctattctct agcctattcc ttaccacctg taatcttgac catatacctt ggagttgaat 420
attgttttca tactgctgtg gtttgaatgt tccctccaac actcatgttg agacttaatc 480
cctaattgtg caaactgaa aggtggggcc tttgagatgt gattggatcg taaggctgtg 540
ccttcattca tgggttaatg gattaatggg ttatcacagg aatgggactg gtggctttat 600
aagaagagga aaagagaact gagctagcat gccagccca cagagagcct ccactagagt 660
gatgctaagt ggaaatgtga ggtgcagctg ccacagaggg cccccaccag ggaaatgtct 720

agtgtctagt ggatccaggc cacaggagag agtgccttgt ggagcgctgg gagcaggacc 780
tgaccaccac caggacccca gaactgtgga gtcagtggca gcatgcagcg ccccttggg 840
aaagcttttag gcaccagcct gcaaccatt cgagcagcca cgtaggctgc acccagcaaa 900
gccacaggca cggggctacc tgaggccttg gggggccaat ccctgctcca gtgtgtccgt 960
gaggcagcac acgaagtcaa aagagattat tctcttccca cagatacctt ttctctccca 1020
tgacccttta acagcatctg cttcattccc ctcaccttcc caggctgac tgaggtaaac 1080
tttgaagtaa aataaaaagct gtgtttgagc atcaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaaaaaaa aangggnggc cgttttaaaag ga 1232

<210> 152

<211> 999

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (917)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (951)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (995)

<223> n equals a,t,g, or c

<400> 152

gcgcagctcc gccggcgcct ggtcccagcg ccgcggcgcg cgcgtccccg gcccacccat 60
ggertcctcc gcggccggct gcgtggtgat cgttggcagt ggagtcattg ggygaagctg 120
ggccatgctg tttgccagtg gaggcttcca ggtgaaactc tatgacattg agcaacagca 180
gataaggaay gccctggaay acatcagaaa ggagatgaag ttgctggagc aggcaggttc 240
tctgaaaggc tccctgagtg tggaagagca gctgtcactc atcagtgggt gtcccaatat 300
ccaagaagca gtagagggg ccatgcwcat ycwgtgaatc cgccatacta catcccctg 360
gttgagctgg tccccacccc ggagacggcc cctacgacag tggacagaac ccacgccctg 420
atgaagaaga ttggacagtg ccccatgcga gtccagaagg aggtggccgg ctctggtctg 480
aaccgcctgc aatatgcaat catcagcgag gcctggcggc tagtggagga aggaatcgtg 540
tctcctagtg acctggacct tgtcatgtca gaagggttgg gcatgcggtg tgcattcatt 600
ggacccctgg aaaccatgca tctcaatgca gaaggtatgt taagctactg cgacagatac 660
agcgaaggca taaaacatgt cctacagact tttggaccca ttccagagtt ttccagggcc 720
actgctgaga aggttaacca ggacatgtgc atgaaggctc ctgatgacct ggagcactta 780
gctgccagga ggcagtggag ggacgagtg ctcagtgagc tcgccaagtt gaaragtcaa 840
gtgcagmyaa sctggaratt tccacctttt cttttcagct tgattgcatt tgactatatt 900
ttacagccag tgattgnagt ttcattgtaa tatgtggcaa aatatttttg naattatttt 960
ctaataccctt tcttaagtac tctggggccc tgcanttat 999

<210> 153

<211> 1212

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (794)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1047)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1146)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1176)
<223> n equals a,t,g, or c

<400> 153
cacgcccacc tctctcctgg agcgctgggc cttcgctggc cgcaccggca gccatgagct 60
cggagatgga gccgctgctc tgggcctgga gctattttag gcgcaggaag ttccagctct 120
ggccgatcta tgcagcaga tgctggagaa gtccccttat gaccaggcgg ctgctgcaga 180
tgggcattta taacggccag ctttttaaca atctggggct gtgttgcttc tatgccagc 240
agtatgatat gactctgacc tcatttgaac gtgccctttc tttggctgaa aatgaagaag 300
aggcagctga tgtctggtac aacttgggac atgtagctgt gggaatagga gatacaaat 360
tggcccatca gtgcttcagg ctggctctgg tcaacaacaa caaccacgcc gaggcctaca 420
acaacctggc tgtgctggag atgcggaagg gccacgttga acaggcaagg gcactattac 480
aaactgcac atcattagca ccccatatgt atgaaccgca ttttaatttt gcaacaatct 540
ctgataagat tggagatctg cagagaagct atgttgctgc gcagaagtct gaagcagcat 600
ttccagacca tgtggacaca caacatttaa ttaacaatt aaggcagcat tttgctatgc 660
tctgattggt ccttagacca catatgttct tatgaagcmg cattatgcaa ggggaaaaaa 720
gcactatgtc tgtgtatgta tgtatatagt gtaatacgtat tattttaaca aacctgtcct 780
tgatattagt taanggtgac acataagggt gacacagaat gtgtaatgca aatttcatag 840
taatagtaac tttataaaat aatattataa aatacaggat ttaaaccttt ctaaatagat 900
cctgaaactg tctctcacat tatatagtag atgtttgttt ataattgttt caaaacattt 960
tgggtgaattt cctcaatgtt ttataaatgt acatttttta agtccttaag ctgactctta 1020
gccatcatgt agcttaagga gtctggnaat ctggccattg aaaactggca cctttgaagc 1080
cagggtgtgt tagcatgtgc ctatagtcac agcttacttg gggaggtggg aggtggggag 1140
ggattnatga aattaggagg acttttcctt ttaagnactt ttaaaaaaat ggtattttga 1200
aaacctattt tt 1212

<210> 154
<211> 2361
<212> DNA
<213> Homo sapiens

<220>

<221> misc feature

<222> (92)

<223> n equals a,t,g, or c

<400> 154

```
ggagccttgc cttgctgctc tacctccacc atgccaagtg gtcccaggct gcacccatgg 60
cagaaggagg agggcagaat catcacgaag gntgaagttc atggatgtct atcagcgcac 120
actgccatcc aatcgagacc ctggtggaca tcttccagga gtaccctgat gagatcgagt 180
acatcttcaa gccatcctgt gtgcccctga tgcgatgcgg gggctgctgc aatgacgagg 240
gcctggagtg tgtgcccact gaggagtcca acatcaccat gcagattatg cggatcaaac 300
ctcaccaagg ccagcacata ggagagatga gcttcctaca gcacaacaaa tgtgaatgca 360
gaccaaagaa agatagagca agacaagaaa aatgtgacaa gccgaggcgg tgagccgkgc 420
aggagggaagg agcctccctc aggggttcgg gaaccagatc tctcaccagg aaagactgat 480
acagaacgat cgatacagaa accacgctgc cgccaccaca ccatcaccat cgacagaaca 540
gtccttaatc cagaaacctg aaatgaagga agaggagact ctgcgcagag cactttgggt 600
ccggagggcg agactccggc ggaagcattc ccggggcggg gaccagcac ggtccctctt 660
ggaattggat tcgccatttt attttcttg ctgctaaatc accgagcccg gaagattaga 720
gagttttatt tctgggattc ctgtagacac acccaccac atacatacat ttatatatat 780
atatattata tatatataaa aataaatatc tctattttat atatataaaa tatatatatt 840
ctttttttaa attaacagtg ctaatgttat tgggtgtctc actggatgta tttgactgct 900
gtggacttga gttgggaggg gaatgttccc actcagatcc tgacagggaaggaggaggaga 960
tgagagactc tggcatgatc ttttttttgt cccacttggg ggggccaggg tcctctcccc 1020
tgcccaggaa tgtgcaaggc cagggcattg gggcaaatat gaccagttt tgggaacacc 1080
gacaaaccca gccctggcgc tgagcctctc taccacaggc cagacggaca gaaagacaga 1140
tcacaggtag agggatgagg acaccggctc tgaccaggag tttggggagc ttcaggacat 1200
tgctgtgctt tggggattcc ctccacatgc tgcacgcgca tctcgcccc aggggcactg 1260
cctggaagat tcaggagcct gggcgccctt cgcttactct cactgtctt tgagttgccc 1320
aggagccac tggcagatgt cccggcgaag agaagagaca cattgttga agaagcagcc 1380
catgacagct ccccttcctg ggactcgccc tcactctctt cctgtctccc ttcctgggg 1440
gcagcctaaa aggacctatg tctcacacc attgaaacca ctagtcttgt ccccccagga 1500
gacctgggtg tgtgtgtgtg agtgggtgac ctctctccat cccctgggtc ttccttccc 1560
ttcccagggc acagagagac agggcaggat ccacgtgccc attgtggagg cagagaaaag 1620
agaaagtgtt ttatatacgg tacttattta atatccctt ttaattagaa attaaaacag 1680
ttaatttaat taaagagtag ggtttttttt cagtattctt ggttaatat taatttcaac 1740
tatttatgag atgtatcttt tgctctctct tgctctctta tttgtaccgg tttttgtata 1800
taaaattcat gtttccaatc tctctctccc tgatcgggtg cagtcaactag cttatcttga 1860
acagatatatt aattttgcta acactcagct ctgcctccc cgatcccctg gctccccagc 1920
acacattcct ttgaaataag gtttcaatat acatctacat actatatata tatttggcaa 1980
cttgattttg tgtgtatata tatatatata tgtttatgta tatatgtgat tctgataaaa 2040
tagacattgc tattctggtt ttatatatga aaaacaaaac aagaaaaaat agagaattct 2100
acatactaaa tctctctcct tttttaattt taatatgtgt tatca'tttat ttattggtgc 2160
tactgtttat ccgtaataat tgtggggaaa agatattaac atcacgtctt tgtctctagt 2220
gcagtttttc gagatatattc gtagtacata tttattttta aacaacgaca aagaaatata 2280
gatatatctt aaaaaaaaaa aagcattttg tattaagaa ttttaattctg atctcaaaaa 2340
aaaaaaaaaa aaaaaggggg g 2361
```

<210> 155

<211> 1831

<212> DNA

<213> Homo sapiens

<400> 155
aaaggtgaga atgcacaaag acagctctgg gttgggtacc acagttttgc ttggtagaaa 60
gaaaccagtg taggaaagga gacgccacca gacatcttca acagacaaga ttctttctgc 120
ctttttcaaa agatgctctc tgcagcagta agactataga tagagttgat tggaaatatca 180
tgtgaccag tatgctactg ctaggcataa ttatcaaaaa ttcatttttc tcattaaata 240
ttgttaattg ctcgccacat aaagagaagc tagagctcac cagtcttggt ggtgtcctag 300
accttcctct aaagcagctc tgggaagctg gatcatcagw tctttagcct agacagagtg 360
tcgctggtaa ataaaggaga cacaggtaac ccagagtga cagtgatttg cgtggggagw 420
cacagtggat ctggggcctc tgatacttg yttcckaaaa cagccccag ttttcggctt 480
gcctatgaga tgatgttcat gtgcttcctt gaaaccaggt ggaaagaaag ggaagaatt 540
aattttctca ttctgttgct gttgaacgta atgtaatctt aatactgtag cttccttaga 600
agcccttccc tctttttcat gctgtaaagt caaatatttg atatccttaa cataaatttt 660
aaaaattaag gtcattaggr agcaaatgtc tatttccaaa gcaatgagct tgttgtgact 720
gtgattttat tcttctatag tatttttttc ctcattttaa ctgagaggag aaaataatac 780
tcttttgcaa tatccttagg ttctccctt ccccttggtg ccccttctag tgtcttaaga 840
ctttgtctta acaagtataa cattacattt tgttggtaaa acctttcgaa actgtattca 900
gtgattcttc caagtttatc tgctctgcac tatttcacta ataaacctg gctaccacgt 960
agcccttgat ctccaagtag ttacctatg caagacctgt gacactctga attcacttct 1020
ctttctttca gaaagtagtc ataaatggag cttaattata aaggtaaaac ttgtctccaa 1080
ccagtttcat tttggccatt tctttttcaa aatgtcagct gttttcctcc aagatttttc 1140
accaaaacaa tgatcataag tgctggaata tataatactt tgcaggcata aaataacca 1200
gacatactct catatttctt tgggtgattt tgggtggtaa aacttaccag cattaatgt 1260
aaaatataat gaggagttaa ttccttacct agaactattt cttcctttta agattcataa 1320
gtaacctttt atttttacag agctacgtat aacttcaca ttacagtcag ggacctgagg 1380
tgtaacttac taagtgaacc ccaaggttat tttatcttgc aaaagaaacc taaaccaaac 1440
taagggcctt acagtttatg gttagactga atcaaaagct ataacctcaa tttttccaaa 1500
aacagcttct gactgcaaaa gcaagtcata cagttgtag gtatgaaata gcaactgatca 1560
ggaaatgcat cttcgcagat ggtatttcct tcagaaaaga cttttctact tttaataata 1620
attaagccat aacagtttca tgctgtggaa agagggtgaa aagggttcatt ttaagagatt 1680
atataatatg aactttcaca tttactgtga aatgtctaac tttgccagt cttcagcaag 1740
tttttttggg ggggtgatgg gggaggggta gtattgggtt ttwggaggkt ttccaaatct 1800
gtggaacttt gggagagggg acagttgttg g 1831

<210> 156

<211> 1186

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1045)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1078)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1118)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1134)

<223> n equals a,t,g, or c

<400> 156

```
ggcacgagcg ggcctggccg aggttcgggc tccgttggcc gaggggggcg tacggaggtg 60
gcagctgtgg gaggaggcgg cgtggaaggc cgaggagctc aagcccgac caatcccccac 120
gttcggggcc gcgacctga ccctgcagcg taccgggaag cgaaaccggc cgcatgggccc 180
gctgagcccg aatcgggcac tgtgtggagc cccctggagc tgagatcagg atgttccgct 240
tcatgaggga cgtggagcct gaggatccca tgttcctgat ggatcccttt gctattcacc 300
gtcagcatat gagccgtatg ttgtcaggtg gctttggata tagcccttc ctcagcatca 360
cagatggcaa catgccagg accaggcctg ccagccggcg gatgcagcag gctggagctg 420
tctccccctt tgggatgctg ggaatgtcgg gtggtttcat ggacatgttt gggatgatga 480
atgacatgat tggaaacatg gaacacatga cagctggagg caattgccag accttctcat 540
cttccactgt catctcctac tccaatacgg gtgatggtgc cccaaggctc taccaagaga 600
catcagagat gcgctcgga ccaggcggga tccgggagac acggaggact gtccgggatt 660
cagacagtgg actggagcag atgtccattg ggcacacat ccgggacagg gctcacatcc 720
tccagcgctc ccgaaacat cgcacgggg accaggagga gcggcaggac tatatcaacc 780
tggatgagag tgaggccgca gcgtttgatg acgagtggcg gcgggagacc tcccgattcc 840
ggcagcagcg tccctggag ttccggcggc ttgagtcctc aggggctggg ggacgaaggc 900
ggaggggcct ccccgctgg ccacccaggg acctgaggay tcccctttcc cgacaktccc 960
gccgytatga ctggtgaggg ccccgggcct caagcctctt ctttgtaaca ggctgagagg 1020
ctgagaaatc aatccctga ataantttc cctctccgat tccccatccc ccaatttnat 1080
aattaaattt aaacagggaa agccgggccc ccaactgntt ccttgggggt cttnagggga 1140
gaacttttaa cgggaccttt ttccctaact ttttccttct ttaaaa 1186
```

<210> 157

<211> 1448

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (456)

<223> n equals a,t,g, or c

<400> 157

```
ggcctgaacc cggggctggt gggcctttct gtgtcctact ccttgaggt gacatttgct 60
ctgaactgga tgatacgaat gatgtcagat ttggaatcta acatcgtggc tgtggagagg 120
gtcaaggagt actccaagac agagacagag gcgccctggg tggggaagg cagccgccct 180
cccgaagggt gggcccccac tggggaggtg gaggctccga attattctgt gcgctaccg 240
ccgggcttag acctggtgct gagagacctg agtctgcatg tgcacgggtg cgaraagggt 300
gggatcgtgg gccgcactgg ggctggcaak tcttccatga ccytttgctt gttccgcac 360
ctggaggcgg caaagggtga aatccgcatt gatggcctca atgtggcaga catcgccctc 420
catgacctgc gctctcagct gacctcatc ccgcangacc ccatcctgtt ctgggggacc 480
ctgcgcatga acctggaccc cttcggcagc tactcagagg aggacatttg gtgggctttg 540
gagctgtccc acctgcacac gtttgtgagc tcccagccgg cagcctggga cttccagtgc 600
```

tcagagggcg gggagaatct cagcgtgggc cagagggcagc tcgtgtgcct ggcccagacc 660
ctgctccgca agascgcat cctgggttta gacgaggcca cagctgccat cgacctggag 720
actgacaacc tcattccaggc taccatccgc acccagtttg atacctgcac tgcctgacc 780
atcgacacac ggcttaacac tatcatggac tacaccaggg tcctggtcct ggacaaagga 840
gtagtagctg aatttgattc tccagccaac ctcatcgag cttagaggcat cttctacggg 900
atggccagag atgctggact tgcctaaaat atattcctga gatttcctcc tggcctttcc 960
tggttttcat caggaaggaa atgacaccaa atatgtccgc agaagtgact tgatagcaaa 1020
cactgggggc accttaagat tttgcacctg taaagtgcct tacagggtaa ctgtgctgaa 1080
tgcttttagat gaggaatga tccccaagtg gtgaatgaca cgcctaaggt cacagctagt 1140
ttgagccagt tagactagtc cccgggtctcc cgattcccaa ctgagtgtta tttgcacact 1200
gcactgtttt caaataacga ttttatgaaa tgacctctgt cctccctctg atttttcata 1260
ttttctaaag tttcgtttct gttttttaat aaaaagcttt ttctccttg aacagaagac 1320
agctgctggg tcaggccacc cctaggaact cagtcctgta ctctgggggtg ctgcctgaat 1380
ccattaaaaa tgggagtact gatgaaataa aactacatgg tcaacagtaa aaaaaawaa 1440
aaaaaaaa 1448

<210> 158

<211> 1004

<212> DNA

<213> Homo sapiens

<400> 158

ggcagagctc cagactactt taaatgtctc tsctctgact gaattgaagg aattgtaagt 60
ttcagttgct gaatatatca gtatatctga actccggatc tttttatttc ctatttttgg 120
cttctacat aaattaataa agtctgtcta tatttaagggt ttctgggagt agttttgtct 180
atatttaatc catagatgtg ctcttagtca tcagtcttat ttcagtaaaa ccatctttct 240
ggccaaatct ctaacacgtt ttttacttca gtagtgaggg caatgagaag ctttattaag 300
aaagatccta tttaaagggc taatttggtg ttgaaagcat gctgatataa agtccagtat 360
ttgaggatat aaagaggatg tgcactgagc tgtgctgatg ttcagcattt attatacttt 420
aatgggatag tgcttcttga ccattataga acaacaaact gtcaaagggt taatacagat 480
gaccagacc tgacattaaa tcctcttgat tagaaataat taaatgaatt atttattcat 540
ttctctctc tgccaaatct acttagcccc acaaaaatgc cccagctcag ttaacatttt 600
tttgtttttc actttaggta tctcttaggc aaatagagat ataaattgta ataggaaaca 660
gtctctaaat gtgaggctga atacagcttc agtctggctc tctaatagga caaaagccag 720
tccatgtggt gtctctmact gatttagctt tttcttttct ttgaacaatt cggcatgaat 780
tgaaaccagg ttttcctgtg gaaagttata gcttgattgg gagatagaag ttgaattgag 840
ttctttcttg caactcttag tgtttatttt tatatctcag taagacgagg ataccttcag 900
tttgaatctg cataatgttc actgccaac tccttctcat ttaatgctta tggccttcac 960
atttctgtat aataaagatc aattatcagc tmaaaaaaaa waaa 1004

<210> 159

<211> 1509

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1420)

<223> n equals a,t,g, or c

<400> 159

agagagagag actagttctc tcttactcct aggcctttcg gtttgcgga cgggcaggaa 60
agcgtgctg cggtctaaag agtggcgct ctgcggccg ctgacgatgg aagaactgga 120
gcaaggcctg ttgatgcagc catggcgctg gctacagctt gcagagaact cctctttggc 180
caagggtttt atcaccaagc agggctatgc cttgttggtt tcagatcttc aacagggtgtg 240
gcatgaacag gtggacacta gtgtggtcag ccagcgagcc aaggagctga acaagcggct 300
cactgctcct cctgcagctt tcctctgtca tttggataat ctcttcgcc cattgttgaa 360
ggacgctgct caccctagcg amgctacctt ctctgtgat tgtgtggcag atgactgat 420
tctacgggtg cgaagtgagc tctctggcct ccccttctat tggaatttcc actgcatgct 480
agctagtctt tccctggtct cccaacattt gattcgtctt ctgatgggca tgagtctggc 540
attacagtgc caagtgaggg agctagcaac gktacttcat atgaaagacc tagagatcca 600
agactaccag gagagtggg ctacgctgat tcgagatcga ttgaagacag aaccatttga 660
agaaaatttc ttcttgaas aatttatgat agagaaactg ccagaggcat gcagcattgg 720
tgatggaaaag ccctttgkca tgaatctgca ggatctgtat atggcagtca csacacaaga 780
ggtccaagtg ggacagaagc atcaaggcgc tggagatcct catacctcaa acagtgtctc 840
cctgcaagga atcgatagcc aatgtgtaaa ccagccagaa caactggtct cctcagcccc 900
aaccctctca gcacctgaga aagagtccac gggtaacttca ggccctctgc agagacctca 960
gctgtcaaa gtcagagga agaagccaag gggctctctt agttaatctg ttgtggcctc 1020
agctgctgag gatggacttg gagaayagct tccaagcttc accttgaaa aagcttacat 1080
ggcagcaata ttctaaaat agtgatacag tcagaggcct cctgtaaggg cgagagaact 1140
gaagttgatg ttgacaggcc cacagggaat tggccttccc tgttcaagtg gaagccagtc 1200
tctgagaatc ccgtgctctc ctctcttttg gtggagggtc tgtaggttca ggtttctacc 1260
atggacttta ggtatatagg gcaagtcagc aagaaagcac cacacactca ggaagccttg 1320
tctacctttc ctagcgtct ctagccagcc agccccagat actcctcaga gaccacttc 1380
tctcttttgc atggaataaa aagcactcac agtgccctgn cttttgggat tacttatggc 1440
tgtggaactt catgaaccca atttcacttt ccttcactgg gttccacccc cattgttgc 1500
tctgggatg 1509

<210> 160

<211> 2160

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (470)

<223> n equals a,t,g, or c

<400> 160

gctcgtgccg tggatgcggc tctgcaggat gaagatgaga tacagggtga tgcacatca 60
tttgaagca ctgcaataa gaaatcctcc tcctcttcct caggaaacccc tgcacatcag 120
ctttatccac agtctcgggg gctggttcca aagtaaagcc agcttctcct ctcccagggc 180
ggaaacagca ttgctcttct gagagaagag actagcaaaa agctgcagag aggattcggc 240
ccaaactcag aactgttccc ctgaggagaa gcggtggcct ctttgcagat caaccaactt 300
aatctggttg aacgtgctgt tcctaactctg gcaactcagcc cctctgggaa acatctttta 360
attagcatct cagaaatgca tgggtaagggt aaagtgcgat agttcaagtg gaaagcaaga 420
gaatgaccag tgacctgtct tccttcccc ttgcttctt ctcccccttn ccctgtgctc 480
cctttctctc ctctctcctt ttctagcctg ttcttwacat ggggctccct tcttgttgaa 540
caatagggca gaatcaggrg tcaccttagc aggaccacat ctttggagcc tcgggataaa 600
atgacagtga ggttgaaaag tgaaaaccct aggaactga ataggtgcct gttctttag 660
ggagaaatga gaaatcgcat ttggatccag gccccagggt ggccacatca gcagtcttgc 720
ttccatgcac ctcaagtaaga agtggatctg cctttgggac ctgctcagtg aggaaatctc 780

ttccaatttc tgcttctgaa tgattcaatg ttgggagcaa tagaaataac attccctttg 840
ccttctctga gtgttttagg aaatagcttc tttaaaacct caaaaccatg accatcctgt 900
caaagacctt agtctgtaag ctgggtgcat gtccatacac catgtcactt tactcttcat 960
ttgtcaccat cttttcccat gcacgcatac tctgaacatc cttgtgtggg cccatcctct 1020
gcatccagag catgctctgc agtgggcctg ttttgtggaa gaaaggaggc tgtctctgcc 1080
ttctctgatg ggactggagt tgagggaagg agctgtattg tggcacttct gaattccccc 1140
ttttgttcca tattggtata gagagcagaa gagtagctag gcagatgcag agatggagac 1200
atgagactca gtgcagtggg caggggaagac ataacagatg gaagcaaagg aatcctgcct 1260
gccttcagca gagaattcac cgaatcctag aactgtggct ccctccaggc agagcctaag 1320
atgctgggtg agaattagctg tgtgattgaa taggctcaaa ggagagttca gaattcccat 1380
ttacatatta ctagtgttgg ttgtaagttt tagttccttg tattattgag attcagagct 1440
tcattttatg ttggtcatta ggtgaatatt actcatttct cctcaagaga agctcataag 1500
tgtgtgtggg tgtgagagca cgatgggtgcc tgtgttctgt gaatgtgtcc atatgtgtct 1560
gtaagagaga cagagaccaa gaacttgccc aattttagaa atacactaat gtgcagtgtg 1620
tgccttttgt ctgtattgaa ggccattga atgactaatc caggctggaa gcattcccat 1680
gtgggtgtct gagtccatga gccaaagcctg aggggacagt gagtctccag gtctgccaca 1740
ctggtgcacc ttgctggcac ggtgcctcag gaaggtggcg actcargtgg gccttgagtt 1800
atattttaac tcagctgctc agtccaccag gcacatttct ggatcagaac ccatgggaaa 1860
caggaggtag taagtgaat gtcttagcat tctgcaaaat ggagatctgt tgtccagcgg 1920
cttatctcct ttttagtaac cttcttttct gaaccagggg cccttttcag ccttccctca 1980
tattttcttg agatcaaaact ttacttcttt cttatttact aagaatttgc ctgtttgaat 2040
aagaacaaaa cgctaagggtg ggtagcctaa gctgatttct tgctgggtac acgtgtctct 2100
cacaccacat ttcctcaaa gctaactctgaa ttctgttagc taaaaatatt catgtagcaa 2160

<210> 161

<211> 3609

<212> DNA

<213> Homo sapiens

<400> 161

ctggtaatcg aaaatgttaa catgcctgag sagattgtta ttcacgcact gcagtgtact 60
cactatgtaa tcctttggca acttgctaag ataactgaaa gcagctctac aaaggaggac 120
ttgctgcgtt taaagaaaca aatgagagta tttgtcaga tatgtcaaca ttacctgacc 180
aacgtgaata ctactgttaa ggaacaggcc ttcactattc tgtgtgatat tttgatgatc 240
ttcagccatc agattatgtc aggagggcgt gacatgttag agccattagt gtatacccct 300
gattcttcat tgcagtctga gttgctcagc tttatttttg atcatgtctt cattgaacag 360
gatgatgata ataatagtgc agatggtcag caagaggatg aagccagtaa aattgaagct 420
ctgcacaaga gaagaaattt acttgacgca ttttgttaag taattgtata tactgtggtg 480
gagatgaata cagctgcaga tatcttcaaa cagtatatga agtattataa tgactatgga 540
gatatcatca aagaaacaat gagtaaaaca aggcagatag acaaaattca gtgtgctaag 600
acccttattc tcagtctgca acagctttta atgaaatgat acaagaaaat ggctataatt 660
ttgatagatc atcctctaca tttagtggca taaaagaact tgctcgacgt tttgctttta 720
cttttggaact tgatcagttg aaaacaagag aagccattgc catgctacac aaagatggca 780
tagaatttgc ttttaaagag cctaataccgc aaggggagag ccattccacct ttaaatttgg 840
catttcttga tattctgagt gaattttctt ctaaactact tcgacaagac aaaagaacag 900
tstgtgttta cttggaaaag ttcattgacct ttcagatgtc actccgaaga gaggatgtgt 960
ggcttccact gatgtcttac cgaaattctt tgctagctgg tggatgatgac gacaccatgt 1020
cagtcattag tggatcagc agccgggggt caacagtacg gagtaaaaaa tcaaaacat 1080
ctacaggaaa acggaaagtg gttgagggca tgcagcttct actcactgaa gaaagtagta 1140
ttagtgacag tatgtgtgta agcagagaaac aaacactgca caccctgtt atgatgcaga 1200
caccacaact caccctcact attatgagag agcccaaaag attacggcct gaggatagct 1260

```
tcatgagtgt ttatccaatg cagactgaac atcatcaaac acctcttgat tataatcggc 1320
gtggcacaag cctaattggaa gatgatgaag agccaattgt ggaagatggt atgatgtcct 1380
cagaaggag gattgaggat cttaatgagg gaatggattt tgacaccatg gatatagatt 1440
tgccaccatc aaagaacaga cgagagagaa cagaactgaa gcctgatttc ttgatccag 1500
cttcaattat ggatgaatca gttcttgagg tgtcaatggt ttaataccag tacacaatta 1560
aatctgtggt gaagtcattt tctaagtggg agaggaaatt ttaaagtgtg gtagatacag 1620
tgaaattctg tacagatttt tctctaagga gaatatgaca tgcttatgct taccaagatc 1680
aagtgcattg aggggcagtt ttgtttgcct gaataaacgt aaaggacaag taaacaattt 1740
gatgataagc tacagttttt cttagaaagt aaatatttta tttatgctgt gttagtggc 1800
ttttgaatcg attatttcat gctttttttt aaaaaaaaaa aaaaacaaat aacaatctga 1860
agaggcattt ggtacagata tgaattctct tacatttatt tactgggtgt actaaataat 1920
gatgacctct gctggatttc tgtttacatc cagaaaacaa tgttaaggat gtattttatc 1980
ccctaccctg aagaaaagtgt aggatagaat tgtttttagc attctaaatt taaatgctta 2040
aaacgtcaat caacaaaact ttgtttttaa tattgtaatt gtggagaaaa gtaacttat 2100
aagcagaact tttaacaattt ttcatctaa aagtatttta agatattttt aaaatccaag 2160
agcttctcta tacttttcag aaatatccag atgcagtga ctgccagaag gtaaccagtc 2220
tcaaacatgc ttatcccatc atcaaccctg aaagtttgct tgccttttaa gataaaaatg 2280
taatgttggt atattccttc cagtaatgcc actgtatttt gtctccaaat aaaagaagct 2340
tattgtagta tgtttgcaga aaaattctaa acaaaaatta tacagcttat tagagtgtgg 2400
gaatagggat ctaaatttta aataaaatta tatatatata taaattggtg ctgattttat 2460
aattgctgag tttgtttagt tttttcttac ttttaaatc caacttaaaa ttatgagggt 2520
tcagaaatat attgaaagt taacaatggt taaaaataga aaagcatgag tgttcagct 2580
ttaaatgat ttttaaatgt gtattttata ttgttttata tatctgtctt tgcaagcagt 2640
cttcaggtta aagatacttc taacaggtta cagtacattt cctctgtatg taaattagat 2700
gggataatag aattcataac ccataatatt ctttgaaagc taagctttta acttcatttt 2760
atgtccttcc acaataaat tagttttaa cagaaagtgg ctacttgcca ttttgacatc 2820
aactcatttt gcgaggctta ggcagctaga catcgtttaa acaaaaatat taacttatat 2880
tacatgtgta tctatctatt gtcagtcgtc tctcagttct tgaggatat tattttaatc 2940
attccatgcc ttaatatgct tgcaatacaa gaatatcttc agatgggtga ataccaaaag 3000
gctttcagtt tttagtcaga aatcaagcat tgggctgtgg tagccaaaaa ccatagggtta 3060
gctaaaaaga tcatgatata attattttat taagtcatgg ttaataacaa atgaatccag 3120
acttgtctaa cagattttcc atcaacaaat attgttatgt gcaaaagtat tgcctatggt 3180
gttttacaca ccactgcatt aactagaact gctgagagga ctgtatatat gattttaaac 3240
ctaagttgat tttttttctc actcttgaaa ggagtacttc tttgtgaaag cagttcttac 3300
agctttggtt tcaaccagct aaaaatgttt tatatattac tctaacctgt tgcctccac 3360
attctattgt ctaattgta ctgttttctg atttgtatgt atgtcttgag acagtaactt 3420
tttgaataaa aataaaccta cagtatgttg tatgttttct cttgtactca aagggggagg 3480
gtggctataa atggtttgca aatttatatc tattatcaca tcttttaatg tgtttgggga 3540
ataatttata gagaatacca tcagtttata tttttaataa atcatatgta tttacaatga 3600
aaaaaaaaa 3609
```

<210> 162

<211> 1603

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1600)

<223> n equals a,t,g, or c

<400> 162

```
gttcaggaga gatcacgctg ctgccctccc gggacaaccc ccgtcgctg cgcggtgctgc 60
tgcgctacaa agtgggagac caggaggaga agaccaaaga ctttgccatg gaggactgag 120
cgttgccttt tctcccagct acctcccaaa gcagcctgac ctgcgtggga gaggcgtagm 180
gagtcggagg ggaaaggagg atcccacgtg caagtagggg gaatatctcc cccttttccc 240
tcatagcctc tagggaggga gagtgaactt attctccatt tkaagagatt cttctggtga 300
tgtttactta aaaagtgatc cccctcaaca acggatacag cgtgcttatt attgggcatt 360
tagcctcaaa agcatgtagt accaagcact tgtatttccg tataattttgt ttcgcggggg 420
agtggggggg aagaacacgg atgaaaatgt cagtttttga aggggtccatg cacatccctg 480
acacctcaca ctttatctaa gtctgaagct ggggagaaa ggggttcatt agacttcata 540
catttccagt acgactttag tatctctcca gagccatatt ttctcagtcc gaattaattc 600
ccctcccta ggtgcctgta ggctatggtta cttcttctc attgttttct aggtaaactt 660
cactactggt aattaagggg aaggatatga ggaagcagtt taaatagccc tgttctcatt 720
actctgacca catacatcat agggtgctaa agttgatgaa cacattaatc cgtaagtaa 780
aatggacttt gtaattgtac agcataccta agaaactcag aagggtgcatt taagagagag 840
acctgaaaga aatagtatgg atttttaaaa attcttgtct ctactattat aacccaaaaa 900
tatttcttgt atgtcccata aaaatatttg kgtaattctt atgaaacagg ctggtagagg 960
aggtttctga gcctagccca agggcttatt catcaccatg ggtaaattat taaactcac 1020
ttaattaagg aaaatatttt ccagctaga aaagtatact cattctcatt taaactctct 1080
catttgaggg gatcatgtga gtggcctac ttacaagtag tgaaagtcc ctttctcagt 1140
ttgttttgtt ttgttttgtt ttctctttc actcagccaa atgtgaaagt tgtgaattta 1200
ggaaaatcac ttgtaatgaa gtgtgaatct tgttatcaaa tttatttctc tgatgtttcc 1260
ttccttatcc ttgtagccaa taaaacattg acattctcac gttttataga tgaggtaaaa 1320
agtcttgtgt gctgtgagtt ataatgcttt tgccctttta atattattag ttcttaagt 1380
ttacagcccc ttcagaatat aacttcagga caattcaaac tatgcttaat gtatgatttt 1440
cgagcttctg tatgctaaga aaataggtgt gaaaaactgg tgttctgaaa tagcctaaca 1500
tttattgtaa ttctgaattt tctgcccttt tattcattgc atattaaagt attagagtat 1560
aaaaactaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ccc 1603
```

<210> 163

<211> 853

<212> DNA

<213> Homo sapiens

<400> 163

```
gaataaaccc aacctacaga gcatcatagc ttagcctagc ctgctttaa tgtgctcaga 60
aaacttccat tagcctgcaa ttaggcaaaa tcatcaaaca taaaaccatc aaacataaaa 120
tatttataaa gtgttgaata tctcatatag ttatttgaat acctgcatcc aaaagatgct 180
ggcaacacag cacactttag agcattgggt gtttactctc ttgatgggtat ggctgcccag 240
catcaagagt tatcatactg caaatcgata gccaggaaga agagcaaaat tcaaagttca 300
aagtagagtt ttactgaat gcttgctttt gcaccgtcgt aaagttgaaa agaattttaa 360
ttgaaccatc ataagctgca gactgtgcat ttatatttga aaagttaata tttttaattt 420
ttaatgcaga gaagtaccca aagcataaga acacaacaca ttttcacaaa gcaaacacag 480
ccatggaacc agcaccata tcaactaaca aaatactagt ttgggctttt ttgtacttta 540
tacaaatgga ctcatataat gtctcatctt ttgggtctgcc tgctttcatt caatattagg 600
tttggtgggt catctctgct gtgtgtagtt ctttctctgt ctttatacag tgttccaaag 660
tatagtatat tacagtttac ccattctact cttgatagta aatgttttca catttgggct 720
attacaaata gtgctgcagt gaacattcac atacacatct tttggtgaac atgtgttaca 780
tttccaagta caattgctgg gtgatgagta tgcatactct taaaacatgg ttgtaccaat 840
ttacacctct acg 853
```

<210> 164
<211> 1917
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1433)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1856)
<223> n equals a,t,g, or c

<400> 164
gcccacgcgt ccgcggacgc gtgggaaact tttgtaaaaa aagtttattc ttaaattgtac 60
aacagctcca agacaacatt aattccagct atacgagaag tttcttcaca ggtataagat 120
gatctctgaa ttcacctggc ccaaccatga ccttccttca gacaaagagg ctgtcaagaa 180
actaattgaa cgggtgtggtt ttcaggatga tgtagcttat gggaaagacca aaattttcat 240
tcgaacaccc cgaacattgt ttaccttggg agaactccgt gcccagatgc tcataaggat 300
tgtcctcttt ctacaaaasg tgtggcgggg caccckggcc cgcattgcgt acaaaaagaac 360
caaggcagct ctgacaataa tcagggtacta ccggcgctac aaagtgaagt cgtacatyca 420
cgargtggtc agacgcttcc atggcgctca gaccatgcga gactacggga agcacgtgaa 480
gtggccaagc cctcctaaag ttcttcgccc ttttgaggag gccctgcaga cgattttcaa 540
tagatggaga gcatcccagc tcatcaagag cattccggcc tcagacctgc cccagggtcag 600
ggcaaagggt gcagccgtgg aaatgttgaa ggggtcaaagg gctgacctcg ggctccagag 660
ggcctgggag ggcaactatc ttgcttcaaa gccagataca cctcagacct caggcacttt 720
tgtcctctgt gctaataaat tgaaacggaa ggacaaatac atgaatgtcc tcttttctctg 780
tcacgtccgt aagtaaatcg atttagtaag gtggaagaca gagcaatttt tgtcactgac 840
cgtcacctgt ataaaatgga tcccactaaa cagtacaagg tgatgaagac tatccctcta 900
tacaatttga ctgggtctgag tgtctccaat ggaaaggacc aacttgtagt gttccatacg 960
aaagacaaca aagacctcat tgtctgcctc ttcagcaaac agccaacca tgagagtcga 1020
attggagaac ttgttgaggt gctgggtgaat catttcaaga gtgagaagcg ccaccttcaa 1080
gtgaacgtca ccaaccagct acagtgcagc ctgcacggga agaagtgcac cgtctccgtg 1140
gagacgcggc tcaaccagcc ccagcccagc ttcaccaaga atcgctcggg cttcatcctc 1200
agcgtgcccc ggaactgacg ccccgcgagg gcctggcccc gagccccggc aactccagag 1260
tcctgggtcc cagtcagct gctgcctccc aaccatgcc cgtagaac ctgctgcgag 1320
ggccccctcc agaggagccc cggccctgta agatttcctt cctgggtttc tgcctttggt 1380
atcatcttcc tctgtcctta ctgtccacgg tccctgttca ataagccaaa gancctggtg 1440
ccccgcccag accctgggc tgacgtccag accaatctca ccccarargc aactggatgg 1500
tgcttttagt tgggtcggat gcccctggc aggtcaagtc agagcacctg gacagggtgc 1560
ctccctgtc ctgacctgc agagggaagg ggtggggatg caggaccccg ctctgcccga 1620
gcccatagc cacctctytt gcccaagggt agccagccct ggggaccag ctcaggaggag 1680
ctctgtcag agttcggcg acccaacctc cagccgcccag ccaaagccc 1740
tggtgtgagc aggtcaccca cagggccagc cctcagcca ggagccagtg gaggagcga 1800
acagcccac cagcagagtg agtccatcct tcccagggtc tcccctggga gactccttt 1860
gccaccaagg cccccaccag ggctctgacc accgctctgg agaggacagt gtggcat 1917

<210> 165

<211> 2420

<212> DNA

<213> Homo sapiens

<400> 165

```
gcacctccgt gcgcccgggt gcagcgtgga cgccggatga gttgctttta ggcttgctgg 60
cccgcggggc tgtccaggca cgcgaggccc ctcagcaaca aaatgcttca acaagttcca 120
gaaaacataa attttcctgc tgaagaagag aaaatccttg agttttggac tgaatttaat 180
tgttttcagg aatgcttaaa gcaatcaaaa cataaaccaa aatttacctt ctatgatggg 240
cctccttttg caactggact gcctcactat ggacatatac ttgcgggtac aattaaagat 300
atagttacaa gatatgctca ccagagtggg tttcatgttg acagaagatt tggatgggat 360
tgccatggct tacctgtgga atatgaaatt gataagacac tgggaatcag aggaccagag 420
gatgtggcca aaatggggat tacagagtat aacaatcagt gccgagcaat tgtgatgaga 480
tattctgctg agtggaaagtc tactgttagc agacttggcc gatggattga ctttgacaat 540
gactataaaa ctctgtatcc acaattcatg gaatcagtct ggtgggtctt caaacaactc 600
tatgataaag gccttgctta tagaggtgtg aaagtcatgc cttctcttac ggcatgtaac 660
actccacttt ccaacttcga gtcacaccag aattataagg atgttcaaga tccttcagta 720
tttgtaactt tccctttgga agaagatgaa actgtatctt tagttgcttg gacaaccact 780
ccctggactc tacctagtaa ccttgctgtg tgtgttaatc cagaaatgca atatgtgaaa 840
attaagatg ttgccagagg acgattactc attttaatgg aagccagatt gtcagccctc 900
tataaattgg agagtgacta tgagatcctt gaaagatttc ctggtgccta tcttaaagge 960
aagaagtaca ggcccctgtt tgactatttc ctgaagtgtg aagagaatgg cgctttcact 1020
gtgcttggtg acaactatgt gaaggaagaa gaaggcacag gggttgtcca ccaagctcct 1080
tacttcgggt ctgaggacta tcgggtctgt atggacttta acattattcg gaaagactca 1140
ctccctgttt gccctgtgga tgcttcaggc tgcttcacaa cggaggtgac agatttcgca 1200
ggacagtatg tgaaggatgc tgacaaaagt atcatcagga ctttgaagga acaaggccga 1260
cttctggttg ccaccacett cactcacagc tacccttttt gctggagatc agacactcct 1320
ctaatttaca aagcagtgcc cagctgggtt gtgcagtggt agaactggt ggaccagctc 1380
ctaaggaaca atgacctgtg ctactgggtc ccagagtggg tacgagaaaa acgatttgga 1440
aattggctga aagatgcacg tgactggaca atttccagaa acagatactg gggcaccctc 1500
atcccactgt gggtcagcga tgactttgag gaggtggtat gcattgggtc agtggcgga 1560
cttgaagaac tgtcaggagc aaagatctca gatctccaca gagagagtgt tgaccacctg 1620
accattcctt cagctgtgtg gaagggatcc ttgcaccgca tctctgaagt gtttgactgt 1680
tggtttgaga gtggcagcat gccctatgct cagggttcatt acccgtttga aaacaagagg 1740
gagtttgagg atgcttttcc tgcagatttc attgcccagg gcacgcacca aaccagagga 1800
tggttttata ccctgctggt gctggccacg gccctctttg gacaaccgcc tttcaagaac 1860
gtaattgtga atgggcttgt cctggcaagt gatggccaaa aaatgagcaa acggaaaaag 1920
aattatccag atccagtttc catcatccag aagtatgggt ctgatgccct cagattatat 1980
ctgattaact cccctgtggt gagagcagaa aacctccgct ttaaagaaga ggtgtgctgg 2040
gacgtcctta aggatgtact gctcccattg tacaatgcct atcgcttctt aatccagaac 2100
gttctgaggc tccagaagga ggaagaaata gaatttctct acaatgagaa cacggttaga 2160
gaaagcccca acattacaga ccggtggatc ctgtccttca tgcagtctct cattggcttc 2220
tttgagactg aaatggcagg tgagtctctc ttggtctgtc ctcccaggaa taaggactat 2280
tctctttgta ackgcccttt tgatatttaa tagtgaaat attgagagat acagamaatt 2340
aaataattgt ttccttttac ctctcatata gttttgttaa ggcttgtaa ttgtcgtgat 2400
atctttgttt aagacctggg 2420
```

<210> 166

<211> 2061

<212> DNA

<213> Homo sapiens

<400> 166
gccggcaccg cagcagcccc aggagggcgc gggcrgcrgg cccggtgcgt gcagcctgca 60
cctcagcgag cgcgcgcgact ggcagtactc gcagcgcgag ctggacgccg tcgaggtctt 120
cttctcgcgc acggcccgagg acaaccggct cggtgcgatg ttctgctcgt gcgcgccctc 180
cagccgctac acgctgctct tctcgcacgg caacgccgtg gacctgggcc agatgtgcag 240
cttctacatt ggctcggtc cccgcacaa ctgcaacatc ttctcctacg actactcggg 300
atacggcgtc agtcggggcaa gccctccgag aagaacctct acgccgacat cgacgccgcg 360
tggmagggcg tgcgcacccg gtatggcggt agtcccgaga acattatcct ctatggtcag 420
agcattggga ctgtccccac ggtagacttg gcctcgaggt atgaatgcgc agggtaattc 480
tccattcccc tctgatgtct ggtttgctg tggttttcc ggataccagg aaaacatact 540
gctttgatgc tttccccagc attgacaaga tatctaaagt cacctctcct gtgttggtca 600
ttcatgggtac agaggatgag gtcacgatt tctcccatgg cctagcgatg tacgagcgct 660
gtccccgagc cgtggagccc ctttggkttg aaggggctgg gcataatgac atagagcttt 720
atgcacaata cctagaaaaga ctaaaacagt tcatactca cgaacttcct aattcctgaa 780
gacaacaact tgatcttacc tcatttactg tgaacagaag agtcctctgt tttgcacatg 840
ctttaactgg gtagctgtaa aggcttgata accatgaaga agtgcccaac ctttaggggt 900
ttctaataca agagctgatg aaatctcagt cttttgtatc tagagggtgt tctgctaatt 960
cacacaacac gttaaactga acagtcgtga tccccagctt cattaccttg caggaaatggg 1020
aatgagagct gaatgtaggg acaattttct agtgctgtat aaagtagcct cgcactctgtt 1080
tctcaacctt atccatcatt tctgacattc atgcaggact tgccctgttg ccaccaatgt 1140
tctcggtatt tcacatgcag ctctctttct gccactggat acatgggttc aatccatttg 1200
tgaagctgtg atagtgtaac tggaaagcta gtgtggtgaa aattccttta ttattttttg 1260
ttaacatgct gatctttccc ggacaaatga actgaagggt aatttactgg aactctcgtg 1320
tacagcttca tcaactgtaa ccatataaat ataactggaa tattctttaa caaaagaaa 1380
ctaggggttt ttttaagtgt aaatttatta ctagccaaca gagttttact attttgattg 1440
tctgggtggg ttaacaaaga gcctagctga ctttccttct gtaaagtcct ccttgtaggc 1500
ttttttaaag tactgtacat atttgcaatc acattgtgca tagattctta atggtagata 1560
tgattttctt tgtcaggcta caacaatgaa ctgcagattc cttgtttgta atgtaaatga 1620
ttgaatacat tttgttaata tgtttttatt cctatgtttt gctattaaaa attttataac 1680
atttccaaga caaaaattcc aagtttatgc tttgaagaat ttatgtaatt aaaatttcac 1740
taaactaatc tttttagttt aggaattatt tgggttttga cactggaagt tgcgccaaat 1800
aagcatcaga aataggagat gcttaacatt gctatactac ttgtgttggt taggggtttg 1860
gatttggggg ttctttgggt ttaatttttt ttccacatt taaaagcctt aaatgtactg 1920
taagcctcag atcgttgtac aactggactg cggttgattg ccagtttggt tactgttgct 1980
tggatgcggc acagtgggtg gtaatggaat aaaggatgca tggatcagaa aaaaaaaaaa 2040
aaaaaaaaa aaaaaaaaaa a 2061

<210> 167

<211> 2567

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (16)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (55)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (74)

<223> n equals a,t,g, or c

<400> 167

```
ggagaaggcg gcggtnggcg gcgtggacgg cccgcagcgg gacggcccg gcgangggcg 60
acagccgggc gcancggcg yacaggggccc cccgccggcg cctcagctca cagagacgct 120
gggcttctac gagagcgacc ggcggcggga gaggcgscgg gccgcacaga gctttcattg 180
ttgaggttcc tcagtgtga actacaaga gggtaacttcc ttgaacataa tgaggccaag 240
tatacagaaa gaagagaaa agtatacact tgtttgcgaa taccaagaga attgaaaaag 300
ctgatggttt ttggaatcct tctgtgcctg gatgcgtttt tgtatgtgtt caccctgctt 360
cctttaagag ttttcctggc actattcagg ctctcactt tgccttgcta tggcttaagg 420
gacagacgtt tgcttcagcc tgcccagggtg tgtgacattt tgaagggtgt cattttggta 480
atctgtctatt ttatgatgca ctatgttgac tactccatga tgtaccacct gataaggggg 540
cagtcctgca tcaagctcta catcatctac aacatgctgg aggtagctga tcgtctgttt 600
tcattctttg gacaagacat attagatgct ctctattgga cagcaacaga gcctaaagaa 660
agaaaaagag cccacattgg ggtgattcct cactttttca tggctgttct ctatgtcttt 720
ttgatgcaa ttcttataat gttcaagca acaactctca atgtagcttt taactcacac 780
aacaagtctt tgtctactat catgatgtct aataatyttg ttgaaattaa aggaagtgtt 840
ttcaagaagt ttgaaaagaa caatctcttt caaatgtcaa atagcgatat taaggaacga 900
ttcacaayt atgtgctytt actgatagtg tgtctaagaa acatggaaca gttttcttgg 960
aatccasatc atctctgggt gttgtttcca gatgtctgta tggtaattgc atcagaaatt 1020
gccgtggata ttgtaaaaca tgcctttatt actaaattca atgacattac tgcagatgtc 1080
tacagtgaat atagagccag tcttgctttt gacctgtta gcagccgaca gaaaaatgca 1140
tactactgatt acagtgactc tgtagcacgg aggatgggct ttattcctct cccactagct 1200
gttttactca tcagagttgt aacaagctca attaaagtgc aaggaatcct gtcttatgcc 1260
tgtgtcatalc tcttctatct tgggttgata tccctgaaag tacttaatag catcgtgctg 1320
ttggggaaat cgtgccagta tgtgaaggaa gccaaaaatg aagagaagct gtcgaatcct 1380
cccgcacact gcactccagg caagccgtcc agtaaatcac agaacaaatg taaacctctc 1440
caaggccctt ccacagaaga aaacctgtct gcctccatca ccaaacaacc tattcatcaa 1500
aaggaaaaata tcataccatt acttgtgaca agcaattctg atcagttttt gacaactcca 1560
gatggtgacg agaaggacat aacgcaggac aattctgaat taaaacacag atcctcaaag 1620
aaagatttgt tagagataga caggttcaca atttgtggaa accgaattga ctgaatctgt 1680
ggcttcatgc gctgaagaag ctgggtcctg gggcaacaag tgctgtgttg ccaggacaaa 1740
tagatgctaa acatggcact taaatattta tttaaaaact taaattatta ttggcaagca 1800
aatcttagta tctttcttcc agtaatatgg cctggctgag ggtcagacca caggacagga 1860
gcgacctccg gccttgactg tctgggaagc ttgatggatt ataaaacttc ctctgcctg 1920
gccaaagcagc agcatcattt ccaggacca acaaaggcaa catcaaaatc tgttttgctt 1980
tgtcagctcg ggcttcaga atgttgaatt tgctcaagg cctcttcagt ataaggaaat 2040
acctggaaaa ctgtgaaact tttaccacga cgtaatcttt ccagtctcat actattttca 2100
caaacagttt tcaaacgtta cttcatctgc caaagcatta aaaaaattaa acataagtca 2160
agataaatgt tcttaccacc agaataaccc ttaaagatgt atcttaatta atcagaataa 2220
aaggctacct taaataagac atgatgaata gtagcatttt gtaggggtta aaaaaaatg 2280
ctaaactagt ttatttatta aaaaaataat taatgaaggc tatgcattac tgggaaaaat 2340
ccttcataat ttttcagttt actctttaaa gcaaaatgtg agctgtatgt catttaattg 2400
gtggataaaa caaatcttct tacaaaaggg cactttttac accaagggaag cagagcagtg 2460
ttaactttaa ggtatactta actgattttg ctttaaaact aattacytcc mtaaaattatt 2520
acmgctaaag ggttycgrtg ttgarccgg gcyygaaaat tgtccat 2567
```

<210> 168
<211> 2324
<212> DNA
<213> Homo sapiens

<400> 168
cccacgcgtc cggtccttcc ggtatcgcgt tgctcagggg cttttcaacc ctctgtcagt 60
cggcgatcgc aagcaggctc ttctaagttc ccgacgcctc ttggcccgga aaactccggg 120
aaccacacac tgctttcctc tgcccagccc gagactcggg tcagttactg gacgaaactg 180
ctctcccagc tccttgccgc gctccccgga ttgcttcaga aggtgctaata ttggagccaa 240
cttttcgggtg gaatgtttcc gaccagatgg ctagattttg ctggagtcta cagcgccctg 300
agagccctga agggacggga gaaaccagcc gcccccacag cgcagaaatc tttagttcg 360
ctgcagctcg actcctcaga cccctcggtc accagtcccc ttgattggct agaggagggg 420
atccactggc aatactcgcc ccagaccta aaattggagc ttaaggccaa gggaagtgc 480
ttggaccctg cagcacaggc ttttctctta gagcagcagc tgtggggagt ggagctgttg 540
cccagtagcc ttcaatcccg tctgtactct aaccgggaac ttggctcttc gccctctggg 600
cttctaaaca ttcaacgcat agacaatttc agtgtggtat cctatttgct gaaccttcc 660
tacctggact gctttcctag gctagaagtc agctatcaga acagtgatgg aaatagcgag 720
gtagtcggct tccagacact aaccccagag agcagctgcc tgagagagga ccattgtcat 780
ccccagccgy tgartgcaga actcattccg gscctcgtggc agggatgtcc acctctttct 840
acggaaggcc taccagaaat tcaccatctt cgcataaaac ggctggaatt ccttcaacag 900
gctagcaagg ggcaagatwt acccaccctt gaccaggata atggctacca cagcctggag 960
gaggaacaca gccttctccg gatggatcca aaactactga gagataacct aacacagttt 1020
gttcctgctg ctggagacat tcctggaaac acccaggaat ccactgaaga aaaaatagaa 1080
ttattaacta cagaggttcc acttgctttg gaagaagaga gcccttctga gggctgtcca 1140
tctagtgaga tacctatgga aaaggagcct ggagagggcc gaataagtgt agttgattac 1200
tcatacctag aaggtgacct tcccatttct gccagaccag cttgtagtaa caaactgata 1260
gattatattt tgggaggtgc atccagtgc ctggaaacaa gttctgatcc agaaggtgag 1320
gattgggatg aggaagctga ggatgatggt tttgatagt atagctcact gtcagactca 1380
gacctgaac aagaccctga agggcttcac ctttggaaact ctttctgcag ttagatcct 1440
tataatcccc agaactttac agcaacaatt cagactgctg ccagaattgt tcctgaagag 1500
ccttctgatt cagagaagga tttgtctggc aagtctgatc tagagaattc ctcccagctc 1560
ggaagccttc ctgagacccc tgagcatagt tctggggagg aagatgactg ggaatctagt 1620
gcagatgaag cagagagtct caaactgtgg aactcattct gtaattctga tgaccctac 1680
aaccctttaa attttaaggc tccttttcaa acatcagggg aaaatgagaa aggctgtcgt 1740
gactcaaaga ccccatctga gtccattgtg gccatttctg agtgtcacac cttactttct 1800
tgtaagggtg agctgttggg gagccaagaa agtgaatgtc cagactcggg acagcgtgac 1860
gttctttctg gaggaagaca cacacatgtc aaaagaaaaa aggtaacctt ccttgaagaa 1920
gttactgagt attatataag tggatgatg gatcgcaaag gacctggga agaatttgca 1980
agggatggat gcaggttcca gaaacgaatt caagaaacag aagatgctat tggatattgc 2040
ttgacatttg aacacagaga aagaatgttt aatagactcc agggaacatg cttcaaagga 2100
cttaattgtc tcaagcaatg ttgagttggc agcctgtagt cctagctagc atacactacc 2160
tcttacctga gaggtgtctt ttaaaaacaa atcttggcag ctgtcctttg acattttttt 2220
tttagrggaa tgtaactggm ctrgtaattt ttttttttgc aacatatccc cactccagaa 2280
acattccagg tttgaagcca gccctggata atgaaggatg aact 2324

<210> 169
<211> 1784
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (200)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (215)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1759)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1778)
<223> n equals a,t,g, or c

<400> 169
gagaattacc ttcagctgta tcaattacag tacacaggaa ttgttacgat ttcctaaact 60
tcatgatgcc atagttgaag tggtagacttg tcttcttcgt aaaagggttg ctgttacaaa 120
tgaaatgggc cataacttag tggcaattga actggcttat atcaacacaa aacatccaga 180
ctttgctgat gcttgtgggn taatgaacaa taatntagag gaacaaagga gaaacaggct 240
agccagagaa ttaccttcag ctgtatcacg agacaagggt gcatctggag gtggtggggt 300
tgagagatggt gttcaagaac caaccacagg caactggaga ggaatgctga aaacttcaa 360
agctgaagag ttattagcag aagaaaaatc aaaaccatt ccaattatgc cagccagtcc 420
acaaaaagggt catgccgtga acctgctaga tgtgccagtt cctgttgac gaaaactatc 480
tgctcgggaa cagcgagatt gtgaggttat tgaacgactc attaaatcat attttctcat 540
tgtcagaaag aatattcaag acagtgtgcc aaaggcagta atgcattttt tggttaatca 600
tgtgaaagac actcttcaga gtgagctagt aggccagctg tataaatcat ccttattgga 660
tgatctcttg acagaatctg aggacatggc acagcgcagg aaagaagcag ctgatatgct 720
aaaggcatta caaggagcca gtcaaattat tgctgaaatc cgggagactc atctttggtg 780
aagagaacta tgtaatactg agactttgtt gactcaaaac ttgctagtta ctgcctacct 840
gagtagaatc ttatttatga actcctgtgt attgcaatgg tatgaatctg ctcatgtgga 900
gactggctat aaactgaaaa gtgtattcca aattgcagaa cacatcacac atttaatcca 960
aataataaat ggctgtttct aaagttccc agtatatata aaatacatca agtctgtctt 1020
gtgacagttt catctgaact taacttaaaa acaactgtta atgttctagt tgtgcaaagc 1080
agtttgccctg tggataagat gacctgtgta ataacttttg ttagtagtct taaagctgct 1140
gccatagtcc tccaagaaga aagcaccaag acaacatttc atatgactat aatgcatgta 1200
ctatataagc tgatctggct ttgaaagatg tgagttggca agttcctcac atagagtcat 1260
tgtattccac ctgtccttca atttagtttt ttctgagctt ctttgacgcc tttgatgtgt 1320
ttttaagaaa gctgaatgca caagaggatc tgtgacactg acatggctgt ggtgtgcata 1380
ctgtgtagtt acatagccct tccaattctg ggtccatttg cactagcaaa ttaaaatatg 1440
ctttgattca tacttaaac tgaaagcagg aatgcctaca ttaattccta cattaataaac 1500
agccatctac ccttgattat ctgwaagac ttggtaatga tggtcagttc cttttagatt 1560
tcagaaaaatc aaatgatgac cttaaattcc cttaatttgc aaatacagta gtaattaagg 1620
tacatctcta aagtggagca cttacaccag gctctaagat tcactttgag gtggaactta 1680
aaaccagtgt actgtatgta tgcattggta atagctactt tkgcttcata gcttccatac 1740
caacmaata tatttrttna gratagtatg aaagtcangg gagg 1784

<210> 170
<211> 1296
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1261)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1276)
<223> n equals a,t,g, or c

<400> 170
ggcgggtgcag aggcggcgagg aagatggagt tggggagttg cctggagggc gggagggagg 60
cggcggaggga agagggcgag cctgaggtga aaaagcggcg acttctgtgt gtggagtgtg 120
cctcggtcgc aagctgcgat gccgcagtgg ctgagtgctt cctggccgag aacgactggg 180
agatggaaaag ggctctgaac tcctacttcg agcctccggt ggaggagagc gccttggaac 240
gccgacctga aaccatctct gagcccaaga cctatgttga cctaaccaat gaagaaacaa 300
ctgattccac cacttctaaa atcagcccat ctgaagatac tcagcaagaa aatggcagca 360
tgttctctct cattacctgg aatattgatg gattagatct aaacaatctg tcagagaggg 420
ctcggaggggt gtgttcctac ttagctttgt acagcccaga tgtgatattt ctacaggaa 480
ttattcccc atattatagc tacctaaaga agagatcaag taattatgag attattacag 540
gtcatgaaga aggakatttc acagctataa tgggtgaagaa atcaagagt aaattaaaaa 600
gccaagagat tattcctttt ccaagtacca aaatgatgag aaacctttta tgtgtgcatg 660
tgaaygtgtc aggaaatgag ctttgcccta tgacatccca tttggagagc accagagggc 720
atgtgcgga acgaatgaat cagttaaaaa tggtttttaa gaaaatgcaa gaggtccag 780
agtcagctac agttatattt gcaggagata caaatctaag ggatcgagag gttaccagat 840
gtggtgggtt acccaacaac attgtggatg tctgggagtt tttgggyaaa cctaaacatt 900
gccagtatac atgggatata caaatgaact ctaatcttgg aataactgct gcttgtaaac 960
ttcgttttga tcgaatattt ttcagagcag cagcagaaga gggacacatt attccccgaa 1020
gtttggacct tcttgatta gaaaaactgg actgtggtag atttcctagt gatcactggg 1080
gtcttctgtg caacttagat ataattattt aaaatgcttt tcaagtgtgg gttttgccct 1140
gattgttgca aatacaattt ccaccttctg gaaaggtagg tttgctgtgg aggaataat 1200
gwactagatc attgtcacag aaaaaccaac tatgatttat ggkkggtgtt tcmccaryat 1260
nagacggaaa gaccncccc ttcttcccc cccct 1296

<210> 171
<211> 1897
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (952)
<223> n equals a,t,g, or c

<400> 171

gagaaaactg acttcgagga attctttcta aggtgtcaga tgcctgatac attcaattca 60
tggtttctta taacctact ccacgtctgg atgtgtctag tccgaatgaa gcaggaaggc 120
cggagtggga agtacatgtg tcgtatcata gttcatttta tgtgggagga tgttcagcag 180
cgcgagagt catgggggtt aatccctata tcctgaagaa gaacatgac ctcagacaa 240
atcatttcta tgcagcgatc ttgggatatg atgaggggat ctttcagat gatcatgggc 300
tggccgctgc cctctggaga accttcttca accggaaatg tgaagaccct cgacatcttg 360
aattgctggt agagtatgtg aggaaacaga tacagtacct ggactccatg aacggggagg 420
atctgcttct gacaggggag gtgagctggc gccctctagt ggagaagaat cctcagagca 480
tcctgaagcc ccattctccg acctacaacg acgagggact ttgatgggct gggcctccgc 540
acggcccgcg agctggcttc gaggaacctc caggagagaa gtgcctgttg gtccaggacc 600
ctgcagaaag tggcctgaac tgacctctga acagcatctg tcaataacct ggccccattt 660
gtgttgagtt tcctcttagt gtgcccagga gtctgatctg ctgggtgaca gggtgggag 720
aaccctagc tctcccgagg tgtcctctcc cttaggggaa gcccgagtg agagtcccc 780
agcacacact ccccaacccc ctccagcaac tacatgtgac tgatagcttt tcccaaaggc 840
caaggaaagg atggtgtagg ttcaaaagg aaacccccca gggcctgctg tggcctagga 900
gcagattgta atgctgccga gtccggtcgg tgaccacggg ttgtccctcg gntttcagcc 960
atggggttga gttggccatt aaaagaaaca gagactctc tctgccatgg cccttcttta 1020
ttccagggac ttagaaactt gcctgagatg gtggacgcag taatgagggc accgcgcast 1080
cagttagaga cggagaaagg gaagaggctg ggatggtctc tgcctgcttt gcctctagtt 1140
catggagatg tgtctctgtt caggccaaga tacagccagc caggcctgtc gtctgggacc 1200
caggaggcct ctgatgacca agggctttca catcctaagt catttggaag gaggccttga 1260
gaacaaagtc acctttgtca ctcccagtga actgaatgag gaacatgctg tctcctgtct 1320
tggcctcccc ttctatgaga tactggggag aagagaacat tcctcctggc ttagttgtag 1380
cagaccagga cctgtgcca gctttggctc cccttcccaa ctctgaagc acgtgctgca 1440
gagccacctt ggtctgagca cctgaggacc agcccctcct ccctcagtgc gggtcatctc 1500
ttgggggatt ttcttaaagt gaagaaagg ggtggggaac catattgccc ctccctcccc 1560
catcaaaact ctttatttta acttgctata aaatgagtc tataaagaaa ctctatatgg 1620
gtgaggtata tcccacttct gtgaaaacat tacaaatcaa accgcttctc tcagtttatt 1680
taagatgctt ttgttgcgag cggagctcta gagtgaagcc tcctgtgtgt gtgtgagata 1740
ataacacctt gtaactcatt acagctgggc actatttaca taaaccagag ctgagccagg 1800
caggaatttg ctgattaatt tatttttaat ggagtgaagt ataccatgca ccaaaataaa 1860
ctttactgtg tgtacctaaa aaaaaaaaaa aaaaaaa 1897

<210> 172

<211> 1723

<212> DNA

<213> Homo sapiens

<400> 172

cggggagttc acctccgcc gtcacctcct ccccttgtcg cctagggtcca cccgagcccc 60
ctccccggg ccgccccga gcacgaagtt ggcgggagcc tataaaagct ggtgccggcg 120
cgaccgsgg acacacagt caggcgcccc agccgcccgc gccagatcgg tgccgattcc 180
tgccctgccc cgaccgccag cgcgaccatg tcccatcact gggggtacgg caaacacaac 240
ggacctgagc actggcataa ggacttcccc attgccaaag gagagcgcca gtcccctgtt 300
gacatcgaca ctcatacagc caagtatgac ccttccctga agcccctgtc tgtttcctat 360
gatcaagcaa ctccctgag gatcctcaac aatggtcatg ctttcaacgt ggagtttgat 420
gactctcagg acaaagcagt gctcaaggga ggacccctgg atggcactta cagattgatt 480
cagtttcaact ttcactgggg ttcaacttgat ggacaagggt cagagcatac tgtggataaa 540
aagaaatatg ctgcagaact tcacttgggt cactggaaca ccaaatatgg ggattttggg 600
aaagctgtgc agcaacctga tggactggcc gttctaggta ttttttgaa ggttggcagc 660
gctaaaccgg gccttcagaa agttgttgat gtgctggatt ccattaaaac aaagggcaag 720

```

agtgctgact tcactaactt cgatcctcgt ggcctccttc ctgaatccct ggattactgg 780
acctaccag gctcactgac caccctcct cttctggaat gtgtgacctg gattgtgctc 840
aaggaaccca tcagcgtcag cagcgagcag gtgttgaaat tccgtaaact taacttcaat 900
ggggaggggtg aacccgaaga actgatgggtg gacaactggc gccagctca gccactgaag 960
aacaggcaaa tcaaagcttc cttcaaataa gatgggtcca tagtctgtat ccaaataatg 1020
aatcttcggg tgtttccctt tagctaagca cagatctacc ttggtgattt ggaccctggg 1080
tgctttgtgt ctagtcttct agacccttca tctcttactt gatagactta ctaataaaat 1140
gtgaagacta gaccaattgt catgcttgac acaactgctg tggctgggtg gtgctttgtt 1200
tatggtagta gtttttctgt aacacagaat ataggataag aaataagaat aaagtacctt 1260
gactttgttc acagcatgta ggggtgatgag cactcacaat tgttgactaa aatgctgctt 1320
ttaaaccata ggaaagtaga atggttgagt gcaaaccat agcacaagat aaattgagct 1380
agttaaggca aatcaggtaa aatagtcattg attctatgta atgtaaacca gaaaaataa 1440
atgttcattg tttcaagatg ttatattaaa gaaaaacttt aaaaattatt atatatttat 1500
agcaaagtta tcttaaatat gaattctgtt gtaatttaaa gacttttgaa ttacagagat 1560
ataaatgaag tattatctgt aaaaattgtt ataattagag ttgtgatata gagtatat 1620
ccattcagac aatatatcat aacttaataa atattgtatt ttagatata tctctaataa 1680
aattcagaat tctaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 1723

```

<210> 173

<211> 1416

<212> DNA

<213> Homo sapiens

<400> 173

```

ccagtagctt ggaaagtaga gatgactaat gttttagcct tttcttgag aaaaggaaga 60
actcttcttg aatattttca cagatgattg tgattgcttt aaatgacctc tgtggcaatt 120
taaattagat ggatttaatc tcagtaattg gctggtcgca taaatgtcat gttttaatag 180
gaaaagttac ttgtaaatct ttagaccttt gttgtcactt aggctgggga gtcactacc 240
tatttgcat cttactagtt ggggggacct tttccgtgta cagtgatggg acttttgtga 300
cctttactct cactatgcaa tagagggttt catgtagtta atctgacatg tcaaaattgg 360
gaagactgta accttttttt ttttttttta agatttctct tttttgtgtc cctcaatact 420
tagcagatgt tcatttggtg gaaattctta ttacttacat gaatgagttt gaatttagtg 480
gcaaggaaga aaaaaaaaaac tcaaatattt gttttaaaag aagaaaactt gcaaagtaca 540
taagtatttt ttaaaaatca atcgaaacaga aagggaatgca tgctgttttt caatggctta 600
gacatgcttt ttattcactg actagtattc acttttttac aacttgatc aaaacaaatg 660
atctttgttt ttgtcacagg caaaaacagg ttgacactgg tgggttggtt ttattaatta 720
attttttttc tattaggttt tctttaataa tgtaaattt ctaaattata gcatatgttt 780
tagttaattc tgaaatcagt tacttctatt gtttaattat cctcatatc atgaatattg 840
ttttttaaat ktccctatac maatttgcat cacttctttt cttacagctt ttgcagttwa 900
tatattctaa acttgaaaat gtggtatcaa tcaataatag aagtatcact ggaggattta 960
tttagctttg tatttcttaa ttttagtctt agctactaaa gtatgtaagc cttaaagttt 1020
aaaatgtttt tcttaaatta gctttatata caaacatttt catttacttt atgaaatggg 1080
aggagatagt cactgtgct tatgtttttt tgtttaattt ctatattctg aagcagtga 1140
gatatagggt atgctaatac agtgagcaag gtggaacatg tacaataataa ggagaagctg 1200
taaaaatcac agtataaaat tatgaagttt ggtaactgta aaatgtactg tatttatatg 1260
taactctcat tctaaaagtt gccacaaaag ctgaattgga agcttcatgt ctgcatgaaa 1320
tttcctatat ttttaatgtg tatgatgaaa ttaatttttc ttgaatatta aagctctgcca 1380
attgctatga aaaaaaaaaa aaaaaaaaaa actcga 1416

```

<210> 174

<211> 1956

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (4)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (6)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<400> 174
nctnantaag ggaacaaagc tggngctcca ccgcggtgcc gaccgctcta gaactagtgg 60
atcccccggt ctgcaggaat tcggcacgag gatcaaccgg attttttagga tatgcaatct 120
aacaagacca caggaagggt atctgatggt gcaacagttt cagtacctag gatgggcttc 180
tcatcgagaa gtgcctgggt ccaaaagggtc attcttgaaa ctgatacttc aggtggaaaa 240
gtggcaggag gaatgcgagg aagggggaagg ccggacgatt atccactgcc taaatgggtg 300
cgggcgaagt ggcattgtct gtgctatagg catcgttggt gaaatggtga aacgggcaaa 360
atgttgctga tgttttccat gcagtaaaag cactgagggg acagcaagcc aaacatggtg 420
ggaagccccg gagcaatacc gtttctgcta tgatgtagct ttggagtacc tgggaatcat 480
cttagttggg tgagactctt taaagtgcac ccatgaagra acctgtccat ctattgagcc 540
agcagctggt gtacctgtta cacttggtga gaaagatttt aatgtggggg gtgggagact 600
tttacatttg agaggtaaaa gtattttttt tatgaagttg tgtatcttaa taaaaaggac 660
tgaattagtt tttattacta tattaagca tcaacatttc atgccacata aattatattt 720
aataagaacc agattgaaat gagaacgtat tgggtgttgt acagtgaaca tgccaccttt 780
tttctcatgg tttcagtaga gcagctacca catgttgcac gagttcatac tttctacgtg 840
gcatttttct ccctttctaa aatgaaagct gatgaatctt aaaaggaaga agaaaagaaa 900
agctgtgcaa attcatagta aagttcgttt tttatatggt tccagtgtag cagatctcta 960
tataaatata taaatatata taactggctt attttctttt aatgtgcaat gatggctgga 1020
tcatttttaa aaaaaagggt tctttttaga aaataacata agccaaagac tcaagtgtaa 1080
atatgtctat atggagaaag cacattatat ttattgggtta cttacattcc ttttttgatg 1140
gctaaaatac taccaccaca caatcatctt ttttttcctg aagaaagctt tttctttagc 1200
taaaatcaat tgtaaacgat tttttagat tattttttgt atgttttagt gtaagtagaa 1260
gataaacttt ttattcataa accaggaagc aatgttcttt atagtgattc tcttgtgtac 1320
atgcttgtga attaaatttg tgtaaaatcc cttggcaatt gggctcttta atataggacc 1380
aaatttaaac attttgctga atatgtatag tttttcacaa tttcattagg taaataatgg 1440
tttggtgatc atacatgaga aatgtacaca ttaaaaggcc ttgctgacaa cttgcacaa 1500
gttgacata gcctttaagc atcatttaaa ttttaaggga atggagtttt tcagcctgtg 1560
gcccgact ggtcaagaaa acaagatggc aacatatatg ctttcagggt caaatttgag 1620

caaactgtaa actgtcaggg tgataaaatg tttctcttga tgtttacatg cacaagcttt 1680
gcgttctgac tataaaaagt gtgaacaaat caatgccaga ttcctgtttt gcgcattgtc 1740
atgggattct taagtgaacc tttctaaatg tggctctgtt cacatgctcc acgtagctgt 1800
aacttcacat catcagcttg cagtttgtaa ttgactaaag cattccagtg tcctctttct 1860
agattgccag ctcatgacat ggtgcttata aagatttaat taaagtaaga atgaaataaa 1920
gtttttataa ttataaaaaa aaaaaaaaaa aaactc 1956

<210> 175

<211> 1689

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1688)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1689)

<223> n equals a,t,g, or c

<400> 175

ggctgcaggt cgacactagt ggatccaaag aattcggcas akctcttatt cctcactttc 60
caccgccgta ttttgtaatg accatgagca atgtttttac tttttgtata atgggggtggg 120
gtggagtggg ggcttctgag agtcagcctg aggtcttttag aggaccagct attgtagcac 180
cttggatact tgaagtttaa tgctcagttg ggtcgggtgg cagttgactt ggtggctggc 240
atgttcagca gtgcctgggg ccctgtttct gggcagcctt tgaggatttt ctatgatatt 300
gaatgcacagt ttaagtggc aactcaggcc cagctcatgc ccttttttgc ctggacatgt 360
gctattttta ttcacttata tatcaattac ttgtaagggt taaactttca aacagggaagt 420
atattgggac aaaagggctc ttggggatta gatatccctt taatctgtga ccattgggca 480
aaaaattttc ctgcagcaaa agtctgaggc tggttggacc atttttgcag ctttaatcct 540
tagcctcttt tgactgtata tttgtgttta aaatgcagag ctcaactgaa tatttccttt 600
tttgttttgt tttgttttgt ttaagaagt aggttgtttt cctgaaccgt aaacttgtat 660
cattttaact tgcacaaagg aagtctgttc ttggtattgc tcttgcaactt gggttttttg 720
ttattgtttt gtgtggattt ttaaaagctt ttctgttcac cctcctgccca ggaaaatccc 780
agaaagctta atgatacccc aaaatgatta caccagggga ggaaaaaag gagcgctttc 840
tagggtcaga atcgtggaga gaatactcag aaatgaacct ctttaaagcc ttgcaggaat 900
gagtcactct tacttaatga aatgttaaag ccaattaaaa agcatgctgt gatgccacgc 960
ttccctttcc acagggtgca tgcgtctcct gctgggtgaat cacatgcggc aagaggcaac 1020
tggtccaca gcctgggatg ctgccgtacc aagaggaaaag aagcagcaaa atgcctttac 1080
gttgttctaa acccccagcg cataaagtgt agaggaggga tggccaaggg tgggtggtag 1140
aaagtgtgtt caggctgaca ctggcaatga gtacagataa tttcactttc ctcttcagg 1200
ggcaaaggct gatggcctct acctttgtat ccaggagaaa ctgcagagca gccctgtgac 1260
tttcaaaaat atgctacctc aaagtgtctc cgataaacct ttctaattgt aagkscctta 1320
ctaagggcac atgtcttaat caaagttagt tttttgtttt ctggtttgtt tttttttttt 1380
tgtatattga tgaatgagat cttacctatt aaatatatta ttggattatg gttcctgaag 1440
gtcattagag tgtgtgtgtg tgtgtgtgtg tgtgtgtgtg ttttatgact taaatatctt 1500
tacgtgtgtt ttttagagct tggttcttta aagatttgga gaagatatgt aaattaccaa 1560
ggcacttggt ttttctgttt tatatactaa taatcagggc ctaagttaaa taaaaatatg 1620
tgtgcatgta ttttaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1680

aaaaaaaaann

1689

<210> 176

<211> 1016

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (895)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (928)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (970)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (992)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1001)

<223> n equals a,t,g, or c

<400> 176

taagagagct ttgcaaaaaa aaaaagaaaa aagaaaaaaa aaaaaccacg tttctttgtt 60
gagctgtgtc ttgaaggcaa aagaaaaaaa atttctacag tagtctttct tgtttctagt 120
tgagctgctg gcgtgaatgc ttattttctt ttgtttatga taatttcact taactttaaa 180
gacatatattg cacaaaacct ttgtttaaag atctgcaata ttatatatat aaatatatat 240
aagataagag aaactgtatg tgcgaggcca ggagtatttt tgtattagaa gaggcctatt 300
aaaaaaaaaa gttgttttct gaactagaag aggaaaaaaa tggcaatttt tgagtgccaa 360
gtcagaaaagt gtgtattacc ttgtaaagaa aaaaattaca aagcaggggt ttagagttat 420
ttatataaat gttgagattt tgcactattt tttaataata atatgtcagt gcttgcttga 480
tggaaacttc tcttgtgtct gttgagactt taaggagaaa atgtcggaa ttcagagtcg 540
cctgacggca gaggggtgag ccccggtggag tctgcagaga ggccttggcc aggagcggcg 600
ggctttcccg aggggccact gtccctgcag agtggatgct tctgcctagt gacaggttat 660
caccacgtta tatattccct accgaaggag acaccttttc ccccttgacc cagaacagcc 720
tttaaatcac aagcaaaata ggaaagttaa ccacggaggc accgagttcc aggtagtggt 780
tttgcccttc ccaaaaatga aaataaactg ttaccgaagg aattagtttt tcctcttctt 840
ttttccaact gtgaagggtc ccgtggggtg gagcatggtg cccctcacia gccgnacggc 900
tggtgcccgg gctaccaggg acatgccnga gggctcgatg acttgtctct gcaggcgct 960
ttggtggtgn ttaactggct aaaggttacc gntgaaggca ngcgcggtta ctggcc 1016

<210> 177
<211> 1364
<212> DNA
<213> Homo sapiens

<400> 177
gaatcccaact cccttctccc acttggttaat tagttacata cttttttgta attgttttatt 60
tggttgctgt ctccctctca agaatgcagg gaccatgtct gcattctgca gtaatcacta 120
ctgcacacccc agaatctatt acagatcctg gcatgtagct gatgcataaa tatttggtga 180
atgaaagtct gtacattgta tttatgctat tgggtattgct atgacctgaa actaaaagga 240
gttggtggaaa agatttctta tggaacagaa atatcccttt tgattaatat cacaatctcg 300
taaattgaga aaacaaawaa tatatactac tggagcattc atgtatagtt ggagattatg 360
actcatttat tgggtgtgtt ttggactcag aacaaagatg agggaaatatt ccttaaagct 420
ctgtattgaa ataacgaaaa gcagtcacat tttataata gaagcttcct agcttactct 480
ttctgtaatc ttcttttctt aaatgtaaga gagcctcata attatgaggc ttattactag 540
agtaaggctg tcaaaggcag caaaatgtct ttctgttttg aagaataaca taaacttgac 600
atgtatggtg ggggacagaa ggtttcaaaa gttaagaat ctgtgttgct ttaacaaata 660
gatgcttctc aaggasstta cgytagtggg tactctgtcc agtcagggtt tttcttctt 720
taacttgggt tcatttcctg atggcacaca tgaagtttg atcatatggt ttgactttag 780
ctatggtcct tagctatggg gagcagcatc agcgacctgt gacatgtaaa ttaaaaaatac 840
aatgccaggg cccttcccca gccctctga tagagaacct cttggccatc tgtattttta 900
gatgttccag gttagtctga ttaacaccct tgggttaagaa ccattgggag gatctgattg 960
ccagtttaag gggaccttca agcctgtagg tctttatagt taaaaaaaaaaa aaaagatttt 1020
aaaaatcatg catatgttgt ggctgaawtc tggtttagca catactgctt ttaatggcct 1080
gaaatgtttt tcccaaataa attstcttgt tatagcttct atgtgtgatt tgggtccagct 1140
tcttgttttg aagatactta cgggggggaa cacttttgta tttctcttag taacatatta 1200
accacttaaa aaacccttct tattacaggt cttcacattt aggcctaatg tgcttaattc 1260
aaatgtaaaa atacacctgc ctttgttctc agtgaaaagta tgtaataaat aaatgagggg 1320
ttggcaaaact actgccacc atctgttttt ttatggccta tgaa 1364

<210> 178
<211> 740
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (38)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (171)
<223> n equals a,t,g, or c

<400> 178
tggctgggac tccaggcgca taccaccatg actggctnat ttttgtgttt ttagtagaga 60
cggggtttca ccatgttggc caggetggtc ttgaacttct ggccccaagt catctgcctg 120
cctcagcctc ccaaggtgct gggattacaa gtgtgagcca ctgtgtacag nctttgtgtt 180
aacttacttt gaattagccc cttattccaa agctgcctcc tgaaaggasg cttcctaaaa 240
ggtgttgctt tggggtaaat aagaggtgct ggtgaagtga gactagcttg cgcactcttc 300

```
tgccatgaag gaaactagct ccgcagacgt tttaactaacc acagcagacc ccctccttga 360
ggaacccaga agggcagaga gcatcagatg aaatcaactg ctgagagagg gaagtgaagt 420
ctgtgttggc accgctgccg gaaactcttg agaatgagca tcgtgtgccg cttatccact 480
gcttgccgac ttcagtagca agagcctcca gaatgctcat ggataaacat catctttgtg 540
gccacacatt gttttatgag ctgagtcatc agactggctg gctccttaaa tgctgctgtg 600
tgattctcta ggcagcagaa aacaaggaga gaagatatga agtgatatctc atcgagcttt 660
gtagagggcc agtatcgatt caccaaaggt gatgggggat ttaccattat ggagtttgct 720
tgtgataaaa aattattgat                                     740
```

<210> 179

<211> 1410

<212> DNA

<213> Homo sapiens

<400> 179

```
gaaaatgtat ataataggca aggaaagaaa tacagtactg tttctggacc cttataaaaat 60
cctgtgcaat agacacatac atgtcacatt tagctgtgct cagaagggct atcatcaycc 120
tacaactcac attagagaac atcctggctt ttgagcactt ttcaaacaat caagttgact 180
cacgtgggtc ctgaggcctg cagcacgtcg gatgctaccc cactatgaca gaggattgtg 240
gtcacaactt gatggctgcg aagacctacc ctccgttttt ctactagata ggaggatggg 300
agaagtttgg ctgctgtcat aacatccaga gcyttgtcgt atttggcaca cagcagaggc 360
ccagatatta gaaaggctct attccaataa actatgagga ctgccttatg gatgatttaa 420
gtgtctcact aaagcatgaa atgtgaattt ttattgttgt acatacgatt taaggtattt 480
aaagtatttt ctctctgtg agaaggttta ttgttaatac aaggtataat aaaattatcg 540
caacccctct ccttccagta taaccagctg aagttgcaga tgttagatat ttttcataaa 600
caagttcgag tcaaagttga aaattcatag taagattgat atctataaaa tagatataaa 660
tttttaagag aaagaattta gtattatcaa agggataaaag aaaaaaatac tatttaagat 720
gtgaaaatta cagtccaaaa tactgttctt tccaggctat gtataaaata catagtgaag 780
attgtttagt gatattacat ttatttatcc agaaaactgt gatttcagga gaacctaaca 840
tgctggtgaa tattttcaac tttttccctc actaattggt acttttaaaa acataacata 900
aattttttga agtctttaat aaataaccca taattgaagt gtataatata aaaaatttta 960
aaaatctaag cagcttattg tttctctgaa agtggtgtga gttttacttt cctaaggaat 1020
taccaagaat atcctttaaa atttaaaagg atggcaagtt gcatcagaaa gctttatttt 1080
gagatgtaaa aagattccca aacgtggtta cattagccat tcatgtatgt cagaagtga 1140
gaattggggc acttaatggt caccttgtaa cagttttgtg taactcccag tgatgctgta 1200
cacatatttg aagggtcttt ctcaaagaaa tattaagcat gttttgttgc tcagtgtttt 1260
tgtgaattgc ttggttgtaa tttaattctg agcctgatat tgatatggtt ttaagaagca 1320
gttgtaacca gtgaaattat tttggagatt ataataata tatacattca aaaaaaaaaa 1380
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa                                     1410
```

<210> 180

<211> 1493

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1328)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (1352)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1406)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1436)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1484)
<223> n equals a,t,g, or c

<400> 180
ggccaaatct tcccgggctg ctgtgggcaa caagatgttt gtcaaggggtg cccctgaggg 60
cgctcatcgac cgctgtaact atgtgcgagt tggcaccacc cgggtgccac tgacggggcc 120
ggtgaaggaa aagatcatgg cgggtgatcaa ggagtggggc actggccggg acaccctgcg 180
ctgcttggcc ctggccaccc gggacacccc cccgaagcga gaggaaatgg tcctggatga 240
ctctgccagg ttcttgagat atgagacgga cctgacattc gtgggtgtag tgggcatgct 300
ggaccctccg cgcaaggagg tcacgggctc catccagctg tgccgtgacg ccgggatccg 360
ggtgatcatg atcactgggg acaacaaggg cacagccatt gccatctgcc ggcgaattgg 420
catctttggg gagaacgagg aggtggccga tcgcgccctac acgggcccag agttcgacga 480
cctgcccctg gctgaacagc ggggaagcctg ccgacgtgcc tgctgcttcg cccgtgtgga 540
gccctcgcac aagtccaaga ttgtggagta cctgcagtcc tacgatgaga tcacagccat 600
gacaggtgat ggcgtcaatg acgcccctgc cctgaagaag ctgagattgg cattgccatg 660
ggatctggca ctgcccgtggc caagactgcc tctgagatgg tgctggctga cgacaacttc 720
tccaccatcg tarctgctgt ggaggarggc cgcgccatct acaacaacat gaagcagttc 780
atccgctacc tcatttcctc caacgtgggc gaggtggctc gtatcttcct gaccgctgcc 840
ctggggctgc ctgaggccct gatcccgggt cagctgctat gggtgaaactt ggtgaccgac 900
gggctcccag ccacagccct gggcttcaac ccaccagacc tggacatcat ggaccgcccc 960
ccccggagcc ccaaggagcc cctcatcagt ggctggctct tcttcgcta catggcaatc 1020
gggggctatg tgggtgcagc caccgtggga gcagctgcct ggtggttcct gtacgctgag 1080
gatgggcctc atgtcaacta cagccagctg actcacttca tgcagtgcac cgaggacaac 1140
accactttg agggcataka ctgtgaggtc ttcgaggccc ccgagcccat gaccatggcc 1200
ctgtccgtgc tggtagccat cgagatgtgc aatgcaactga acagcctgtc cgaraaccag 1260
tccttgcctg ggaattgcma cccctgggggt traacaattc ttgggctggc tgggsttcca 1320
ttcttggnct tcttccaatg tccccctggc ancttccctc aattccctcc taatgnntga 1380
accccccttg cccgaatgaa tctttnaagg cttccggggc ccttggaacc ttcaanccaa 1440
gttgggttaa tggttcctcc aagattttca attggccagt taantggggg ttc 1493

<210> 181
<211> 2040
<212> DNA
<213> Homo sapiens

<400> 181
gcatgagctc atgcaagagg caggggatga gtgtgagccc gagtgggtgtg atgccgagga 60
cccactcttc atcctgtaca ccagtggctc cacaggcaaa cccaaggggtg tggttcacac 120
agttgggggc tacatgctct atgtagccac aaccttcaag tatgtgtttg acttccatgc 180
agaggatgtg ttctgggtgca cggcagacat tggttggatc actggtcatt cctacgtcac 240
ctatggggcca ctggccaatg gtgccaccag tgttttgttt gaggggatc ccacatatcc 300
ggacgtgaac cgcctgtgga gcattgtgga caaatacaag gtgaccaagt tctacacagc 360
accacagcc atccgtctgc tcatgaagtt tggagatgag cctgtcacca agcatagccg 420
ggcatccttg caggtgttag gcacagtggg tgaacccatc aacctgagg cctggctatg 480
gtaccaccgg gtggtaggtg ccagcgtg cccatcgtg gacaccttct ggcaaacaga 540
gacaggtggc cacatgttga ctccccttcc tgggtgccaca cccatgaaac ccggttctgc 600
tactttccca ttctttgtg tagctcctgc aatcctgaat gagtccgggg aagagttgga 660
aggtgaagct gaaggttata tgggtgtcaa gcagccctgg ccagggatca tgcgcacagt 720
ctatgggaac cacgaacgct ttgagacaac ctactttaag aagtttcctg gatactatgt 780
tacaggagat ggctgccagc gggaccagga tggctattac tggatcactg gcaggattga 840
tgacatgctc aatgtatctg gacacctgct gagtacagca gaggtggagt cagcacttgt 900
ggaacatgag gctgttgca aggcagctgt ggtggggccac cctcatcctg tgaaggggtga 960
atgcctctac tgctttgtca ccttgtgtga tggccacacc ttcagcccca agctcaccga 1020
ggagctcaag aagcagatta gagaaaagat tggcccccatt gccacaccag actacatcca 1080
gaatgcacct ggcttgcccta aaacccgctc agggaaaatc atgaggcgag tgcttcggaa 1140
gattgctcag aatgaccatg acctcgggga catgtctact gtggctgacc catctgtcat 1200
cagtcacctc ttcagccacc gctgcctgac catccagtga acatgatcct gacctttacc 1260
taggattcct cctgtcccaa actttgcca tcctctttgc cccctcagga gtgctgaggg 1320
ccagtgttga cccacactac cctcccttga ccagtgtct gggaccggaa accagctttg 1380
tctccaggta gagacaacat cctgtgactg ccaggcagaa aggacagggc ccaggtcagc 1440
ctcagctctg tgtgcctcca gactgcagag ctctcagaac ccagaacaga gacgaaaagg 1500
ctacctctcc tacccaagtt aagtgttcaa aggggatgtg agggcctcca ctgaagcagg 1560
gaggcagctg tgtaatccta tgtcagctct cttaggaagc cccagtactt atattgggca 1620
tgcacttgcc cttaaaaaa atgattttgt agtccaggaa caatttacta tttttaaaat 1680
atthttgtgc ttctgttctg ggtctgaatt cccttttgtg ccagatgcca gtactgtctg 1740
ccatttggt ccaggggctg tatgggcaga ttcagtctcc agagggtatt cagatcatct 1800
gcttctttga aggagtaaat gtgttttgtt cctagggcca gaggagcttg tcttccttgt 1860
cctctgttcc caccctcccc tgaacagaa ccagcccata agagacattc tcagatgaaa 1920
ctctgttttc ttgccccagt caggctcaag ccctgtggtt gtaggaataa agcctgtgat 1980
ctcaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaactcgaa 2040

<210> 182
<211> 969
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (128)

<223> n equals a,t,g, or c

<400> 182

```
agacagttct natttctccc agggcagggg gcaggttatg accaggacta aggtcccaaa 60
gtccccaccc tgaacccctc cctgctgttc caccgctccc tcatatccac ccctgcccc 120
tctcctgnac tttggtcacg ctagcatctt ctgctgatcc tgaaattgta ccagcggcaa 180
gatgtggcct ggaaggggac tttaagttct ccacaactgc cagcaatcct tccaccaggc 240
aaaacacatc atctaaggaa aagaagttag gtttgcttag ggcgtggcag cttcggataa 300
acgcaggact ccgcctggca gcccgaattc tcccgaacc tctgctcagc ctggtgaacc 360
acacaggcca gcgctctgac atgcagaagg tgaccctggg cctgcttggt ttcctggcag 420
gctttcctgt cctggacgcc aatgacctag aagataaaaa cagtcctttc tactatgact 480
ggcacagcct ccaggttggc gggctcatct gcgctggggt tctgtgcgcc atgggcatca 540
tcctcgtcat gagtgaagtgg aggagctcgg gggagcaggc gggccggggc tggggctccc 600
ctcccctgac cactcagctc tcccacacag gtgcaaatg caaatgcaag tttggccaga 660
agtccggtca ccatccaggg gagactccac ctctcatcac cccaggctca gcccagaagt 720
gatgaggaca gaccagctga aattgggtgg aggaccgttc tctgtcccca ggtcctgtct 780
ctgcacagaa acttgaactc caggatggaa ttcttctctc tctgctggga ctcccttgca 840
tggcagggcc tcatctcacc tctcgcaaga gggctctctt gttcaatttt ttttwtacta 900
aaatgattgt gcctctgccc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 960
aaaaaaaga
```

<210> 183

<211> 1452

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (18)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1430)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1441)

<223> n equals a,t,g, or c

<400> 183

```
gtcgaccac gcgtccnncc acggtccggt ggggagaagc cgggaggact ggggtgcgct 60
```

gcagggatcg gaagccgggtt ggggtgtgag aggttttctc gctctagga gattcttcaa 120
gcaatcacta tgtcaacaga cacaggtgtt tcccttcctt catatgagga agatcaggga 180
tcaaaactca ttcgaaaagc taaagaggca ccattcgtac ccgttggaat agcgggtttt 240
gcagcaattg ttgcatatgg attatataaa ctgaagagca ggggaaatac taaaatgtcc 300
attcatctga tccacatgcg tgtggcagcc caaggctttg ttgtaggagc aatgactgtt 360
ggtatgggct attccatgta tcgggaattc tgggcaaaac ctaagcctta gaagaagaga 420
tgctgtcttg gtcttgttgg aggagcttgc tttagttaga tgtcttatta ttaaagttag 480
ctattattgt tggaaataaa ctaatttgta tgggtttaga tggtaacatg gcattttgaa 540
tattggcttc ctttcttgca ggcttgattt gcttgggtgac cgaattacta gtgactagt 600
tactaactag gtcattcaag gaagtcaagt taacttaaac atgtcaccta aatgcacttg 660
atgggtgttg aatgtccacc ttcttaaatt ttttaagatga acttagttct aaagaagata 720
acaggccaat cctgaaggta ctccctgttt gctgcagaat gtcagatatt ttggatgttg 780
cataagagtc ctatttggcc cagttaattc aacttttgtc tgccctgttt gtggactggc 840
tggctctgtt agaactctgt ccaaaaagtg catggaatat aacttgtaaa gcttcccaca 900
attgacaata tatatgcatg tgtttaaacc aaatccagaa agcttaaaaca atagagctgc 960
ataatagtat ttattaaaga atcacactg taaacatgag aataacttaa ggattctagt 1020
ttagtttttt gtaattgcaa attatatttt tgctgctgat atattagaat aattttttaa 1080
tgtcatcttg aaatagaaat atgtatttta agcactcacg caaaggtaaa tgaacacgtt 1140
ttaaatgtgt gtgttgctaa tttttccat aagaattgta aacattgaac tgaacaaatt 1200
acctataatg gatttgggta atgacttatg agcaagctgg tttggccaga cagtataccc 1260
aaacttttat ataataatac gaaggctatc acacttgtga aattctcttg tctaacttga 1320
atttgcattc catgggtgta acatgggtata tgtattgtta ttaaagtaag tgacccatgt 1380
caaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1440
naaaaaaaaa aa 1452

<210> 184

<211> 2119

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2090)

<223> n equals a,t,g, or c

<400> 184

aaattgaaac ttctaataaa aatgatatga ctatagatat attacatgct gatggtgaaa 60
gacctaatgt tctagaaaac ctagacaact caaaggaaaa gactggttga tcagaagcag 120
caaaaactga agatacagtt ctctgcagca gtgatacaga tgaggagtgt ttaatcattk 180
wtacagaatg taaaaataat agtgatggaa agacagctgt tgtgggttct aacttaagtt 240
ccagaccagc tagtccaaat tcttctcag gacaggcttc tgtaggaaac cagactaata 300
ctgcttgtwg tcttgaagag tcatgtgttt taaaaaaacc tatcaaacga gtatataaaa 360
aatttgatcc agttggagag attttaaaaa tgcaggatga gctcttwaag ccaattttcca 420
gaaaagtacc agaattgccc ttaatgaatt tagaaaattc taaacagcct tctgtttctg 480
agcaattgtc tggctcctca gactcctcta gttggccgaa atctggatgg ccttctgcat 540
ttcagaagcc aaaaggacga ttgccatag aacttcagga ctatgttgaa gatacatcgg 600
aatacctagc tcttcaggaa ggaaattttg tttataagtt atttagcctg caagacctgt 660
tgttactcgt acgctgcagt gtccagagga tagagacaag accacgttct aaaaaacgga 720
agawwatyg aagacaattt ccagtttatg tactaccaaa agtagagtat caagcttgtt 780
atggagtgtg agctctgact gaaagtgaac tttgtcgctt atggactgaa agtttattgc 840
attccaacag ctcattttat gttgggcata tcgatgcatt tacttcaaaa ctttttctac 900

tggaagaaat tacctcagaa gaattaaaaag aaaagctttc agcactcaag atttccaatt 960
tatttaacat cctccaacac attctaaaga aactaagtag cttgcaggag gggttcctact 1020
tgttatctca tgcagcagaa gattcttcac tcctgattta taaggcctct gatggaaaag 1080
ttactaggac agcatacaat ttgtataaaa cacattgcgg ccttcctggt gtaccttcca 1140
gtctctcagt tccctgggtc ccattagatc ccagcctggt attaccatat catatccatc 1200
atggaagaat accttggtact tttccaccga aatcactgga taccacaaca caacaaaaa 1260
ttggtggaac gagaatgcct acacgcagcc acaggaatcc agtttccatg gaaacaaaa 1320
gcagttgctt gcctgctcag caagttgaaa ctgaaggagt ggctccacat aaaagaaaa 1380
taacttgagg actgtacat ggaaaaactaa atttaaaaa mcagttataa cagtgtttaa 1440
tttagataag tttgaggga aataatcagt aggcaaggag aacatttttc ctgtagtagc 1500
tagagtgcct tgaaaaaatg tgttggtat gtgaaggagt atttcaacta aaatggaatg 1560
gtatgctttt cacccttraa gtttgaggag gatcttgata tgttttaaca ttatcatggc 1620
agggaaatat ataaagaaga aaaatatattt tacattaaac cttttctaaa aattgtaaat 1680
agaaaaataa tttggtttt tatcaagaac aacacttatc gttatgtatt gtgttagtta 1740
tattgccagt ctgttgcgac tgactcaaaa agttaaatgt tgccactgct gaagatgatt 1800
atgagcatcg caaactttgt ttctgacca ttttgacagt ttttatatac tccttataaa 1860
tgatgaatgt tacaggttaa taaagttaat accttataaa acttggtgaa attccattac 1920
agaagccaaa aataaaaact cctgcctct gaaaagtcag attactgact tcttgtttg 1980
caaccatcag tttgtttaat aaaagaaaa atttggtggt ataacatgtt tgatgacaga 2040
tgcctctatc tctagattca agctgagtgt tgaaatacac tgctgaaagn aagagatagg 2100
tatgttttcc agaaaaaaa 2119

<210> 185

<211> 1325

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1319)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1320)

<223> n equals a,t,g, or c

<400> 185

atgactgggg agcgacccag ncacggcgct cccgggacag acgctggggg ccscggaatc 60
tctccctgga tattggcact gaagtgtttg cccctggacc gggctctggt attcagaaac 120
agcgggagcc acggaagggc cgctcatcg tgttgggcca tgggacgctg gagcgggacg 180
gagtcctctg tctctcagc gatgatcatg gtgcctcctg gcgctacgga agtgggggtca 240
gcggcatccc ctacggtcag cccaagcagg aaaatgattt caatcctgat gaatgccagc 300
cctatgagct ccagatggc tcagtcgtca tcaatgccc aaaccagaac aactaccact 360
gccactgccg aattgtcctc cgcagctatg atgcctgtga tacactaagg ccccgatgatg 420
tgaccttcga ccctgagctc gtggaccctg tggtagctgc aggagctgta gtcaccagct 480

ccggcattgt cttcttctcc aaccacagcac atccagagtt ccgagtgaac ctgaccctgc 540
gatggagcct cagcaatggt acctcatggc ggaaagagac agtccagcta tggccaggcc 600
ccagtggcta ttcacccctg gcaaccctgg agggcagcat ggatggagag gagcaggccc 660
cccagctcta cgtcctgtat gaaaaaggcc ggaaccacta cacagagagc atctccgtgg 720
ccaaaatcag tgtctatggg acactctgag ctgtgccact gccacagggg tattctgcct 780
tcaggactct gccttcagga acacgggtct gtagagggtc tgctggagac gcctgaaaga 840
cagttccatc ttcctttaga ctccagcctt ggcaaaatca ccttcccttt accagggaaa 900
tcacttcctt taggactgaa agctaggcgt cctctccac aaaaaagtcc tgccctcatc 960
tgagaatact gtctttccat atggctaagt gtggcccccac caccctctct gcctcccggt 1020
acattgattg gtcctgtctt gggcaggtct agtgagctgt agaattgaat caatgtgaac 1080
tcagggaact ggggaaggct gagcctcctc ttgggtgttg cggtaagata accgacaggg 1140
ctggtgaaaag tcccagatg gcaggatatt tggtttcaga gtaaggacta ggtgcaccac 1200
catgactgac tatcaatcaa aatgtttgta acttaaaatt tttaatgaag gataatgaat 1260
attaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaann 1320
aaaaa 1325

<210> 186

<211> 433

<212> DNA

<213> Homo sapiens

<400> 186

aattcgccac gagggcgag gccccagcca gctcaggcta cactatccca ggatcagcat 60
ggccgtccgc cagtgggtaa tcgccctggc cttggctgcc ctcttctgtg tggacaggga 120
agtgccagtg gcagcaggaa agctcccttt ctcaagaatg cccatctgtg aacacatggt 180
agagtctcca acctgttccc agatgtccaa cctgggtctgc ggcaactgat ggctcacata 240
tacgaatgaa tgccagctct gcttggcccc gataaaaacc aaacaggaca tccagatcat 300
gaaagatggc aaatgctgat cccacaggag cacctcaagc catgaagtgt cagctggaga 360
acagtgggtg gcatggagag gatatgacat gaaataaaag atccagccca aaaaaaaaaa 420
aaaaaaaaaa aaa 433

<210> 187

<211> 859

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (803)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (853)

<223> n equals a,t,g, or c

<400> 187

cccacgcgtc cgcccacgcg tccgcccacg cgtccgaccg gggcttttcg tgagctcgtc 60
tctgatctcg cgcaagagtg acacacaggt gttcaaagac gcttctgggg agtgagggaa 120
gcggtttacg agtgacttgg ctggagcctc aggggcgggc actggcacgg aacacaccct 180
gaggccagcc ctggctgccc aggcggagct gcctcttctc ccgcgggttg gtggaccgcg 240

tcagtacgga gttggggaag ctctttcact tcggaggatt gctcaacaac catgctgggc 300
atctggaccc tcctacctct ggttcttacg tctgttrcta gattatcgtc caaaagtgtt 360
aatgccaag tgactgacat caactccaag ggattggaat tgaggaagac tggtactaca 420
gttgagactc agaacttgga aggcctgcat catgatggcc aattctgcca taagccctgt 480
cctccagggtg aaaggaaagc tagggactgc acagtcaatg gggatgaacc agactgcgtg 540
ccctgccaaag aagggaagga gtacacagac aaagcccatt tttcttcaa atgcagaaga 600
tgtagattgt gtgatgaagg acatggctta kaagtggaaa taaactgcac cgggaccag 660
aatmccaagt gcagatgtaa accaaacttt ttttgkaact ctactgtatg tgaacactgt 720
gacccttgca ccaaattgtga acatggaatc atcaaggaat gcacactcac cagcaacacc 780
aagtgc aaag aggaaggatc canatctaac ttgggggtggc ttgggcttct tcttttgcca 840
attccctaatt tgn ttgggt 859

<210> 188

<211> 833

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (65)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (798)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (803)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (809)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (812)

<223> n equals a,t,g, or c

<400> 188

tcgcagagat tacaccactg caacgagact cgggccccaga cgagatctgc ctgggcgaca 60
cagtnagact ctgtctcaaa ataaataaat aagtaaatag aacctcctga gctaccctgg 120
cttataagga ttcagattac tttaggaaat taggcattag cactaggagg ggggaagatg 180
gctggcaaaa cttctagcta gcaacctatg tctagcacat gggtaactaa ccatagcgag 240
atcctgaaca catatcctct aggtgcagggt ggagggaaacg atgtccaata cctgaagcag 300
aatctcacat ggacggaacg tctgtatttc cctctcctgc acgaatcact catcattctt 360
ggaggggcttc tctgcattcc tcttttctt ctctccctc ccctgccttt tgtcttttct 420
aaagagtctg aactccgatt tccatgctct cctgctacat taataagcaa aacctgcttg 480

```

tgtgtacggt tctttactgg aaacatgact ttttgtttct gtattggttt tactgtcatc 540
cagttttcta gtttaatatc tagcaaaact aaatctgaat gtactcgctt tttccgttaa 600
gtatgccaaag cacctcggtt gaaaggcttg actcaaacca gaagtcttgc tggaattcgt 660
ctctgaacac ttgaaaaaca gaaaccctga gccgcaacaa acatgcctct gtgtgtcggg 720
attgcctttg tctctgcttc catgtggctt ttccttttgt agctatgctt agtgacataa 780
tcttccctct tcaagcantc tgngttggnc cnttcccggt ccccccccg tgc 833

```

<210> 189

<211> 2211

<212> DNA

<213> Homo sapiens

<400> 189

```

ggcacgagga gtacagggcc cgcgcatgcy tggattgtcg tcttctgtcc aagttggctg 60
cttccctgcy ccaaagttag cagtagccaa catgtcaggg tgggagtcac attacaaaac 120
cgaggcgcat gaagaagcag aggaagaaca agaagagaac cttgaagcaa gtggagacta 180
taaataattca ggaagagata gtttgatttt tttggttgat gcctccaagg ctatgtttga 240
atctcagagt gaagatgagt tgacaccttt tgacatgagc atccagtgtg tccaaagtgt 300
gtacatcagc aagatcataa gcagtgatcg agatctcttg gctgtggtgt tctatggtac 360
cgagaaagac aaaaattcag tgaattttta aaatatctac gtcttacagg agctggataa 420
tccaggtgca aaacgaattc tagagcttga ccagtttaag gggcagcagg gacaaaaacg 480
tttccaagac atgatgggcc acggatctga ctactcactc agtgaagtgc tgtgggtctg 540
tgccaacctc tttagtgtatg tccaattcaa gatgagtcac aagaggatca tgctgttcac 600
caatgaagac aacccccatg gcaatgacag tgccaaagcc agccgggcca ggaccaaagc 660
cgggtgatctc cgagatacag gcactcttct tgacttgatg cacctgaaga aacctggggg 720
ctttgacata tctttgttct acagagatat catcagcata gcagaggatg aggacctcag 780
gggttcacttt gaggaatcca gcaagctaga agacctgttg cggaagggtc gcgccaagga 840
gaccaggaag cgagcactca gcagggttaa gctgaagctc aacaaagata tagtgatctc 900
tgtgggcatt tataatctgg tccagaaggc tctcaagcct cctccaataa agctctatcg 960
ggaaacaaat gaaccagtga aaaccaagac ccggaccttt aatacaagta caggcggttt 1020
gcttctgcct agcgatacca agaggtctca gatctatggg agtcgtcaga ttatactgga 1080
gaaagaggaa acagaagagc taaaacggtt tgatgatcca ggtttgatgc tcatgggttt 1140
caagccgttg gtactgctga agaaacacca ttacctgagg cctccctgtc tctgtacctc 1200
agaggagtcg ctgggtgatt ggagctcaac cctgttcagt gctctgctca tcaagtgtct 1260
ggagaaggag gttgcagcat tgtgcagata cacacccgcg aggaacatcc ctcttattt 1320
tgtggctttg gtgccacagg aagaagagtt ggatgaccag aaaattcagg tgactcctcc 1380
aggcttccag ctgggtctttt taccctttgc tgatgataaa aggaagatgc cctttactga 1440
aaaaatcatg gcaactccag agcaggtggg caagatgaag gctatcgttg agaagcttcg 1500
cttcacatac agaagtgaca gctttgagaa ccccgctgctg cagcagcact tcaggaaacct 1560
ggaggccttg gccttggtt tgatggagcc ggaacaagca gtggacctga cattgcccga 1620
ggttgaagca atgaataaaa gactgggctc cttggtggat gagtttaagg agcttgttta 1680
cccaccagat tacaatcctg aagggaaggt taccaagaga aaacacgata atgaagggtc 1740
tggaagcaaa aggcccaagg tgagtatctc agaagaggag ctgaagacct acatcagcaa 1800
gggtacgctg ggcaagttca ctgtgcccat gctgaaagag gcctgcccgg cttacgggct 1860
gaagagtggt ctgaagaagc aggagctgct ggaagccctc accaagcact tccaggactg 1920
accagaggcc gcgcgtccag ctgcccttcc gcagtgtggc caggctgcct ggccttctcc 1980
tcagccagtt aaaaatgtgtt tctcctgagc taggaagagt ctaccgcaga taagtcgagg 2040
gactttatgt ttttgaggct ttctgttgcc atggtgatgg tgtagccctc ccactttgct 2100
gttcccttact ttactgcctg aataaagagc cctaaagttt tactataaaa aaaaaaaaaa 2160
aaaaaaaaaa aaagtcgacc ggccgttaat ttagtagtag taggcggccg c 2211

```

<210> 190
<211> 1659
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1582)
<223> n equals a,t,g, or c

<400> 190
atcgggttca agaagaaaat gctaggctta agaagaaaa agagcagttg cagcaggaaa 60
tcgaagactg gagtaaatca catgctgagc tcagttagca aatcaaatca tttgagaagt 120
ctcagaaga tttggaagta gctcttactc acaaggatga taatattaat gctttgacta 180
actgcattac acagttgaat ctgttagagt gtgaatctga atctgagggt caaaataaag 240
gtggaaatga ttcagatgaa ttagcaaatg gagaagtggg aggtgaccgg aatgagaaga 300
tgaaaaatca aattaagcag atgatggatg tctctcggac acagactgca atatcggtag 360
ttgaagagga tctaaagctt ttacagctta agctaagagc ctccgtgtcc actaaatgta 420
acctggaaga ccaggtaaaag aaattggaag atgaccgcaa ctactacaa gctgccaaag 480
ctggactgga agatgaatgc aaaaccttga ggcagaaagt ggagattctg aatgagctct 540
atcagcagaa ggagatggct ttgcaaaaaga aactgagtca agaagagtat gaacggcaag 600
aaagagagca caggctgtca gctgcagatg aaaagggcagt ttcggctgca gagggaagtaa 660
aaacttacaa gcggaagaatt gaagaaatgg aggatgaatt acagaagaca gagcgggtcat 720
ttaaaaacca gatcgctacc catgagaaga aagctcatga aaactggctc aaagctcgtg 780
ctgcagaaag agctatagct gaagagaaaa ggggaagctgc caatttraga cacaattat 840
tagawttaac acaaaagatg gcaatgctgc aagaagaacc tgtgattgta aaaccaatgc 900
caggaaaacc aaatacacaa aaccctccac ggagagggtcc tctgagccag aatgtctttt 960
ggcccatccc ctgtgagtgg tggagaatgc tcccctccat tgacagtgga gccaccctg 1020
agacctctct ctgctactct caatcgaaga gatatgccta gaagtgaatt tggatcagtg 1080
gacgggcttc tacctcatcc tcgatggtea gctgaggcat ctgggaaacc ctctccttct 1140
gatccaggat ctggtacagc taccatgatg aacagcagct caagaggctc tccccctacc 1200
aggggtactc atgaaggcaa ggttaatatg gctccaaaag ggccccctcc tttcccagga 1260
gtccctctca tgagcaccac catgggaggg cctgtaccac caccattctg atatggacca 1320
ccacctcagc tctgcggacc ttttgggctt cggcacttcc tccacccttt ggccctggta 1380
tgcgctccacc actaggctta agagaatttg caccaggcgt tccaccagga agacgggacc 1440
tgccctctcca ccctcgggga tttttacctg gacacgcacc atttagacct ttaggttcac 1500
ttggcccaag agagtacttt attcctggta cccgattacc accccaacc catggtcccc 1560
aggaataccc accamcacct gnttgtaar gagratttat kgcgttcagt gtttagagatg 1620
aggcctcaac tggcttctta gaggcattag ccagggatt 1659

<210> 191
<211> 3894
<212> DNA
<213> Homo sapiens

<400> 191
cacacttgaa gctgaaaaag aaagaagaaa atctgggcta tcctcaagag ttcagtttcg 60
aaaccaaggt tctgagccca aatatactca agaactaact ctgaagaggc agaaacagaa 120
agtgtgcatg gaggaacccc tgtggctaca ggataatata agagataaac tgcgtcccat 180
tcccataact gcctcagtg agatccaaga gccaaagctct cgtaggcgag tgaattcact 240
tccagaagtt cttccaattc tgaattcaga tgaacccaag acagctcata ttgatgttca 300

cttcttaaaa gagggatgtg gagacgacaa tgtatgtaac agcaacctta aactagaata 360
taaatTTTgc acccgagaag gaaatcaaga caaatTTwct tatttaccaa ttcaaaaagg 420
tgtaccagaa ctagttctaa aagatcagaa ggatattgct ttagaaataa cagtgcacaa 480
cagcccttcc aacccaagga atcccacaaa agatggcgat gaygcccatg aggctaaact 540
gattgcaacg tttccagaca cttaacctta ttctgcatat agagaactga gggctttccc 600
tgagaaacag ttgagttgtg ttgccaacca gaatggctcg caagctgact gtgagctcgg 660
aaatcctttt aaaagaaatt caaatgtcac tttttatttg gttttaagta caactgaagt 720
cacctttgac accccagatc tggatattaa tctgaagtta gaaacaacaa gcaatcaaga 780
taatttggct ccaattacag ctaaagcaaa agtgggttatt gaactgcttt tatcggctctc 840
gggagttgct aaaccttccc aggtgtattt tggaggtaca gttgttgccg agcaagctat 900
gaaatctgaa gatgaagtgg gaagtttaat agagtatgaa ttcagggtaa taaacttagg 960
taaacctctt acaaacctcg gcacagcaac ctgaacatt cagtggccaa aagaaattag 1020
caattgggaaa tggttgcttt atttggtgaa agtagaatcc aaaggattgg aaaaggtaac 1080
ttgtgagcca caaaaggaga taaactccct gaacctaacg gagkctcaca actcaagaaa 1140
gaaacgggaa attactgaaa aacagataga tgataacaga aaattttctt tatttgctga 1200
aagaaaatac cagactctta actgtagcgt gaacgtgaac tgtgtgaaca tcagatgcc 1260
sctgcggggg ctggacagca aggcgtctct tattttgcgc tcgaggttat ggmacagcac 1320
atttctagag gaatatcca aactgaacta ctggacatt ctcatgcgag ccttcattga 1380
tgtgactgct gctgccgaaa atatcaggct gccaaatgca ggcactcagg ttcgagtgc 1440
tgtgtttccc tcaaagactg tagctcagta ttcgggagta ccttggtgga tcatcctagt 1500
ggctattctc gctgggatct tgatgcttgc tttatttagt tttatactat ggaagtgtgg 1560
tttcttcaag agaaataaga aagatcatta tgatgccaca tatcacaagg ctgagatyca 1620
tgctcagcca tctgataaag agaggsttac ttcygatgca tagtattgat ctacttctgt 1680
aattgtgtgg attcyttaa cgctctaggt acgatgacag tgttccccga taccatgctg 1740
taaggatccg gaaagaagag cgagagatca aagatgaaaa gtatattgat aaccttgaaa 1800
aaaaacagtg gatcacaagg tggaacgaaa atgaaagcta ctcatagcgg gggcctaata 1860
aaaaaaagct tcacagtacc caaactgctt tttccaactc agaaattcaa tttggattta 1920
aaagcctgct caatccctga ggactgattt cagagtgcact acacacagta cgaacctaca 1980
gttttaactg tggatattgt tacgtagcct aaggctcctg ttttgcacag ccaaatttaa 2040
aactgttgga atggatTTTT cttaactgc cgtaatTTaa ctttctgggt tgcctttrtt 2100
tttggcgtag ctgacttaca tcatgtgttg gggaaaggcc tgcccagttg cactcagggtg 2160
acatccctca gatagtgtag ctgaggaggc acctacactc acctgcacta acagagtggc 2220
cgtcctaacc tcgggcctgc tgcgcagacg tccatcacgt tagctgtccc acatcacaag 2280
actatgccat tggggtagtt gtgtttcaac ggaaagtgtc gtcttaaaact aaatgtgcaa 2340
tagaagggtg tggtgccatc ctaccgtctt ttccgtgttc ctactgtgtg gaatacctgc 2400
tcacgtcaaa tgcatacaag tttcattctc ctttctacta aaacacacag gtgcaacaga 2460
cttgaatgct agttatactt atttgtatat ggtatttatt ttttcttttc tttacaaacc 2520
attttgttat tgactaacag gccaaagagt ctccagttta cccttcagggt tggtttaatc 2580
aatcagaatt agagcatggg aggtcatcac tttgacctaa attatttact gcaaaaagaa 2640
aatctttata aatgtaccag agagagtgtt tttataaact tatctataaa ctataacctc 2700
tccttcatga cagcctccac cccacaaccc aaaaggTTta agaaatagaa ttataactgt 2760
aaagatgttt atttcaggca ttggatattt ttacttttag aagcctgcat aatgtttctg 2820
gatttcatac tgtaacattc aggaattctt ggagaaaatg ggtttattca ctgaactcta 2880
gtgcggttta ctactgctg caaatactgt atattcagga ctgaaagaa atggtgaatg 2940
cctatgggtg atccaaactg atccagtata agactactga atctgctacc aaaacagtta 3000
atcagttagt cgatgttcta tttttgttt tgtttctctc cctatctgta ttcccaaaaa 3060
ttactttggg gctaatttaa caagaacttt aaattgtgtt ttaattgtaa aaatggcagg 3120
gggtggaatt attactctat acattcaaca gagactgaat agatatgaaa gctgattttt 3180
tttaattacc atgcttcaca atgttaagtt atatggggag caacagcaaa caggtgctaa 3240
tttgttttgg atatagtata agcagtgtct gtgttttgaa agaatagaac acagtttgta 3300
tgccactgt tgttttgggg gggctttttt cttttcgaa atcttaaacc ttaagatact 3360

aaggacgttg ttttggttgt actttggaat tcttagtcac aaaatatatt ttgtttacaa 3420
aaatttctgt aaaacaggtt ataacagtgt ttaaagtctc agtttcttgc ttggggaact 3480
tgtgtcccta atgtgttttag attgctagat tgctaaggag ctgatacttt gacagtgttt 3540
ttagacctgt gttactaaaa aaaagatgaa tgcctgaaa aggggtgttg gaggggtgtt 3600
caacaaagaa acaaagatgt tatggtgttt agatttatgg ttgttaaaaa tgcatctca 3660
agtcaagtca ctggtctgtt tgcatttgat acatttttgt actaactagc attgtaaaat 3720
tatttcatga ttagaaatta cctgtggata tttgtataaa agtgtgaaat aaatttttta 3780
taaaagtgtt cattgtttcg taacacagca ttgtatatgt gaagcaaact ctaaaattat 3840
aaatgacaac ctgaattatc tatttcatca aaaaaaaaaa aaaaaaaaaa aaaa 3894

<210> 192

<211> 695

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (23)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (28)

<223> n equals a,t,g, or c

<400> 192

annctataca ncatacaggg aanggtanac tgacagtacg gtcggattcc cgggtcgacc 60
cacgcgtccg ctgagccatt agaaaatcca tttaaaaaga tgaaaaacaa cattgttgat 120
gctgcaaaca atcacagtgc cccagaagta ctgtatgggt ccttgcttaa ccaggaagag 180
ctgaaattta gcagaaatga tcttgaattt aaatatcctg ctgggtcatgg ctcagccagc 240
graaagtgaac acaggagtgt ggccagagag agcaaaagct ttaatgttct gaaacagctg 300
cttctctcag aaaactgtgt gcgagatttg tccccgcaca gaagtaactc tgtggctgac 360
agtaaaaaga aaggacacaa aaataatgtg accaacagca aacctgaatt tagcatttct 420
tctttaaatg gactgatgta cagttccact cagcccagca gttgcatgga taacaggaca 480
ttttcatacc cagggtgtagt aaaaactcct gtgagtccta ctttcctga gcacttgggc 540
tgtgcagggt ctagaccaga atctgggctt ttgaatgggt gttccatgcc cagtgaagaa 600

ggacccatta agtgggttat cactgatgcg gagaagatga gtatgaaaag cctctccaga 660
ttgaccaaac caccacatac twtattacat gcttc 695

<210> 193

<211> 3131

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (10)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (26)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3128)

<223> n equals a,t,g, or c

<400> 193

gggnncaaan gctggagctc caccgnggtg cgtccgctct agaactagtg gatcccccg 60
gctgcaggat tcggcacgag gccacttcct ggggccgccg gcggggccgc tggctgcact 120
cagcgccgga gccgggagct agcggccgcc gccatgtccc accagaccgg catccaagca 180
agtgaagatg ttaaagagat ctttgccaga gccagaaatg gaaagtacag acttctgaaa 240
atatctattg aaaatgagca acttggtgatt ggatcatata gtcagccttc agattcctgg 300
gataaggatt atgattcctt tgttttacct ctgttgagg acaaacaacc atgctatata 360
ttattcaggt tagattctca gaatgccag ggatatgaat ggatattcat tgcattggtct 420
ccagatcatt ctcatgttcg tcaaaaaatg ttgtatgcag caacaagagc aactctgaag 480
aaggaatttg gaggtggcca cattaagat gaagtatttg gaacagtaaa ggaagatgta 540
tcattacatg gatataaaaa atacttgctg tcacaatctt cccctgcccc actgactgca 600
gctgaggaag aactacgaca gattaaaatc aatgaggtac agactgacgt ggggtgggac 660
actaagcatc aaacactaca aggagtagca tttccattt ctcgagaarc ctttcaggct 720
ttggaaaaat tgaataatag acagctcaac tatgtgcagt tggaaataga tataaaaaat 780
gaaattataa ttttgccaa cacaacaaat acagaactga aagatttgcc aaagaggatt 840
cccaaggatt cagctcgtaa ccatttcttt ctgtataaac attcccatga aggagactat 900
ttagagcca tagtttttat ttattcaatg cctggataca catgcagtat aagagagcgg 960
atgctgtatt ctagctgcaa gagccgtctg ctagaaattg tagaaagaca actacaaatg 1020

gatgtaatta gaaagatcga gatagacaat ggggatgagt tgactgcaga ctctctttat 1080
gaagaagtac atccaagca gcatgcacac aagcaaagtt ttgcaaaacc aaaaggtcct 1140
gcaggaaaaa gaggaattcg aagactaatt agggggccag cggaactga agctactact 1200
gattaaagtc atcacattaa acattgtaat actagttttt taaaagtcca gcttttagta 1260
caggagaact gaaatcattc catgttgata taaagtaggg aaaaaaattg tacttttttg 1320
aaaatagcac ttttcacttc tgtgtgtttt taaaattaat gttatagaag actcatgatt 1380
tctatttttg agttaagct agaaaagggt tcaacataat gtttaatttt gtcacactgt 1440
tttcatagcg ttgattccac acttcaaata ctcttataaa ttttatacag ttgggccagt 1500
tctagaaagt ctgatgtctc aaagggtaaa ctactactt tctgtggga cagaaagacc 1560
ttaaaatatt catattactt aatgaatatg ttaaggacca ggctagagta ttttctaagc 1620
tggaactta gtgtgccttg gaaaaggccg caagttgctt actccgagta gctgtgctag 1680
ctctgtcaga ctgtaggatc atgtctgcaa cttttagaaa tagtgcttta tattgcagca 1740
gtctttttata tttgactttt ttttaatagc attaaaattg cagatcagct cactctgaaa 1800
ctttaagggt accgatattt ttctatactg caggatttct gatgacattg aaagacttta 1860
aacagcctta gtaaatatc tttctaattc tctgtgaggc caaacattta tgttcagatt 1920
gaaattttaaa ttaatatcat tcaaaaggaa acaaaaaatg ttgagtttta aaaatcagga 1980
ttgacttttt tctccaaaac catacattta tgggcaaatt gtgttcttta tcaactccga 2040
gcaaatactc agattttaaa ttactttaaa gtccgtgtac ttaacaggct aacgtagata 2100
aacaccttaa taatctcagt taatactgta tttcaaaaca catttaactg ttttctaattg 2160
ctttgcatta tcagttacaa cctagagaga ttttgagcct catatttctt tgatacttga 2220
aatagaggga gctagaacac ttaattgttta atctgttaaa cctgctgcaa gagccataac 2280
tttgaggcat tttctaaatg aactgtgggg atccaggatt tgtaatttct tgatctaaac 2340
tttatgtctc ataaatcact tatcggaat gcacatttca tagtgtgaag cactcatttc 2400
taaacccttat tatctaaggt aatatatgca cctttcagaa atttgtgttc gagtaagtaa 2460
agcatattag aataattgtg ggtgacaga tttttaaaat agaatttaga gtatttgggg 2520
ttttgtttgt ttacaaataa tcagactata atatttaaac atgcaaaata actgacaata 2580
atgttgctact tgtttactaa agatataagt tgttccatgg gtgtacacgt agacagacac 2640
acatacaccc aaattattgc attaagaatc ctggagcaga ccatagctga agctgttatt 2700
ttcagtcagg aagactacct gtcatgaagg tataaaataa tttagaagtg aatgtttttc 2760
tgtaccatct atgtgcaatt atactctaaa ttccactaca ctacattaaa gtaaatggac 2820
attccagaat atagatgtga ttatagtctt aaactaatta ttattaaacc aatgattgct 2880
gaaaaatcagt gatgcatttg ttatagagta taactcatcg tttacagtat gttttagttg 2940
gcagtatcat acctagatgg tgaataacat attcccagta aatttatata gcagtgaaga 3000
attacatgcc ttctgttgga catttttataa gtgcatttta tatcacaata aaaatttttt 3060
ctcttttaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3120
aaaaaaaaaa a 3131

<210> 194

<211> 2058

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2045)

<223> n equals a,t,g, or c

<400> 194

gccactccgg gtttatttgt ttacaagcgg attacgtcag ctctccctc tcttccctat 60
ctctggaccg gccctctgaa ctcttttccc gcccttttcg gctccgaacc ggcttgcgct 120
acaatgggtgc gatattcgga ttggctggag tcggccatca cgctccacgt acgccacttc 180

```
cttttcgtgg cactataaag ggtgctgcac ggcgcttgca tctcttcgcc tctcggagct 240
ggaaatgcag ctattgagrt cttcgaatgc tgcggagctg gaggcggagg cagctgggga 300
gggccgagcg atgtgaccag gccgccatcg ctcgtctctt cctctctcct gccgcctcct 360
gtctcgaaaa taactttttt agtctaaaga aagaaagaca aaagtagtcg tccgccccctc 420
acgccctctc ttctctcag ccttcgcccc ggtgaggaag cccggggtgg ytgctccgcc 480
gtcggggccg cgccgccgag ccccggccgc cccgggccgc ccccgcacgc cgcccccatg 540
catcccttct acaccgggc cgccaccatg ataggcgaga tcgcgccgc cgtgtccttc 600
atctccaagt ttctccgcac caaggggctc acgagcgagc gacagctgca gaccttcagc 660
cagagcctgc aggagctgct ggcagaacat tataaacatc actggttccc agaaaagcca 720
tgcaagggat cgggttaccc ttgtattcgc atcaaccata aaatggatcc tctgattgga 780
caggcagcac agcggattgg actgagcagt caggagctgt tcaggcttct cccaagttaa 840
ctcacactct ggggtgacct ctatgaagtg tcctacagaa ttggagagga tggctccatc 900
tgtgtgctgt atgaagcctc accagcagga ggtagcactc aaaacagcac caacgtgcaa 960
atggtagaca gccgaatcag ctgtaaggag gaacttctct tgggcagaac gagcccttcc 1020
aaaaactaca atatgatgac tgtatcaggt taagatatag tctgtggatg gatcatctga 1080
tgatgatgga taaatttgat ttttgctttg ggtgggctcc tcttggggat ggattatgga 1140
atttaaacca tgtcacagct gtgaagatct ggcacaagat agaattgtaa aaaaaaaaaa 1200
aaattttaag tgacagtgcc atagtttgga cagtaccttt caatgattaa ttttaatagc 1260
ctgtgagtc aagtaaatga tcactttatt tgctagggag ggaagtccta ggggtggttc 1320
agtttctccc agacatacct aaatttttac atcaatcctt ttaaagaaaa tctgtatttc 1380
aaagaatctt tctctgcagt aaatctcgca ggggaatttg cactattaca cttgaaagtt 1440
gttattgtta accttttcgg cagcttttaa taggaaagtt aaacgtttta aacatggtag 1500
tactggaaat tttaacaagac ttttacctag cacttaaata tgtataaatg tacataaaga 1560
caaactagta agcatgacct ggggaaatgg tcagaccttg tattgtgttt ttggccttga 1620
aagtagcaag tgaccagaat ctgccatggc aacaggcttt aaaaaagacc cttaaaaaga 1680
cactgtctca actgtggtgt tagcaccagc cagctctctg tacatttgct agcttgtagt 1740
tttctaagac tgagtaaaact tcttattttt agaaagtggg ggtctggttt gtaactttcc 1800
ttgtacttaa ttgggtaaaa gtcttttcca caaaccacca tctattttgt gaactttgtt 1860
agtcactctt tatttggtaa attatgaact ggtgtaaatt tgtacagttc atgtatattg 1920
attgtggcaa agttgtacag atttctatat ttggatgag aaatttttct tctctctata 1980
ataaatcgtt tcttatcttg gcattttaaa aaaaaaaaaa aaaaaaaaaa ggcggggcgc 2040
tctanaagat ccaagctt 2058
```

<210> 195

<211> 831

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (28)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (44)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (791)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (793)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (806)

<223> n equals a,t,g, or c

<400> 195

```
anccctcact aagggaaaca aagctggnag ctccaccgcg gtgncgaccg ctctagaact 60
agtggatccc ccgggctgca ggaattcggc agagttcgac ctctgctgca gcccggtgccg 120
ccgccgcctc ctgggaagag aggaagcggg agaggagccc acgtcgccctg tcaccaata 180
tctccagccg cgcagtcctg aagagtgtaa gatgttcgcc tgcgccaagc tcgcctgcac 240
cccctctctg atccgagctg gatccagagt tgcatacaga ccaatttctg catcagtgtt 300
atctcgacca gaggctagta ggactggaga gggctctacg gtattttaatg gggcccagaa 360
tggtgtgtct cagctaattc aaagggagtt tcagaccagt gcaatcagca gagacattga 420
tactgctgcc aaatttattg gtgcaggtgc tgcaacagta ggagtggctg gttctggtgc 480
tggtattgga acagtctttg gcagccttat cattggttat gccagaaacc cttcgctgaa 540
gcagcagctg ttctcatatg ctatcctggg atttgccttg tctgaagcta tgggtctctt 600
ttgtttgatg gttgctttct tgattttgtt tgccatgtaa caaattactg cttgacatgt 660
tggcattcat attaatcag gatgtaattc tgtgtatctt actgtgactc cgaaaactgt 720
agtattggtg tcatgggaat gtacgttatt tccaaagtca tttcattaaa gatgaaaact 780
ttaattcctc ngngattgga ctacanaagt tagattaccc caagagacgg c 831
```

<210> 196

<211> 961

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (32)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (923)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (947)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (953)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (954)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (960)

<223> n equals a,t,g, or c

<400> 196

tagatagaac agatgttttg tgtgaaattt nntatcttta acttaatawaa ccagcaggaa 60
ctgtatgaac acaacacacc caactgacaa acagagagaa ctaacatgtt tatttagctg 120
tatgtatata tgcttaacta cacccgagga agctgtagag ttagaaaaac atgaaccatt 180
aacagatgtg gcctccctgc agaactttta ctttgaaaaa gaagtacgtc tgaaccagat 240
tcacatgttt gatatttgga tgcagagaaa atggggcaga aagcatcgca acagttggct 300
ctgaaggaca gcaaagaggt gcccgctcgtc tgtgargtggt tcagtgaagc tatagtccat 360
gcagctcaga aactgaagga gtaccttgga tttgaatatc ctccaagtaa actctgcccc 420
gctgcaaata ctctgaatga gatcttctta atccatttca tcactttctg ccaagaaaag 480
ggagttgatg agtggctgac caccaccaag atgaccaagc accaagcctt cctgtttggg 540
gcagactgga tttggacctt ttggggatcy gacaagcaaa taaagcttca gctcgcagta 600
cagactctgc agatgtcttc acctcctcct gtggaatcta agccttggtga cctttccaat 660
ccagaatcaa rggtaragga rtcttccttg aagaaaagta gatttgataa gctggaagaa 720
ttctgtaact taataggaga ggattgcctg ggtctgttta tcacttttgg tatgccagga 780
aagcctaaag acatcagggg agttgtcctg gacagtgtca aaagtcagat ggtgaggagc 840
catctgccag gaggaaggc tgtggctcas tttgtcctgg aaactgaaga ttgtgtgttc 900
atcaaagagc tgctcaaaat tgnctgagta agaaagacgg gctgganaga agnnggcaan 960
g 961

<210> 197

<211> 606

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (488)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (500)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (603)
<223> n equals a,t,g, or c

<400> 197
gcttgggtct gtcccgggtg gacgacgccg tggtgcca caccgccag tgygcccaga 60
ggagggacag gagaggcggg gaaggaagag ggcaggggat cgagccttca cccgcctctg 120
ccacccacag gactcgggga gtatgccgca tgccagtcac acgccttcat gaagggcgtt 180
ttcaccttcg tcacaggcac cggcatggcc tttggcttgc agatgttcat tcagaggaag 240
tttccatacc ctttgcagtg gagcctccta gtggccgtgg ttgcaggctc tgtggtcagc 300
tacgggggtga cgagagtgga gtcggagaaa tgcaacaacc tctggctctt cctggagacc 360
gggcagctcc ccaaagacag gagcacagat cagagaagct aggagagctc cagcaggggc 420
acagaggatt gggggcagga ggagtctgga acacagcctt catgccccct gacccacagg 480
cgaccctncc cacaccctan ggtacccag tcgtatcctc tgtccgcatg tgtggccagg 540
cctgacaaaac acctgcagat ggctgctgcc caacctggga ccttgcccag gaggttggag 600
canaaa 606

<210> 198
<211> 393
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (354)
<223> n equals a,t,g, or c

<400> 198
cccacgcgtc cgcccacgcg tcctggacct cctgtgcaag aacatgaaac acctgtggtt 60
cttctctctg ctggtggcag ctcccagatg ggtcctgtcc caggtgcagc tgcaggagtc 120
gggcccagga ctggtgaagc cttcacagac cctgtccctc acctgcactg tctctggtgg 180
ctccatcagc agtgggtgctt actactggag ctggatccgc cagcaccag ggaagggcct 240
ggagtggatt ggntwcactt attacagtgg gascacctac tacaayccgt ccctcaarag 300
tctagtasc atatcastag acacgtctaa gaaccasttc tccctgaasc tganctctgt 360
gackgcccgc gacacggmcg tgtattactg tgc 393

<210> 199
<211> 1061
<212> DNA
<213> Homo sapiens

<220>

<221> misc feature

<222> (1035)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1056)

<223> n equals a,t,g, or c

<400> 199

```
gccccacgct gaccaccggc gtggaggga ggcctctgag actctgcact gagcaccatt 60
ttggacttgt gtgtccagga gtcattgtccc ccgacgagta ccaytcaggg gtcaacaact 120
ctgtgtacac caacgtcctg gtccagaaca ggtagacac aagatcccc caccctaccc 180
cacccccgcc ccagctggag acctgcccc gtgccccac taggcaggca gcagctggaa 240
gtgtaggggt tgcagcctcc cccamctacc tccacctcca gcctgcgctt tgctgctgcc 300
ctggcccagg acctgggtct tcccatcccc agccaktggc tggcgggtggc tgacaagatc 360
aagggtacct ttgacgtkga gcagaacttc cacccgaggt tcgatgggta tgagcctggg 420
gaktggaccc cttcaagggy tcctccccctg ccgtcgagac cctcgagtct gtcctggaac 480
accttccagt cagcggcacc tccctgtagg agaggtgggt aagcaggcag acgtcgtgct 540
cctgggatac cmagtcccct tctccctgag tctgatgtt cgcaggaaaa atctggagat 600
ttacgaggct gtgacgtccc ccagggccc cgccatgacc tggagcatgt ttgctgtggg 660
ctggatggag ctgaaggacg cagtgcgggc ccggggcctc ctggacagga gctttgcca 720
catggctgaa ccttcaagg tgtggacgga gaatgcagac gggtcaggcg ctgtgaactt 780
cctgacaggc atggggggct tctgcaggcg gtggtcttcg ggtgcacggg gttcagggtc 840
acccgagcgg gtgtgacctt tgacctgtg tgtctgtcgg ggatctccag artgagcgt 900
tccggcatct tctaccagg gaacaagctc aacttctctt tttccgagga ctccgtgacc 960
gtggaggtea cagctcgagc agggccctgg gctcctcacc tggaggctga gctgtggcca 1020
ttccagtcce ggmntttcct tgtkcmggam macaangggc t 1061
```

<210> 200

<211> 1359

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1005)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1128)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1342)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1343)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1344)
<223> n equals a,t,g, or c

<400> 200
tcacgcgtcc ggggctgcag taggtcccg caaccgcagg ctgcgcggcg gcgctgggcg 60
cgggatccga ctctagtcgt aatggaggcg ggcggcttct tggactcgct catttacgga 120
gcatgcgtgg tcttcaccct tggcatgttc tccgccggcc tctcggacct caggcacatg 180
cgaatgaccc ggagtgtgga caacgtccag ttccctgccct ttctcaccac ggaagtcaac 240
aacctgggct ggctgagtta tggggctttg aaggagagac ggatccctcat cgtcgtcaac 300
acagtgggtg ctgcgcttca gacctgtat atcttggcat atctgcatta ctgccctcgg 360
aagcgtgttg tgctcctaca gactgcaacc ctgctagggg tccttctcct gggttatggc 420
tacttttggc tcctgggtacc caaccctgag gcccggttc agcagttggg cctcttctgc 480
agtgtcttca ccacagcat gtacctctca ccactggctg acttggctaa ggtgattcaa 540
actaaatcaa cccaatgtct ctccatccca ctaccattg ctacccttct cacctctgcc 600
tcctgggtgcc tctatgggtt tcgactcaga gatccctata tcatgggtgc caactttcca 660
ggaatcgta ccagctttat ccgcttcttg cttttctgga agtaccacca ggagcaagac 720
aggaactact ggctcctgca aacctgagge tgctcatctg accactgggc accttagtgc 780
caacctgaac caaagagacc tccttgtttc agctgggcct gctgtccagc tccccaggtg 840
cagtgggttg tgggaacaag agatgacttt gaggataaaa ggaccaaaga aaaagcttta 900
cttagatgat tgattgggac ctaggagatg aaatcacttt ttatttttta gagatttttt 960
tttttaattt tggagggttg ggtgcaatct ttagaatatg ccttnaaagg ccgggcgcgg 1020
tgctcacgcc tgtaatccca gcactttggg aggccaaggt ggcgggatcg cctgagggtca 1080
ggagtccaag accaacctga ctaacatggt gaaaccccat ctctactnaa aatacaaaat 1140
tagccaaggca tgatggcaca tgctgtaat ccagatact tgggaggctg aggcaggaga 1200
attgcttgaa ccagggaggt ggaggttgca gtgagctgag atcgtgccat tgtgatatga 1260
atatgcctta tatgctgata tgaatatgcc ttaaaataaa gtgttcccca cccctgaaaa 1320
aaaaaaaaa aaaaaaaaaa annnaaaaag ggcggccgc 1359

<210> 201
<211> 726
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (7)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (30)
<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (179)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (724)

<223> n equals a,t,g, or c

<400> 201

```
ttcgaantta cccttcatta aagggaaan aagttggagc ttccaccgcg gtggcgggccg 60
ctctagaact agtggatccc ccgggctgca ggaattcggc acgaggagag aactagtctc 120
gcgcagagcg cagcgmacgg ccacagacag ccctgggcat ccaccgacgg cgcacggang 180
ccagcagagc cggaaggcgc gccccgggca gagaaagccg agcagagctg ggtggcgtct 240
ccggggccgccc gctccgacgg gccagcgccc tccccatgtc cctgctccca cgcgcgcccc 300
ctccggtcag catgaggctc ctggcgggcg cgctgctcct gctgctgctg gcgctgtaca 360
ccgcgcgtgt ggacgggtcc aaatgcaagt gctcccgaa ggaccaaga tccgctacag 420
cgacgtgaag aagctggaaa tgaagccaaa gtaccgcac tgcgaggaga agatggttat 480
catcaccacc aagagcgtgt ccaggtaccg aggtcaggag cactgcctgc accccaagct 540
gcagagcacc aagcgcttca tcaagtggta caacgcctgg aacgagaagc gcagggctta 600
cgaagaatag ggtgaaaaac ctcaagaggg aaaactccaa accagttggg agacttgtgc 660
aaaggacttt gcagattaaa aaaaaaaaaa aaaaaaaaaa aaaagccttt ctttctcaca 720
ggcnta 726
```

<210> 202

<211> 2714

<212> DNA

<213> Homo sapiens

<400> 202

```
gctttgtttg ccatttccta caggtgaaac gccatcatta ggattcactg taacgttagt 60
gctattaaac tctactagcat ttttattaat ggccgttatc tacactaagc tatactgcaa 120
cttggaaaaa gaggacctct cagaaaaactc acaatctagc atgattaagc atgtcgcttg 180
gctaactctc accaattgca tctttttctg ccctgtggcg tttttttcat ttgcaccatt 240
gatcactgca atctctatca gccccgaaat aatgaagtct gttactctga tattttttcc 300
attgcctgct tgccctgaatc cagtccctgta tgttttcttc aacccaaagt ttaaagaaga 360
ctggaagtta ctgaagcgac gtgttaccac gaaaagtggc tcagtttcag tttccatcag 420
tagccaaggt ggttgtcttg aacaggattt ctactacgac tgtggcatgt actcacattt 480
gcagggcaac ctgactgttt gcgactgctg cgaatcgttt cttttaacaa agccagtatc 540
atgcaaacac ttgataaaat cacacagctg tcctgcattg gcagtggctt cttgccaag 600
acctgagggc tactgggtccg actgtggcac acagtssgcc cactctgatt atgcagatga 660
agaagattcc tttgtctcag acagttctga ccaggtgcag gcctgtggac gagcctgctt 720
ctaccagagt agaggattcc ctttgggtcg ctatgcttac aatctaccaa gagttaaaga 780
ctgaactact gtgtgtgtaa ccgtttcccc cgtcaaccaa aatcagtgtt tatagagtga 840
accctattct catctttcat ctgggaagca cttctgtaat cactgcctgg tgtcacttag 900
aagaaggaga ggtggcagtt tatttctcaa accagtcatt ttcaaagaac aggtgcctaa 960
attataaatt ggtgaaaaat gcaatgtcca agcaatgtat gatctgtttg aaacaatat 1020
atgacttgaa aaggatctta ggtgtagtag agcaatataa tgtagttttt ttctgatcca 1080
taagaagcaa atttataacct atttgtgtat taagcacaag ataaagaaca gctgttaata 1140
ttttttaaaa atctatttta aaatgtgatt ttctataact gaagaaaata tcttgctaata 1200
tttaccta atgtttcatcct taatctcagg acaacttact gcagggccaa aaaagggact 1260
```


gtcccgagcta gaactgtgag agtatacata ggcattactt tattatgttt tcacttgcca 1320
tccttgacat aagagaacta taaattttgt ttaagcaatt tataaatcta aaacctgaag 1380
atgtttttaa aacaatatta acagctgtta ggtaaaaaa atagctggac atttgttttc 1440
agtcattata cattgctttg gtccaatcag taattttttc ttaagtgttt tgtgattaca 1500
ctactagaaa aaaagtaaaa ggctaattgc tgtgtgggtt tagtcgattt ggctaaacta 1560
ctaactaatg tgggggttta atagtatctg agggatttgg tggcttcag taatgttctc 1620
attaatgaat acttccta atcgttggct ctactaatat tttccaattt gctgggatgt 1680
cacctagcaa tagcttgat tatatagaaa gttaaactgtg gtcaatactt gcatttaatt 1740
agacgaaacg gggagtaatt atgacacgaa gtacttatgt ttatttctta gtgagctgga 1800
ttatcttgaa cctgtgctat taaatggaaa tttccataca tcttccccat actatttttt 1860
ataaaagagc ctattcaata gctcagaggt tgaactctgg ttaaacaaga taatatgtta 1920
ttaataaaaa tagaagaaga aagaataaag cttagtcctg tgtcttttaa aattaaaaat 1980
tttacttgat tcccatctat gggctttaga cctattactg ggtggagtct taaagtata 2040
attgttcaat atgttttttg aacagtgtgc taaatcaata gcaaaccac tgccatatta 2100
gttattctga atatactaaa aaaatccagc tagattgcag ttaataatt aaactgtaca 2160
tactgtgcat ataatgaatt tttatcttat gttaaattatt tttagaacac aagttgggaa 2220
atgtggcttc tgttcatttc gttaattaa agctacctcc taaactatag tggctgccag 2280
tagcagactg ttaaatgtg gtttatatac ttttgcatt gtaaatagtc tttgtgttac 2340
attgtcagtg taataaaaac agaactcttg tatatcaaaa tcatgtagtt tgtataaaat 2400
gtgggaagga tttatttaca gtgtgttgta attttgtaag gccaaactatt tacaagtttt 2460
aaaaattgct atcatgtata tttacacatc tgataaatat taaatcataa cttggtaaga 2520
aactccta at taaaagggtt tttccaaaat tcaggttatt gaaaayttt cattttatc 2580
atttaaaaac tagaataaca gatataataa agtgtaatac tttgtgctat atggtatgaa 2640
atacaatatt gtactcagtg ttttgaatta ttaaagtttc tagaaagcaa aaaaaaaaaa 2700
aaaaaaaaactc gtag 2714

<210> 203

<211> 422

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (419)

<223> n equals a,t,g, or c

<400> 203

gacggagccg gaagcaggaa gcagcctgk ctccccagga cctgcctggt tgggggaatt 60
ggaggcttct aggaggtggc acggtgcacg ccaagatggc tgtgtccaca gaggagctgg 120
aggccacggt tcaggaagtc ctggggagac tgaagagcca ccagttttcc cagtccacat 180
gggacactgt tgccttcatt gttttctca ccttcattgg caccgtgctg ctctgctgc 240
tgctggctgt cggccactgc tgctgctgca gctccccgg gccccgcagg gaaagcccca 300
ggaaggaaag acccaaggga gtggataact tggccctgga accctgacc tggtgtctct 360
gcccgtggc agtaacaaag ccttctgtct gccagaaaa aaaaaaaaaa aaaaaaang 420
gg 422

<210> 204

<211> 2339

<212> DNA

<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (18)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (19)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (47)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (91)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (125)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (199)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2238)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2321)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2329)
<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2334)

<223> n equals a,t,g, or c

<400> 204

```
anacatggcc aatcaatnnt ctttcagcgt caagaatacg ggacatncac caggaccctg 60
tgaactgacc gctgtgcctg cccttcaagt ncaaggccaa ggattgcact gtgccccatca 120
acacntgcat ccagaacccc tgtcagcatg aggcacctgc cacctgagtg acagccccc 180
ggatgggttc agctgytcnt gccctctggg ctttragggg cagcgggtgtg agatcaaccc 240
agatgactgt gaggacaacg actgcgaaam caatgccacc tgcgtggacg ggatcaacaa 300
ctacgtgtgt atctgtccgc ctaactacac aggtgagcta tgcgacgagg tgattgacca 360
ctgtgtgcct gagctgaacc tctgtcagca tgaggccaa ggcaccccc tggaacaaagg 420
attcagctgc gagtgtgtcc ctggctacag cgggaagctc tgtgagacag acaatgatga 480
ctgtgtggcc cacaagtgc gccacggggc ccagtgcgtg gacacaatca atggctacac 540
atgcacctgc cccagggct tcagtggacc cttctgtgaa cccccccac ccattgtcct 600
actgcagacc agcccatgcg accagtacga gtgccagaac ggggcccagt gcatcgtggt 660
gcagcaggag cccacctgcc gctgcccacc aggcctcgcc ggccccagat gcgagaagct 720
catcactgtc aacttcgtgg gcaaagactc mtacgtggaa ctggcctccg ccaagggtccg 780
accccaggcc aacatctccc tgyaggtggc cctgacaagg acaacggmat ctttctctac 840
aaaggagaca atgacccctt ggcactggag ctgtaccagg gccacgtgcg gctggtctat 900
gacagcctga gttccctccc aaccacagtg tacagtgtgg agacagtga trattgggag 960
tttcacagtg tggagctggt gacgctaacc cagaccctga acctagtagt ggacaaagg 1020
actccaaaga gcctggggaa gctccagaag cagccagcag tgggcatcaa cagccccctc 1080
taccttgag gcatccccc ctcaccggc ctctccgcct tgcgccaggg cagggaccgg 1140
cctctaggcg gcttccacgg atgcattcat gaggtgcgca tcaacaacga gctgcaggac 1200
ttcaaggccc tcccaccaca gtccctgggg gtgtcaccag gctgcaagtc ctgcaccgtg 1260
tgcaagcacg gcctgtgcg ctccgtggag aaggacagc tgggtgtgca gtgcccacca 1320
ggctggaccg gccactctg cgatcaggag gcccgggacc cctgcctcgg ccacagatgc 1380
caccatggaa aatgtgtggc aactgggacc tcatacatgt gcaagtgtgc cgagggttat 1440
ggaggggact tgtgtgacaa caagaatgac tctgccaatg cctgctcagc cttcaagtgt 1500
caccatgggc agtgccacat ctacagacaa ggggagccct actgcctgtg ccagcccgcc 1560
tttagcggcg agcactgcca acaagagaat ccgtgcctgg gacaagtagt ccgagagggtg 1620
atccgcccgc agaaagggtt tgcatcatgt gccacagcct ccaagggtgcc catcatggaa 1680
tgtcgtgggg gctgtggggc ccagtgcctg cagccacccc gcagcaagcg gcggaaatac 1740
gtcttccagt gcacggacgg ctccctggtt gtagaaggag tggagagaca cttagagtgc 1800
ggctgcctcg cgtgttcta agccctgcc cgctgcctg ccacctctcg gactccagct 1860
tgatggagtt gggacagcca tgtgggaccc cctggtgatt cagcatgaag gaaatgaagc 1920
tggagaggaa ggtaaagaag aagagaatat taagtatat gtaaaataaa caaaaaatag 1980
aacttatttt tattatggaa agtgactatt ttcattttt attatataaa tatattacac 2040
catctgcgta tatgtacat atagtgaatt atttttacca agtttttgtg tgtgtatttg 2100
ttgtgttttt aaaaatagct gtttaaaaat ttaagaaaaa aatagactaa taaaaatgct 2160
ttaaacaaca aggataagaa taaagaatga tagcctgtct gaggaaggag gaagatgttt 2220
tcatgtttat gaaacggnat tatgtcggca gtagaccaag agctgctcat gagatcagaa 2280
aaaggaataa gaggaagaga gaggaaggaa aagacctgt ntctgtttnt taantccct 2339
```

<210> 205

<211> 1655

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature
<222> (1548)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1559)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1564)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1623)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1643)
<223> n equals a,t,g, or c

<400> 205
ttcggggcta gcgtcggcga ggcttgagct tgcagcgcgc ggcttccctg ctttctcgcg 60
gccaccccg ctcggcgccg ctcggcgccg gaggggctgg aggtgcggga gccgctctcc 120
gccggtcggg ccccgcgccg ctgagcccag gccgccagcg ccgcgccmc gtgcggtgtc 180
cctgagctcc tgctcccgcg cgggctgctc cgagcaacgg tgcttcggag ctccaaactc 240
gggctgcggg ggcaagtgtc ttcataaacc cagaggatgt ccgggaagca ctacaagggt 300
cctgaagtca gttgttgcat caaatacttc atatttggtc tcaatgtcat attttggttt 360
ttgggaataa catttcttgg aattggactg tgggcatgga atgaaaaagg agttctgtcc 420
aacatctctt ccatcaccca tctcgcgccg tttgacccag tttggctctt ccttgggtg 480
ggaggagtga tgttcatttt gggatttgca ggggtgcattg gagcgctacg ggaaaacact 540
ttccttctca agtttttttc tgtgttcctg ggaattattt tcttctgga gctcactgcc 600
ggagttctag catttgtttt caaagactgg atcaaagacc agctgtattt ctttataaac 660
aacaacatca gagcatatcg ggatgacatt gatttgcaaa acctcataga cttcacccag 720
gaatattggc agtgctgtgg ggcttttggg gctgatgatt ggaacctaaa tatttacttc 780
aattgcacag attccaatgc aagtcgagag cgatgtggcg ttccattctc ctgctgact 840
aaagatcccg cagaagatgt catcaacact cagtgtggct atgatgccag gcaaaaacca 900
gaagttgacc agcagattgt aatctacacg aaaggctgtg tgccccagtt tgagaagtgg 960
ttgcaggaca atttaaccat cgttgctggt attttcatag gcattgcatt gctgcagata 1020
tttgggatat gcctggccca gaatttggtt agcgatatcg aagctgtcag ggcgagctgg 1080
tagacccctt gcaaccgctg ctgcaagaca ctggacagac ccagctttcg ggaccctccc 1140
gcgtgccgaa ctgactctcg agctgcatgg acctaatcac agatgcagcc tgcagtctcg 1200
cctaattggg ctgcccattg gggagtgtaa aactgggaaa tgctgctcac tgacagaatt 1260
aaaaaaaaaa ataaccagta tgaaagtcgt tgcgccgtga atctctactg tagccatgaa 1320
tttatggaca gttagatgct taccataaaa gaaaaaaagg gagggtaggg gacccagatg 1380
tacttgaatg tgcagaaaat acattcttgt cctcatcttc cgtaattgga gggctgggag 1440
argcagcttt gctcttcacc acaccttgga cggaccacct tctttctgtt ccatggcctg 1500
aaggagtggc atctcctcaa aggactcagc ccctcacctg ggaggggancg agtttctgnc 1560

cccncttaga tttttaatcc agcttggtgg tccacccgtt tgttctgccc cccctgggg 1620
agncaagatt atcaagcctt gcncgcactt cttgt 1655

<210> 206

<211> 5145

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (5126)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (5143)

<223> n equals a,t,g, or c

<400> 206

gtcngattcg cgcccgatc ggaggtttcg gctaagttgg aggtactggc cacgactgca 60
tgccccgcgc cgccaggtga tacctccgcc ggtgaccag gggctctgcg acacaaggag 120
tctgcatgtc taagtgtctag acatgctcag ctttgtggat acgcggaactt tgttgctgct 180
tgcagtaacc ttatgcctag caacatgccca atctttacaa gaggaactg taagaaagg 240
cccagccgga gatagaggac cacgtggaga aaggggtcca ccaggccccc caggcagaga 300
tgggtgaagat ggtcccacag gccctcctgg tccacctggt cctcctggcc cccctggtct 360
cgggtgggaac ttgtctgctc agtatgatgg aaaaggagtt ggacttggcc ctggaccaat 420
gggcttaatg ggacctagag gcccacctgg tgcagctgga gcccagggc ctcaagggtt 480
ccaaggacct gctggtgagc ctgggtgaacc tgggtcaaact ggtcctgcag gtgctcgtgg 540
tccagctggc cctcctggca aggtctggtga agatgggtcac cctggaaaac ccggacgacc 600
tgggtgagaga ggagtgttg gaccacaggg tgctcgtggt ttccctggaa ctctggact 660
tcctggcttc aaaggcatta ggggacacaa tggctctggat ggattgaagg gacagcccg 720
tgctcctggt gtgaagggtg aacctggtgc ccctggtgaa aatggaactc cagggtcaaac 780
aggagcccgt gggcttcctg gtgagagagg acgtgttggg gccctggcc cagctggtgc 840
ccgtggcagt gatggaagtg tgggtcccgt gggctcctgct ggtccattg ggtctgctgg 900
ccctccaggc ttcccagggtg cccctggccc caagggtgaa attggarctg ttggtaacgc 960
tggctcctgct ggtcccgcgc gtccccgtgg tgaagtgggt cttccaggcc tctccggccc 1020
cgttggaact cctggtaatc ctggagcaaa cggccttact ggtgcaaagg gtgctgctgg 1080
ccttcccgcc gttgctgggg ctcccgccct ccctggacc cgcggtattc ctggccctgt 1140
tgggtgctgcc ggtgctactg gtgccagagg acttggtggg gagcctgggc cagctggctc 1200
caaaggagag agcggtaaca aggggtgagc cggctcctgct gggcccaag gtcctcctgg 1260
tcccagtggt gaagaaggaa agagaggccc taatggggaa gctggatctg ccggccctcc 1320

aggacctcct gggctgagag gtagtcctgg ttctcgtggt cttcctggag ctgatggcag 1380
agctggcgctc atggggccctc ctggtagtcg tgggtgcaagt ggccctgctg gagtccgagg 1440
acctaattgga gatgctggtc gccctgggga gccctggtctc atgggacca gaggtcttcc 1500
tgggtccccct ggaaatatcg gccccgctgg aaaagaaggt cctgtcggcc tccctggcat 1560
cgacggcagg cctggcccaa ttggcccmgc tggagcaaga ggagagcctg gcaacattgg 1620
attccctgga cccaaaggcc ccactggtga tcctggcaaa aacggtgata aaggctatgc 1680
tggctcttgc ggtgctcggg gtgctccagg tcctgatgga aacaatggtg ctcagggacc 1740
tcctggacca cagggtgttc aagggtgaaa aggtgaacag ggtcccgtg gtccctcagg 1800
cttcagggt ctgcctggcc cctcagggtcc cgctggtgaa gttggcaaac caggagaaaag 1860
gggtctccat ggtgagtttg gtctccctgg tcctgctggt ccaagagggg aacgcgggtcc 1920
cccaggtag agtggtgctg ccggtccctac tggctctatt ggaagccgag gtccttctg 1980
acccccaggg cctgatggaa acaagggtga acctggtgtg gttggtgctg tgggcactgc 2040
tggctccatct ggtcctagt gactcccagg agagaggggt gctgctggca tacctggagg 2100
caaggagaaa aagggtgaac ctggtctcag aggtgaaatt ggtaaccctg gcagagatgg 2160
tgctcgtggt gtcctggtg ctgtagggtgc ccctggtcct gctggagcca caggtgaccg 2220
gggcgaactg gggctgctgg tcctgctggt cctgctggtc ctgggggaa cctggtgaa 2280
cgtggtgagg tcggtcctgc tggccccaat ggatttgcgt gtcctgctgg tgctgctggt 2340
caacctggtg ctaaaggaga aagaggagcc aaaggcccta agggtgaaaa cgggtgtgtt 2400
ggtcccacag gcccggttg agctgctggc ccagctggtc caaatggtcc ccccggtcct 2460
gctggaagtc gtggtgatgg aggccccct ggtatgactg gtttccctgg tgctgctgga 2520
cggactggtc cccaggacc ctctggtatt tctggccctc ctgggtcccc tggctcctgt 2580
gggaaagaag ggcttcgtgg tcctcgtggt gaccaaggtc cagttggccg aactggaga 2640
gtagggtcag ttgggtcccc tggctcgtc ggtgagaagg gtccctctgg agaggctggt 2700
actgctggac ctgctggac tccaggtcct cagggtcttc ttggtgctcc tggattctg 2760
gggtctcctg gctcgagagg tgaacgtggt ctaccaggtg ttgctggtgc tgtgggtgaa 2820
cctggtcctc ttggcattgc cggccctcct ggggcccgtg gtcctcctgg tgctgtgggt 2880
agtctgtag tcaacggtgc tcctggtgaa gctggtcgtg atggcaacc tgggaacgat 2940
ggtccccag gtcgcatgg tcaaccgga cacaaggag agcgcggtta ccctggcaat 3000
attggtccc ttggtgctgc aggtgcacct ggtcctcatg gcccggtgg tcctgctggc 3060
aaacatggaa accgtgtgga aactggtcct tctggtcctg ttggtcctgc tgggtgctgtt 3120
ggcccaagag gtccctagtgg ccacaaggc attcgtggcg ataagggaga gcccggtgaa 3180
aaggggccca gaggtcttcc tggcttaaag ggacacaatg gattgcaagg tctgcctggt 3240
atcgctggtc accatggtga tcaagggtgct cctggtccg tgggtcctgc tggctcctagg 3300
ggccctgctg gtccctctg ccctgctgga aaagatggtc gactggaca tcctggtaca 3360
gttggacctg ctgctgcatg aggcctcag ggtcaccag gccctgctgg cccctggt 3420
ccccctggcc ctccctggacc tccaggtgta agcggtggtg gttatgactt tggttacgat 3480
ggagacttct acagggtga ccagcctgc tcagcacctt ctctcagacc caaggactat 3540
gaagttagt ctactctgaa gtctctcaac aaccagattg agacccttct tactcctgaa 3600
ggctctagaa agaaccagc tcgcacatgc cgtgacttga gactcagcca cccagagtgg 3660
agcagtgggt actactgat tgaccctaac caaggatgca ctatggatgc tatcaaagta 3720
tactgtgatt tctctactgg cgaaacctgt atccggggcc aacctgaaaa catcccagcc 3780
aagaactggt ataggagctc caaggacaag aaacacgtct ggctaggaga aactatcaat 3840
gctggcagcc agtttgaata taatgtagaa ggagtgaact ccaaggaaat ggctacccaa 3900
cttgccttca tgcgctgct ggccaactat gcctctcaga acatcaccta cactgcaag 3960
aacagcattg catacatgga tgaggagact ggcaacctga aaaaggctgt cattctacag 4020
ggctctaatt atgttgaaact tgttgcgtgag ggcaacagca ggttcactta cactgtctt 4080
gtagatggct gctctaaaaa gacaaatgaa tggggaaaaga caatcattga atacaaaaca 4140
aataagccat cagcctgcc cttccttgat attgcacctt tggacatcg tgggtgctgac 4200
cagggaattct ttgtggacat tggcccagtc tgtttcaaat aaatgaactc aatctaaatt 4260
aaaaaagaaa gaaatttgaa aaaactttct ctttgccatt tcttcttctt cttttttaac 4320
tgaaagctga atccttccat ttcttctgca catctacttg cttaaattgt gggcaaaaaga 4380

gaaaaagaag gattgatcag agcattgtgc aatacagttt cattaactcc ttccccgct 4440
cccccaaaaa tttgaatttt tttttcaaca ctcttacacc tggtatggaa aatgtcaacc 4500
tttgtaagaa aaccaaaata aaaattgaaa aataaaaaacc ataaacattt gcaccacttg 4560
tggtctttga atatcttcca cagaggggaag tttaaaaccc aaacttccaa aggttttaac 4620
tacctcaaaa cactttccca tgagtgtgat ccacattgtt aggtgctgac ctagacagag 4680
atgaactgag gtccttgttt tgttttgttc ataatacaaa ggtgctaatt aatagtattt 4740
cagatacttg aagaatgttg atggtgctag aagaatttga gaagaaatac tcctgtattg 4800
agttgtatcg tgtggtgtat tttttaaaaa atttgattta gcattcatat tttccatctt 4860
attcccaatt aaaagtatgc agattatttg cccaaatctt cttcagattc agcatttggt 4920
ctttgccagt ctcatcttca tcttcttcca tggttccaca gaagctttgt ttcttgggca 4980
agcagaaaaa ttaaattgta cctattttgt atatgtgaga tgtttaataa aattgtgaaa 5040
aaaatgaaat aaagcatgtt tggttttcca aaagaaaaaa aaaaaaaaaa aacycgrggg 5100
ggggcccgw acccaattgg ccccaanggg ggccgtttac atnca 5145

<210> 207

<211> 487

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (470)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (483)

<223> n equals a,t,g, or c

<400> 207

tccgccctcc cgctcgctccc ctgctcgggg tcagctagtg tckctctgc tcggccgcgg 60
gtccccggag gactgcaggc aggatgacgc aaaacacggg gattgtgaat ggagttgcta 120
tggcctctag gccatcccag cccacccacg tcaacgtcca catccaccag gagtcagctt 180
tgacacaact gctgaaagct ggaggttctc tgaagaagtt tctttttcac cctggggaca 240
ctgtgccttc cacagccagg attggttatg agcagctggc tctaggggtg actcagatat 300
tgctgggggt tgtgagttgt gttcttgag tgtgtctcag cttggggccc tggactgtgc 360
tgagtgcctc agctgtgcct tctgggcggg gtctgtggtg atcgcarcar gastkgggcc 420
attgtccatg araarcaccc gggcaaaatk gytggctata ttscagccgn tcaccgcgca 480
agntttg 487

<210> 208

<211> 2296

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (202)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (252)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2258)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2285)
<223> n equals a,t,g, or c

<400> 208
gaagaattag gcttggttcc aggttacata aataatagag cagatgggtg cctttggggtc 60
ttgaaccaaataaagactta ctaaaggaag agtgaggttt tagaagagaa ctttacgagc 120
gtgacattat gcattactcc acatccaccc tttaagaatt gatttattac aaaattatgt 180
tctattttaa agtccactcc cncaaaatac tgatttcttc catggattat ttttatatga 240
tttgtaaaaa tnttttctga aatttaaatt gacaagcaat gttataaaaa tattgctaata 300
cttctgctta tttttcagcc tttaaagtct gaaatactat aaaacactga gagtatctat 360
gtctttctga atttaaaatg aaaattcatg tccatatgga gacttacata tcccccaatc 420
tttccatgaa ttgcattttt ttctaatact tatctctgtt aagactccat tgaaagttaa 480
acaaaaagcc gatataatag ttggggtttt ttttttttgc ttatttttagt attttgaatc 540
agagaatact tggccaagct cgtatttgta aactaaaacc ccttgttatt tgtatcttct 600
gcatttccagt ttactgtaa gaacatacat yttgaaatta tgaacatttt tttcataaaa 660
attaatgctt ttaagaaatg gagaaggaaa aatagctttt atcaattttt ttctcctcaa 720
gattctgtaa taaacctcaa agtctggtgt agaatggtat tttcccgttt gaggtcttgg 780
tagttttttt ttttttttta aacatggtgc cacttttaat taactaaatg ttacaaaatt 840
ctcagcattt gtttttgtct ggtagtatta ttgagtcttt tcatacactt cttaaaatgt 900
gaaaatctat gttttcagta ttatgacagt atctgcagat actataataa tatgctttgg 960
ccaaatttgc tktttataga gtgaaccaag atgtcagttt tgtattgggg atatgtttta 1020
cataaataga actcatttaa aagtatggtt caagtactgc tcttcttttt atgggggaaat 1080
tgatttttaa ggagcgattc ttttcctata taaaagctat tgaggcttag ctttattcat 1140
acatgagtga ttgggttata actcaatcaa aatttttatc tttaaactctg agacaagtat 1200
tatattttaa actagatcta ttaatacatt catcactgcc tgaaaatttg ctgttctaata 1260
gwtgtggaat ttctactcat tacctagagt aaacttattg catattttca taatcagtca 1320
aaataataaa actaaaaaca aaaagtaaaa attagttccc aagtaaatat atcagtgttt 1380
cctgtgtttg ggcagaagtg caaagtagtt ttaatttctc acgttgtaga aaaagaggcc 1440
aaaaaggtca ccttagttta tttttccaaa acataacttt aggtattcta atgtattttg 1500
ttagttgctt agaaatggaa ataaaatata tgtgcttatt aaataatgca cagtacacta 1560
gattatagag tttatttttg cttttatgct ttaaagaatg catgccccaa acactactac 1620
caacaagagg attttttaaa atcacagaat tgatatttat ctcaagacta tcttaagcta 1680
atgtacttct gttacataat taaactattt ctgctattat ttttaaagca gtattatttt 1740
ataaccaatt ctaaaattca gctttaaaaa tattgracat accttcctac cacttacgct 1800
tcaagtttca ttattgaaat accagggtca tatcttagaa ttgtatataa aacaacatgt 1860
aaccatttca tgaaaaacgt ttttaagtat tgctttttgc tgctatgttc tgcatgtgac 1920
ctagttttgc cattgtcccc tgaatgtagt attatctata gattatacat aactacatct 1980
attgcttttg tgggtaaatc aagatttagt tgtaattttc cagctgtgaa aatgttgctc 2040
tgtattttaa agggtttcat gaatggaac ctaagtaaaa ctaagctcat tagtgacaga 2100
cttgttttct tcttggtatt cctccagcaa ctccctcacc accacgcctc cctgcctacc 2160

atccccggaa ggggtgcttat tctttaacaa agagaatcta aaaaaaaaaac aaaaaactgt 2220
tgtctatatt cacgtaatgt cctggcttta ttccctgngt ttagagagca ttcctggttg 2280
tacantgact ttgatg 2296

<210> 209

<211> 625

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (10)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (25)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (26)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (600)

<223> n equals a,t,g, or c

<400> 209

ntctcattcn acacccaacc ttcannttca actacaactt ctccgctttg ggcaaacacc 60
gtcactcttg ctggaggcaa gcttcattcc aaagggttga aatacttcca tcactttacc 120
ctcagtctct gtggaaacca gggtaggaaa atgtctgtgt gcaccgacaa tgtcactgac 180
ctccggattc ctgagggtga gtcagggttc tccaaatcta tcacagccta cgtctgccag 240
gcagtcacat tccccccaga ggtgacaggc tacaaggccg gggtttcctc acagcctgtc 300
agccttgctg atcgacttat tggggtgaca acagatatga ctctggatgg aatcacctcc 360
ccagctgaac ttttcacct ggagtccttg ggaataccgg acgtgatctt cttttatagg 420
tccaatgatg tgaccagtc ctgcagttct gggagatcaa ccaccatccg cgtcagggtgc 480
agtccacaga aaactgtccc tggaagttag ctgctgccag gaacgtgctc agatgggagc 540
tgtgatggct gcaacttcca cttcctgtgg gaaaamsgs tgsttgccc cttctgctcan 600
tggtgacta ccatgctatc gtcac 625

<210> 210

<211> 1551

<212> DNA

<213> Homo sapiens

<220>
<221> misc feature
<222> (760)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1543)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1544)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1545)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1546)
<223> n equals a,t,g, or c

<400> 210
ttaaggctgt ccgctctggg ctccgcggcc tcggcccgt gcactccacc tccgccccct 60
cggaactccct cccctctgct tctactctc ctgctccagt gcggatcggt tcgcaactgc 120
ttgccactcg tcccgctgct ggctgtttty ccatttccc gccccctctt cttgagtact 180
ttacccctcg catttgggga cagggaactgg aaaaggggag ggtggagcgt ccagtggaga 240
agaaggaagc gagggcccga gagggaggag atcggcgagc tgtggggagg agaccccacg 300
ccaccctttc tggcatctc cctcccgc ccgcccctgc gcacactccc tcgcgggcga 360
gctactttcg gaccaggaag gtaagagcgg ccctgggtga cagcgccgcg gggccagtc 420
cggggttagc cgcgcgtctg ctgcgttctg gtccgtcgcg ctcccagcca gggcacagcg 480
gaccgaggat ggcttcgacc acaacctgca ccagggtcac ggacgagtat cagcttttcg 540
aggagcttgg aaagggggca ttctcagtgg tgagaagatg tatgaaaatt cctactggac 600
aagaatatgc tgccaaaatt atcaacacca aaaagctttc tgctagggat catcagaaac 660
tagaaaagaga agctagaatc tgccgtcttt tgaagcacc taatattgtg cgacttcatg 720
atagcatatc agaagagggc ttctactact ttggtgggtt gattaagtta ctggagggtga 780
actgtttgaa gacatagtgg caagagaata ctacagtga gctgatgcca gtcattgcat 840
tcagcagatc ctggaggctg tgctacactg ccatcagatg ggcgtgggtc atcgggacct 900
gaagcctgag aatttgcttt tagctagcaa atccaaggga gcagctgtga aattggcaga 960
ctttggctta gccatagaag ttcaagggga ccagcaggcg tggtttggtt ttgctggcac 1020
acctggatat ctttctycar aagttttacg taaagatcct tatggaaagc cagtggatat 1080
gtgggcatgt ggtgtcatc tctatattct acttggtggg tatccaccct tctgggatga 1140
agaccaacac agactctatc agcagatcaa ggctggagct tatgattttc catcaccaga 1200
atgggacacg gtgactcctg aagccaaaaga cctcatcaat aaaatgctta ctatcaacc 1260
tgccaaacgc atcacagcct cagaggcact gaagcaccca tggatctgtc aacgttctac 1320
tgttgcttcc atgatgcaca gacaggagac tgtagactgc ttgaagaaat ttaatgctag 1380
aagaaaacta aagggtgcca tcttgacaac tatgctggct acaaggaatt tctcagcagc 1440

caagagtttg ttgaagaaac cagatggagt aaaggagtca actgagagtt caaatacaac 1500
aattgaggat gaattcagct tggacttaac caggttgact cannnngggg g 1551

<210> 211

<211> 1011

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (309)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (801)

<223> n equals a,t,g, or c

<400> 211

tgcacccacg cgtccgggag aatggcggcg gaaggctgga tttggcgttg gggctggggc 60
cgggcgtgcc tgggaaggcc tgggcttctc ggccccggcc ctggccccac tacacctctc 120
tttcttcttt tggtgttggg gtctgtgact gcggatataa ctgacggcaa cagtgaacat 180
ctcaagcggg agcattcget cattaagccc tacciaagggg tcggttccag ctctatgccc 240
ctctgggact tccagggcag cactatgctc acgagccagt acgtacgtct gacctctgac 300
gagcgcagna aagagggctc tatctggaac caccagccgt gcttcctcaa agactgggaa 360
atgcacgtcc acttcaaagt ccacggcaca gggaagaaga acctccatgg agacggcatc 420
gccttgtggg acacccggga ccgcctcgtg ccagggcctg tgtttggaag caaagataac 480
ttccacggct tagccatctt cctggacacc taccccaatg atgagaccac tgagcgcgtg 540
ttcccgctaca tctcggtgat ggtgaacaat ggctccctgt cctacgacca cagcaaggat 600
gggcgctgga ccgagctggc gggctgcacg gctgacttcc gcaaccgcga tcacgacacc 660
ttcctggctg tgcgctactc ccggggccgt ctgacggtga tgaccgacct ggaggacaag 720
aacgagtgga agaactgcat tgacatcacg ggagtgcgcc tgcccaccgg ctactacttc 780
ggggcctccg ccggcacccg nracctgtct gacaatcatg acatcatctc catgaaagct 840
gttccaagct gatggtggaa gcacacgccc gacgaggaga agcatcgact tggaccaaga 900
tcgagcccag cgtcaacttc tcaagtcgcc caaagacaac gtggacgacc ccacggggaa 960
ctttcgcagc gggcccctga mggggtggcg ggtgtttcct gctgctgctg t 1011

<210> 212

<211> 1639

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1630)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1637)

<223> n equals a,t,g, or c

<400> 212

```
ggcacgagga cgcgccgagg tactaggcag agccgtggaa ccgccgccag gtcgctgttg 60
gtccacgccc cccgtcgcgc cgcgccccc ctcagcgtcc gccgccgcca tgggagtgca 120
ggtggaaacc atctcccag gagacgggcg caccttcccc aagcgcggcc agacctgcgt 180
ggtgcactac accgggatgc ttgaagatgg aaagaaattt gattcctccc gggacagaaa 240
caagcccttt aagtttatgc taggcaagca ggaggtgatc cgaggctggg aagaaggggt 300
tgcccagatg agtgtgggtc agagagccaa actgactata tctccagatt atgcctatgg 360
tgccactggg caccaggca tcatccacc acatgccact ctcgtcttcg atgtggagct 420
tctaaaactg gaatgacagg aatggcctcc tcccttagct ccctgttctt ggatctgccc 480
tggaggatgc tgggtgcctcc agacatgtgc acatgartcc atatggagct tttcctgatg 540
ttccactcca ctttgtatag acatctgccc tgactgaatg tgttctgtca ctcagctttg 600
cttcgcagac cctgttttcc tcttcccctt tctcctcgta tgtgtgttta cctaaactat 660
atgccataaa cctcaagtta ytcattttat tttgttttca ttttggggtg aagattcagt 720
ttcagtcctt tggatatagg tttccaatta agtacatggt caagtattaa cagcacaagt 780
ggtagggtta cattagaata ggaattggtg ttgggggggg gtttgcaaga atattttatt 840
ttaatttttt ggatgaaatt tttatctatt atataataaa cattcttgct gctgcgctgc 900
aaagccatag cagatttgag gcgctgttga ggactgaatt actctccaag ttgagagatg 960
tctttgggtt aaattaaaag ccctacctaa aactgagggt gggatgggga gaggcctttg 1020
ctccaccatt cccaccacc ctcctcttaa accctctgcc tttgaaagta gatcatgttc 1080
actgcaatgc tggacactac aggtatctgt ccctgggcca gcagggacct ctgaagcctt 1140
ctttgtggcc tttttttttt ttcacctctg ggtttttcta atggactttc aggaattttg 1200
taatctcata actttccaag ctccaccact tcctaaatct taagaacttt aattgacagt 1260
ttcaattgaa ggtgctgttt gtagacttaa caccagtgta aagccagcc atcatgacaa 1320
atccttgaat gttctcttaa gaaaatgatg ctggtcatcg cagcttcagc atctcctgtt 1380
ttttgatgct tggctccctc tgctgatctc agtttcctgg cttttcctcc ctcagccctt 1440
tctcaccctt ttgctgtcct gtgtagtgtt ttggtgagaa atcgttgetg cacccttccc 1500
ccagcaccat ttatgagtct caagttttat tattgcaata aaagtgttt atgccggctt 1560
ttctcaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1620
aaaaaaaaan ggggggncc 1639
```

<210> 213

<211> 2127

<212> DNA

<213> Homo sapiens

<400> 213

```
ttgaggttct gggggtcctg gagacttacc attgagccat gcaatctggg aagcacagga 60
ataagtagac actttgaaaa tggatttgaa tgttctcatc ctttttgca cttttctttt 120
tggtctctct atgtccttgg cttgtcctc tattctacct ctctttctcc agcaataata 180
tgcaaatgaa gacatgtatc cataagaagg agtgccttc atcaactaat agagcaccta 240
ccacagtgtc atacctggtg gaggtgagca attcatattc aaaggttgca aagtgtttgt 300
aatatattca tgaggctgga akkaagaaga attaaaaatt tgtcctaatt acaatgagaa 360
ccattctagg tagtgatctt ggagcacaca tgaataactt tctgaagggt caaccaaatc 420
catttttatt tctgcctggc ttggtcacct ctgtaaagggt ttaacttagt gttgtcaagt 480
aacagttact gaaagagctg agaaaaagaa caatgaacag caacgatctt gactgtgcaa 540
ctcagacatt cctgcagaaa agacatatgt tgctttacaa gaaggccaaa gaactatggg 600
gccttcccag catttgactg ttcattgcat agaatgaatt aaatatccag ttacttgaat 660
gggtataacg catgaatatt tgtgtgtctg tgtgtgtgtc tgagtgtgtt gattttatta 720
ggggcatctg ccaattctct cactgtgggt ccttctctga ctttgctgtt tcatcatcta 780
aggaggctag atccttcgct gacttcacca ttctcaaac ctgtaagttt ctcacttctt 840
```

ccaaattggc tttggctctt tcttcaacct ttccattcaa gagcaatctt tgctaaggag 900
taagtgaatg tgaagagtac caactacaac aattctacag ataattagtg gattgtgttg 960
tttggtgaga gtgaagggtt cttggcatct ggtgcctgat taaggcttga gtattaagtt 1020
ctcagcatat ctctctattg tcttgacttg agtttgctgc attttctatg tgctgttcgt 1080
gacttgagga acttaaagta atcgagctat gccaaacttg ggtggtaaca gagtacttcc 1140
caccacagtg ttgaaaggga gagcaaagtc ttatggataa accctccttt cttttgggga 1200
cacatggctc tcaattgaga agctcacctg tgctgaatgt ccacatggtc actaaacatg 1260
ttatccttaa accccccgta tgcctgagtt gaaagggctc tctcttatta ggttttcatg 1320
ggaacatgag gcagcaaatc tattgctaag actttaccag gctcaaatca tctgaggctg 1380
atagatattt gacttggtaa gacttaagta aggctctggc tcccaggggc ataascaaca 1440
gtttcttgaa tgtgccatct garaaggag acccagggtta tgagttttcc ttgaaacaca 1500
ttggctcttt ctcaaagtcc ctgccttgct agactgttag ctctttgagg acagggacta 1560
tgtcttatca atcactatta tttcctgtt acctagcatg ggacaagtac acaacacata 1620
tttgttcaat gaatgaatga atgtcttcta aaagactcct ctgattggga gaccatatct 1680
ataattggga tgtgaatcat ttcttcagt gaataagagc acaacggcac aaccttcaag 1740
gacatattat ctactatgaa ctttttactg tgagactctt tttttgcct tctacttgcg 1800
ctgaaatgaa accaaaacag gccgttggtg tccacaagtc aatatatggt ggatgaggat 1860
tctgttgctt tattgggaac tgtgagactt atctggtatg agaagccagt aataaacctt 1920
tgacctgttt taaccaatga agattatgaa tatgttaata tgatgtaaat tgctatttaa 1980
gtgtaaagca gttctaagtt ttagtatttg ggggattggt ttttattatt tttttccttt 2040
ttgaaaaata ctgagggatc ttttgataaa gttagtaatg catgttagat tttagttttg 2100
cagagcatgt tgtttaacaa ataaagg 2127

<210> 214

<211> 1166

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1163)

<223> n equals a,t,g, or c

<400> 214

gcggacgcgt gggcgacgc gtggcgacgc gcgtggcggt gtcgcttggt ggctccgtct 60
gtctgtccgt ccgcccgcgt gtgccatcat ggcggacgcg gccagtcagg tgctcctggg 120
ctccggtctc accatcctgt ccagccgct catgtacgtg aaagtgtca tccagggtgg 180
atatgagcct ctctctcaa caataggacg aaatatcttt gggcggaag tgtgtcagct 240
tcttggtctc tttagttatg ctcagcacat tgccagtatc gatgggaggc gcgggttggt 300
cacaggctta actccaagac tgtgttcggg agtccttggg actgtggtcc atggtaaagt 360
tttacagcat taccaggaga gtgacaaggg tgaggagtta ggacctggaa atgtacagaa 420
agaagtctca tcttctttg accacgttat caaggagaca actcgagaga tgatcgctcg 480
ttctgtgct accctcatca cacatccctt ccatgtgatc actctgagat ctatggtaca 540
gttcattggc agagaatcca agtactgtgg actttgtgat tccataataa ccatctatcg 600
ggaagagggc attctaggat ttttcgctgg tcttggtcct cgccttctag gtgacatcct 660
tctttgttg ctgtgtaact cactggccta cctcgtaaat acctatgcac tggacagtgg 720
ggtttctacc atgaatgaaa tgaagagtta ttctcaagct gtcacaggat tttttgcgag 780
tatgttgacc taccctttg tgcttgctc caatcttatg gctgtcaaca actgtggtct 840
tgctggtgga tgccctcctt actccccaat atatacgtct tggatagact gttggtgcat 900
gtacaaaaa gaggggaata tgagccgagg aaatagctta tttttccgga aggtccctt 960
tgggaaagact tattgttgtg acctgaaaat gttaatttga agatgtgggg caggacagct 1020

gacattttctg tagtcccaga tgcacagaat tatgggagag aatgttgatt tctatacagt 1080
gtggcgcgct tttttaataa tcatttaatc ttgggaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aacggggggg gcccgaacc canttc 1166

<210> 215

<211> 3323

<212> DNA

<213> Homo sapiens

<400> 215

ggggcaaaaa ggccgagggg gcccagaatc agggcaaaaa ggccgagggg gcccagaacc 60
agggcaagaa ggcmgagggg gcccagaacc agggcargaa ggccgagggg gcccagaacc 120
aggrcaagaa ggccgagggg rcccmkaacc agggcargaa ggccgagggg gcccagaacc 180
agggcargaa ggccgagggg gcccagaacc agggcaagaa ggccgagggg gcccagaayc 240
agggcaagaa ggccgagggg gcccagaayc agggcaagaa ggccgagggg gcccagaayc 300
agggcaagaa ggccgagggg gcccagaacc agggcaagaa ggccgagggg gctcagaacc 360
agggcaaaaa agtakaaggg gcccagaacc agggcaagaa ggctgagggg gcccagaacc 420
agggcaagaa ggccgagggg gctcagaacc agggcaaaaa ggccgagggg gcccagaacc 480
agggcaaaaa aggagagggg gcccagaatc agggtaaaaa gacagaaggg gctcagggca 540
aaaaggcaga aaggagtccc aaccaaggca aaaaggaga gggagctccc atccagggca 600
aaaaggcaga ttcggttgct aatcagggca caaaggtaga gggattaca aaccagggga 660
aaaaggcaga aggggtcccc agtgaaggca aaaaggcaga aggggtcccc aaccaaggca 720
aaaaggcaga cgcagctgcc aatcagggta aaaagacaga gtcagcttct gtcagggca 780
gaaatacaga tgtggcccag agcccagagg caccaaagca agaggctcct gccaagaaga 840
agtctggttc aaagaaaaaa ggtgagcctg ggcccccaga tgccgacggc cctctctacc 900
tcccctacaa gacgctggtc tccacggttg ggagcatggt gttcaacgag ggcgaggccc 960
agcggctcat cgagatcctg tctgagaagg ctggcatcat tcaggacacc tggcacaagg 1020
ccactcagaa gggtgaccct gtggcgattc tgaacgcca gytggaagag aaggaaaaac 1080
tgctggccac agaacaggaa gatgcggtg tgcaccaagag caaaytgagg gagtcaaca 1140
aggagatggc agcagaaaaag gccaaagcag cagccgggga ggccaaagtg aaaaagcagc 1200
tggtggcccc ggagcaggag atcacggctg tgacggcacg catgcaggcc agctaccggg 1260
agcacgtgaa ggaggtgcag cagctgcagg gcaagatccg gactcttcag gagcagctgg 1320
agaatggccc caacacgcag ctggcccgcc tgacgcaggga gaactccatc ctgcccggatg 1380
ccttgaacca ggccacgagc caggtggaga gcaagcagaa cgacagctg gccaaagctc 1440
ggcaggagct cagcaaggct agcaaaagc tgggtggagaa gtcagaggct gtgcggcaag 1500
atgagcagca gcgaaagct ctggaagcca aggcagctgc cttcgagaag caggtcctgc 1560
agctgcaggc gtcccacagg gagagtgagg aggcctgca gaagcgctg gacgaggta 1620
gccgggagct gtgccacacg cagagcagcc acgccagcct ccggggcgat gccgagaagg 1680
cccaggagca acagcagcag atggccgagc tgcacagcaa gttacagtcc tccgaggcgg 1740
agggtgcgag caaatgcgag gagctgagtg gcctccacgg gcagctccag gaggccaggg 1800
cggagaactc ccagctcaca gagagaatcc gttccattga ggccctgctg gaggcggggc 1860
agggcgggga tgcccaggac gtccaggcca gccagscgga ggctgaccag cagcagactc 1920
gcctcaagga gctggagtcc caggtgtcgg gtctggagaa ggaggccatc gagctcaggg 1980
agggcgctga gcagcagaaa gtgaagaaca atgacctccg ggagaagaac tggaaaggcca 2040
tgagggcact ggccacggcc gagcagcct gcaaggagaa gctgcactcc ctgacccagg 2100
ccaaggagga atcggagaag cagctctgtc tgattgaggc gcagaccatg gaggcctgtc 2160
tggtctgtct cccagaactc tctgtcttgg cacaacagaa ttacaccgag tggctgcagg 2220
atctcaaaga gaaaggcccc acgctgctga agcaccggcc agctcccgcg gagccctcct 2280
cggacctggc ctccaagttg agggaggccg aggagacgca gagcactcag caggccgagt 2340
gtgaccagta ccgacgcatc ctggcgagga cggagggcag gctcagagac ctgcagaaga 2400
gcgtggagga ggaggagcag gtgtggaggg ccaaggtggg cggcgagag gaggagctcc 2460

agaagtcccg ggtcacagtg aagcatctcg aagagattgt agagaagcta aaaggagaac 2520
ttgaaagtgc ggaccagggtg agggagcaca cgtygcattt ggaggcagag ctggaaaagc 2580
acatggcggc cgccagcgcc gagtgccaga actacgcca ggaggtggca gggctgaggc 2640
aacttctcct agaattctcaa tctcagctcg atgccgcca gagcgaagcc agaaacagag 2700
cgatgagctt gccctgggtca ggcagcagtt gagtgaatg aagagccacg tagaggatgg 2760
tgacatagct ggggccccag cttcctcccc agaggcgccc ccagccgagc aggaccccg 2820
tcagctgaag acgcagctgg agtggacaga agccatcctg gaggatgagc agacacagcg 2880
gcagaagctc acggccgagt ttgaggaggc tcagacctcg gcatgtcggg tacaagaaga 2940
attggagaag ctccgcacag ccggccccct agagtcttca gaaacagagg aggcctcaca 3000
gctgaaggag agactagaaa aagagaagaa gttaacaagt gacctggggc gcgccgccac 3060
gagactgcag gagcttctga agacgaccca ggagcagctg gcaagggaga aggacacggg 3120
gaagaagctg caggaacagc tggaaaaggc agaggacggc agcagctcaa aggagggcac 3180
ctctgtctga gtttctctt tggaaaaaga agttactgtt caacttacca aaatgcctta 3240
cacattcctt acaaatataac caaccaacct acaaaaaaaaa aaaaaaaaaa aactcgaggg 3300
ggggcccggt acccaattcg ccc 3323

<210> 216

<211> 1408

<212> DNA

<213> Homo sapiens

<400> 216

taaactattt atcttgtgtg tgtacatttg tgggtggagt ttgtgcgcct ggtttttttg 60
tttgaaaaac actgcgtggt caatgtggtt atggggggga gtgatgcatt tttttctagt 120
cttaaaacta aaaacttgag tctaccattt ctgtgttgca ctgaaaatac cgcccagcct 180
gatggtgttc ccgtgctgtc cctccccctt cccttctccc cgctctacc tccccacccc 240
gttctgttcc cctccctcc ttctccctct ccctcaaatc cgtgagtttt ggaagcccca 300
gggcctctct cccccgccc tcttgatga ggccaccatc ccccaaaccg gcttgttttg 360
cagtttcccc aggatcctgg aagctcgtg gcgctcgagg gtggcgggga cacggggggg 420
tgggtgaagg ttctgtacct tttctagtgc gttctatcat agttaacggg tgcacacttt 480
tttaaaaaaa gtaaatggat ttgccacaat taaatgtcat aacatttatg acagaatata 540
aaatattaac atattttaag ccaagtttta ggtgtatttt ttgaatcttg gttataaacc 600
caatttttaa ggagcagatga tccagcgttg tgaaggcaac agagtgtacc catatttata 660
tttttataaa atacctataa gactgtgaat ctcttgtgct aatggctgag ttaattgaag 720
gatcgttttg ccctttttta gcctcccaga gcttcgagga ctcaattcga acccgaaatc 780
ctgccgtggg ggagggggtg cgtcgagacc tgggcccggg gaggttctcc tgcgtcactt 840
tctgtcctga aaggcgccct tcctgggttc tgtggctcca attttctatg cagccccaca 900
ccccttgttg ttttgatcct gagaataaaa agggaggctg aattattcaa atttaaatga 960
ggtttccctc tcatggaagt gctgctgacc cttcgtgcag aaatggggag cacttgagga 1020
cacagtgagg tggaggccct ttgtgcgtgg ctggctcgat tcgggcagcc ctccgtcgct 1080
ttttataaaa ctttgtgtga gaagaatata ttgataatgt cagtgaaca agcagacatt 1140
gaaatggagg cacagattac tccacaagga gttcttctgt atattttttc tagatgcaa 1200
taccttttta attatgttaa ttaatgttaa gactttctag gcttatatcg aagctgtgtg 1260
tgggtcacgg ggtgatcact gctaactgga taaagtttgt gcagcacatt cctgagtgta 1320
cgatattgac ctgtagccca gcgtgaaaaa tttataataa aatttttcat tgatcttttt 1380
atattaaaaa aaaaaaaaaa aaaaaaaaaa 1408

<210> 217

<211> 2111

<212> DNA

<213> Homo sapiens

<220>
<221> misc feature
<222> (2102)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2104)
<223> n equals a,t,g, or c

<400> 217
ggcacgagga gcattccata tagaaactgc tgaaactgcc acaggtgctt ctccgaaaac 60
cttacagttg tggcattgaa tggtcagtat cgcttccttt ctgcacacga ggcatactgt 120
tgagcctgag aaatgccagt ggcctgacag cagcagacat tgcacaaacc cagggtttcc 180
aagagtgtgc ccagtttctc ttgaacctcc agaattgtca tctgaaccat ttctataaca 240
atggcatctt aaatgggggt catcagaatg tatttcctaa tcatattagt gtgggaacaa 300
atcgaaaagag atgcttgagg agactcagaa gactttggag taaagaaagc tagaactgaa 360
gctcaaagct tggattctgc cgtgccactc acgaatggcg acacagaaga cgatgctgac 420
aaaatgcacg ttgataggga gtttgcgtgt gtaacagata tgaaaaacag tagctccgta 480
tcgaatacat tgacaatgga tgtgtcatca atggacattt ggacttcccc tccacgaccc 540
cgctcagtgg gatggaaaagc aggaatggcc agtgcttgac aggaactaac ggaattagca 600
gtggattagc cccaggacag ccgtttccga gttagccaggg ttctctctgc attagtggga 660
ctgaggagcc agagaagacc ctgagagcta accctgagtt gtgcggttct ctgcacctga 720
acgggagtc aagtagctgc atagccagta ggccttcctg ggtggaagac attgggggata 780
acctgtacta tggacactac cacgggtttg gggacactgc tgaaagcatg ccagaaactg 840
aacagtgtgg tcgagcattc caagtccgtg aaggtgcagg agcgggtacga cagtgccgtg 900
ctgggcacca tgcacctgca ccacggctcc tagagacgct gacctggctc tcggaaaagc 960
aggagtcctt cctggttagc agctcagaat acccatgtag cagcaacttg aacgaatgtc 1020
acaacttgta ctttttttat atacttcaac tttctgaaaa agtaaaacttc gacaagtcc 1080
cagcaactgc ttgtttgtgc atgagtaggg ctactaagt gcatagatgt ttctacagt 1140
agggtgcctt ttataaggt gcacttttg agttttctg atgccaatct caacattgtc 1200
ttttaatac tgtcaccaga tattgccatt ttctttttg ttaaaagatt atatgatcaa 1260
gataaattgg ggtggtaaat cagggtgcctg gtaatttatc tctttgcaca tgggcatcat 1320
tttaaaaagc ttgcttcac tcttttctgt agaatttgac ggaacacagc tatttcccta 1380
tgcaaggtag agccttacia agatttctgc agtgatttgt gtgaagaaga gaatgtttgt 1440
ctttttcaat gaagctttgc agatcccat gtggttgaag gtttttagttg tggacacagt 1500
ggtccctcct taatgatgaa gatcactgcc ttgggcttca tggaaaacag gccagacctg 1560
gggctgcgtt tggatttatt gtttttattc cacacttcct acttgggtctc tggaaagttt 1620
accacatgta acagattcct ttatatgtag tgaaaaatcac tatttgtaga aactgtcagg 1680
tcaaaaatatt taactgactg ttgacatgta ttttctttt tccttgtttt tgttttttg 1740
ggttttctgc ttaagatat ataccactat gtatatccag ttaactgaga gaattttgac 1800
tctcttaata aaactgcatt aagtttttga tttttagtaa attagctttt gtctaggcaa 1860
ctagtgggta tactctgcaa atattgtaat gaatttttac ttttttgatt tttgtaataa 1920
aaattggtgc agataaaatg tcaaatgaac aaaccagtgt tctaagagt ttactaacat 1980
tttgttctaa aactgtcctt cacaattga ataaaaaact ctacactca aaaaaaaaaa 2040
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2100
ananaaaaaa a 2111

<210> 218

<211> 2493

<212> DNA

<213> Homo sapiens

<400> 218

```
ggcccggtgg ggccggccgg gactcgccgc tcgcacgccc ttgggcccgg gccgggccc 60
cgctcttctt tccgcttgcg ctgtgagctg aggcgggtgta tgtgcggcaa taacatgtca 120
accccgctgc ccgccatcgt gcccgccgcc cggaggcca ccgctgcggg gatcttcctg 180
catggattgg gagatactgg gcacggatgg gcagaagcct ttgcaggat cagaagtcca 240
catatcaaat atatctgccc gcatgcgcct gttaggcctg ttacattaaa tatgaacgtg 300
gctatgcctt catggtttga tattattggg ctttcaccag attcacagga ggatgaatct 360
gggattaaac aggcagcaga aaatataaaa gctttgattg atcaagaagt gaagaatggc 420
attccttcta acagaattat ttggggaggg tttctcagg gaggagcttt atctttatat 480
actgccctta ccacacagca gaaactggca ggtgtcactg cactcagttg ctggcttcca 540
cttcgggctt cctttccaca gggctctatc ggtggtgcta atagagatat ttctattctc 600
cagtgccacg gggattgtga ccctttggtt cccctgatgt ttggttctct tacggtggaa 660
aaactaaaaa cattggtgaa tccagccaat gtgaccttta aaacctatga aggtatgatg 720
cacagtctgt gtcaacagga aatgatggat gtcaagcaat tcattgataa actcctacct 780
ccaattgatt gacgtcacta agaggccttg tgtagaagta caccagcatc attgtagtag 840
agtgtaaacc ttttcccatg cccagctctc aaatttctaa tgttttgtag tgttaaaatg 900
ttttgcaaat acatgccaat aacacagatc aaataatatc tcctcatgag aaatttatga 960
tcttttaagt ttctatacat gtattcttat aagacgaccc aggatctact atattagaat 1020
agatgaagca ggtagcttct ttttctcaa atgtaattca gcaaaataat acagtactgc 1080
caccagattt ttattacat catttgaaaa tagcagtat gcttaatgaa aatttggtca 1140
ggtataaatg agcagttaag atataaacia tttatgcatt ctgtgactta gtctatggat 1200
ttattccaaa attgcttagt caccatgcag tgtctgtatt ttatatatg tgttcatata 1260
tacataatga ttataatata taataagaat gaggtggtat tacattatc ctaataatag 1320
ggataatgct gtttattgtc aagaaaaagt aaaatcgctt tcttcaatta atggcccttt 1380
tattttggga ccaggctttt attttccctg atattatttc tatttaatac tcttttctct 1440
caagaaaaaa aaaaaagttt gtttttctt tattgtcctt catagcaggc caagtattgc 1500
ctctctgcaa tagacagcta ctgtcaatc atgtgtaat ttgacattct gggtcacaga 1560
tataaggat ttaaaatcta ttatgcttt atagagaaac cagacattaa aacttcatgc 1620
actactatt tcgaattact gtacctatc caaatttaca cctagctatt aggatcttca 1680
accaggttaa caggaataat tctgtggtt catttttctg taaacaactg aaagaataat 1740
tagatcatat tctagtatgt tctgaaatat ctttaagact gatcttaaaa actaacttct 1800
aagatgattt catcttctca tagtatagag tttactttgt acacgtttga aaccaactac 1860
tgtagaagat gaggaatcta ttgtaatttt ttgctttatt ttcattctgc agtggaacta 1920
tttgaaattt tcaacttagt caaattattt ttgtatttag tttttgatgc agacataaaa 1980
atagcaatca ttttaaattg tcaaaatttc cagattactg gtaaaaatta tttgaaaaca 2040
aacttatggg taataaaggc tagtcagaac cctataccat aaagtgtagt taccatacag 2100
attaatatgt agcaaaaatg tatgcttgat atttctcaac tgtgttaatt tttctgctgt 2160
attccagctg accaaaacaa tattaagaat gcatctttat aaatgggtgc taattgataa 2220
tggaataaat ttagtaatgg actatacagg atgttaataa tgaagccata tgtttatgtc 2280
tggttttaaa aattttaaac aatcatttac tatgtcattt ttctttacct tgaagaacat 2340
aaactgttat ttcacttcta caaatcagca agatattatt tatggcaaga aatattccat 2400
tgaaatattg tgctgtaaca tgggaaagtg taaatgtttt tcatggtttc tatcaatgtg 2460
aaataaaatt taattctgaa aaaaaaaaaa aaa 2493
```

<210> 219

<211> 1259

<212> DNA

<213> Homo sapiens

<400> 219

gcgcgcgggt gccggaaccg ctgggcggga gcgagcggt gcggctgcag ctgcagggcg 60
aggagctgcg gctgcaggag gagagcgtgc ggctgcacca gattaacatc tacctcagcg 120
accgcatctc actgcaccgc cgcctgcccg wgcgctggaa cccgctgtgc aaagagaaga 180
aatatgatta tgataatttg cccaggacat ctgttatcat agcattttat aatgaagcct 240
ggtcaactct ccttcggaca gtttacagtg tccttgagac atccccggat atcctgctag 300
aagaagtgat cctttagat gactacagtg atagagagca cctgaaggag cgcttgcca 360
atgagcttcc gggactgccc aagggtgcgc tgatccgcgc caacaagaga gagggcctgg 420
tgcgagcccg gctgctgggg gcgtctgcgg cgargggcga tgttctgacc ttcctggact 480
gtcactgtga gtgccacgaa ggggtgctgga gccgctgctg cagaggatcc atgaagagga 540
gtcggcagtgt gtgtgcccgt tgattgatgt satcgactgg aacaccttcg aatacctggg 600
gaactccggg gagccccaga tcggcgggtt cgactggagg ctgggtgtca cgtggcacac 660
agttcctgag agggagagga tacggatgca atccccgctc gatgtcatca ggtctccaac 720
aatggctggt gggctgtttg ctgtgagtaa gaaatatattt gaatatctgg ggtcttatga 780
tacaggaatg gaagtttggt gaggagaaaa cctcgaattt tccttttagga tctggcagtgt 840
tggtggggtt ctggaaacac acccatgttc ccatgttggt catgttttcc ccaagcaagc 900
tccctactcc cgcaacaagg ctctggccaa cagtgttcgt gcagctgaag tatggatgga 960
tgaatttaaa gagctctact accatcgcaa cccccgtgcc cgcttggaac cttttgggga 1020
tgtgacagag aggaagcagc tccgggacaa gctccagtgt aaagacttca agtggttcct 1080
ggagactgtg tatccagaac tgcagtgtcc tgaggacagg cctggcttct tcgggatgct 1140
ccagaacaaa ggactaacag actactgctt tgactwtaac cctcccgatg aaaaccagat 1200
tgtgggacac caggtcatte tgtacctctg tcattgggatg ggccagaatg acctggtgc 1259

<210> 220

<211> 1849

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (920)

<223> n equals a,t,g, or c

<400> 220

ctgttgatg gagcaggggt tgtgggtttt ctgggcccat cattatggct gcttcagagt 60
cagaagaaag ccatagggca gtaggggagc tcctattgcc tagcccctct ccctttgtgg 120
ctcccactct agctgcctat ttttgctcat cagctggtga gtcagtatgg gccagcagtt 180
ctccctccct aagcccttgc tactttatgg gttagctttg caggtttggg ggcttgaggg 240
gtggggggcaa ctcaccactg ccaggtaact ccctgaaggg tgggagtggg ttatcttcta 300
ggctcttacc cgcggtaggg aagggcacat acactgtctt ccttccattc tcctttcccc 360
catcccattt agtgctgcca cagggcagaa gcacacaaac caaccacaca gtctctgact 420
tctcctaagc actttgagtt gttgaatggg gctcaggggc aagagttttt gctgccctcc 480
ccagcgtggt cacagggtta ttgaactgcc tgcacttgtt tctcatgcaa ctccagcatt 540
ttccccagaa gttgaactat ggatagcagc ttgggtatgga ttctctaaat cttaacattt 600
gaagcagctt cttgaggctg gcaactatcc tgggttctgt cttggagggg gtgggttgtt 660
tgctggggcc caacgtctgt cccaagtggg ggggtgagag taagttaact ttgggtgccg 720
gtgagagggt ggggctcttt gcttagactc cctatcatgg aaagattgga gttttctatg 780
cagggcactg ggaaaagga ttgctgattc tgactgacct tgatcagaga gattaggatt 840
gtattttgac ataggatttg gaacccatct aaatgttgaa gttccctgag acagctctcc 900
agctgctgag cctgcgccan gggctaagca gccctaatg agaggctctg ctccctttcc 960

```
cacctcgcca atgttggtgt tgctgccttt ttgatttgta tcctctgtta tagacatttt 1020
ttaaaaacga tttcctcttt cattgtgcac aagtgcctgag agtctgaggc cccatttctg 1080
ctgtgtatat atacctgac tcggggcctt tattcagcaa actgttcatt cttctgtcag 1140
acaatgtcat attcaactct gttcatatta aaccactgtg aagcaagcct ctgttttcct 1200
gcttaagttg taaatttagt attcttttagt gtctaggata tgctgggtat tatgcagaaa 1260
tcatacagtg tggccagtgt cctgaggtaa tgttttgcat ttaaattttt ttagaaagca 1320
gaatcttaac ttatcttaat gatatttacc tacccttttt gcaactcaca actgactttg 1380
tcacagaggt aatgcatctg cttgcaggaa gtarctgtag gctcagtacc tgttgkttga 1440
gtcagattta gcagatttgg tttttaagct tgtgggtttg kgctaatttg ggcagaatat 1500
atattattata tatgtgtgtg tgtatgtktg tatgtgtgtg tctgcatatg taatacatgt 1560
acataaacac acatgcatgt gttcatcctc tgacacaccc acacaacacc aacaaacatt 1620
tcttctatag gctttttatc tcaactgaca ctgttttttt tcccaaataa atttgacaca 1680
ggcagaaagg tgggtgaact ctcagaactt ttggtgggtg gatattcatc tgaccagtga 1740
gctctgaaat gggttcccta cacagagtgg gttttggcaa ggggttgaat gaggggaggt 1800
agcagtcttg tcatttagaa aatcaaagct agttttgatg tagctcaac 1849
```

<210> 221

<211> 1267

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (26)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1244)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1252)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1264)

<223> n equals a,t,g, or c

<400> 221

```
cggattatgc ngamacnccc cagggntttrt gctatgacgt cgcattgcacg cgtaagcttg 60
ggccccctga gggatcctct agagcggccg ccgcgccatt cggggaatct gcagggcaga 120
tgagtaacga aagaggcttt gaaaatgtag aactgggagt cataggaaaa aagaagaaaag 180
tcccaaggag agtcatccac tttgttagtg gtgaaacaat ggaagaatat agcacagatg 240
aagacgaagt tgatggcctg gagaagaaag atgttttgcc tactgttgat ccgacaaaac 300
ttacctgggg tccctactta tggttttaca tgcttcgggc tgctacatca actctctcag 360
tgtgtgactt ccttgagag aagattgcat ctgttttggg tatcagcacc ccaaagtacc 420
aatatgccat tgatgaatat tatcgatga agaaggagga agaagaagaa gaagaagaaa 480
acaggatgtc tgaagaagca gaaaaacaat atcaacagaa taaattgcag actgattcca 540
ttgttcagac agatcaacca gagacagtga tatccagctc atttgtgaat gtcaattttg 600
aaatggaggg agacagtga gtaattatgg aaagcaagca aaatccagtc tctgtcccac 660
cataaaatga aatgactatc aagcttcaaa ctcttaagtt tttttttttt aatacaaaaa 720
ctttcacatt ctttattcag tgggacttaa tacaattatt tatattttta attattaaag 780
tatctggaaa gggaaaatgt tttcttcatt tttaggatct atctagcaaa agccagatct 840
gaaattcaga tatttgtact gtttttactg tgtatagaaa ttagtgcttt ggttttaaaa 900
tgatctttta aaaaagttaa ggacatccta gagccttaat agttaagaag agttaaatta 960
tcaagcctat ttgtgcattt gctttttttg aaaaaggtaa gttgctgatt aagtctaatt 1020
ggaattgata attccatagt cttagattaa aatgaggata ttttctccta gattttctca 1080
tgttatgcca tgcatttata tatctaacca ttaatttcac actaaggatg cttcaccata 1140
taataaaaag agcaagatgg aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaggggag gacgctggg tccgaccgg gaattcccg atcngtcact gncgggctga 1260
cttntct 1267
```

<210> 222

<211> 754

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (702)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (710)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (754)

<223> n equals a,t,g, or c

<400> 222

```
cagagaactg agatgaaagc aatgatgaag ggaccctggg accctgcctt tcctctccct 60
actcagcagg agcttatgaa gtcattctac ttctctttct gtcctccctc tcctcttcat 120
gcaggatata agaaatatga ggcccccaatt ggaacagcag catggagggt cccagttggg 180
tccccaacgt gctcccatg ggcaagacaa tggaaacttc cacaagcagg gaaggcaaac 240
cctctttatt gaacattagc cagcccgacc cagacccag ggctgctaag gacacagaga 300
ttctccatgg gaaggggact gccaaagcatg aggaaataga agattcaggg gcctgarctc 360
```

tggaagctgc aagcaaaagg gatgggacta gggctgagtt gtgtctycat ttgataagg 420
aaaggatatg ctcacactct tgcttggtca gattccaaga cagaaggctt cacaaggcca 480
acgcctggaa aatgggcatc tctccctccc atgttaagct ttaacctctg taatctgcct 540
gtatctatag gtgggcatct cactccatca aaggagccca gcctcttttg tcctctacca 600
tgcacagtct ttctgtgcat ttccccaagc tgggccctct tctactctca tttaggcctg 660
tgataactca ttaccacccc atcacttggtg tcttcagggc anacttggcn aagcagggaa 720
tgcccttga cataaattga agggcactcc tttt 754

<210> 223

<211> 1258

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (41)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1205)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1241)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1247)

<223> n equals a,t,g, or c

<400> 223

aacgatacag ccaaggmgat caccatcaac aagattctkt nccaacgwga gccttggtgca 60
ggagaacctg tacttccagc gctgcctaga ctggaaccgt gacatcctca agaaggagct 120
gggactgaca gagcaggaca tcattgacct gcccgctctg ttcaagatgg acgaggacca 180
ccgtgccaga gccttcttcc caaacatggt gaacatgac gtgctggaca aggacctggg 240
catccccaag ccattcgggc cacagggtga ggaggaatgc tgcctggaga tgcacgtgcg 300
tggcctcctg gagccccctg gcctcgaatg caccttcac gacgacattt ctgcctacca 360
caaatttctg ggggaagtcc actgtggcac caacgtccgc aggaagccct tcaccttcaa 420
gtggtggcac atggtgccct gacctccag gggccctggc gtttgccctc ttmgcttart 480
tctccagacc ctgcctcaca ygccagagc cttctgctga catggactgg acagccccgc 540
tgggagacct ttgggacgtg gggtggaatt tggggtatct gtgccttgcc ctccctgaga 600
ggggcctcag tgwcctctga agccatcccc agtgagcctc gactctgtcc ctgctgaaaa 660
tagctgggcc agtgtctctg tagccctgac ataaggaaca gaacacaaca aaacacagca 720
aacctgtgc ccaactgct ccccaaaaaa ttttgagtct ctaatctgac actgaatgag 780
gggagaaggg aaggagattc tgggattgcc agttcttcca gcagccatgc tctgaaaatc 840
aaggtagaat ccatggaaaag ggaccccagg accccgggac cctagacgta tcttgaactg 900
ccatcgctcat ttcaaataca tctccctcag ggtttccagg tggccacccc caatwattca 960
ttccttacca acctctcaaa tcctcttggc tttctctctg cagtgtggac actgttggct 1020

agtcctcccc actccctgag ggtccagtaa gttaccttag aaccttcctg ggaaacattt 1080
catctggagg caggkttccc cacgtgtggg atgctccttt ttgcctcatc tgtcttcagg 1140
ggatggcagg ctcccccgca ttgcatgggg gattttttcc ccagaccagg cataatttgt 1200
ggacntgagg agtttcaatg tgttaaagat ggccctgggt naagccntat tccatctt 1258

<210> 224

<211> 584

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (494)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (495)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (577)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (583)

<223> n equals a,t,g, or c

<400> 224

cttaccacaga aggcaacgct tctctttctg gtcaaaatgg ctggtaagca ggccgtttca 60
gcatcaggca agtggctgga tggatttcga aaatgggtatt acaatgctgc aggattcaat 120
aaactggggg taatcgcgaga tgatacaata tacgaggatg aagatgtaaa agaagccata 180
agaagacttc ctgagaacct ttataatgac aggatgtttc gcattaagag ggcaactggac 240
ctgaacttga agcatcagat cttgcctaaa gaggcagtga ccaaatatga agaggaaaat 300
ttctaccttg aaccgtatct gaaagagggt attcgggaaa gaaaagaaa agagaatgg 360
gcaaagaagt aatcatgtag ttgaagtctg tggatgcagc tgttatgaag atgggttaaac 420
ttgaaacaaa caattttaag aattatttgg tctgaagatg ttttacttta aataaatgtc 480
tattgtaatg gcwnnaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 540
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaggggnggc cgnt 584

<210> 225

<211> 3449

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2330)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3434)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3443)

<223> n equals a,t,g, or c

<400> 225

gcgagcacgt tagaaagcct gacatggcct gactcgggac agctcagagc agggcagaac 60
tggggacact ctggggccggc cttctgcctg catggagcct ctgaagccac cctgtctctg 120
gaggaaccac gagcgaggga agaaggacag ggactcgtgt ggacaggaaga actcagagcc 180
gggaagcccc cattcactag aagcactgag agatgcggcc ccctcgcagg gtctgaattt 240
cctgtgctg ttcacaaaga tgctttttat ctttaacttt ttgttttccc cacttccgac 300
cccggcggtg atctgcatcc tgacatttgg agctgccatc ttcttgtggc tgatcaccag 360
acctcaaccc gtcttacctc ttcttgacct gaacaatcag tctgtgggaa ttgagggagg 420
agcacggaag ggggtttccc agaagaacaa tgacctaacca agttgctgct tctcagatgc 480
caagactatg tatgaggttt tccaaagagg actcgcctgtg tctgacaatg ggccctgctt 540
gggatataga aaaccaaacc agccctacag atggctatct tacaacacagg tgtctgatag 600
agcagagtac ctgggttcct gtctcttgca taaaggttat aaatcatcac cagaccagtt 660
tgtcggcatc tttgctcaga ataggccaga gtggatcatc tccgaattgg cttgttacac 720
gtactctatg ttagctgtac ctctgtatga caccctggga ccagaagcca tctgacatat 780
tgtcaacaag gctgatatcg ccatggtgat ctgtgacaca ccccaaaagg cattggtgct 840
gatagggaa ttagagaaag gcttcacccc gagcctgaag gtgatcatcc ttatggaccc 900
ctttgatgat gacctgaagc aaagagggga gaagagtga attgagatct tatccctata 960
tgatgctgag aacctaggca aagagcactt cagaaaacct gtgcctccta gcccagaaga 1020
cctgagcgtc atctgcttca ccagtgggac cacaggtgac cccaaaggag ccatgataac 1080
ccatcaaaat attgtttcaa atgctgctgc ctttctcaaa tgtgtggagc atgcttatga 1140
gccactcct gatgatgtgg ccatatccta cctccctctg gctcatatgt ttgagaggat 1200
tgtacaggct gttgtgtaca gctgtggagc cagagttgga ttcttccaag gggatattcg 1260
gttgctggct gacgacatga agactttgaa gccacattg tttcccgcgg tgcctcgact 1320
ccttaacagg atctacgata aggtacaaaa tgaggccaag acaccctga agaagttctt 1380
gttgaagctg gctgtttcca gtaaattcaa agagcttcaa aagggtatca tcaggcatga 1440
tagtttctgg gacaagctca tctttgcaaa gatccaggac agcctgggag gaagggttcg 1500
tgtaatgtc actggagctg ccccatgtc cacttcagtc atgacattct tccgggcagc 1560
aatgggatgt caggtgtatg aagcttatgg tcaaacagaa tgcacaggtg gctgtacatt 1620
tacattacct ggggactgga catcagggtca cgttgggggtg cccctggctt gcaattacct 1680
gaagctggaa gatgtggctg acatgaacta ctttacagtg aataatgaag gagaggtctg 1740
catcaagggt acaaacgtgt tcaaaaggata cctgaaggac cctgagaaga cacaggaagc 1800
cctggacagt gatggctggc ttcacacagg agacattggt cgctggctcc cgaatggaac 1860
tctgaagatc atcgaccgt aaaaagaacat tttcaagctg gcccaaggag aatacattgc 1920
accagagaag atagaaaata tctacaacag gagtcaacca gtgttacaaa tttttgtaca 1980
cggggagagc ttacggtcat ccttagtagg agtgggtggt cctgacacag atgtacttcc 2040
ctcatttgca gccaagcttg ggggtgaagg ctcctttgag gaactgtgcc aaaaccaagt 2100
tgtaaggga gccattttag aagacttgca gaaaattggg aaagaaagtg gccttaaac 2160
ttttgaacag gtcaaaagcca tttttcttca tccagagcca ttttccattg aaaatgggct 2220
cttgacacca acattgaaag caaagcgagg agagctttcc aaatactttc ggacccaaat 2280
tgacagcctg tatgagcaca tccaggatta ggataaggta cttaagtacn tgccggccca 2340

ctgtgcactg cttgtgagaa aatggattaa aaactattct tacatttggt ttgcctttcc 2400
tcctattttt ttttaacctg ttaaaactcta aagccatagc ttttgtttta tattgagaca 2460
tataatgtgt aaacttagtt cccaaataaa tcaatcctgt ctttcccatc ttcgatgttg 2520
ctaataattaa ggcttcaggg ctacttttat caacatgcct gtcttcaaga tcccagttta 2580
tgttctgtgt ccttctcat gatttccaac cttaatacta ttagtaacca caagttcaag 2640
ggcacaaggg accctctgtg ccttcttctt tgtttgtga taaacataac ttgccaacag 2700
tctctatgct tatttacatc ttctactgtt caaactaaga gattttttaa ttctgaaaaa 2760
ctgcttacia ttcattgttt ctagccactc cacaaccac taaaatttta gttttagcct 2820
atcactcatg tcaatcatat ctatgagaca aatgtctcgc atgctcttct gcgtaaatta 2880
aattgtgtac tgaagggaag agtttgatca taccaaacat ttcctaaact ctctagttag 2940
atatctgact tgggagtatt aaaaattggg tctatgacat actgtccaaa aggaatgctg 3000
ttcttaaagc attatttaca gtaggaactg gggagtaaat ctgttcccta cagtttgctg 3060
ctgagctgga agctgtggg gaaggagtgt acaggtgggc ccagtgaact tttccagtaa 3120
atgaagcaag cactgaataa aaacctcctg aactgggaac aaagatctac aggcaagcaa 3180
gatgccaca caacaggctt attttctgtg aaggaaacca ctgatctccc ccaccctgg 3240
attagagttc ctgctctacc ttaccacag ataacacatg ttgtttctac ttgtaaatgt 3300
aaagtcttta aaataaacta ttacagatam ttaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3360
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3420
aaaaaaaaaa aaangggggg cnttttaa 3449

<210> 226

<211> 1866

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1859)

<223> n equals a,t,g, or c

<400> 226

caccaatgtt gataaactgg tgaaggacat ttacggagga gactatgaac gatttggcct 60
tcaaggatct gctgtagcat caagctttgg caacatgatg agtaaagaaa agcgagattc 120
catcagcaag gaagacctcg cccgggccac attggtcacc atcaccaaca acattggctc 180
cattgctcgg atgtgtgctg tgaatgagaa catagacaga gttgtgttg ttggaaattt 240
tctcagaatc aatatggtct ccatgaagct gctggcatat gccatggatt tttgggtccaa 300
aggacaactg aaagctctgt ttttgaaca tgagggttat tttggagccg ttggggcact 360
gttgggaactg ttcaaatga ctgatgayaa gtagagacga gcagtggagg aaacagcctc 420
ccaaaaggac agagaactaa aaaattgctg ctggagaagg tgaaagtcgc tttgggacgg 480
aagccaagcc attatggcag atgaacctgc tggatttgta aataatttaa aatccttcca 540
gatgatcttt tactcttagg ttttgagcta atgattcaaa acgggggaat ataaaaggtt 600
tttttctgt atactgtatt tttttaaaaa aatggtgcag cgtggccaaa cctaccaatt 660
gtatgcatta actttgaaaa gttgtttgat gtttaagaag gacctgatat gtaagcgctg 720
gtcatttttc ttctggggtt tactgatcag tgtgtgtgatt ttaacttcat ttagtaatta 780
ctctaggaga ttttacctg acttatattt ttcatgacgt ttcattgatt gctgttggtt 840
tcaaatgaaa ctacaaatct ggcattgttt actgtgaaca cttttgttat ttgttttgta 900
cccttttttg tcttgktttt ctgktttagt tgkcttctga aaaaagagtc gttccctctg 960
ttctgtcct cagatgatgt cctccccct acctgtaacc tttctttgac ataattgttc 1020
atatcaatga aggtgctgac cagctcaata caaagttaag cacaagatct aaagctcttg 1080
aaaatgcccg tgaagagaag actgaatgtg ttaatgaatt taatgagtct ggcaaaagtt 1140
gcaattata tgcaagtttgc tctatcgt tataaatgta gtgtttcatt ggatttattt 1200

tatgctaggt tatattaagt tgaaatagtc tgtgattaaa tgcctcatc catgcacaga 1260
atatgaatgg cagcaaatct ttgtgcaaga aatttgaaac ttattgggaa aagcctccca 1320
gtagattaat tgttcataatc aggagattta gggtaagtca tgggttgagg tgcagatag 1380
taatatctat ttgttttga catgtatata tctaggaact ttgtaacaac acatctttaa 1440
taatgttaaa ggttttttca tttttaatat tttaactaa aaactgtact tcaatctcag 1500
tttctaaaat taaaaataat ttatactgat ctatatattt tttctttttg aaagatttca 1560
ttaagactga tgggtaactt tcaaatgagg gtcatgtaca aatattggga tgcagagat 1620
cccatgatct tgtgtattga gcttattgtt gaaagggatt tttgaaggac agaacaatta 1680
ctccatgatg aatcttcctt tctctgcctt ctgagcaccg tctttaattt ccatactctc 1740
aagtcttgaa gaagttagtg ttaattgaag aattcacttg tctggttgaa ataaagcctg 1800
tttctgttgt gaaaaaaaaa aaaaaaact sgtggggggc ccggacccaa ttgccctang 1860
ggagcc 1866

<210> 227

<211> 1064

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (209)

<223> n equals a,t,g, or c

<400> 227

ggcaacagca tggccacaat tttatagcag caaaatgaag caaaaatcaa aggagttagc 60
aaaggcagaa acatatgtgt agtgtgctgc cagcataaga tggaagaact taaagaaggc 120
ctgcggaaca gagatgagct tattgaggag aagcagcgca tgcagcagaa aatagacacc 180
atgacaaaag aggtgtttga cctccaggng acacttcttt ggaaagataa aaaaattgrg 240
aaacatggct tagttataat ccccgatggc actcccaatg gtgatgtcag tcatgaacca 300
gtggctggag ccatcactgt tgtgtctcag gaagctgctc aggtcttgga gtcagcagga 360
gaagggccat tagatgtaag gctacgaaaa ctgctggag agaaggaaga actactgtca 420
cagattagaa aactgaagct tcagttagag gaggaacgac agaaatgctc caggaatgat 480
ggcacagtgg gtgacctggc aggactgcag aatggctcag acttgcagtt catcgaaatg 540
cagagagatg ccaatagaca aattagcgaa tacaaattta agctttcaaa agcagaacag 600
gatataacta ccttggagca aagtattagc cggcttgagg gacaggttct gagatataaa 660
actgtgctg agaatgtga gaaagttgaa gatgaattga aagcagaaaa acggaagcta 720
caacgagagt tacgaacagc actggacaag attgaggaga tggagatgac caacagccac 780
ctggccaagc ggctggagaa gatgaaggcc aataggacag cacttctggc ccagcagtag 840
gaaaaccacc cttcaacctg ggtgatgctc cttggggccc tacctagagg gactgacttt 900
tgtccattga cacaaccacc ttttagtact gttttgagtt ttgtcattaa aacagccacc 960
tttgtatttt ataatttatg acagratgaa gtcattttga atctacatga atgaacactt 1020
tggatttttg ttggagtttg attctagggt agaaccagtc catg 1064

<210> 228

<211> 373

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (329)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (332)

<223> n equals a,t,g, or c

<400> 228

```
gctgcactct caggtattcc ctgctcttac tccaaaaaga tggacccagg tccgaagggg 60
cactgccact gtggggggca tggccatcct ccaggtcact gcggggccacc ccctggccat 120
ggcccagggc cctgcggggc accccccccac catggcccag gccctgagg gccaccccc 180
caccatgggc cagggccctg cggggccacc cctggccatg gcccaggggc ctgaggggcca 240
ccccccacc atggtccagg gccctgagg cctccccctg gccatggccc aggtcaccca 300
ccccctgggc cacatcactg aggaagtana anaaaacagg acacaagatg gcaagcctga 360
gagaaattgc cca 373
```

<210> 229

<211> 2844

<212> DNA

<213> Homo sapiens

<400> 229

```
tcgaccacg cgtccgcagg tgttcaagat ggcggtagct gaggggttga ccgagagacc 60
cagttgamgg cctttacgaa gtgaaaggag ccgggaatcg cccctaccc gcttctcgta 120
gtcctgggag cacagcagaa gtgtttttct ttttttaatg aacaagtaaa ccatacaaat 180
tgtcaacatg ggacggagat ctacatcatc caccaagagt ggaaaattta tgaacccac 240
agaccaagcc cgaagggaag cccggaagag agaattaaag aagaacaaaa aacagcgcat 300
gatggttcga gctgcagttt taaagatgaa ggatccaaaa cagataatcc gagacatgga 360
gaaattggat gaaatggagt ttaacccagt gcaacagcca caattaaatg agaaagtact 420
gaaagacaag cgtaaaaaag tcggtgaaac ctttgaacgt attctacgac tctatgaaaa 480
agagaatcca gatatttaca aagaattgag aaagctagaa gtagaatatg aacagaagag 540
ggctcaactt agccaatatt ttgatgctgt caagaatgct cagcatgtgg aagtggagag 600
tattcctttg ccagatatgc cacatgctcc ttccaacatt ttgatccagg acattccact 660
tcctggtgcc cagccaccct ctatcctaaa gaaaacctca gcctatggac ctccaactcg 720
ggcagtttct atccttcctc ttcttggaac tgggtgtcca cgtttgcccc ctggcagaaa 780
acctcctggc cctccccctg gtccacctcc tcctcaagtc gtgcagatgt atggccgtaa 840
agtgggtttt gccctagatc ttccccctcg taggcgagat gaagacatgt tatatagtcc 900
tgaacttgcc cagcgaggtc atgatgatga tgtttctagc accagtgaag atgatggcta 960
tcctgaggac atggatcaag ataagcatga tgacagtact gatgacagtg acaccgacaa 1020
atcagatgga gaaagtgcag gggatgaatt tgtgcaccgt gataatggtg agagagacaa 1080
caatgaagaa aagaagtcag gtctgagtgt acggtttgca gatatgcctg gaaaatcaag 1140
gaagaaaaag aagaacatga aggaactgac tcctcttcaa gccatgatgc ttcgtatggc 1200
aggtaagaaa atccctgagg agggacggga agtagaggaa ttttcagagg acgatgatga 1260
agatgattct gatgactctg aagcagaaaa gcaatcacia aagcagcata aagaggaatc 1320
ccattctgat ggcacatcca ctgcttcttc acagcagcag gctccgccc agtctgttcc 1380
tccttctcag atacaagcac ctcccatgcc aggaccacca cctcttgagg caccacctgc 1440
tccaccatta cggcctcctg ggccacctac aggccttcc cctgggtccac ctccaggagc 1500
tcctccattc ctgagaccac ctggaatgcc aggactccga gggcccttac ccgactttt 1560
acctccagga ccaccaccag gccgaccccc tggccctccc ccagggtccac ctccaggtct 1620
gcctcctggt cccctcctc gtggaccccc accaaggcta cctccccctg caccctcagg 1680
tattcctcca cctcgtcctg gcatgatgag cccacctttg gtgcctcccc ttggacctgc 1740
```

```
ccccctggg ctgttccac cagctccctt gccaaaccct ggggttttaa gtgccccacc 1800
caacttgatt cagcgaccca aggcggatga tacaagtgca gccaccattg agaagaaagc 1860
cacagcaacc atcagtgcc aagcacagat cactaatccc aaggcagaga ttactcgatt 1920
tgtgcccact gcaactgagag tacgtcggga gaataaagg gctactgctg ctccccaag 1980
aaagtcagag gatgattctg ctgtgcctct tgccaaagca gcacccaaat ctggtccttc 2040
tgttcctgtc tcagtacaaa ctaaggatga tgtctatgag gctttcatga aagagatgga 2100
agggtactg tgacagcttt tgatgccaga aaaggcttct gttcacaaca gtggcccatg 2160
gagaaagagg ctcttattaa acttagatga aagagctgct tccattgtca gggatatttc 2220
taatttcagt tcaaggaata tcctaaaatt tagccttggt cagaatttac tgcacataaa 2280
aaagggtatt tcatccagaa tagatcagtt attgaagcag tgctgctaac atccattccc 2340
tttcatacca ccattttcac cctgtttctt cccctcctcc agttctttgg aaatttgtga 2400
tcgggggagc ttagttgctt atttgttttg actcttggt gctgtgggca ctggagtaga 2460
gatttctgga gaaaaaaaaa cagtttattt catcttgctt tttgtgtttg agttattttt 2520
aatattttcc tgtaaatatt ttgtaaatatt ttacttgtaa tgaaatggat cacaatgtca 2580
tttcctaata caaggcagga tatgtgggaa gaatatgtac aattatttga ttaaaattat 2640
ttcccaactga cctaaacttt cagtgttttg tgggaaaaat aaataaatgt tctacaccaa 2700
gatactctgt gttactgata atatgtaaac ataataccta ttttatggga caggaaaaag 2760
ttaaaatatt taaaaacttg ggggttttaa gggaactgcc tttctgggta cagatgagga 2820
tcttactgga aaactgggta aaaa 2844
```

<210> 230

<211> 1798

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (15)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (24)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (501)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (1798)
<223> n equals a,t,g, or c

<400> 230
ngcaaatata cgtgnacact tatnaaaagt nacgcctgac acgtgaccgg ttccgggaat 60
tccccgggtcg acccacgcgt ccggcccagc actagaagtc ggcgggtgtt ccattcggtg 120
atcagcactg aacacagagg actcaccatg gagtttgggc tgagctgggt tttcctcgtt 180
gctcttttaa gaggtgtcca gtgtcaggtg cagctggtgg agtctggggg aggcgtggtc 240
cagcctggga ggtccctgag actctcctgt gcagcstctg gattcacctt cagtagctat 300
ggcatgcact gggctccgcca ggctccaggc aaggggctgg agtgggtggc agttatatsr 360
tatgatggaa gtaataaata ctatgcagac tccgtgaagg gccgattcac catctccaga 420
gacaattcca agaacacgct gtatctgcaa atgaacagcc tgagagctga ggacacggct 480
gtktattact gtgcgaraga ngttactatg gttcggaaag catctactac tactttgact 540
ctggggccag ggaacmctgg tcaccgtctc ctacgcctcc accaaggggc catcgggtctt 600
ccccctggca cctcctcca agagcacctc tgggggcaca gcggccctgg gctgcctggg 660
caaggactac tccccgaac cgggtgacgtg gtcgtggaac tcaggcgccc tgaccagcgg 720
cgtgcacacc tccccggtg tcctacagtc ctacggactc tactccctca gcagcgtggt 780
gaccgtgccc tccagcagct tgggcaccca gacctacatc tgcaacgtga atcacaagcc 840
cagcaacacc aaggtggaca agaragttga gcccaaactc tgtgacaaaa ctcacacatg 900
cccaccgtgc ccagcacctg aactcctggg gggaccgtca gtcttctctt tccccccaaa 960
acccaaggac accctcatga tctcccgac ccctgaggtc acatgcgtgg tgggtggacgt 1020
gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa 1080
tgccaagaca aagccgaggg aggagcagta caacagcacg taccgtgtgg tcagcgtcct 1140
caccgtctcg caccaggact ggctgaatgg caaggagtac aagtgcgaag tctccaacaa 1200
agccctccca gccccatcg agaaaacat ctccaaagcc aaagggcagc cccgagaacc 1260
acaggtgtac accctgcccc catcccgga kgagmtgacc aagaaccagg tcagcctgac 1320
ctgcctggtc aaaggcttct atcccagcga catcgccgtg gagtgggaga gcaatgggca 1380
gccggagaac aactacaaga ccacgcctcc cgtgctggac tccgacggct ccttcttctt 1440
ctayagcaag ctaccgtgg acaagagcag gtggcagcag gggaacgtct tctcatgctc 1500
cgtgatgcat gaggtctctg acaaccacta cagcgagaag agcctctccc tgctcyccgg 1560
taaatgagtg cgacggccgg caagcccccg ctccccgggc tctcgcggtc gcacgaggat 1620
gcttggcacg taccctgtst acatacttcc crggcrcca gcatggaaat aaagcaccca 1680
scrctgcctt gggcccctgc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1740
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaawaa aaaaaaaaaa aaaaaaan 1798

<210> 231
<211> 1823
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (82)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1593)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (1714)
<223> n equals a,t,g, or c

<400> 231
gtcgccctgcc acccgcttcg gccacggcca gacgcccgcg gccttcctca gcgctctgct 60
gccctcgcag ccgcccgcg gncgggtcaac gccctggggc tgcccaaggg tgtcaccccc 120
gcagatttcc aaagaaggcc agtagaactg ctagaatagc ctccgatgag gaaattcaag 180
gcacaaagga tgctgttatt caagacctgg aacgaaaact tcgcttcaag gaggacctcc 240
tgaacaatgg ccagccgagg ttaacatacg aagaaagaat ggctcgtcga ctgctagggtg 300
ctgacagtgc aactgtcttt aatattcagg agccagaaga ggaaacagct aatcaggaat 360
acaaagtctc cagctgtgaa cagagactca tcagtgaat agagtacagg ctagaagggt 420
ctcctgtgga tgaatcaggt gatgaagttc agtatggaga tgtgcctgtg gaaaatggaa 480
tggcaccatt ctttgagatg aagctgaaac attacaagat ctttgaggga atgccagtaa 540
ctttcacatg tagagtggct ggaaatccaa agccaaagat ctattgggtt aaagatggga 600
agcagatctc tccaaagagt gatcactaca ccattcaaag agatctcgat gggacctgct 660
ccctccatac cacagcctcc accctagatg atgatgggaa ttatacaatt atggctgcaa 720
accctcaggg ccgcatcagt tgtactggac ggctaattgt acaggctgtc aaccaaagag 780
gtcgaagtcc ccggtctccc tcaggccatc ctcatgtcag aaggcctcgt tctagatcaa 840
gggacagtgg agacgaaaat gaaccaatc aggagcgatt cttcagacct cacttcttgc 900
aggctcctgg agatctgact gttcaagaag gaaaactctg cagaatggac tgcaaagtca 960
gtgggttacc aaccctgat ctaagctggc aactagatgg aaagcccgtc cgccctgaca 1020
gtgctcacia gatgctgtg cgtgagaacg ggtgacactc tctgatcata gagccagtca 1080
cgtcacgtga tgccggcatc tacacatgta tagctacca cagagcagga cagaactcat 1140
tcagcctgga gcttgtggtt gctgctaaag aagcacacia accccctgtg ttatttgaga 1200
agctccaaaa cacaggagtt gctgatgggt acccagtgcg gctggaatgt cgtgtattgg 1260
gagtgccacc acctcagata ttttggaaga aagaaaatga atcactcact cacagcactg 1320
accgagttag catgcaccag gacaaccacg gctacatctg cctgtctcatt caggagacca 1380
caaaagaaga tgctgggttg tatactgtgt cagccaagaa tgaagcaggg attgtgtcct 1440
gtactgccag gctggacgtt tacacccagt ggcacagca gtcacagagc accaagccaa 1500
aaaaagtacg gccctcagcc agtcgctatg cagcactttc ggaccaggga ctagacatca 1560
aagcagcggt ccaacctgag gccaaacct ctnacctgac actgaatact gccttggtag 1620
aaagttagga cctgtaatcc agcattcttg ktaaaagctga aacacttgaa acagscattg 1680
scttgaccaa catattcctt tgtcacatta tgtnaaaggc agaamcatac ctttgctatt 1740
agaaattaaa aaaacaccaa aatatatttt ctacttgaat accaacttag ttaaagtaga 1800
taatgctaata acaaatatac aca 1823

<210> 232
<211> 970
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (936)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (960)

<223> n equals a,t,g, or c

<400> 232

```
aattcgggac gagggcagcc cgcttgccaa tcagagcgcg gctgagcggc cccgcagcca 60
acccccgagg agcgcccggc tggcgtccgc cgcgcccagg agttggggat gtcctacaaa 120
cccatcgccc ctgctcccag cagcaccctt ggctccagca cccctggggc gggcaccccg 180
gtccctacag gaagcgctcc gtcccgctcg ggctcagtgc caggagcccg cgctcctttc 240
agaccgctgt ttaacgactt tggaccgcct tccatgggct acgtgcaggc gatgaagcca 300
cccggcgccc agggctccca gagcacctac acggacctgc tgtcagtcac agaggagatg 360
ggcaaagaga tccggcctac ctatgctggc agcaagagcg ccatggagcg cytgaagaga 420
ggaagcgccct cggcctcagc gtctggacct atccggccac tgcagagcac ccgcttctcc 480
ctggccttca tcccgagttg cactaaccat cctgggcttc ctgtcctgtg tcccttggtg 540
ggtcccccctc aggaaccaag gagtggccct ccaggtggca gactaagga cccccccca 600
caacaagagt tagcagcgag gtcccatga gtcccacca tgacctgccg acagtgttgc 660
ccaccggaac ttttgtggcc cctaccgctc agcccttccc agcacttctc ccactttgtc 720
ccgagcctcc ttctccccc gcaggggcac aggcctggca cctccctgcc ttgtgtcctg 780
agccatagtg actcttttat ctgtgtgtct tttgctaaat atgccctttt tatattaata 840
aaagatgatt tggagttgtg ctctcaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 900
aaaaaaaggc cggccgctct agaggatcca tgcttnacgt acgcgtgcat gcgtacgtcn 960
ttgctcttct                                     970
```

<210> 233

<211> 967

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (923)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (926)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (955)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (957)

<223> n equals a,t,g, or c

<400> 233

```
ggcagagcac ttcctggcca ggaaacctga gcggtgagac tcccagctgc ctacatcaag 60
gccccaggac atgcagaacc ttcctctaga acccgaccca ccaccatgag gtccctgcctg 120
tggagatgca ggcacctgag ccaaggcgct cagtggctct tgcttctggc tgccctggtc 180
ttctttctct tcgccttgcc ctcttttatt aaggagcctc aaacaaagcc ttccaggcat 240
```

caacgcacag agaacattaa agaaaggtct ctacagtccc tggcaaagcc taagtcccag 300
gcacccacaa gggcraggag gacaaccatc tatgcagagc cagygccaga gaacaatgcc 360
ctcaacacac aaaccacagc caaggccac accaccggag acagaggaaa ggaggccaac 420
caggcaccgc cggaggagca ggacaaggtg cccacacag cacagagggc agcatggaag 480
agcccagaaa aagagaaaac catggtgaac aactgtcac ccagagggca agatgcaggg 540
atggcctctg gcaggacaga ggcacaatca tggaaagagc aggacacaaa gacgaccaa 600
ggaaatggg gccagaccag gaagctgacg gcctccagga cgggtgcaga gaagcaccag 660
ggcaaagcgg caaccacagc caagacgstc attyccaaa gtcagcacag aatgctggct 720
yccacaggag cagtgtcaac aaggacgaga cagaaaggar tkaccacagc agtcatcca 780
cctaaggaga agaaacctca ggccaccca cccctgccc cttccagag cccacgacg 840
cagagaaacc aaagactgaa gggcggcaac ttcaaactct agcctcgggtg ggattttgag 900
gaaaaataca agcttcgaaa tanganggtc tcagacgact tggccttgac tctgngnaga 960
tcaaaag 967

<210> 234

<211> 2163

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1140)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1157)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1158)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1166)

<223> n equals a,t,g, or c

<400> 234

gcccagacct gcgatggctc ttccccaccg gaggtcagc ccgtggctgc gtcaaaggca 60
tcaaggccct gggcaagtat gtggacctca agcggctgaa cagcacaggc gtgacgccgg 120
ctgcacygcc gacctgctgg tggggcgcgc catgactttc catggccacg gcttccttcg 180
cctggcgctc tcgaacgtgg caccgctcac tggcaacgtc tactccggct tcggcttcca 240
cagcgcaccg gacagtgcc tgctctacta ccgggcgtcc ccggatgggc tatgccaggt 300
gtccctgcag cagggccgtg tgagcctaca gtcctgagg actgaagtga aaactcaagc 360
gggcttcgcc gatgggtgcc cccattacgt cgcctctac agcaatgcc cgggagctctg 420
gctgtatgtc gatgaccagc tccagcagat gaagccccac cggggaccac ccccgagct 480
ccagccgcag cctgaggggc ccccgaggct cctcctggga ggcctgcctg agtctggcac 540
catttacaac ttcagtggct gcatcagcaa cgtcttcgtg cagcggctcc tgggcccaca 600
gcgcgtatct gatctgcagc agaacctggg cagcgtcaat gtgagcacgg gctgtgcacc 660

cgccctgcaa gccagaccc cgggcctggg gcctagagga ctgcaggcca ccgcccggaa 720
ggcctcccgc cgagccgctc agcccgcgcg gcctcctgcc tgcctgctgc cccacacct 780
caggaccacc cgagactcct accagtttgg gggttccctg tccagtcacc tggagtttgt 840
gggcatcctg gcccacata ggaactggcc cagtctctcc atgcacgtcc tcccgcgaag 900
ctcccagggc ctctctctct tcaactgccc tctgaggccc ggcagccctt ccctggcgct 960
cttctgagc aatggccact tcgttgacac gatggaaggc ctccggactc ggctccgcgc 1020
ccagagccgc cagcgtctcc ggcctggcgc tggcacaagg tctccgtgcg ctgggagaag 1080
aaccggatcc tgctgggtgac ggacggggcc cgggscgtga gccaggaggg gccgcaccgn 1140
agcaccaggg ggcagannac cccangccc cacacctctt ttgtggggcg cctcccggcm 1200
agcagccaca gctccaaact tccggtgacc gtcgggttca gcggctgtgt gaagagactg 1260
aggctgcacg ggaggccctt gggggccccc acacggatgg caggggtcac accctgcatc 1320
ttgggcccc tggaggcggg cctgttcttc ccaggcagcg ggggagttat cactttagac 1380
ctcccaggag ctacactgcc tgatgtgggc ctggaactgg aggtgcggcc cctggcagtc 1440
accggactga tcttccactt gggccaggcc cggacgcccc cctacttgca gttgcagggtg 1500
accgagaagc aagtctctgt gcggcgcatg gacggagcag gggagttctc cacgtcataa 1560
cccggccctc agtctctgtg gatggccagt ggcaccggct agcgggtgatg aaaagcggga 1620
atgtgctccg gctggaggtg gacgcgcaga gcaaccacac cgtgggcccc ttgctggcgg 1680
ctgcagctgg tggccagcc cctctgtacc tcgggggcct gcctgagccc atggccgtgc 1740
agccctggcc cccgcctac tgcggctgca tgaggaggct ggcggtgaac cgggtccccg 1800
tcgcatgac tcgctctgtg gaggtccacg gggcagtggt ggccagtggc tgccagccg 1860
cctaggacac agccaacccc gggccctggt caggccctg cagctgcctc acaccgcccc 1920
ttgtgctcgc ctcataggtg tctatttggc ctctaagctc tacgggtgac agatcttgtt 1980
tctgaagatg gtttaagtta tagcttctta aacgaaagaa taaaatactg caaatgttt 2040
ttatatttgg cccttccacc catttttaat tgtgagagat ttgtcaccaa tcatcactg 2100
ttcctcctta aaaattaaaa agtaacttct gtgtaaaaaa aaaaaaaaaa aaaaaaaaaa 2160
aaa 2163

<210> 235

<211> 1321

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1313)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1320)

<223> n equals a,t,g, or c

<400> 235

gcgcgacgcc gccacctccg gaacaagcca tgggtggcggc gacgggtggca gcggcgtggc 60
tgctcctgtg ggctgcggcc tgcgcgcagc aggagcagga cttctacgac ttcaaggcgg 120
tcaacatccg gggcaaacctg gtgtcgtctg agaagtaccg cggatcgggtg tccctgggtg 180
tgaatgtggc cagcagatgc ggcttcacag accagcacta ccgagccctg cagcagctgc 240
agcgagacct gggcccccac cacttcaacg tgctcgctt cccctgcaac cagtttggcc 300
aacaggagcc tgacagcaac aaggagattg agagctttgc ccgccgcacc tacagtgtct 360
cattccccat gtttagcaag attgcagtca ccggtactgg tgcccatcct gccttcaagt 420
acctggccca gacttctggg aaggagccca cctggaactt ctggaagtac ctagtagccc 480

cagatggaaa ggtggtaggg gcttgggacc caactgtgtc agtggaggag gtcagacccc 540
agatcacagc gctcgtgagg aagctcatcc tactgaagcg agaagactta taaccaccgc 600
gtctcctcct ccaccacctc atcccgccca cctgtgtggg gctgaccaat gcaaactcaa 660
atgggtgcttc aaaggggagag acccactgac tctccttcct ttactcttat gccattggc 720
ccatcattct tgtgggggaa aaattctagt attttgatta tttgaatctt acagcaacaa 780
ataggaaactc ctggccaatg agagctcttg accagtgaat caccagccga tacgaacgctc 840
ttgccaacaa aaatgtgtgg caaatagaag tatatcaagc aataatctcc caccacaaggc 900
ttctgtaaac tgggaccaat gattacctca tagggctgtt gtgaggatta ggatgaaata 960
cctgtgaaag tgcctaggca gtgccagcca aataggaggc attcaatgaa cattttttgc 1020
atataaamca aaaaataact tgttatcaat aaaaacttgc atccaacatg aatttccagc 1080
cgatgataat ccaggccaaa ggttttagttg ttgttatttc ctctgtatta tttcttctat 1140
tacaaaagaa atgcaagttc attgtaacaa tccaaacaat acctcacgat ataaaaataa 1200
aatgaaagta tcctcctcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1260
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aanggggggn 1320
c 1321

<210> 236

<211> 683

<212> DNA

<213> Homo sapiens

<400> 236

tgcacccacg cgtccgagca gctgctgcag ggcccatggc ggacaccag tacatcctgc 60
ccaatgacat cggcgtgtct agcctggact gccgtgaggc ctccgcctg ctgtcaccca 120
cagagcgctt ctatgcctac cacctgtccc gtgccgcctg gtacggaggc ctggctgtgc 180
tgcttcagac ctcccttgag gccccctaca tctatgtctt gctcagccgc ctcttccgcg 240
cccaggaccc cgaccagctg cgccaacatg ccctggctga aggccttacc gaggaggagt 300
atcaggcggt cctggtctat gccgcgggtg tttactccaa catgggcaac tacaagtcct 360
ttggtgacac caagtttgtt cccaacttgc ccaaggaaaa gctggaacgg gtgatcctag 420
ggagtgaggc tgctcagcag caccagaag aagtcagggg cctctggcak acctgcgggg 480
agcttatgtt ctctctggag ccaaggcttc gacacctcgg actgggggag gagggaaatca 540
ccacctattt ctctgggaat tgtaccatgg aagatgccaa attggccmag atttctggac 600
tcacagaacc tyartgccta caacaaccgg stcttyaaag aagtcsatgg garaaaggaa 660
rccctamtac gaagggtgcgg gtt 683

<210> 237

<211> 2115

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2112)

<223> n equals a,t,g, or c

<400> 237

attcccgggt cgaccacgc gtccgggctc cggcaaccgc tccggcaacg ccaaccgctc 60
cgctgcgcgc aggtggtgct gcaggctctc ggctgcagcg ctggggtctt gctttgtctc 120
ccaggctggt gtgcagtggt gcgaccacgg ctacagcgag cctcagccac ccagatgtaa 180
gcgatctggt tcccacctca gcctcccag tagtgatctt aggatccggc ttccaacatg 240
tggcagctct gggcctccct ctgctgcctg ctggtgttgg ccaatgcccg gagcaggccc 300

tctttccatc ccstgtcggg tgagctgggc aactatgtca acaaacggaa taccacgtgg 360
caggccgggc acaacttcta caacgtggac atgagctact tgaagaggct atgtggtacc 420
ttcctgggtg ggcccaagcc accccagaga gttatgttta ccgaggacct gaagctgcct 480
gcaagcttcg atgcacggga acaatggcca cagtgtccca ccatcaaaga gatcagagac 540
cagggtcctt gtggctcctg ctgggccttc ggggctgtgg aagccatctc tgaccggatc 600
tgcattccaca ccaatgcgca cgtcagcgtg gaggtgtcgg cggaggacct gctcacmtgc 660
tgtggcagca tgtgtgggga cggctgtaat ggtggctatc ctgctgaagc ttggaacttc 720
tggacaagaa aaggcctggt ttctggtggc ctctatgaat cccatgtagg gtgcagaccg 780
tactccatcc ctccctgtga gcaccacgtc aacggctccc ggcccccatg cacgggggag 840
ggagataccc ccaagtgtag caagatctgt gagcctggct acagcccagc ctacaaacag 900
gacaagcact acggatataa ttcctacagc gtctccaata gcgagaagga catcatggcc 960
gagatctaca aaaacggccc cgtggaggga gctttctctg tgtattcgga ctctctgctc 1020
tacaagtca ggtgtacca acacgtcacc ggagagatga tgggtggcca tgccatccgc 1080
atcctgggct ggggagtgga gaatggcaca ccctactggc tgggtgcca ctcttggaac 1140
actgactggg gtgacaatgg cttctttaa atactcagag gacaggatca ctgtggaatc 1200
gaatcagaag tgggtggctg aattccacgc accgatcagt actgggaaaa gatctaattc 1260
gcccgtgggc tgtcgtgcca gtcctggggg cgagatcggg gtagaaatgc attttattct 1320
ttaagttcac gtaagataca agtttcagrc agggctctgaa ggactggatt ggccaaacat 1380
cagacctgtc ttccaaggag accaagtcct ggctacatcc cagcctgtgg ttacagtga 1440
gacaggccat gtgagccacc gctgccagca cagagcgtcc ttccccctgt agactagtgc 1500
cgtagggagt acctgtgccc ccagctgact gtggccccct ccgtgatcca tccatctcca 1560
gggagcaaga cagagacgca ggaatggaaa gcggagttcc taacaggatg aaagtcccc 1620
catcagttcc ccagttacct ccaagcaagt agctttccac atttgtcaca gaaatcagag 1680
gagagatggt gttgggagcc ctttgagaa cgccagtctc ccaggcccc tgcatctatc 1740
gagtttgcaa tgtcacaacc tctctgatct tgtgctcagc atgattcttt aatagaagtt 1800
ttattttttc gtgcactctg ctaatcatgt gggtagacca gtggaacagc gggagacctg 1860
tgctagtttt acagattgcc tcctaataac gcggctcaaa aggaaaccaa gtggtcagga 1920
gttggtttctg acccactgat ctctactacc acaaggaaaa tagtttagga gaaaccagct 1980
tttactgttt ttgaaaaatt acagcttcac cctgtcaagt taacaaggaa tgctgtgccc 2040
aataaaaggt ttctccaact tgaaaaaaa aaaaaaaaa aaamwmgggg ggggcccgtg 2100
cccatggcc tntgg 2115

<210> 238

<211> 1642

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1633)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1638)

<223> n equals a,t,g, or c

<400> 238

taagcaggca ctagaggacc tggagcgcac atcaggccgt ggcaccctga tgcgccacc 60
atgctcaaca tccttgcttt ggtgtatcgt gaccagaata agtataagga agctgccac 120
ctgctgaatg atgcccttag catccgggag agcaccttg gacctgacca tcctgctgtg 180

gctgccacac tcaacaattt ggctgtgctc tatggcaaaa ggggcaagta caaggaggca 240
gagcctctgt gccagcgggc actggagatt cgagaaaagg tcctgggcac gaatcatcca 300
gatgtggcaa aacagctgaa caacctggcc ctcttggtcc aaaaccaggg caagtatgag 360
gccgtggaac gctactacca gcgagcactg gccatctacg aggggcagct ggggcccggac 420
aaccctaatt tagcccgac caagaacaac ctggcttcct gttacctgaa acaggggcaa 480
tatgctgagg ctgagacact atacaaagag atcctgacct gtgcccattg acaggagttt 540
gggtctgtgg atgatgacca caagcccatc tggatgcatg cagaggagcg ggaggaaatg 600
agcaaaagcc ggcaccatga ggggtggaca ccctatgctg agtatggagg ctggtacaag 660
gcctgcaaaag tgagcagccc cacagtgaac actactctga gaaacctggg agctctgtat 720
aggcgccagg gaaagctgga ggctgctgag accctggagg aatgtgccct gcggtcccgg 780
agacagggca ctgaccctat cagccagacg aaggtggcag agctgcttgg ggagagtgat 840
ggtagaagga cctcccagga gggccctgga gacagtgtga aattcgaggg aggtgaagat 900
gcttctgtgg ctgtggagtg gtccggggat ggcagtggga ccctgcagag gagtggctct 960
cttggcaaga tccgggatgt gctccgcaga agcagtgaac tcttggtgag gaagctccar 1020
gggactgagc ctcgccctc cagcagcaac atgaagcgag cagcctcctt gaactatctg 1080
aaccaacctt gtgcagcacc cctccagtc tcccggggcc tcagtgccag caccatggac 1140
ctctcttcaa gcagctgaca ttcaaccggg ccccaggtc tgctgggtcc ccccacccc 1200
acagccctca cagcattccc cattgtcctt ggctcttccc caccctaggg tgggacagt 1260
aaggggagca gtttaaccag aagattgctg ctgcccttag ggtctcagct ccctcctcag 1320
gaatccctct taggaaggac cctcaggaca ccctctctgc accctgtggg cctctagagt 1380
agctagctct gaggcccaa ggtgggtaca aagcaggtat ggccctcaga gatgcagcct 1440
gctgctggct ttctagtcag agggttgggg gctggccagc caagctgcct tgccctggcc 1500
gctcttactc cctccctctg ctgtctcact tcagggtccat gtatttctact tttcttaaat 1560
aaaagaatca ggtaacctta aaaaaaaaaa aaaaaaaaaa cttggggggg gccccgaacc 1620
caatttgccc canagggnng cg 1642

<210> 239

<211> 468

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (461)

<223> n equals a,t,g, or c

<400> 239

gcggacgcgt gggcctggtc tcagtatggc gctgtccttg gttcttacag tcctgagcct 60
cctacctctg ctggaagcca gatccattg tgtgccaacc tagtaccggt gcccatcacc 120
aacgccaccc tggaccggat cactggcaag tggttttata tcgcatcggc ctttcgaaac 180
gaggagtaca ataagtcggt tcaggagatc caagcaacct tcttttactt taccaccaac 240
aagacagagg acacgatctt tctcagagag taccagacct gccagaacca gtgcttctat 300
aactccagtt acctgaatgt ccagcgggag aatgggaccg tctccagata cgagggaggc 360
cgagaaacat gttgctcacc tgctgttcct tagggacacc aagaccttga tgtttggtty 420
ctacctggac gatgagaaga actgggggtg tctttctatg ntgacaag 468

<210> 240

<211> 1329

<212> DNA

<213> Homo sapiens

<400> 240

tacatcttat aaaacctaac tggcatttaa aaaatactgt ggccgggctg ggtggctcat 60
gcctgtaatc ccagcacttt gggaggccga ggtgggagga ttgcttgagt ccaggaattt 120
gagaccagcc tggacaacac agtgagacct catctctatc aaaaaataaa aattagctag 180
atgtgggtggc atgagcctgt gttcccagct gcttaggagg ctgaagcagg aggattgatt 240
gagcctgcga ggccaaggct gcagcaggct gtgattgcac cactgcactt cagcttgggc 300
aacagagcaa gaccctgtct ccgaaacaaa taaaaaatat tgtaataaaa gtacttataa 360
acatactaata cctcttttcag gaccctaaag ttgcaggtta gtaggtcttc aaggacaaat 420
ctgtaagttt cttattttctg tagtgcaagt aaaatttcac tttttgaaac tatagagaga 480
tcccctttctg attagcctac agaacttaaa gtgagggaac catttcctct cacagacaaa 540
gaggcctggg atattaggac tttgggggtt gagagcatca tggggcagac agatggtgga 600
tggtctggac aagaagcag taagccactg cggttggtca tactgaaggg aattgatggc 660
aagaggatcc cctgagcaag tcagaagtta ctctcatcag tcgttcctgg tcacaacctg 720
aggtagctctg ctgagtgggc aaggctgaag aagaggcctg tggaatgcag cattacctgc 780
tggacagagc agggcaggca gttctatgcc ttggagctcc tgactgcagg gactctgtcc 840
ccacactcar aaagactcag ctactcaat gagagaatgt gatttacttt atagaacgta 900
taatcaactt tgttgaataa tttgttctat taaggctgtc taaagtatgt gatgtcttca 960
tcatagtatg aagtgttgaa aattaataac gagcctagtt taggaaaaag ctgcttaaaa 1020
ctgtggctct aagagagtaa tcataaaata ccttagataa aattgacta tggaattttc 1080
attgagtatg tttaaattat tggcttgtct actaatcac atctgcttca aaatgaacat 1140
atttcataaa aattggcatca attttaatga cgctcctggt atggaacctc agatataccc 1200
tattggagac aatcctttga tcataaattc tcccacaata taaatcattt tatgtcttta 1260
tcawaaaaaa aaaaaggaat tcgatatcaa gcttatcgat accgtcgacc tcgagggggg 1320
gcccgtacc 1329

<210> 241

<211> 1652

<212> DNA

<213> Homo sapiens

<400> 241

gagcayttcg gcaactggca aaaattaggt gtacagggat ctaggtaata ctgtttattt 60
gagcaataat atattgtgct aacgttcagg catcctatta ctgagaaata agggaaaatg 120
agtgtaaaagt acaactaaga gtctcggcta cagggaataa taccatcagt taaatatcca 180
tagtccctaga gcatattatgt aaaactgcaa tttgaatcct gcaatacatt ttggcttttt 240
cctcagtgat accatgtgtg ggaagttgtt ctgtcaaggt gggtcggata atttgccctg 300
gaaaggacgg atagtgactt tcctgacatg taraacattt gatcctgaag acacaagtca 360
agaaatargc atgggtggcca atggaactaa gtgtggcgat aacaagggtt gcatkaatgc 420
agaatgtgtg gatattgaga aagcctacaa atcaaccaat tgctcatcta agtgcaaagg 480
acatgtgtgt tgtgacctg agctccagt tcaatgtgag gaaggatgga tccctcccga 540
ctgcatgagay tcctcagtgg tcttccactt ctccattgtg gttgggggtc tgttcccaat 600
ggcgggtcatt tttgtgtgtg ttgctatggt aatccggcac cagagctcca gagaaaagca 660
gaagaagatg cagaggccay tatctaccac tggcaccagg ccacacaaac agaagaggaa 720
acccacagatg gtaaaggctg ttcaacccca agagatgagt cagatgaagc cccatgtgta 780
tgatctgccca gtagaaggca atgagccccc agcctctttt cataaagaca caaacgcact 840
tccccctact gttttcaagg ataattcaat gtctacacct aaggactcaa atccaaaagc 900
atgaagcaac agctaagcaa gaactaatgg ctaaattatc aacttgaaa actggaaaat 960
ctggatggca gagaaatata ctatctcacc agtatttggc ctgcactcaa gaagggttaac 1020
atcttctgat tcatgttaga ctttgaagag actaaagaaa attttcaaga ggaacatatg 1080
cctgagaacc tttgatgaa tttaaaattt caattatcca ttcttataag aagggaagatg 1140
attgtaaaga aatatctccg aagttaaaaat ctgtaatagg aattgattca ttctctaagt 1200

aaaacaaaac ataaaaacat cacactaatc ttggaggaat aagaaaaatt gtacatccat 1260
taaatgtaca attgattgca acatcttgat tgttttaacc attaacttgt caaattacaa 1320
tcacagttaa gaaaatgatg taaaattctg ttttgtggat ctctttccta gattagcttc 1380
tgaaatcatt attagctata tcatttgagg ttttctacaa tttgggtataa ctaagaattt 1440
aaaaatgttt tatcatatat atttgtataa ttaattactg gcatgggttaa agtgggtttc 1500
acttttttaa tggagaaaaa ttcagttaaa ttaataggat aaaccagggt gcgaactggt 1560
gacctgtagg ccatgtttgc actgcaaata tatttgggtc gaatgatatt gaaaaaaaaa 1620
aaaaaaaaaa ctcgaggggg ggcccgtacc ca 1652

<210> 242

<211> 1946

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (84)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (538)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1932)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (1941)

<223> n equals a,t,g, or c

<400> 242

ggcacgagta caaggtggcg gcgctgggmc tggccaccgg cwtcgtcttg gtgctgctgc 60
tgctctgcct ctaccgcgtg ctangcccgc gcaactacgg gcagctgggt ggtgggcccg 120
ggcggcgagg gcgcggggag ctgccctgcg acgactacgg ctatgcgcca cccgagacgg 180
agatcgctgcc gcttgtgctg cgcggccacc tcatggacat cgagtgcctg gccagcgacg 240
gcatgctgct ggtgagctgc tgcctggcag gccacatctg cgtgtgggac gcgcagaccg 300
gggattgcct aacgcgcatt ccgcgcccag gcaggcagcg ccgggacagt ggcgtgggca 360
gcgggcttga ggctcaggag agctgggaac gactttcaga tgggtgggaag gctggtccag 420
aggagccttg ggacagccct cccctgagac accgccccg gggccctccg ccgccttccc 480
tcttcgggga ccagcctgac ctcacctgct taattgacac caacttttca gcgcagcnc 540
ggtccctaca gcccactcag cccgagcccc ggcaccgggc ggtctgtggc cgtctcggg 600
actccccagg ctatgacttc agctgcctgg tgcagcgggt gtaccaggag gaggggctgg 660
cggcctctg cacaccagcc ctgcgcccac cctcgcctgg gccggtgctg tcccaggccc 720
ctgaggacga ggggtggctcc cccgagaaa gctccccttc cctcgcctgg gccccagtg 780
ccgagggttc catctggagc ttggagctgc agggcaacct catcgtgggt gggcgaggca 840
gcggccggct ggaggtgtgg gacgccattg aagggtgtgt gtgctgcagc agcgaggagg 900
tctcctcagg cattaccgct ctggtgttct tggacaaaag gattgtggct gcacggctca 960

```
acgggtccct tgatttcttc tccttggaga cccacactgc cctcagcccc ctgcagttta 1020
gagggacccc agggcggggc agttcccctg cctctccagt gtacagcagc agcgacacag 1080
tggcctgtca cctgacccac acagtgccct gtgcacacca aaaacccatc acagccctga 1140
aagccgctgc tgggcgcttg gtgactggga gccaaagacca cactctgaga gtgttccgctc 1200
tggaggactc gtgctgcctc ttcacccttc agggccactc aggggccatc acgaccgtgt 1260
acattgacca gaccatggtg ctggccagtg gaggacaaga tggggccatc tgcctgtggg 1320
atgtactgac tggcagccgg gtcagccatg tgtttgctca ccgtggggat gtcacctccc 1380
ttacctgtac cacctcctgt gtcacagca gtggcctgga tgacctcatc agcatctggg 1440
accgcagcac aggcataaag ttctactcca ttcagcagga cctgggctgt ggtgcaagct 1500
tgggtgtcat ctacagacaac ctgctggtga ctggcggcca gggctgtgtc tccttttggg 1560
acctaaacta cggggacctg ttacagacag tctacctggg gaagaacagt gaggcccagc 1620
ctgcccgcca gatcctggtg ctggacaacg ctgccattgt ctgcaacttt ggcagtgagc 1680
tcagcctggt gtatgtgccc tctgtgctgg agaagctgga ctgagcgagc ggcctccttg 1740
cccaggcagg aggctggggg gctgtgtggg ggccaatgca ctgaacctgg acttggggga 1800
aagagccgag tatcttccag ccgctgcctc ctgactgtaa taatattaaa cttttttaa 1860
aaaccataaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1920
aaaaaaaaa anaaaaaaaaa nggggg 1946
```

<210> 243

<211> 518

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (298)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (472)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (494)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (500)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (507)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (513)

<223> n equals a,t,g, or c

<400> 243

```
ggacgcgtgg gcagctgcag aaatccaaaa atattgacct tataagctca agatgctgta 60
tgctgggttt ttacaaggat attcgggtatt tgatgaaacc cagttaggca tgacatatct 120
cccttctccc cattcctacc tctcttttat taaaaatttt atagttacag tgtctatgtt 180
gcccagcctg gtcttgaacc ctttggtca agtgatccac ctgccttggc ctcccaaagt 240
gctgggatta cagatgggag ccatcgtgtc tggcccatc ctgcctctta aagccctnca 300
cccaacaggc agacttgaga ctcaaaccgt ggtcccagct cttgcatcag gggtagtttg 360
ggaattcaaa atctttcaaa ctggggtaaa aaccaggccc agaattcaga gttcaaccaa 420
caagctttcc aagaatgccg gcccaaagga aaaaagtggg ttcttggaaa anccggggaa 480
aaggggggtg caanccttn gggagtnaca agngcttg 518
```

<210> 244

<211> 621

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (460)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (569)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (593)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (609)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (612)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (621)

<223> n equals a,t,g, or c

<400> 244

```
gggaagggtcc atgggttgat acctcaggtc aaaaatgtgt ttactctgtt gattgctgtt 60
tcactttact tgtatatcag atatataagc tatgaacaca agtttgtagt aaaagtatct 120
```

tctgtctggg caatggctca cacctgtaat tccaacactt tggggggctc aggtgggagg 180
atctctagtc cccaggagtt tgagaccagc ctgggcaata aactagaccc catgtctcta 240
aaaaatgtaa aaaacatcaa gaggtgagt caggaggatc atttgagctt aggagttaa 300
ggctgcagta agttatgatt gtgccactgc attccagcct gggtgacaag agtgagaccc 360
tgacccaaaa aaagtatctt ctgataaagt ggcatttgaa aattgcagaa aattttaata 420
ttttttcttt taatattaag aactttattg atctctctan gacaccttg tgaggaaatt 480
aatggcactt ttcttgacat tcaacctggg actttgggtg attctgggtt ctggtgaatt 540
taaagtttca acatttcttt ggggcaccnt ttacaagggg tttggccctt ggnaccttat 600
gaccggtcna anggtccaaa n 621

<210> 245

<211> 480

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (340)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (366)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (431)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (447)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (458)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (460)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (469)

<223> n equals a,t,g, or c

<400> 245

ggcacgaggt gaaaccctgt ctctactaaa aatacaaaaa ttagctgggt gtggtggtgg 60
gcgcctgtag tcccagctac tcgggaggtt gaggcgggag tatgacttga acctgggagg 120
cggaggttgc agtgagctga gattgtgcca ctgcactcca gcctgggtga cagagcaaga 180
ctccgtttca gaaaaaaaaa aaaactgtgt ctcatcattg cactgttctt ttaaaattgt 240
atatatatat ttttcttgca gatacactct ggaaatgatt aaactagtac cacataatga 300
aagtgcattg aactatttga aagggatttt gcaggatcgn ggtctttcca aatatcccaa 360
tctgtnaaat caattacctt gatttacaac caagtcaaag ttccccctac ctaattgcct 420
ttcttgggga naccatgaag acatgcnaga aaaccagnn gacaataang gagacatcct 480

<210> 246

<211> 451

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (102)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (107)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (110)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (135)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (198)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (309)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (314)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (319)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (324)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (328)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (330)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (332)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (342)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (349)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (379)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (434)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (438)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (439)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (451)

<223> n equals a,t,g, or c

<400> 246

```
agggcgaccca ccaccatgcc ccggctaaat ttagttgtat ttttagtaga gatgggggtt 60
caccgtatta gccaggatgg tctcgatctc ctgacctcgt gntccgngcn cctcggcctc 120
ccaaagtgct ggganttaca ggcgtgagcc accgtgcccg gcctgattct cttaaaattg 180
aagaggtgct gccaaagnct tcagatctaa cgcagatgca tagaccttgt tcatggtact 240
tgttcagcct gtgctgggga gccgtgggtc cgagtttcct gggaggntga cagggtgaag 300
ccaccgtanc cacnaccnc cganttcncn tncgctttct tntcagcant aggattaaag 360
ggattttgaa tgaagcaant tttaaggggt aggaggtgtt ggggaaaata aataattatt 420
caagtaaggc cttnggggnt ttaggggggt n 451
```

<210> 247

<211> 530

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (29)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (404)

<223> n equals a,t,g, or c

<400> 247

aggaaannag tanctgacag taacgggtcng aattcccggg tcgacccacg cgtccgctgg 60
aactttctcct ggtctctcag ctggggccac tgtcggcatc atgattggag tgctggttgg 120
ggttgctctg atatagcagc cctggtgtag tttcttcatt tcaggaagac tgacagtgtg 180
tttgcttctt ccttaaagca ttgcaacag ctacagtcta aaattgcttc tttaccaagg 240
atatttacag aaaagactct gaccagagat cgagaccatc ctagccaaca tcgtgaaacc 300
ccatctctac taaaaataca aaaatgagct gggcttggtg gcgcacacct gtagtccag 360
ttactcgggg aagctgaggc aggagaatcg cttgaaccg ggantggaga ttgcagtgg 420
cccagatcgc accactgcac tccagtctgg ccaacagagc aagactccat ctcaaaaaag 480
aaaagaaaag gagactctga cctggtactc ttgaatacaa gtttctgata 530

<210> 248

<211> 635

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (217)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (224)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (235)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (408)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (437)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (484)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (513)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (516)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (560)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (568)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (576)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (603)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (635)
<223> n equals a,t,g, or c

<400> 248
ggtggcaggt gcttataatc ccagctactc tggggggccaa ggcaggagaa ttgcttgagc 60
ctgggagatg gaggttgagc tgagctgaga tcatgccact gcactccagc ctgggcaaca 120
gagcaagact ctgcctcaaa aaaaaattaa aataaattta aatacaaaaa aaaatagcca 180
ggtgtgggggt gcatgcctgg aatcccagct acttganaag gctnaaggca ccaanaattg 240
cttgaaccca ggaggtggag gttgcagtga gccaaagatca caggagccac tgcactccag 300
cctgggtgac agagtggagac tctgtctcaa aaaaaaatt aaataaatta ttataacctt 360
tcaaaaatgc tgtgtgcatt ttcattgttct tttttttaac attactgnca ctctccctaa 420
tgaaatggac ttcaanaaag cagtattttg ttaaataaat acataacctc attctgaata 480
atgnccctca ttttgactat aactgggctt ggnttnaaaa gcaaaattta acaaaaattt 540
aatccccttc caaaggaacn ttgggtancc tgggtnaaaa aatgccaaat ggggtacttt 600
tanttaaaat ttggaattta acatcttttt tggan 635

<210> 249

<211> 360

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (118)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (280)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (282)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (305)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (360)
<223> n equals a,t,g, or c

<400> 249
ctcaggaaac catgtccaaa accctagcag cggtagcagca tgctgtctcc aacccttacc 60
cccaggttta aggggtggtt atggccatac gtggaggttt tttgttggtg tttttganac 120
tgagtttcac tcttgttgcc caggctggag tgcaatggca ccactctggc tcaactgcaac 180
ctccacctcc tggttcaagc gatctcaggc ctcagcttcc caagtagttg ggattacagg 240
cgcttgccac cacacctggc taattttgta tttttagtat anatgggggt ctccatgttg 300
gtcangctgg tctcaaaact ccgaactcaa gtgatctgcc cgccttgggc tcccaaaatn 360

<210> 250
<211> 464
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (397)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (436)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (446)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (459)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (460)
<223> n equals a,t,g, or c

<400> 250
gctgtgagga aggggaatcg cttgaacccg ggaggcggag gttgcggtga gccgagatag 60
cgccattgca ctccagcctg ggcaacaaga gcgaaaccct gtctcaaaaa aaaaacaaaa 120
aaccaccaca aaactaggag cctggaaggc cggactgggt ctccgtggga ggggcctggg 180
tctggagagc agggcagggt ctccctgggc taggggatgg ggatggggct ggggtctcaga 240
ggaggcaggg ttacgtgca gaagagcggg cttggtctcc ggggtcccga gtgggtgacg 300
cggcccgccca cagggtgctc ctgaaggtga gccggctgga ggcacaactg ctccctggagc 360
gctaccccga gtgcgggaac ctgctgctgc ggcccancgg ggacggcgcc gacggcgtgt 420
cggtcaccac tcggcngatg ccaacnggaa cgcacgtgnn ccgg 464

<210> 251
<211> 315
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (28)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (37)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (77)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (113)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (202)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (210)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (220)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (265)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (302)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (305)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (311)
<223> n equals a,t,g, or c

<400> 251
aaagtaaaca tcggggaagg tttnccgnaa cgctccnagg taccgggtccg gaattcccgg 60
gtcgacccac gcgtttingct ccttggtcag gctgggtctgg aactggcgac ttnggggtgat 120
ccgcccgcct cggcctccca aagtgggtggg attacaggtg tgagccaccg tgcccagccc 180
tgaaatagtc ttaattgctt gnttttcttn ttgtctgan gtgtgctttt taaaatctct 240
atggagatgg agaagactga cattntctgg cctgatgtga aaaacctctc attaaaaacc 300
gngtntgtag ntttg 315

<210> 252
<211> 333
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (79)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (196)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (200)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (233)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (254)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (302)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (303)
<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (325)

<223> n equals a,t,g, or c

<400> 252

```
gggctaattt gtgtattttt agtggaaacg gggttttgcc acgttgcca ngctgggtctc 60
gaactcctga cctcaggtna tccacccacc ccagcctccc aaagtgctgg gattacaggc 120
gtgagccact gagcccagcc nacttttcag tttttaacat aatttttggt ttatccacaa 180
cttttcaagt attganagtn caataaaaaa catgggttct tagtctgtaa ctntctgtta 240
aagcctatga atnncttctg aaaatcatgt ttttaaatgc ataaaatata tacgattacg 300
anngaattctt attattgtcg aaatncagtt att                                     333
```

<210> 253

<211> 307

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (42)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (48)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (250)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (260)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (281)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (299)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (302)

<223> n equals a,t,g, or c

<400> 253

```
gaaaaanttt taaaggaaga agggaaggca ggctggggag gntggggnaa ggaagcaggg 60
agtcagatc actccccttc catgagttgt tttcttaaga tgctagaact tggccaggca 120
tggtaggctca cccctgtaat ccagcactt tgggaggctg aggctggcag atcacttgag 180
gtcaggagtt ccagaccagc ctggccaaca tggtgaaacc ccatctctac taataatata 240
aaatttagcn ggggtgtgggn ggtgtgcacc tgtggcccta nctacttggg aaggctgana 300
cnaggag                                           307
```

<210> 254

<211> 401

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (256)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (340)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (350)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (359)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (378)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (379)

<223> n equals a,t,g, or c

<400> 254

```
gtataaatgt tacttaaaaa aaatatattt ggccgagcac agtggtcac gcctgtaatc 60
cccacacttt gggaggccga ggcgggcaga ttgcctgagg tcaggaattt gagatcagcc 120
tggccaacat gatgaaaccc cgtctctact aaaaatacaa aaattagccg ggcattggtgg 180
caggcacctg taatcccggc tactcgggag gctgaggcag gaaaatcgct tgaacccggg 240
agtcggaagt tgcagngagc caaggatcatg tcatcattgc actccagctt gggcaacaag 300
agtgaagact tcgtctcaaa aaaaaaactt acagactttn ttttcctttn gtaaattgnt 360
agtttaactt tgaattcnng atgaaggatt tattcttatt t 401
```

<210> 255

<211> 406

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (149)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (243)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (369)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (370)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (376)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (390)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (392)

<223> n equals a,t,g, or c

<400> 255

```
gcttatcatt tgtattaaaa tatggtttta tgtaaagagt tgcctttctc cattcccccc 60
tcccctcccc tcccctcctc tcccctcctc tcttaccctc aattaaaaaa tagcacatcc 120
```

ctaattgtggg aaacagtaag aaagcatana ttacatttta ggcagggcac agtggctcac 180
gcctgtaatc ccagcacttt cggaggctga ggtgggtgga tcacttgagg tcaggagttt 240
ganaccagcc tggccaacat ggtgaaaccc tgtctctact aaaagtacaa aaattagcca 300
ggcatggagg cgggtgcctg taatcccagc tacttgggag gctgaggcag gagaattgct 360
tgagccann aggcanaagg tgcagtgagn anagatcggg ccattg 406

<210> 256

<211> 415

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (27)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (324)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (378)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (379)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (411)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (412)

<223> n equals a,t,g, or c

<400> 256

gacatgaatt ccttaatgat gggagangat aaaatcaagt tcaaacacat cccccccctg 60
caggagcaga gcaaagaggt ggccatccgc atctttcagg gctgccagtt tcgctccgtg 120
gaggctgtgc aggagatcac agagtatgcc aaaagcattc ctggttttgt aaatcttgac 180
ttgaacgacc aagtaactct cctcaaatat ggagtccacg agatcattta cacaatgctg 240
gcctccttga tgaataaaga tggggttctc atatccgagg ggccaagctt catgacaagg 300
gagtttctaa agagcctgcg aaancctttg gtgactttat gggagcccag tttgagtttg 360
ccgtgaaatt ccatggcnng ggattatatg gacagcggct tgggcaaatt nnagc 415

<210> 257

<211> 414
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (102)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (297)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (375)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (380)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (397)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (411)
<223> n equals a,t,g, or c

<400> 257
cgcagaggaa accccgcacc cctggcagaa gttccggacc aagccccagg gggaccagga 60
caccggcaag gaggtgatg acggatgtgc ccttgggggc angtgatggg agcacagctg 120
gaacaatgtg ctcggccccc agtgctctgt gggagcccca ggacaagtga gctggtgtca 180
cctcctgcct gggggaagag ccaggccctg aagaacagcc gcagcgtgtc acaggtgttg 240
gtgaggacac acactaggcc caaggtgcct gtgctcccag caggttccaa gtgcaanttc 300
aagccaactt tgcgtgttaa cttcaacggg gacttccaag cttccaagct aacttttgtg 360
gtgttaacct tcaanaaaan caanaaaggg gggcttnggg gcattttggt ncct 414

<210> 258
<211> 373
<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (321)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (324)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (328)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (351)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (361)

<223> n equals a,t,g, or c

<400> 258

```
ctcccagtcagggtgcagctctccgggtggacacagaatgcagccatgctgaggg 60
cagccggagc caaggccctg agaaagccttcagcccggtcttccatgtgcctggaacgt 120
gtgtgtcacc aggaaggccc cctgctggcctctgacagt agctcctctg ggggtccca 180
cagcaggatggcgaccagaaggcagcgagtccatggatgcggtgagcaggggtcccg 240
ccgggaggcc ccccgctgcc cacagtggcc aagacagaag aagcttttgg caagggttcgg 300
ttttctgaca accggctttt naancctncc ttgccccgcgccaaaagga nttaatttct 360
ngcccaagcgttg 373
```

<210> 259

<211> 529

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (377)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (438)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (443)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (444)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (495)
<223> n equals a,t,g, or c

<400> 259
aaactaacag atgaagaagt agatgaaatg atcagagaag cagatattga tggagacgga 60
caagtcaact atgaagaatt cgtacagaat gatgactgca aaatgaagac ctactttcaa 120
ctcctttttc cccctcttta gaaagaatca aatttgaatc ttttacttac ctccttgcaa 180
aaaaaagaaa aaagaaaaaa ttccatttat tcatccttgt ttcctatata gccaaactga 240
atgttcaaaa gtaccttcct tgtcccacac acacaaatct gcatgtntgg ttgggggggt 300
ccctgtcccc cttaaagatca agctacactt cccattttta caatataata cttgttctac 360
cttatgatag atccctnaaa agtttcccct ttgctaataa ttataacctgt ttgggtggcc 420
agttttccct gcttgcantt gannaatgac ccagccggcc tttgttttaa attgaaatga 480
aaacaattcc aaccttccat ggttcccgtc cattgttaat taatggcca 529

<210> 260
<211> 566
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (361)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (413)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (437)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (460)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (473)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (509)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (562)
<223> n equals a,t,g, or c

<400> 260
actgtctgta tattacggaa gctcttttca cataatatga ctcgtctgcg aaagtttatg 60
gtttactttg gaaagaacca aagtttgagc aagatacaga aaactcctct ctttgtggcg 120
gcgatctgtg ctcattggtt tcagtatcct ttgacccat ctttgatga tgtggctgtt 180
ttcaagtcct atatggaacg cctttcctta aggaacaaag cgacactgaa aattctcaaa 240
gcaactgtgt cctcctgtgg tgagctggcc ttgaaagggt ttttttcatg ttgctttgag 300
tttaatggat ggatggatct cgcagaagca gggggtggat ggaagatgaa gatctaacca 360
ngtgccttga tgagcaaatt ttacagccca gagactaaga ccattctacc ggntttttaa 420
gttccggcct tccaagnaatt tcttggcggg ggatgaggcn ggattgaact ccngggatcc 480
agataggcag gaacatcaag attggggang gtaaccattg gaacaaacca ctcaccagc 540
tgacggtaag cgctacaac antttt 566

<210> 261
<211> 502
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (25)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (44)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (47)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (61)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (72)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (75)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (89)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (98)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (146)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (149)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (151)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (157)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (169)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (176)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (180)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (190)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (194)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (211)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (224)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (231)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (235)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (246)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (247)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (262)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (273)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (334)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (339)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (362)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (375)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (446)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (454)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (477)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (484)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (497)
<223> n equals a,t,g, or c

<400> 261
gggttaaaagt ggantgtatg ttgtnataga agttaaagtt gcanctnatt atggaataga 60
nataacctgt cnaanttatc tgatgacana ttaccaangt gctcccccat cccacagta 120
tagaangatt atttgcatgg gtgcanaana naatggnttg ccgctggant atcaanagan 180
gttaanagcn ttanaaccaa atgactatac ntgaaaggtc tcanaagaaa ntgangacat 240
catcannaag ggggaaacac anactcttta gancataaca gaatatatct aagggtattc 300
tatgtgctaa tatanaatat tattaacact tganaacang gatctggggg atctccacgt 360
tngatccatt ttcannagtg ctctgagagg agtatcttac ttgggggtgac tccttgtttt 420
tagactatac tcagaaactg ggatangggag ttanaccatt taaaacgggt gtatganggc 480
ctgnaatatg tgacaantga at 502

<210> 262
<211> 489
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (3)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n. equals a,t,g, or c

<220>
<221> misc feature
<222> (440)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (464)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (477)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (483)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (486)
<223> n equals a,t,g, or c

<400> 262
gnngacgcgt ggggccgga tgaggagtgg ggtggccact ccccccggag tcccagggga 60
tgggaccagg agcccgccag ggagcaggca ggcgggggct ggcgggccag gcggccccgg 120
gcccgctccg tggacgccct ggacgacctc accccgccga gcaccgccga gtcaggggagc 180
aggtctccca cgagtaatgg tgggaggaga agccgggcct acatgcccc gcggagccgc 240
agccgggacg acctctatga ccaagacgac tcgagggact tcccacgctc ccgggacccc 300
cactacgacg acttcaggtc tcgggagcgc cctcctgccc accccaggtc ccaccaccac 360
cgtacccggg accctcgga caacgntcc aggtccgggg acctccccta tgatgggagg 420
ctactggagg aggtgtgan gaagaagggg tcggacgaga ggangagacc ccacaangag 480
gangangaa 489

<210> 263
<211> 93
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature

<222> (22)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (75)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (76)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (93)
<223> n equals a,t,g, or c

<400> 263
gcggctgcc gccctgcgcg tntccctgc ccgggtctcc tcgctgtcgc ngccgatgcc 60
acaccatggc cttcnnccacc aggcagttca tgn 93

<210> 264
<211> 389
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (45)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (72)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (106)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (178)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (241)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (249)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (309)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (327)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (333)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (344)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (358)
<223> n equals a,t,g, or c

<400> 264
ggcagagtnt gctcttggga agtncaggct cggttgcttt ttggnagcca tggagagtga 60
ctttaatctg cnttactacg tggggcacaa gggcaagtcc ggccangagt tcctggagtt 120
tgaaagtttcg accggacgga gcttgaaatc gtcattggag atgaacacat ttcttttnac 180
aacatcaaaa attggttccc ttattgatgt caatcatgcc aaggatccag aaggcttacg 240

nagtatttna ttatcttgtc caggacctga agtgtttggg cttcagtcctt attgggttac 300
acttcaagnt taaacctct agactgnata ttngtgtggg acangggggg ggggtggngt 360
aggaaatttt gtgtattttc aggggcatt 389

<210> 265

<211> 400

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (296)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (308)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (342)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (343)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (389)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (390)

<223> n equals a,t,g, or c

<400> 265

tttgcccacg cgtccgcccc cgcgtccgcc cagcgcgtcg cccacgcgtc cgcacccccca 60
tcccagtcct agcctacttc cgccacgcag aacctggatt ctccctcaag agggccaggg 120
ggctcagccg gagcctcccc cctccgcccc ctgccaaggg cagcattccc atcagccgcc 180
tcttccctcc tcggaccccc ggctggcacc agctgcagcc ccgcgggtgt cattccgggc 240
gaaggcctag agactctgca gagccctggg tatgacccaa gccgcaaaag tcttntttca 300
agcaaagntt caaaaggtta agccgcttgg gcaatggttt annggaaagg ttttaagggg 360
gcctccaagg gaggcagggc ggtttttttn ggaaaagggg 400

<210> 266

<211> 382

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (10)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (42)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (44)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (49)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (88)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (103)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (109)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (162)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (166)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (185)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (220)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (248)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (278)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (301)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (326)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (341)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (350)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (362)
<223> n equals a,t,g, or c

<400> 266
ggacctggan cggaggcggg aagaaagtca taccagatc cnanatccnc acgcagaatg 60
cacaggtgag ctgccgctgg gcccgganca tgctgggctg tcncacctng cagactgcac 120
gctccaaccg cccgctccac ctactcttc caggcccggc ancttntgga gaaggaattc 180
agcannctta tctccttagg cacagacagg ctgctggacn aggacatgcg ccaagtcttt 240

cagttcgnlc cccatcctgg cggaagatgt tccggganga aggacctccg aggcgtaact 300
ngccgactca ctgagatggt gcccncnaac tttcgttcgg ntgcagcggn attcctgggc 360
tnttcggggg ctccccttttc cc 382

<210> 267
<211> 449
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (374)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (433)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (446)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (447)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (448)
<223> n equals a,t,g, or c

<400> 267
gggcgatggc tgaaggacca ggagctgagc cctcgggagc cgggtgctgcc cccacagaag 60
atggggccta tggagaaatt ctggaataaa tttttggaga ataaatcccc ttggaggaaa 120
atggtccatg gggatataca aaagagtatc tttgttttca ctcatgtact tgtacctgtc 180
tggattatcc attattacat gaagtatcat gtttctgaaa aaccatattg catagttgaa 240
aagaagtcca gaatattccc tgggtatata attctggaga ctggagaagt aattccacca 300
atgaaagaat ttcctgatca acatcattaa agattatgta aaaaggtaaa aggccttatga 360
gcctaagttt ggtncatatat taccatattt actggaattt tcttgaaaa gtactttaat 420
aaagttaaat cttagaaaaa aaaaannna 449

<210> 268
<211> 594
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<400> 268

```
ngcaagcnga caccaaccct cactaaaggg aacaaaagct ggagctccac cgcggg'gcgg 60
ccgctctaga actagtggat cccccgggct gcaggaattc ggcacgaggt cagtgtcggg 120
cgcagacggc ggcagtgcgg cttgctcttg gaagttcagg ctcggttgtc ttttgggagc 180
catggagagt gacttttata tgcgttacta cgtggggcac aagggcaagt tcggccacga 240
gttcctggag tttgagtttc gaccggacgg gaagttaaga tatgccaaca acagcaatta 300
caagaatgat gtcatgatca gaaaagaggc ttatgtacat aaaagcgtga tggaggaact 360
gaagagaata attgacgaca gtgaaattac caaagaggat gatgcattgt ggcctcctcc 420
tgaccgagtg ggccggcagg agcttgaaat cgtcattgga gatgaacaca tttcttttac 480
aacatcaaaa attggttccc ttattgatgt caatcaatcc aaggatccag aaggcttacg 540
agtattttat tatcttgccc aggacctgaa gtgtttggtc ttcagtcctta ttgg      594
```

<210> 269

<211> 332

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (19)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (240)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (275)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (296)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (321)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (332)
<223> n equals a,t,g, or c

<400> 269
gcgggatttg ggtcgcagnt cttgtttgtg gattgctgtg atcgtcactt gacaatgcag 60
atcttcgtga agactctgac tggtaagacc atcaccctcg aggttgagcc cagtgcacacc 120
atcgagaatg tcaaggcaaa gatccaagat aaggaaggca tccctcctga ccagcagagg 180
ctgatctttg ctggaaaaca gctggaagat gggcgacccc tgtctgacta caacatccan 240
aaagagtcca ccctgcacct ggtgctccgt ctcanagggtg ggatgcaaat cttcgngaag 300
accctgactg gcaagaccat nacccttgag gn 332

<210> 270
<211> 515
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (3)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (4)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (7)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (10)
<223> n equals a,t,g, or c

<400> 270
gcnnnganacn aaccctcact aaagggaaca aaagctggag ctccaccgcg gtgcggccgc 60
tctagaacta gtggatcccc cgggctgcag gaattcggca cgagggatgt tgatgtcctg 120
catctaacgc ggtgtaaccc ccgaagccga gcgagctccg gaggaatttc agtatctgct 180
acggtaactt catcagccc ccaagatggc gatgcaagcg gccaaagagg cgaacattcg 240
acttccacct gaagtaaatc ggatattgta tataagaaat ttgccataca aaatcacagc 300
tgaagaaatg tatgatatat ttgggaaata tggacctatt cgtcaaatca gagtggggaa 360
cacacctgaa actagaggaa cagcttatgt ggtctatgag gacatctttg atgccaagaa 420
tgcatgtgat cacctatcgg gattcaatgt ttgtaacaga taccttgtag ttttgtaacta 480
taatgccaac agggcatttc agaagatgga caca 515

<210> 271
<211> 141
<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (14)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (25)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (39)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (141)

<223> n equals a,t,g, or c

<400> 271

cactccatgg gnnnctctgt gtggnatttg gggcccctnc atacttcatt tattatatta 60
tcaaaactga gagggatagg aaagtaaaaa cttatccagg taggaaaatt ggatcgaaca 120
tttcacctct catattaagg n 141

<210> 272

<211> 433

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (32)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (49)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (52)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (94)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (139)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (166)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (177)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (187)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (198)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (244)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (248)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (259)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (274)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (276)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (288)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (355)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (364)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (396)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (402)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (427)
<223> n equals a,t,g, or c

<400> 272
gagcggcggc ggaccgcagc gaagtggcga gntagtggca aacgttgant tntgaagggg 60
ggcccaagat gaccggttct aacgaagttc aagntgaacc agccaccccg aggatagcat 120
ctcctccgta aagttcagnc ccaacacctc ccagttcctg cttgtntcct cctgggncac 180
gtccgtncca ctctacgnat gtgccggcca actccatgcg gctcaagtac cagcacaccg 240
gagncgtntc ggactgcgnc ttctacggtc caangnatgc ctggagtnga ggactagatc 300
atcagttgaa aatgcatgat ttgacactga tcaagaaaat ttcttggacc catgntgccc 360
tatnagatgt gttgaatact gtcagaagtg aattanaatg gnactggaag ttgggggtag 420
ccagtttacc tgt 433

<210> 273

<211> 432
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (187)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (190)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (332)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (371)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (373)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (380)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (430)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (432)
<223> n equals a,t,g, or c

<400> 273
ggttcatctc cgtctcagaa aagagcaagg atcgcggcag caacacgata ggcgcccgcc 60
tgaaccgagt agaagacaag gtgacgcagc tggaccagag gctggcactc atcacgcaca 120
tgcttcacca gctgctctcc ttgcacggtg gcagcaccoc tgagcccact gtgcgtgggg 180
ctcccgnctn caaccctctg cccagtccca gcagccagcc aaacacacag aaggggactg 240
ccacccttccc ttgccagctg ctgagccgca gagaagtgac ggttcctaca caggacaggg 300
gttccttctg ggcattacat cgcataaaaa tnaataattt gtggtgattt ggatctgggt 360

tttaatgaat ntnacagtgn gacttttgatt attaattgag caagcttttc ctaataaacg 420
tggagaattn cn 432

<210> 274

<211> 276

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (95)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (234)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (255)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (264)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (265)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (269)

<223> n equals a,t,g, or c

<400> 274

gatecgttct ctgggtccaa gtctgccagc acggcctccc tgaccatctc tgggctccag 60
gctgaggacg aggctgatta ttactgcagc tcatntacaa gtagtatctc ttatgtcttc 120
ggaactggga ccaagggtcac cgtcctagtc cagcccaagg ccaacccac tgttcaactcc 180
tgtttccccc cctcctcctt aagaacttcc aagcccaaca aaggcaacta tgtnttttgg 240
aaccattact tctanccggg aaanntttna aaatgc 276

<210> 275

<211> 351

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature
<222> (48)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (78)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (87)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (117)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (150)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (154)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (174)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (178)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (206)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (219)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (241)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (248)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (288)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (292)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (303)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (329)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (349)
<223> n equals a,t,g, or c

<400> 275
ccgcagacta gggcgccctcg ggccagggga acgcggaaga gccatggnc cgggctaacg 60
gggcccgtgga aaacgggnac cggacangaa gccgccggcc ctgccgcgcc ccatccncaa 120
cctggaagtc gagttcacca agatatttan caanaatgga atgggacgaa tccnagantg 180
ggaaaaagtg tgctacatgt taccntcaa ctccggganc aaatatgtga agtggaagaa 240
ngagatangg ccgacgtggg acgaaggctg tggaagctgc acagggtngc tnccaaaaag 300
ggntcccca tgggcgcccg gtgggatgnc ctgaattcct tggccggtng c 351

<210> 276
<211> 463
<212> DNA
<213> Homo sapiens

<400> 276
gctgaattct ggctgaccag ggcagtcacc agagctccag acaatgtctg tctccttcct 60
catcttctcg ccggtgctgg gcctcccatg ggggtgcctg tcacaggtgc agctgcagca 120
gtcaggtcca ggactggtga agccctcgca gaccctctca ctcacctgtg ccatctccgg 180
ggacactgtc tctaggaaca gtgctggttg gaactggatc aggcagtccc catcgagagg 240

```
ccttgagtgg ctggaagga catactacag gtccaaatgg tataatgatt atgccgtatc 300
tgtgaaaagt cgaataacca tcaacgcaga ttcaaccaag aatcagttct ctctgcagct 360
gaactctgtg actcccaggg acacggctct gtattactgt gcaagagatc ggggcagctg 420
gtccgatgaa gccgaggggc tcccgcgcg ttacttctac tac 463
```

<210> 277

<211> 463

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (392)

<223> n equals a,t,g, or c

<400> 277

```
ctgtcctcta gagaaaaccc tgtgagcaca gtcctcacc atggactgga cctggaggat 60
cctcttcttg gtggcagcag ctacaagtgc ccactcccag gtgcagctgg tgcagtctgg 120
tgctgagggtg aagaagcctg gggcctcagt gaaggctctc tgcaaggctt ctggatacac 180
cttcaccagt tatgatatca actgggtgcg acaggccact ggacaggggc ttgagtgggt 240
gggatggatg aaccctaaca gtgctaacac aggctatgca cagaagtcc agggcagagt 300
caccatgacc aggaacacct ccataagcac agcctacatg gagctgagca gcctgagatc 360
tgaggacacg gccgtctatt actgtgagag anggaggcgg tgggagctgc tcgggatgat 420
gtgggacttt gactactggg gccagggaac cctggtcacc gtc 463
```

<210> 278

<211> 343

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (172)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (271)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (284)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (293)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (294)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<400> 278
ggcagagggga tctcagggct tctttttctg tcctccacca tcatgggggc aaccgccatc 60
ctcgccctcc tcctggctgt tctccaagga gtctgtggcg aggtgcagct ggtacatgct 120
ggagggagaga tgaggaaagc ccggggagtc tctgaagatc tcctgtgaag gntgttggga 180
tacacctttg aacatctact gggatcggtt ggggtgcgcca gatgcccggg gaaggcctgg 240
gagtgggtgg ggcatacatat gtcctgggtga ntctgatggc cagngatagc canncttttc 300
gagggccaga tgcancatgt gcagtcgaca agtgccacca aca 343

<210> 279
<211> 436
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (121)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (151)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (211)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (326)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (362)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (373)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (392)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (403)
<223> n equals a,t,g, or c

<400> 279
gcaccatggc ttggacccca ctctctctcc tcaccctcct cctccactgc acaggggtctc 60
tctcccagct tgtgctgact caatcgccct ctgcctctgc ctcccctggga gcctcggtca 120
ngctcacctg cactctgagc agtgggcaca ncgactacgc catcgcatgg catcagcagc 180
agccagagaa gggccctcgg tacttgctga ngcttaacac tgatggcagc cacaggaagg 240
gggacgggat ccctgatcgc ttctcaggct ccagctctgg ggctgagcgc tacctacca 300
tctccagcct ccagtctgag gatgangctg actattactg tcagaactgg ggctttggga 360
tngtattcgg cgaaggggac caanctgaac gncctaagtc agnccaaggc tgccccctcg 420
gtcaatctgt tcccgg 436

<210> 280
<211> 315
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (76)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (138)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (224)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (246)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (280)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (308)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<400> 280
ggcacgagct cccccagcct tgctgaagat ccgttccaag gagggcaggt gtgcgcacca 60
tcccgggcta tacagnccat ctgcctgccc tcgatgtata acgatcccca gtttggcaca 120
agctgtgaga tcaactggncct ttggaaaaag gaattctagt gaagtgaaca attgcgaact 180
gaacttagga aggtcctgga ggagtgtttt gacctggaaa atgnagccca gtgtgggttca 240
agggtnagac tgcagagttt agagggtggg agcactgagn cgggtggcaga ttgggtccca 300
gggttggnntt gaagn 315

<210> 281
<211> 411
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (305)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (339)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (358)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (403)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (411)

<223> n equals a,t,g, or c

<400> 281

```
ggcagggttg agccgctgcc gtcgccatga cccgcggtaa ccagcgtgag ctgcccgc 60
agaagaatat gaaaaagcag agcgactcgg ttaagggaaa gcgccgagat gacgggcttt 120
ctgctgccgc ccgcaagcag agggactcgg agatcatgca gcagaagcag aaaaaggcaa 180
acgagaagaa ggaggaaccc aagtagcttt gtggcttcgt gtccaacctt ctggcccttc 240
gcctgtgtgc ctggagccag tcccaccacg ctgcggtttc ctctgtagt gctcacaggt 300
cccancaccg atggcattcc ctttgccctg agtctgcanc gggtcccttt tgtgcttnc 360
tccctcaag tacctntttc cccctgggcc acttccgggg gtnagggggg n 411
```

<210> 282

<211> 570

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (6)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (217)

<223> n equals a,t,g, or c

<400> 282

```
gcaagnngaa antaaccctc actaaaggga acaaaagctg gagctccacc gcggtgcggc 60
cgctctagaa ctagtgatc ccccgggctg caggaattcg gcacgagtag agacagcgcc 120
ggggcaagtg agagccggac gggcactggg cgactctgtg cctcgctgag gaaaaataac 180
taaacatggg caaaggagat cctaagaagc cgagagnaaa atgtcatcat atgcattttt 240
tgtgcaaact tgtcgggagg agcataagaa gaagcaccca gatgcttcag tcaacttctc 300
agagtthtct aagaagtgtc cagagagggtg gaagaccatg tctgctaaag agaaaggaaa 360
atttgaagat atggcaaaaag cggacaaggc ccgttatgaa agagaaatga aaacctatat 420
ccctcccaaa ggggagacaa aaaagaagtt caaggatccc aatgcacca agaggcctcc 480
ttcggccttc ttctcttct gctctgagta tcgccccaaa atcaaaggag aacatcctg 540
```

gctgtccatt ggtgatgttg cgaagaaact

570

<210> 283

<211> 366

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (10)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (32)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (39)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (327)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (333)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (337)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (348)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (357)

<223> n equals a,t,g, or c

<400> 283

```
gntnatgtgn aattaagaca aataaaaacg tnaagcggna caaatcacag agagccacaa 60
agcggatttc acacatgcct agcagaccag aactctcggc agttgctaca agggaagaaa 120
ggactatgtg gatcccttgt ggctatgcag atacctacct cacagagttg ttgtagaaga 180
ctggtgggttt ggttcaaacc ttgtgattaa agagtttgtc aagcatttta ttcttttgaa 240
taaaagcaac atatctaaaa catttaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 300
aaaaaaaaaa aaaaaaaaaa aaagggnggc cgntttnaag gatccaanct tacgaancgt 360
gcatgc                                     366
```

<210> 284

<211> 414

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (389)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (394)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (395)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (398)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (407)

<223> n equals a,t,g, or c

<400> 284

```
ggaagctgag acaggcattc caaggggact ccatcccggt ttctgacctg ctgatcctgg 60
gggtggggcc cgatggtcac acctgctcac tcttcccaga ccaccccccc ctacaggagc 120
gggagaagat tgtggctccc atcagtgact ccccgaaagcc accgccacag cgtgtgacct 180
tcacgctacc tgtcctgaat gcagcacgaa ctgtcatctt tgtggcaact ggagaaggca 240
aggcagctgt tctgaagcgc attttgagg accaggagga aaaccgcgtg cccgccgcct 300
```

gggccagccc cacaccggga aactgtgctg gtcttggacg aggcggccgc cgcttctgac 360
cgtgcccttc gagaaacatt ccactttgna actnngcnca aaggacnccc aaat 414

<210> 285

<211> 551

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (22)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (24)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (64)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (180)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (186)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (225)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (234)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (271)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (296)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (298)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (319)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (322)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (324)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (368)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (410)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (482)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (526)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (528)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (536)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (545)
<223> n equals a,t,g, or c

<400> 285
gtccngcntg agccgcacgg cngnacgctc gtcttcgccc gccatggccg agagcgactg 60
gganacgggtg acggtgctgc gcaagaaggg ccctacggac gccagcaac atccaagcag 120
gctatcttag cggcacaaag actaggagaa gatgtggaga cttccaataa atgggctgcn 180
ggccanaaca aacaacattc tattaccaag aacacggcca agctngaccg gganacagag 240
tgctgcacca tgacaggtga ccctgaagtg ngtaagtga tccagcaagt cggcananca 300
agggcttaca cataacganc tngnnacgaa aatcatgata agccacagtg atcgagact 360
atganagngg tacggccata cccaataacc agtgcttggc aaaatcgacn ggaccattgg 420
ctccaagctc cgggaaagac attggaaaac ccatcgagaa gggcctaggc gaaatgaaca 480
cnaagcctcg aaatcatgag ctccagctga tctcttcgcg cggttcnctg gccgcnattc 540
cgttntcctc a 551

<210> 286
<211> 615
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (3)
<223> n equals a,t,g, or c

<400> 286
gengacacca accctcacta aagggaacaa aagctggagc tccaccgagg tgcggccgct 60
ctagaactag tggatcccc gggtgcagg aattcggcac gagtattgcc tctgaggag 120

tccaactgta tacctgcac agtgtcattc ctttgtgtga tttcttaatg ctgtatttgt 180
tcattctaaa cctagatgta tacagctctg agttataaat ggttataaag ctccctgttac 240
tcattattagt tatttacatc aaaaagcttt tagaaaatgg tacgaggtaa ccaattcttg 300
tcattggtgaa atctgattga gtaaccaagc agttttacta ttctggtgct gcttcataac 360
aaaaatgaaa agctgcatgc atctacagca ggcatggatt gtttatgtcg tatgatattcc 420
tttattaagt aagttcactt atagtatttc tataatttga ttcattgccg taatagagcc 480
atgtaggaaa tgcactgatt gcatgttatt gtggcaagaa taccctaaat gtcattaaaa 540
tcctccaaca tgatggatct acttatggtc ttgtttgttg acatgacaaa ttaacattct 600
tatagttaca tctgg 615

<210> 287

<211> 302

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (221)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (226)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (237)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (286)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (296)

<223> n equals a,t,g, or c

<400> 287

gcggaatgtc acacacattg atcaggcact ccaggaagct catcggtgtc tgaaaccagg 60
aggacggttt ctctgtctgg aatttagcca agtgaacaat cccctcatat ccaggcttta 120
tgatctatat agcttccagg tcatccctgt cctgggagag gtcacgctg gagactggaa 180
gtcctatcag taccttgtag agagtatccg aaggtttccg nctcangaag agttcangga 240
catgatagaa gatgcaggct ttcacaaggt gacttacgaa agtcnecat caggcnttgt 300
gg 302

<210> 288

<211> 414

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (53)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (57)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (63)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (83)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (86)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (91)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (95)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (96)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (100)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (107)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (116)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (147)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (153)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (157)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (158)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (168)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (201)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (205)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (208)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (212)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (216)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (221)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (227)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (233)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (275)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (281)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (292)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (298)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (299)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (318)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (332)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (341)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (354)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (359)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (363)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (364)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (372)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (379)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (385)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (402)
<223> n equals a,t,g, or c

<400> 288
aattcggcac gagagagaga gagagagaga gagagagaga gagagagaga gtngcgntag 60
ttngtgtgtc ctcggtttct cgnctnctgg ntcgnncccn cccacanctg gggcgntcta 120
tgtcgagtgg cgcccatggc gaagagntct canctcnnat gtggaagnct ctcaccttct 180
tcgtcgcgct ccccggggtg ncagncanca tnetgnatgt ntacctnaag tcncaccacg 240
gagagcacga ganaccgag ttcacgtctt acccntatct ncgtatcagg ancaagcnnt 300
ttccctgggg agatngtnac catacttttt tncataacct ntatgtgnat ccanttcnc 360
ttntacgga gntgtatana gagantttgn ccattaccgg gncaacaggg ccca 414

<210> 289
<211> 184
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (7)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (25)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (35)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (101)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (142)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (159)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (175)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (176)
<223> n equals a,t,g, or c

<400> 289
ggacttnant tctntggaat gcatngaatt gcagngacgc attggccctg cccttggcct 60
gcattgatga cgagatggac tgtgagcctc agggcccccgc ncctggccca gctctccgat 120
gtggccatgc acagcctggg tntggcttcc atctatganc agactgacga catcnnggat 180
gttc 184

<210> 290
<211> 207
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (15)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (45)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (63)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (107)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (133)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (149)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (160)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (176)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (200)
<223> n equals a,t,g, or c

<400> 290
ggacgtacng tctnctagt ngaggaatat gtcgagtttc tctanggctg ccccagcaat 60
gggccacttt tgctaaaata tggatatct tagatgggaa aatgcancca cctggcaaac 120
ttgctgctat ggnatctata agacttcang gattacatan acctgcgtac catgcnctga 180
ctgactgtgg ggatcatgtn tgttata 207

<210> 291
<211> 492
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (33)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (437)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (448)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (472)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (474)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (477)
<223> n equals a,t,g, or c

<400> 291
gaagagacat tgaacctgga gaagaaattt ctngttatta tggagatggg ttctttggag 60
aaaataatga gttctgcgag tgttacactt gcgaaagacg gggcactggg gcttttaaat 120
ccagagtggg actgcctgcg cctgctcctg ttatcaatag caaatatgga ctcagagaaa 180
cagataaacg tttaaatagg cttaaaaagt taggtgacag cagcaaaaat tcagacagtc 240
aatctgtcag ctctaact gatgcagata ccactcagga aaaaaacaat gcaacttcta 300
accgaaaatc ttcagttggc gtaaaaaaga atagcaagag cagaacgta acgaggcaat 360
ctatgtcaag aattccagct tcttccaact ctacctcatc taagcttaac tcatataaat 420
aattccaggg taccaanaga aactgaanaa agtctgcaaa gcctttactt tnanagntaa 480
acattgagaa at 492

<210> 292
<211> 237
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (19)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (42)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (65)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (79)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (111)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (123)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (129)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (167)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (192)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (210)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (226)
<223> n equals a,t,g, or c

<400> 292
gggggacatg ccatancnt ccttgggggc ctatggaacc tncaaagcgg ccgtggcgct 60
tactnatgga cacattcanc tgtaaaactcc ttccctgggg ggtcaaggtn naccatcatc 120
cancctggnt gcttcaagac agagtcagtg aagaaacgtg gggttcantgg gaaaagcgcc 180
aacaanttgc tnctggccca accttgtctn aagagcttct gcaggnccta cgggaaa 237

<210> 293
<211> 630
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (580)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (588)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (594)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (608)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (609)
<223> n equals a,t,g, or c

<400> 293
gnttggaag acaccacttg gagtttggca aaacactgct cagagacgag agcctgaata 60
tctttcaaaa cctcaatcgt cgacagcatg agcatgccat ccacatgatg gacattgcaa 120
tcattgccac agacctcgcc ctgtatttca agaagaggac gatgttccaa aagatcgtgg 180
atcagtctaa gacatatgag agttaacagg agtggacaca gtacatgatg ctggagcaga 240
cacggaagga aatcgttatg gccatgatga tgaccgcctg tgatctctca gccatcacca 300
aacctcgga ggtgcagagc caggtagctc tgctggtggc tgctgaattc tgggaacaag 360
gtgacctgga gcgcacggtg ctgcaacaga atcccattcc catgatggac agaaacaaag 420
cagatgaact ccctaagctt caagtcggct tcattgactt tgtttgcacc ttcgtctaca 480
aggaattctc ccgtttccac gagggagatc ccccaatggt ggacgggatc accaacaatc 540
gcaaggagtg gaaggccgct tgctgatgaa tacgatgccn agatgaangt gcangaagag 600
aagaaacnna aaccagcaag tcggccaagt 630

<210> 294
<211> 346
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (12)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (83)

THIS PAGE BLANK (USPTO)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (113)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (205)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (206)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (207)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (220)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (249)

<223> n equals a,t,g, or c

<400> 294

```
nnagcgctcg cnagtggaaac actnattgga atccaaaaaa ttcggacagt gggaaatact 60
ggggtaaatac ctggttaccc canaactata ccctggttga tatgaagatt gantttggtg 120
ttgatataac caccaaagaa atgggttcttg ctgatgattc ctggagactg gccattacga 180
gtattgaagc caacagtaaa gacannnagt cttactgggn gctcaaagaa gtaactcctg 240
aagggtcna aatggtaaag aaaagctttg aggccgggca cggtgactca tgcctgtaat 300
cccagcactt tgggaggcca aggcaggcag atcacccaag gtcagg 346
```

<210> 295

<211> 531

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (206)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (209)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (277)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (286)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (311)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (319)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (320)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (345)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (353)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (374)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (383)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (421)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (423)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (425)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (427)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (430)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (432)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (438)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (441)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (444)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (448)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (462)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (463)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (464)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (471)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (495)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (508)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (519)

<223> n equals a,t,g, or c

<400> 295

```
gttctgtctc gctgatgtca gacctggagg gcaacaccaa gtcacgtgtc cgcctgttgg 60
tcctcgtgcc accctccaaa ccagaatgcg gcatcgagg agagaccata attgggaaca 120
acatccagct gacctgcca tcaaaggagg gtcaccaac ccctccagta cagctggaaa 180
gaagttacaa catcctgaat caagancanc ccctggcccc acccacctca ggttcaacct 240
gttctccctt aaaaaatatt tcccacagaa cacatcnggt ttatantctt gttcctccca 300
gcaataagaa nggaaaccnn ttcttgcaac ttcacggtgg ccttncaaat ctncceccca 360
taaaantttg ccntttttt tgnctccccg ggggggtttt tttttcacc ccccttatcc 420
ntngngncan cnccttang ngcngcnc caggggaaag annnaacccc naaaaaaagg 480
gaaaaccggg caccnggacc ctttgaanac cccacactnt aaaatttcca a 531
```

<210> 296

<211> 572

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (435)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (446)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (451)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (465)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (489)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (501)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (539)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (546)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (570)

<223> n equals a,t,g, or c

<400> 296

ggttctgccc ccactgctta taatgctggt gatctacatt aagatcttcc tggtaggcctg 60
caggcagctt cagcgactg agctgatgga ccactcgagg accaccctcc agcgggagat 120
ccatgcagcc aagtcactgg ccatgattgt ggggattttt gccctgtgct ggtagcctgt 180

gcatgctggt aactgtgtca ctcttttcca gccagctcag ggtaaaaata agcccaagtg 240
ggcaatgaat atggccattc ttctgtcaca tgccaattca gttgtcaatc ccattgtcta 300
tgcttaccgg aaccgagact tccgctacac ttttcacaaa attatctcca ggtatcttct 360
ctgccaaagca gatgtcaaga gtgggaatgg tcaggctggg gtacagcctg ctctcggtgt 420
gggcctatga tctangctct cgcctnttca ngagaagatc aaatncacaa gaaacaaaga 480
ggacacggnt ggttttattg ngaaagatag ctacacctta caaggaaatg gactggctnt 540
tttgancact tcctggaact accacgtatn ta 572

<210> 297

<211> 334

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (44)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (67)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (74)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (84)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (90)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (106)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (108)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (118)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (131)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (167)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (176)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (177)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (219)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (227)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (263)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (292)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (293)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (315)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (316)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (317)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (319)
<223> n equals a,t,g, or c

<400> 297
ggcagaaaaga aaggccctgc tcctccaggg ctccaacgag atcnaaatcc gcgcccagg 60
gcaacanccg ctnacctac agcntcactn tcgatggctg cagantnac accggagncc 120
tggggcaaga nagtgaattg aattacaaaa ccaccaagac ctcccgnntg cccatnnatc 180
gatgtggccc ccttgaacg ttggtgcccc aaaccaggna attcggnttc gaacgttggg 240
ccctgtttgg ttctctgtaa aantcccttc cattcccaac ctggggtccc tnnccaaccc 300
aaccaaattt ttccnnncna acccgggaaa cagg 334

<210> 298
<211> 293
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (23)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (165)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (199)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (237)
<223> n equals a,t,g, or c

<400> 298
atggaggccg ctgactaccg agnggccagc agccagcagg gcctggccta tgccacagag 60

gctgtctatg aaagcgcaga ggccccgggc cactatcccg cagaggacag cacctacgat 120
gagtacgaga acgatctggg gatcacagcc gtcgccctgt acgantacca ggctgcgggc 180
gatgatgaga tctcatttna ccctgatgac atcatcacca acatcgagat gattgangac 240
ggctggtggc gcgggggtgtg caagggccgc ttcagggagc tcgcattctc ttg 293

<210> 299

<211> 418

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (29)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (59)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (71)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (76)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (78)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (114)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (154)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (168)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (213)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (221)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (241)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (260)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (269)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (278)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (284)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (292)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (294)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (313)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (330)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (334)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (346)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (347)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (352)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (354)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (355)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (358)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (373)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (377)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (379)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (383)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (392)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (393)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (408)
<223> n equals a,t,g, or c

<400> 299
aatcggcac gaggggacctg ggcgttttnt tgttggtctt cgtgctgggt ctgggtctna 60
ccccaccgac nttggntnag gataactcca ggtacacaca ctccctgacc cagnactatg 120
atgccaaacc acagggcccg gatgacagat actntgaaag catcatgnng agacggggcc 180
tgacctgacc ctgcaaagac atcaacacat ttnttcatgg naacaagcgc agattcaagg 240
ncatctgtga aaacaaggan tggaaacctt tacagggnaa cctnagggtt angnaagttt 300
ttttttcca ggnaaccatt tggaaagttt aatngggggg ttcccnnggg tntnnaanga 360
ctttccggg gcnaaangng ggnttaaaaa annttttttt ttttttnga aaaagggtt 418

<210> 300
<211> 346
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (41)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (97)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (107)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (277)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (296)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (317)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (320)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (330)
<223> n equals a,t,g, or c

<400> 300
ccacaaacag ttgcatccc tagaacatgg cattgtgcca nttaccagtg attgccagta 60
cctcttcctt gccaaaggttg tctctcgctt ggtgaantgg gtgacanttg cccatgagga 120
ttacatggag ctgcacttca ccaaagacat tgtggatgcg ggactggctg gggacaccaa 180
tctctactac atggcgctca tcgaaagggg cacagccaaa ctgcaggccg ctgtggtgtt 240
gaaccctggc tactcctcca tcccacctgt tttccanctc tgtttgaact ggaaanggga 300
gaaaaccaac agcaacnatn acaacattcn gggccatgga gggcga 346

<210> 301
<211> 213
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (19)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (46)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (60)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (66)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (80)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (119)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (130)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (165)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (170)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (202)

<223> n equals a,t,g, or c

<400> 301

gctttataat ctgctcatnt ttgcctaata aatgcaacat ccacangaac tgaatttctn 60
aaaaanttgg tactaccagn tattggttcg ttacaatta ttgatggaaa tcaagtcanc 120
ggacaaaatn ctggaaacaa ttcttcctt caaaaaatct tatcngcaan aaccgactga 180
agctgccata gatttcctta cnagaattaa ata 213

<210> 302

<211> 440

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (121)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (324)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (342)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (363)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (382)

<223> n equals a,t,g, or c

<400> 302

```
ggcaccgagcg agaccccagc aggcaccatc ctttaccacg ctcatttaga catcgaggcc 60
ttcaccatgg accgggaagt gcgcaaaatc aaacaaggcc tgggcttgaa atttgctgag 120
ntgggtgtata ccggtttctg gcacagccct gagtgtgaat ttgtccgccca ctgcatcgcc 180
aagtcaccagg agcgagtggg agggaaagtg cagggtgtccg tcctcaaggg ccagggtgtac 240
atcctcggcc gggagtcccc actgtctctc tacaatgagg agctgggtgag catggatgag 300
aacctcatgc acatcagcta cgangctgga atcctggaga ancccaagaa ccaagcgctc 360
cangtcttaa acgaagacct angaccagc caaagcccca acaaccctga catctccgag 420
atcgagttca aaaaagggtc 440
```

<210> 303

<211> 495

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (22)

<223> n equals a,t,g, or c

<400> 303

```
ctcataaggg accaaagctg gngctccacc gcggtggcgg ccgctctagc actagtggat 60
cccccgggct gcaggacgcc atgttggggt ttgtgggtcg ggtggccgct gctccggcct 120
ccggggcctt gcggagactc accccttcag cgtcgtgcc cccagctcag ctcttactgc 180
gggcccgtcc acggcgggtc atcctgggtg gagcacagta aggactattg ctatggatgg 240
tacagaaggc ttggttagag gccagaaagt actggattct ggtgcaccaa tcaaaattcc 300
tgttggtcct gagacttttg gcagaatcat gaatgtcatt ggagaacctt ttgatgaaag 360
aggtcccacg aaaaccaaac aatttgctcc cattcatgct gaggtccag agttcatgga 420
aatgagtgtt gagcaggaaa tcttgtgact ggtatcaagg ttgtcgatct gctagcttcc 480
```

tatgccaaagg gtggc

495

<210> 304

<211> 293

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (284)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (286)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (289)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (292)

<223> n equals a,t,g, or c

<400> 304

ggtcagggac aagatgggtgc caccgggtgca ggtctctccg ctcatcaaga ttacctaaaa 60
cctcgggcag aagaggagag gaggatagca gcagaagaga agaagaagca ggatgaactg 120
aaacggattg ccagagaatt ggcagaagat gacagcatat taaagtgagt gaccctgcga 180
cccactcttt ggaccagcag cggatgaata aagcttcctg tgttgtgtga aaaaaaaaaa 240
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaagggg cggnctntnt tna 293

<210> 305

<211> 261

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (36)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (181)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (190)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (205)
<223> n equals a,t,g, or c

<400> 305
ccggaatcgg cctgtngcc cccgcccga ctccanagcc ggctgagaac ggcagcggcg 60
agggctccac gtgcagcgtc tctccgggca ccttcgggtc caggactaca actcccggca 120
gggtgcgcaa aacgaccgcc caaggcaacg gaggtcacg agaatcagca tgattcttca 180
naggctcttn aggttctcct ctgtnattcg gtcagccgtc tcagtcatt tgcggaggaa 240
cattggtggt acagcagttg g 261

<210> 306
<211> 502
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (41)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (147)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (165)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (173)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (213)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (260)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (270)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (274)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (293)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (316)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (324)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (346)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (423)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (436)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (462)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (468)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (488)
<223> n equals a,t,g, or c

<400> 306
gcaacactaa aggacgaacc tgatctctta tactagtatc nttaatcatt tttattgcca 60
caactaacct cctcggactc ctgcctcact catctacacc aaccacccaa ctatctataa 120
tcctagccat ggccatcccc ttatgancgg gcgcagtgat tatangcttt cgntctaaga 180
ttaaanaatgc cctagcccac ttcttatcaa aangcacacc tacacccctt atccccatac 240
tagttattat ggnaaacatn atcctactcn ttcnaccaat agccctgggc gtnagcctaa 300
tcgettacat tactgnaggg cacntactca tgcacctcat tggaancgct ccctacaata 360
tcaatcatta aacttcctc tacacttato tcttcacatt ctattctaac ttaatatcct 420
aanaaattcc ttttcncttt attccaagcc caacttttct cnactttntt cttaaaccct 480
ttactttncg aaaaaccttt tt 502

<210> 307
<211> 119
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (33)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (67)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (76)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (113)

<223> n equals a,t,g, or c

<400> 307

aaccgactat gganttacag ccacaaagca agnggtgaan gctggaacat tcttttggtc 60
tgttgnatc cctcanttgc ggagaatatt aaccatattg cagtggctca cntaccag 119

<210> 308

<211> 403

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (9)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (344)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (368)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (375)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (388)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (389)
<223> n equals a,t,g, or c

<400> 308
gcggacggnc ngngaagggtg cttaagcggc tgcgacttca gaagcgaggc acaggcggtg 60
tggacacggc tgcggtgggc ggggtcttcg acgtctccaa cgctgaccgc ctgggcttct 120
cagagggtgga gctggtgcag atggtggtgg acggagtga gctgctcatc gagatggagc 180
agcggctgga gcaggggccag gccatcgacg acctcatgcc tgcccagaaa tgaagcccgg 240
cccacacccg acaccagccc tgctgcttcc taacttattg cctgggcagt gccaccatg 300
caccctgat gttcgccgtc tggcgagccc ttagccttgc tgnagagat tccgttcaac 360
cttggtanaa ttanttttt tgatgggna agatatgctg atg 403

<210> 309
<211> 559
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<400> 309
aagcngacac caccctcact aaagggaaca aaagctggag ctccaccgcg gtgcggccgc 60
tctagaacta gtggatcccc cgggctgcag gaattcggca cgaggtttat ctcaaggcct 120
gagtagccgg taacaaacga ggggtcccg gattggaccg acgcagccat gcctctgcga 180
cttgatatca aaagaaagct aactgctaga tctgacgag ttaagagtgt ggatctgcat 240
cctacagagc catggatgtt ggcaagtctt tacaatggca gtgtgtgtgt ttggaatcat 300
gaaacacaga cactggtgaa gacatttgaa gtatgtgatc ttctgttcg agctgcaaag 360
tttgttgcaa ggaagaattg gggtgtgaca ggagcggatg acatgcagat tagagtgttc 420
aattacaata ctctggagag agttcatatg tttgaagcac actcagacta cattcgctgt 480
attgctgttc atccaacca gcctttcatt ctaactagca gtgatgacat gcttattaag 540
ctctgggact gggataaaa 559

<210> 310
<211> 521
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (227)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (245)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (494)
<223> n equals a,t,g, or c

<400> 310
ggagatgtnt cgctccgtgg ccttagctgt gctcgcgcta ctctctcttt ctggcctgga 60
ggctatccag cgtaactcaa agattcaggt ttactcacgt catccagcag agaatggaaa 120
gtcaaatttc ctgaattgct atgtgtcttg gtttcatcca tccgacattg aagttgactt 180
actgaagaat ggagagagaa ttgaaaaagt ggagcattca gacttgnctt tcagcaagga 240
ctggncttct tatctcttgt actacactga attcaccccc actgaaaaag atgagtatgc 300
ctgccgtgtg aacctatgtga ctttgtcaca gcccagata gttaagtggg atcgagacat 360
gtaagcagca tcatggaggt ttgaagatgc cgcatttgga ttggatgaat tccaaattct 420
gcttgcttgc tttttaatat tgatatgctt atacacttac actttatgca caaaatgtag 480
ggttataata atgntaacat ggacatgatc ttctttataa t 521

<210> 311
<211> 591
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (52)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (528)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (537)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (557)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (563)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (571)
<223> n equals a,t,g, or c

<400> 311
ggccccctgc aggtaccggc ccggaattcc cgggtacggc tgcgagaaga nnacagaagg 60
ggattcctga agctgacagc attcgggccc agatgtctcg ctccgtggcc ttagctgtgc 120
tcgcgctact ctctctttct ggcttgagg ctatccagcg tactccaaag attcagggtt 180
actcacgtca tccagcagag agtggaagt caaatttcct gaattgctat gtgtctgggt 240
ttcatccatc cgacattgaa gttgacttac tgaagaatgg agagagaatt gaaaaagtgg 300
agcattcaga cttgtctttc agcaaggact ggtctttcta tctcttgatc tacactgaat 360
tcacccccac tgaaaaagat gaggatgcct gccgtgtgaa ccatgtgact ttgtcacagc 420
ccaagatagt taagtgggat cgagacatgt aagcagcatc atggagggtt gaagatgccc 480
gcatttggat tggatgaatt caaattctgc ttgcttggtt tttaatantg atatgcntat 540
acacttacct ttatgcncaa aanggtaggg ntataataat ggtaacatgg g 591

<210> 312
<211> 584
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (4)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (20)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (33)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (35)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (119)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (259)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (262)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (299)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (310)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (327)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (329)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (330)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (350)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (400)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (424)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (437)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (508)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (527)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (535)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (553)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (576)
<223> n equals a,t,g, or c

<400> 312
gagnttcccg ggtncgacgn cacgcgtccg aantnaaat ttttaaaaaa aaaaaaaaaa 60
aaaaaaaaaa aaaagggcgg ccgttctaga ggatccaagc ttacgtacgc gtgcatgcna 120
cgtcatagct cttctatagt gtcacctaaa ttcaattcac tggccgctcg tttacaacgt 180
cgtgactggg aaaaccctgg cgttacccaa cttaatcgcc ttgcagcaca tcccccttcc 240
gccagctggc gtaatagcna anaggcccg accgatcgcc ctccccaaca gttgcgcanc 300
ctgaatggcn aatgggacgc gccctgnann ggcgcatata gcgcggcggn tgaggtggtt 360
acgcncagcg tgaccgctac acttgccagt gccctagcgn ccgctccttt cgctttcttc 420
cctncctttc tcgccangtt cgccggcttt ccccgtaag ctctaaatcg ggggctccct 480
ttagggttcc gatttagtgc ttacgggnac cttcgacccc aaaaaanttg attanggtga 540
tggttcacgt aantgggcca tcgcccttga tagacnggtt tttc 584

<210> 313
<211> 763
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (12)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (52)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (54)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (264)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (297)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (331)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (372)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (431)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (496)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (528)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (530)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (588)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (641)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (657)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (676)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (714)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (730)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (731)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (744)
<223> n equals a,t,g, or c

<400> 313
gaaaaaaaaa naaaaaaaaa aaaaaaaaaa aaaaaaaaaa aagggcggcc gntntagagg 60
atccaagctt acgtacgcgt gcatgcgacg tcatagctct tctatagtgt cacctaaatt 120
caattcactg gccgtcggtt tacaacgtcg tgactgggaa aaccctggcg ttacccaact 180
taatcgccct gcagcacatc cccctttcgc cagctggcgt aatagcgaag aggcccgcac 240
cgatcgccct tcccaacagt tgcncagcct gaatggcgaa tgggacgcgc cctgtancgg 300
cgcattaagc gcggcggttg tgggtggttac ncgcagcgtg accgttacac ttgccagcgc 360
cctagcgcgc gntcctttcg ctttcttccc ttccctttct gccacgttcg ccggctttcc 420
ccgtcaagct ntaaatcggg ggctcccttt agggttccga tttagtgtt tacggcacct 480
cgaccccaaa aaactngatt aggggtgatg ttcacgtagt gggccatngn cctgatagac 540
ggtttttcgc cctttgacgt tggagtcac gttcttaata gtggactntt gttccaaact 600
ggaacaacac ttaaccctat ctcgggctat tcttttgatt nataagggat tttccnatt 660
tcgggcctat tggttnaaaa aatgaagctt gttttaacaa aaaatttaac gcgnaatttt 720
acaaaaaatn nttaaacggc ttanaattta agggggggcc ctt 763

<210> 314
<211> 761
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (76)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (103)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (109)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (119)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (155)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (225)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (248)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (251)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (255)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (257)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (297)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (339)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (345)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (367)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (371)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (386)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (397)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (437)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (441)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (447)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (460)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (467)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (483)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (510)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (526)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (527)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (528)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (535)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (545)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (557)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (567)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (581)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (596)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (611)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (632)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (644)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (651)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (653)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (657)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (660)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (664)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (675)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (678)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (692)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (723)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (744)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (756)
<223> n equals a,t,g, or c

<400> 314
gccccacgcgt ccgcccacgc gtccgaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
aaaaaaaaaa aaggngggcc gctctaaagg atccaagctt acntacgcnt gcatgcaang 120
tcatanctct cctatagtgt cacctaaatt caatncactg gccgtcgttt tacaacgtcg 180
tgactgggaa aaccctggcg ttaccaact taatcgctt gcaanacatc cccctttcgc 240
cagctggngt natancnaaa aggcccgcac cgatcgccct tcccaacagt tgcgcancct 300
gaatggcaaa tgggacgcgc cctgttacg cgcattaanc ccgnggggtg tgggtggttac 360
ccccaangtt naccgctaca cttgcnagcg ccctagngcc cgctcctttc gctttcttcc 420
cttcctttct cgccaanttc ngccggnttt ccccgtaaan ctctaantcg ggggcccctt 480
tanggttcca atttatgctt tacggcaccn caacccaaaa aacttnnnta aggtnatggt 540
ccacntatgg gccatcncct aataaanggt ttccccctt ngaaattgga atccanttct 600
taaaagggaa nctgttccaa atggaaaaaa anccaaccct atcncgggct ntncctngan 660
ttanaaaggg aattnggnca attccggcct antgggttta aaaaattgaa actgaattta 720
aanaaaaatt taaacgcaaa attntaaaaa aaattnttta a 761

<210> 315
<211> 540
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (40)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (46)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (47)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (109)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (252)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (367)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (394)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (422)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (432)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (478)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (484)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (488)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (504)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (528)
<223> n equals a,t,g, or c

<400> 315
gattttttat aaaccacggt tgcagccaaa aaaaaaaan aaaaannaaa aaaaaaaaaa 60
aaaagggcgg ccgctctaga ggatccaagc ttacgtacgc gtgcatgcna cgtcatagct 120
cttctatagt gtcacctaaa ttcaattcac tgcccgctcg tttacaacgt cgtgactggg 180
aaaaccctgg cgttacccaa cttaatcgcc ttgcagcaca tcccccttc gccagctggc 240
gtaatagcga anaggcccg accgatcgcc ctccaaca gttgcgcagc ctgaatggcg 300
aatgggagcg gccctgtagc ggcgcattaa gcgcggcggg tgtggtggtt acgcgcagcg 360
tgaccgntac acttgnaagc gccetaacgc ccgntccttt cgctttcttc ccttcctttc 420
tnccacgttc gnccggtttc ccgtcaagct ctaaactcgg ggctccttta aggggtccnaa 480
ttantggntt tacgggacct tgancccaaa aaaacttgat ttaggggnga aggggttcacg 540

<210> 316
<211> 513
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (265)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (420)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (468)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (478)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (483)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (496)

<223> n equals a,t,g, or c

<400> 316

```
gtctttctgt taatttcct ctgcctaaag actacatgac agaaatgacc taccactact 60
tattatttct gaagcctaac tgcaagactg atttctgaga acaagtaaag aactggaata 120
cttatttttc atataaaaat ctaaatgtgt taataaatca ttccatacaa aagtacatta 180
ttaaataacc acattattaa aataaaaaaa aaaaaaaag ggcggccgct ctgaggatc 240
ccaagcttac gtacgcgtgc atgcnacgtc atagctcttc tatagtgtca cctaaattca 300
attcactggg ccgtcgtttt acaacgtcgg tgactgggaa aaccctggcg ttacccaact 360
taatccgcct tgcagcacat ccccttttcg ccagctggcg taatagcgaa gaggcccgcn 420
ccgatcgccc ttccaacag ttgcgcgtgc tgaatggcga atgggacncg ccctgtancg 480
ggngcattaa agcgcngcgg gtgtggtggt tac 513
```

<210> 317

<211> 389

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (9)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (60)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (70)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (72)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (75)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (105)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (111)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (248)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (282)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (296)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (342)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (349)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (357)

<223> n equals a,t,g, or c

<400> 317

```
gcccacgcnt ccgaaaaaaaa aaaaaaaaaa naaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
ggcgggccgn tntanaggat ccaagcttac gtacgcgtgc atgcnacgtc ntagctcttc 120
tatagngtca cctaaattca attcactggc cgtcgtttta caacgtcgtg actgggaaaa 180
ccctggcggt acccaactta atcgccctgc agcacatccc cctttcgcca gctggcgtaa 240
tagcgaaanag gcccgcaccc atcgcccttc ccaacagttg cncagcctga atggcnaatg 300
ggacgcgccc tgtagcgggc cattaagcgc ggcgggtgtg gnggttacnc ncagecngac 360
cgttacactt gccagcgccc tagcgccc                                     389
```

<210> 318

<211> 388

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (14)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (143)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (144)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (146)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (150)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (154)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (159)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (160)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (163)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (168)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (171)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (174)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (196)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (198)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (223)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (226)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (227)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (234)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (240)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (242)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (255)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (280)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (314)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (359)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (366)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (368)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (377)
<223> n equals a,t,g, or c

<400> 318
nnnatctctc tggnggaccc aatttacata ggaaacagct atgaccatga ttacgccaaag 60
ctctaatacg actcactata gggaaagctg gtacgcctgc aggtaccggt ccggaattcc 120
cgggtcgacc cagcggtccg gtnnanaccn atgngggggn agntctcngg nggnaaggaa 180
aatcgtgcaa agaggnanta atgaatgtgg atcacgagga tancnntta gtgnaggaan 240
tncatcgttt gggntntaaaa aatgctgatg gaaagttaan tgtgaaaatt ggggtcctct 300
ttcgtgatga taantgagcc aacctcttg aagcattggt aggaactctt aaagctgtna 360
aacgananaa gaatgtncat atccagga 388

<210> 319
<211> 418
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (235)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (240)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (285)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (380)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (391)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (392)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (396)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (401)
<223> n equals a,t,g, or c

<400> 319
gatatttggt acttttagtt ctataagtga ggaagagttt atggcaaaga tttttggcac 60
tttgttttca agatggtggt atcttttgaa ttcttgataa atgactgttt ttttctgcct 120
aatagtaact ggtaaaaaa caaatgttca tatttattga ttaaaaatgt ggttgcttaa 180
ttcctaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaggcg gccgntctan 240
aggatccaag cttacgtacg cgtgcatgcg acgtcatagc tcttntatag tgcacctaa 300
attcaattca ctggccgctg ttttacaacg tcgtgactgg gaaaaccctg gcgttaccca 360
acttaatcgc cttgcagcan atcccccttt nccanctgg ngaataccaa aaggcccg 418

<210> 320
<211> 684
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (41)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (251)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (570)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (594)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (596)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (606)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (670)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (684)
<223> n equals a,t,g, or c

<400> 320
cgggaaattc ncaaagctgt ttccgcctgc aggtcccgggt ncggaattcc cgggtcgacc 60
cacgcgtccg natccccagt gtgaggtaa tgagcaaaat tggctcttgg ggcccttttt 120
gtccaagccc cactgaaagg cctcttcaga aaactattat cttaaagcc ctactttaac 180
tccttaattc cagcatacag ctaaaactgg atgtatatc ttggcaagtaa aggctgagga 240
ctcctcttta ntcctcagat ctagataact catgacattt tatttgacca acatagcaca 300
tgatgagata tcaaggtaat taaaatagca tgcttgaaaa aaaatacgta atctgtttca 360
cctgtaactg ttaagccaa taaacttttc aaaattttaa aaaaaaaaaa aaggcgcgcc 420
gctctagagg atccaagctt acgtacgcgt gcatgcgacg tcatagctct tctatagtgt 480
cacctaaatt caattcactg gccgtcgttt tacaacgtcg tgactgggaa aaccctggcg 540
ttacccaact taatcgctt gcagcacatn cccctttcgc agctggcgta atancnaaga 600
ggcccnaccg atcgcccttt ccaacagttg cgcagcctga atggcgaatg ggacgcgccc 660
tgtagccggn cattaagcgc cggn 684

<210> 321
<211> 216
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (10)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (25)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (108)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (155)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (183)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (190)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (192)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (202)

<223> n equals a,t,g, or c

<400> 321
cncnagaccn nctttggaaa cgccngacta tagggaaagc tggtagcct gcaggtagcg 60
gtccggaatt cccgggtcga ccacgcgc cgcgcgctg gaggctgnc gaaggatgct 120
ggggataagc ccttaggcac cagcttagac acctncaaga accaggcccc gctgatgcaa 180
ganggcagan cngataccca tnagagcccc gagaat 216

<210> 322
<211> 557
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (40)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (98)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (286)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (394)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (415)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (509)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (545)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (546)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (550)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (555)
<223> n equals a,t,g, or c

<400> 322
ggtccagaag cctccccgac cccccaagct atttgetcan attaacaaat taaagtgcct 60
gaagcataat tcatttttgt atctgtggta acaaaacntt aaccaaaaga ttttctgtcc 120
agaagcctcc ccgaccccc aagctatttg ctcacattaa caaattaaag tgcctgaagc 180
ataattcatt ctttacctgt atactaaaaa ccctgttgta ttgatttttt tataataagc 240
ctttttacct ctgtgtaaaa aatataatata caagtgtatg atgtanattt tagttcttaa 300
ctttttttta tggtttctaa tatgtatgac caatgtagcc attgctttta aatgtaccgt 360
gtaaatataa acacatccta tcagaaaaaa aaanaaaaaa gggcggccgc tctanaggat 420
ccaagcttac gtacgcgtgc atgcgacgtc atagctcttc tatagtgtca cctaaattca 480
attcactggc cgctgtttta caacgtcgng actgggaaaa accctgggagc ttaccaact 540
taatnnggcn ttgcngg 557

<210> 323
<211> 145
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (7)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (15)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (19)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (56)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (139)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (140)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (142)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (145)

<223> n equals a,t,g, or c

<400> 323

ggtanengca gtacngtcng aattcccggc tctagagatc caagcttacg tactgngcat 60
gcacgtcata gctcttctat agtgtcacct aaattcaatt cactggccgt cgttttacaa 120
cgtcgtgact gggaaaaacnn tntan 145

<210> 324

<211> 353

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (28)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (43)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (133)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (146)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (216)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (219)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (225)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (232)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (235)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (263)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (332)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (334)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (349)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (353)
<223> n equals a,t,g, or c

<400> 324
nnnattcatc atagccacca tcaccctnct taacctctac ttntacctac gcctaatacta 60
ctccacctca atcacactac tccccatata tagcaacgta aaaataaaan gacagtttga 120
acatacaaaa ccnaccocat tcctcnccac actcatcgcc cttaccacgc tactcctacc 180
tatctcccc tttatactaa taatcttata aaaaaanaana aaaangggcg gncgntctag 240
aggatccaag cttacgtacg cgngcatgcg acgtcatagc tcttctatag ggtcacctaa 300
attcaattca ctggcgcgctg tttacaacg tngngactgg gaaaaccng gcn 353

<210> 325
<211> 553
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (359)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (381)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (406)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (448)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (461)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (481)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (513)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (524)

<223> n equals a,t,g, or c

<400> 325

```
ggcagcagga caaagccttg ctctgttccc caggctggag tgcattggtg caatcatggc 60
tcactgcagt ctcaaccttc cggactcaag tgatgggtccc gcttcagcta cttgggaggc 120
tgaggcagga gaatcgcttg aaccgaggag gcggaggttg cggggagccg ggattgtgcc 180
actgcactcc accctagagt gagactccct ctcaaaaaaa aaaaaaaa actcgagggg 240
gggcccggtg cccaattcgc cctatagtga gtcgtattac aattccactg gccgtcgttt 300
tacaacgtcg tgaactggaa aaccctgggc gttaccaaac ttaaatcgcc ttgcagcana 360
tcccctttcg ccagctgggt naattagcga agaggccgc accgantcgc cttcccaaca 420
gttgcgagtg ggaatggcga atggcaantg taagcttaaa ntttgtttaa aattcgcggtt 480
naaatTTTTT gttaaatacag ctcatTTTTT aancataagg ccgnaatcgg gaaaatccct 540
tattaaatcc aaa                                     553
```

<210> 326

<211> 628

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (434)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (435)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (437)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (464)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (592)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (602)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (613)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (622)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (625)
<223> n equals a,t,g, or c

<400> 326
gcntgcntcc aagtccatag gtcctgcctc ttcaatcctg gctttctagg gcctgggatg 60
atcattgcta gaactgagag accagcctgg ctagaacttc agggcggtcc attcattctt 120
tcagtaaatg tttgcagcac atgtgttaca tgtcaggcag tgaaaccccc cacagcagcc 180
ttccctctca gaggatacat ttgtaacctat tacacagtca tcagaggaat aatttttttt 240
aatcaccagt gtgcacacag tcatggagtt gggatttccc agctaccagg gaggctgagg 300
tgggaggatt gcttgatgcc aggagttagg gaatatagt caccgtgatt ggacttgcca 360
atagccactg cactgcggcc tggacgacgt agtgataccc tgactcttat aaataaataa 420

```
atgaataaac acannanaaa aaaaaaaggg cgcccgctct agangatcca agcttacgta 480
cgccgtgcat gcgacgtcat aactcttcta taggggcacc taaattcaat tcaactggccc 540
gccgtttaca acgtctgact gggaaaaccc tggcgttacc caacttaatc gnettgacaga 600
anacccccctt ttnccaactg gnggnata 628
```

<210> 327

<211> 416

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (22)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (254)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (258)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (260)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (265)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (285)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (297)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (299)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (300)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (305)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (334)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (349)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (416)
<223> n equals a,t,g, or c

<400> 327
gatgactgat gttattatac antatggtaa atacaaagtt gtagtattat gggtcttgat 60
gggtgtgattt acttatgtaa attatgtaca aatttaagtt caatgaaaca ctgatttaaat 120
gtactgaaag gtaaatagta ccatagtgaa ttattttctg tcttgcccta gggttatttt 180
tattatcttc aatattagaa taaatgctaa attattttca aaaaaaaaaa aaaaaaaaaa 240
aaaaaaaaaa gggnggangn tctanaggat ccaagcttac gtacncgtgc atgcaangnn 300
atagntcttc tatagtgtea cctaaattca attnactggc cgtcgattna caacgtcgtg 360
actgggaaaa ccctggcggt acccaactta atcgccctgc agcacatccc cctttn 416

<210> 328
<211> 629
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (227)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (228)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (229)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (233)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (242)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (243)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (244)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (257)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (271)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (275)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (284)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (301)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (394)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (484)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (512)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (544)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (554)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (573)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (598)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (619)
<223> n equals a,t,g, or c

<400> 328
tttttttttt ttttttgggc gctgccgtat cgttccgcct gggcgggatt ctgacttaga 60
ggcggttcagt cataatccca cagatggtag cttcgcccca ttggctcctc agccaagcac 120
atacaccaaa tgtctgaacc tgcggttcct ctcgtactga gcaggattac catggcaaca 180
acacatcatc agtagggtaa aactaacctg tctcacgacg gtctaannna aaaaaaaaaa 240
annnaaaaaa gaattcnaaa agctttctcga nagtncttct aaancggccg cgggcccac 300
nattttccac ccgggtgggg taccaggtaa gtgtacccaa ttcgccctat agtgagtcgt 360
attacaattc actggccgctc gttttacaac gtcntgactg ggaaaaacct ggcgttacct 420
aacttaatcg ccttgacgca catccccctt tcgccagctg gcgtaatanc gaaaaaggcc 480
gcancgatcg cccttcccaa cagtttgccg ancctgaaat ggcgaatgga aatcaatttt 540

taantgttta atgntgttaa actactgaat ccnaattggt tgggtgtttt taaaatcnca 600
gtcccaaggg tcatttcang gccctcaat 629

<210> 329

<211> 458

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (20)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (86)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (96)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (194)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (226)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (238)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (262)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (267)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (277)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (280)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (285)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (295)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (332)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (333)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (339)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (346)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (352)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (356)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (372)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (380)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (405)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (414)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (452)
<223> n equals a,t,g, or c

<400> 329
aatntaaaaa aaaaaaaaaa aaaaaaaaaa naaaaaaaaa agggcggccg ctctaaagga 60
tccaagctta cgtacgcgtg catgcnacgt catagntctg ctatagtgtc acctaaattc 120
aattcactgg ccgtcgtttt acaacgtcgt gactgggaaa accctggcgt taccctaactt 180
aatcgccctt cagnacatcc ccctttcgcc agatggcgta atagcnaaaa ggccccgcnc 240
gatcgccctt cccaacagtt gngcagntctg aatggcnaaa gggangcgcc ctgtngcggc 300
gcattaagcg cggcgggtgt ggtggttacg cnnagggtna ccgctncaact tnccancgcc 360
ctagegcccg gnccttttcgn tttcttccct tcctttctcg ccacnttcgc cggntttccc 420
cgtcaagctc taaatcgggg ggtccctttt anggttcc 458

<210> 330
<211> 453
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (78)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (83)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (113)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (134)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (323)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (327)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (391)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (418)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (427)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (428)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (453)
<223> n equals a,t,g, or c

<400> 330

gccccatatac cttctgcata atgaattaac aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
aaaaaaaaagg gcggccgntc tanaggatcc aagcttacgt acgcgtgcat gcnacgtcat 120
agctcttcta tagngtcacc taaattcaat tcaactggcg tcgttttaca acgtcgtgac 180
tgggaaaacc ctggcggtac ccaacttaat cgccttgacg cacatcccc tttcgccagc 240
tgggcgtata gcgaaaaggc ccgcaccgat cgccttccc aacagttgag cagcctgaat 300
ggcggaatggg acgcgccttg tancggngca ttaagcgagg cgggtgtggt ggttacgcgc 360
agcgtgaccg ttacacttgc cagcgcccta ncgcccgcgc ctttcgtttc ttcccttnc 420
tttcggnnac gttcgccggt ttcccggtta agn 453

<210> 331

<211> 498

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (356)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (361)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (381)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (393)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (424)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (433)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (452)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (483)
<223> n equals a,t,g, or c

<400> 331
gagaaagctg actaatccaa agggcaaggt ggtttacctt aaagaaaaa aaaaaaaaag 60
ggcgcccgct ctagaggatc caagcttacg tacgcgtgca tgcgacgtca tagctcttct 120
atagtgtcac ctaaattcaa ttacttgcc gtcgttttac aacgtcgtga ctgggaaaac 180
cctggcggtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag ctggcgtaat 240
agcgaagagg ccgcaccga tcgcccttcc caacagttgc gcagcctgaa tggcgaatgg 300
gacgcgcctt gtacggcgca ttaagcgcg cgggtgtggt ggttacgccc agcgtnaccg 360
ntacacttgc cagcgcccta ncgccccgct ccntttcgtt ttctttccct ttccttttct 420
tggncacggt tcncggggtt tttccccgt tnaagcctt ttaaaatcng ggggggcttc 480
ccntttaaag ggggttccc 498

<210> 332
<211> 450
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (278)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (317)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (328)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (335)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (338)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (347)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (353)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (358)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (368)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (373)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (374)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (377)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (378)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (386)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (392)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (395)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (397)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (415)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (417)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (426)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (428)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (430)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (446)

<223> n equals a,t,g, or c

<400> 332

aattcggcag aggtactcct ggcttatcaa tggaacattc cagcaaagca cacaagagct 60
ctttatccct aacatcactg tgaataatag tggatcctat acctgccacg ccaataactc 120
agtcactggc tgcaacaggg ccacagtcaa gacgatgcat agtcactgag ctaagtccag 180
tagtagcaaa gcccctaaatc aaagccagca agaccacagt cacaggagat taaggactct 240
gtggaacctg gacctgggtc cacaaaatgg acactgggnat cttccatccc gttggttctt 300
tcaaaaaacc aggggtnttc ccgtcctnng ggggnttnga aggtttntcc ccnggggnaa 360
aaaaccancc tttnngnntt aaancnttt tnaangnggg ggggtttttt gggangngtt 420
ttgggntntn ggggttttta aaccntttg 450

<210> 333
<211> 434
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (58)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (119)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (195)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (214)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (218)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (269)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (291)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (296)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (316)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (333)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (337)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (382)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (399)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (405)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (425)
<223> n equals a,t,g, or c

<400> 333
gntgaacgta aactataatg ctctaccaca agaaaatggc ctctcacctg gggccatngc 60
tggcattgtg attggagtag tggccctggt tgctctgata gcagtagccc tggcatgtnt 120
tctgcatttc gggaagaccg gcagctcagg accactccaa tgacccacct aacaagatga 180
atgaagttac ttatnctacc ctgaactttg aagnccanca acccacacaa ccaacttcag 240
cctccccatc cctaacagcc acagaaatna ttattccag aagtaaaaaa ncagtnatga 300
aacctggtcc tgctcnctgc agtgctgat gtntttncaa gtctctcacc ccccatcact 360
aggaaattcc tttcccctgt tngggtaaag ggtggggana gaaancaatt tctctactc 420
tcctncctat agga 434

<210> 334
<211> 612
<212> DNA
<213> Homo sapiens

<220>

<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (437)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (455)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (474)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (482)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (493)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (497)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (525)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (540)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (565)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (583)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (606)

<223> n equals a,t,g, or c

<400> 334

```
ggcttagcca caacaggggc aatgctcgcc ttgcaggtgc cccagcgag gacccccagt 60
tccccaaggt gcagtggcca ccccgatgaac ttgtttctgc ctgccacaat gaacgcctgg 120
atgtgcccgt gtgggacgtg gaagccaccc tcaacttcct caaggcccac ttctcccca 180
gcaacatcat cctggacttc cctgcagctg ggtcaacttg cccgagggat gtgcagaatg 240
tggaagccg ccccaactg gcgatgggag ccctggagct ggaaagccgg aattcaactc 300
tggaacctgg gaagcctgag atgatgaagt ccccccacaa caccaccca catgtgccg 360
ctganggacc tgaggcaagt cgacccccga agcttgccacc ctggcctaaa acttgaccag 420
gccaggaacc ctctgancac aatggcaaaa gcttnaaagg aatgaaccag gaancaagcc 480
gntttgggca atnggggnact ttgagccaaa gccgaagaac accanggggc ttggatttgn 540
tttgcttta atttccaagg ggttnaaaaa aaaaaacccg gcnttttttg gggggccccc 600
tttttnggaa gg 612
```

<210> 335

<211> 462

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (337)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (358)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (373)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (428)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (456)
<223> n equals a,t,g, or c

<400> 335
gcccacgcgt ccgcatgttt ccaggagggg ccatgggggt cagggtggtt ttgagagagc 60
agggtaagga aggaatgtgc tagaagtgtc cctagtttct tgtaaaggaa gccagagttg 120
acagtacaaa gggtcgtggc cagccctgca gcttcagcac ctgccccacc cagagtggga 180
gtcagggtgga gccacctgct gggctcccc agaactttgc acacatcttg ctatgtatta 240
gccgatgtct ttagtggtga gcctctggat tctgggggtct gggccantgg ccatagttaa 300
acctgggaat gaatggtact tgaacatctg ggcttgncag ccacaaggaa aggccaancc 360
catgtacccc aancattctt gccaacctt gctctgggca ttcccggaag ggtcgatcct 420
taggcttngc tttaaaagcc cttgcccttg cctttntctt gg 462

<210> 336
<211> 336
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (170)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (175)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (177)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (183)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (196)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (204)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (208)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (230)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (262)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (275)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (293)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (318)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (335)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (336)
<223> n equals a,t,g, or c

<400> 336
gcagattttg atcgttttaa agttatgaag gcaaagaaaa tgaggaacag aataatcaag 60
aatgaattaa gaagcttcaa aaggcagctc tcctgaaagc ttctcccaaa aaacacctgg 120
tactaagggt actgctgctg ctgctgctgc tgctgctgct gctgctaaan tccancnaaa 180
aanatcaccg ccganttaa aagntccncc caaaagtcct gcccaaaaan cccaggccaa 240
aaacacgctg ctccaaactc anaggtcaaa actcncccaa aaccctctcc aangctctgc 300
aaaaagctta ttgcatcnta aaattataag gtctnn 336

<210> 337
<211> 303
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (25)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (28)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (37)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (49)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (50)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (64)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (66)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (67)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (70)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (120)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (168)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (189)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (190)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (192)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (223)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (249)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (256)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (257)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (267)
<223> n equals a,t,g, or c

<400> 337
agacaatcct taccatgtga naggnggnat gactgtnatg gcatgagann ngcatcacta 60
tganannngan aactcagtag aagataatgg caagtccaga ctggggatat gatgacaaan 120
atggtcctga acaatggagc aagctgtatc ccattgccaa tggaaatnac cagtcccctg 180
ttgatattnn gnccagtga accaaacatg acacctctct ganacctatt agtgtctcct 240
acaacccanc cacagnnaaa gaaattntcc aatgtggggg cattccttcc atgtaaattt 300
ttg 303

<210> 338
<211> 460
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (13)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (409)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (423)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (424)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (433)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (454)
<223> n equals a,t,g, or c

<400> 338
gattttaaggc atntcctgca tgcacagttg cagttagtta ttccaggtat tatttttgtt 60
ttcagaaaaa gaaaactcag tagaagataa tggcaagtcc agactgggga tatgatgaca 120
aaaatggtcc tgaacaatgg agcaagctgt atcccattgc caatggaaat aaccagtccc 180
ctgttgatat taaaaccagt gaaaccaaac atgacacctc tctgaaacct attagtgtct 240
cctacaaccc agccacagcc aaagaaatta tcaatgtggg gcattccttc catgtaaatt 300
ttgaggacaa cgatnaccga tcaagtgtg aaaggtggtc ctttctctga cagctcaggc 360
tctttcagtt ccattttcac tggggcagtc aaaatgagca tgggttaana acattccagt 420
ggnntgggag tcnaaatatt ctggccgagc ttnnacgtaa 460

<210> 339
<211> 416
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (30)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (33)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (42)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (137)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (147)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (161)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (173)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (258)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (267)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (268)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (378)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (403)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (407)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (411)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (415)
<223> n equals a,t,g, or c

<400> 339

gggtgccgatt cctgccctgc cccgaccgcn agnccggacca tntcccatca ntgggggtac 60
ggcaaacaca acggaccta gacactggcat aaggacttcc ccattgccaa gggagagagcg 120
ccagtncccc tgttganatc gacactncat acagccaagt ntgaaccctt ccntgaaagc 180
ccctgtttgt ttccatgaa tcaagcaact tccctgagga tcctcaacaa tggcatgct 240
ttcaacgtgg gagtttgnat gactctnnag gacaaagcag tgcttcaagg gaaggacccc 300
tgggttgggc actttacaga ttggttttca ttttttcaat tttcaatggg ggtttcaatt 360
tgatgggaca aggtttcnga ggcataattg tgggttaaaa agnaatntgg ntgcng 416

<210> 340

<211> 362

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (76)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (117)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (136)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (170)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (201)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (204)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (233)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (310)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (331)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (332)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (335)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (338)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (339)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (357)

<223> n equals a,t,g, or c

<400> 340

ggcanagtga atgttgctgg agaacaaggc cagcattttt ggaggaggac tgcctgcccc 60
ataccaggtc aaacanttgc acctgcactg gtccgacttg ccatataagg gctcggngca 120
cagcctcgaa tgggngcac ttgccatgg gagatgcaca tagtacatgn gaaaagagaa 180
ggggacatcg aggaatgtga naanagggcc caggaaccct gaagacgaaa ttncggtgct 240
gggccttttt ggtgggaggc tgggaaccca ggtgaaacga gggctttcca gccactggtg 300
ggaggcactn tcttaatat ccccaaacct nnggntgnng cattacgatt ggcaganagg 360
ca 362

<210> 341

<211> 328

<212> DNA

<213> Homo sapiens

<220>
<221> misc feature
<222> (63)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (98)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (152)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (171)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (184)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (195)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (276)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (280)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (303)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (311)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<400> 341
aagcctttca agatgatccc cggagtgggt gatgggggtct tcctgcccag gcacccccag 60
gancgtgttg cctctgccga ctttcagcct gtccctanca ttgttggtgt caacaacaat 120
gaattcgggt ggctcatccc caaggtcatg angatctatg ataccagaa ngaaatggac 180
aganangcct ccangctgc tctgcagaaa atgttaacgc tgetgatttg cctcctacat 240
ttggtgacct gctgaaggaa gaattcattt gggganaatn gggatcccca aaacccccaa 300
acncatttcc nggaaaataa ttgcngaa 328

<210> 342
<211> 140
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (3)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (33)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (70)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (73)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (129)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (130)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (140)
<223> n equals a,t,g, or c

<400> 342
aangggactt nccagtgatg cacccaagag ctntttatcc ccaacatcac tgtgaataat 60
aggggatccn atncgtgcc agcccataac tctacactgg ccttaatagg agcacagtca 120
cgaatatcnn agtctatgcn 140

<210> 343
<211> 477
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (396)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (421)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (435)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (459)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (463)
<223> n equals a,t,g, or c

<400> 343
gggacagcag agctgacagt cacagcagcc ctgacaagag agttcctgga gcccaagctc 60
ttctccacag aggacaagca ggcagcagag accatgggggt ccccttcagc ctgtccatac 120
agagtgtgca ttccctggca ggggctcctg ctcacagcct cgcttttaac cttctggaac 180
ctgccaacaac gtgcccagac caatatgat gtcgtgccgt tcaatgtcgc agaagggaag 240

gaggtccttc tagtagtcca taatgagtcc cagaatcttt atggctacaa ctggtacaaa 300
ggggaaaggg tgcattgcaa ctatcgaatt ataggatatt gtaaaaaata taagtcaaga 360
aaatgcccac ggcccacac aacgtcgaga gacatntacc caatggaacc ttgtgttcca 420
nacgtaccac atgncagga ttttctccta ctataaaana aanttgatg agaatac 477

<210> 344

<211> 389

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (6)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (19)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (22)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (32)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (35)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (144)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (145)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (159)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (201)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (379)
<223> n equals a,t,g, or c

<400> 344
nnnnnnnnnt ntcgcacgnc nngccagcca anccntcact aaagggaaca aaagctggag 60
ctccaccgcg gtgcgaccgc tctagaacta gtggatcccc cgggctgcag gaattcggca 120
cgaggccggc cggggctggc acgnncgccg cggcggggnc tggaggcgcg accgggccc 180
cccgagcggg aatcagaacg ncgccgaggg gaccagatca acgccagcaa gaacgaggag 240
gacgcgggaa aaatgttcgt tgggtggcctg agctgggata ctagcaaaaa agatttaaaa 300
gactatttta ctaaatttg agaggctcgtt gactgtacaa taaaaatgga tccaacact 360
ggacgggtcaa gagggtttng gtttatcct 389

<210> 345
<211> 152
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (34)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (42)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (48)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (61)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (64)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (84)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (87)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (122)
<223> n equals a,t,g, or c

<400> 345
tcgacccacg ngtcgggggc gcaccatggc ccgnggggct gngctggngc tgctgctctt 60
nggnctgctg ggtgttcttg tcgncgnccc ggatggtggt ttcgatttat ccgatgccct 120
tctgacaat gaaaacaaga aaccactgc aa 152

<210> 346
<211> 634
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (32)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (38)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (284)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (491)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (586)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (588)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (598)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (613)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (621)
<223> n equals a,t,g, or c

<400> 346
tntgtctgaa tgtaaatcac cttctgagcc gntgattncc aagagggttg gtcttattca 60
catatcccaa gtcattctcag aaattgatgg taacaggatg accttgagcc aagaaggagc 120
acaagattcc ttccctcttc agcagaagat cttgggtttgc tctttgatgc tcttgatcag 180
gcagttgaaa atcaaagagg tcaactctggg gaagtatat gaagcctaca gtaaagtctg 240
tcgcaaacag caggtggcgg ctgtggacca gtcagagtgt ttgncacttt cagggtcttt 300
ggaagccagg ggcatttttag gattaaagag aaacaaggaa acccgtttga caaagggtgtt 360
tttcaagatt gaagagaaaag aaatagaaca tgctctgaaa gataaagctt taattggaaa 420
tatcttagct actggattgc cttaaattct tctcttacac cccaccgaa agtattcaag 480
ctggcattta nagagctaca ggcttcattt taaggcttta cacattcggg cctgaaaaca 540
aatatgacct tttttacttg aagccaatga attttaatct atagancntt aaaattanca 600
cagaataata tcnttggggc ntactatttt accc 634

<210> 347
<211> 363
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (228)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (233)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (242)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (268)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (319)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (322)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (323)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (324)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (328)
<223> n equals a,t,g, or c

<400> 347
ggcacgagag gaatatggag aggctgactc ttgcttggtg tggggtggcc ctgaattctt 60
ttgaggacct aagtcctgac tgcttgggac atgcaggact tgtatatgag tatacattgg 120
gagaagttca cctttattga gaagtgtaac aaccctcgtt ctatcacatt attgatcaaa 180
ggaccaaata agccacact tagatcaaag atgcagtaaa gggatgggct tgnagggctg 240
tncaaaaatg ctgttgatga tggctgtntg gttccgggtg ctggtgctgt ggaagtggca 300
atggcagaag ccctggatna annnnaantc catgtaaagg gcaaggcaca gttgggggtcc 360
agc 363

<210> 348
<211> 388
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (205)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (339)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (344)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (364)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (382)
<223> n equals a,t,g, or c

<400> 348
aattcggcac gagctgaagg ctagcaaacc gagcgatcat gtcgcacaaa caaatttact 60
attcggacaa atacgacgac gaggagttag agtatcgggt agtgctggcg cgggagcaat 120
agcgaagggtc gtgagctttg gccgctgagg gcacaaggaa ttagtaacag gaactgaggc 180
gatagaattg gcgcatgcgt atganacatg tcatgctgcc caaggacata gccaaagctgg 240
tccctaaaac ccattctgatg tctgaatctg aatggaggaa tcttggcgtt cagcagagtc 300
agggatgggt ccattatatg atccatgaac cagaacctna aatnttgctg ttccggggcc 360
attnccagga accaaggaat tnagtttc 388

<210> 349
<211> 194
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (38)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (57)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (60)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (67)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (69)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (74)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (83)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (91)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (124)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (127)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (140)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (180)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (191)

<223> n equals a,t,g, or c

<400> 349

gagttgttgc ctgggctgga cgtgggttttg tctgctgngc ccgctcttag ngctctngtn 60
taatttnang cagngcgcca gcnggatggt ncacaagcag atatactact cggacaagta 120
cttngangaa cactacgagn accgggatgg tatgttacc agagaacttg acaacaagn 180
acctaaaact natc 194

<210> 350

<211> 524

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (9)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (14)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (17)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (29)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (85)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (98)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (101)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (148)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (240)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (258)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (484)

<223> n equals a,t,g, or c

<400> 350

```
ttaaccttnt naanggnaca aagtctggnt ctccaccagg tggaggaccg ctccctagcaa 60
ctagtgggtc ccccgggcct gcagnaattc ggcagtgnccg naaagcccta cggcgtcact 120
gcaatgtgct ggaactggga gcaagtgnca gcggctggac ggcattcctga gtcgagacca 180
ttccgattca cggcgcgccg aacttcccca cgctcgagct gcagccgagc ctgtatcgtg 240
aaggtggtgc ggcggcgcnct tgccgagaag cgcctcggcg tccgcgacgt gcgcctcaac 300
ggctcggcag ccagccatgt cctgcaccag gacagcggcc tgggctacaa ggacctggac 360
ctcatcttct gcgccgacct gcgcggggaa ggggagtttc agactgtgaa ggacgtcgtg 420
ctggactgcc tgttggactt cttacccgag ggggtgaaca aagagaagat cacaccactc 480
acgntcaagg aagcttatgt gcagaaaatg gttaaagtgt gcaa 524
```

<210> 351

<211> 352

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (261)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (291)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (295)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (328)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (348)

<223> n equals a,t,g, or c

<400> 351

```
gccaaatcct ttgaatactg ngccaggatt tttaagcaac actttatgga cagcagaata 60
ccttgcttaa tcgtagctgc aaagtcagac ctgcatgaag ttaaacaaga atacagtatt 120
tcacctactg atttctgcag gaaacacaaa atgcctccac cacaagcctt cacttgcaat 180
actgctgatg cccccagtaa ggatatcttt ggtaaattga caacaatggc catgtatccc 240
catgcccgtg tacgctgtat ntgcacctgc aacagggtga cattttgcat ntgtnaaaac 300
ttcctcaact tatactttgc tgcaaatntg gtaaaagaac aaaaatcntt tc 352
```

<210> 352

<211> 632

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (8)
<223> n equals a,t,g, or c

<400> 352
agnnaganac caaccctcac taaagggaac aaaagctgga gctccaccgc ggtgcggccg 60
ctctagaact agtggatccc ccgggctgca ggaattcggc acgaggtgtg tcagaacaat 120
cttgaatcat gaagctacta accagagccg gctctttctc gagattttat tccctcaaag 180
ttgcccccaa agttaaagcc acagctgccc ctgcaggagc accgccacaa cctcaggacc 240
ttgagtttac caagttacca aatggcttgg tgattgcttc tttggaaaac tattctcctg 300
tatcaagaat tggtttgctt attaaagcag gcagtagata tgaggacttc agcaatttag 360
gaaccaccca ttgctgctg cttacatcca gtctgacgac aaaaggagct tcattcttca 420
agataacccg tggaaattgaa gcagttggtg gcaaattaag tgtgaccgca acaagggaaa 480
acatggctta tactgtggaa tgcctgcccgt gtgatgttga tattctaag gagttcctgc 540
tcaatgtcac cacagacca gaatttcgtc gttgggaagt agctgacctt cagcctcagc 600
taaagattga caaagctgtg gcctttcaga at 632

<210> 353
<211> 440
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (435)
<223> n equals a,t,g, or c

<400> 353
ctttaactcc accattagca cccaaagcta agattctaatt ttaaactatt ctctgttctt 60
tcattgggaa gcagatttgg gtaccaccca agtattgact caccatcaa caaccgctat 120
gtatttcgta cattactgcc agccaccatg aatattgtac ggtaccataa atacttgacc 180
acctgtagta cataaaaacc caatccacat caaaaccccc tccccatgct tacaagcaag 240
tacagcaatc aaccctcaac tatcacacat caactgcaac tccaaagcca cccctcacc 300
actaggatac caacaaacct acccaccctt aacagtacat agtacataaa gccatttacc 360
gtacatagca cattacagtc aaatcccttc tcgtcccat ggatgacccc cctcagatag 420
gggtcccttg accancatcc 440

<210> 354
<211> 609
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (3)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (8)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (139)
<223> n equals a,t,g, or c

<400> 354
aancnganac naaccctcac taaagggaac aaaagctgga gctccaccgc ggtgcggccg 60
ctctagaact agtggatccc ccgggctgca ggaattcggc acgagccgga ggggtgtgtgt 120
tgggcaaagc cggaggagna ggaggacgat tgttttacgg atcccgtgat cggcccgtcc 180
ttctccctt tccccctcc ctaccgcccc tgtcccgcg gggagcggcg gcggccttg 240
actttgctgt ctttcctcgc ggagacagat ttcaacacta cacttgcaca atgtctttga 300
aaccaagagt agtagatttt gatgaaacat ggaacaaact ttgacgaca ataaaagccg 360
tggtcatggt ggaatacgtc gaaagagcaa catggaatga ccgtttctca gatatttatg 420
ctttatgtgt ggcctatcct gaacccttg gagaaagact ttatacagaa actaagattt 480
ttttggaaaa tcatgttcgg catttgcata agagagtttt ggagtcagaa gaacaagtac 540
ttgttatgta tcataggtac tgggaagaat acagcaaggg tgcagactat atggactgct 600
tatataggt 609

<210> 355
<211> 466
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (368)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (450)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (451)

<223> n equals a,t,g, or c

<400> 355

```
gctccattat tgatcgggtc atgcagaata attgtgtgcc caagaagatg ctgcagctgg 60
ttggtgtcac tgccatgttt attgcaagca aatatgaaga aatgtaccct ccagaaattg 120
gtgacttttc ttttgtgact gacaacactt atactaagca ccaaatcaga cagatggaaa 180
tgaagattct aagagcttta aactttggtc tgggtcggcc tctacctttg cacttccttc 240
ggagagcatc taagattgga gaggttgatg tcgagcagca tactttggcc aaatacctga 300
tggaactaac tatgttggaac tatgacatgg tgcactttcc tccttctnaa attgcagcag 360
gagctttntg cttagcactg aaaattctgg gataatggtg aatggttatg cagcacctgg 420
ctaagaatgt agtcatgggt aaatcaaggn nttacaaagc acatgc 466
```

<210> 356

<211> 447

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (269)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (375)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (380)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (436)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (440)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (446)

<223> n equals a,t,g, or c

<400> 356

```
ggtgatgagg aggcttgtcc agaggacaag ggaccccagg acccacaggc actggcgctg 60
gacaccaga tccctgcaac ccctggaccc aaaccctgg tccgcaccag ccgggagcca 120
```

gggaaggacg tcacgacctc aggttactcc tccgtcagca ccgcaagtcc cacaagtccc 180
gtggacggtg gcttgggggc cctgcccaca cctacctcag tgctgtccct ggacagtgc 240
tcgcacacac agccctgccca ccatcaggnc aggaagtcag gtttacagtg tcgtccccc 300
agtcccccgg agagcagtgt tccccagcaa caggtgaagc ggataaacta tgcatacaca 360
gtgaagagga ggacntgaan ctgggcttgt gaagctgtaa gtgtgtcagc acatttgcgc 420
agtggatttt actgangggn tgaagng 447

<210> 357

<211> 510

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (453)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (486)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (491)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (496)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (500)

<223> n equals a,t,g, or c

<400> 357

tnccacggt ccgcccacgc gtccgcccac gcgtccgcag attccctccg tcgcccgc 60
gatgatgtgc ggggcgccct ccgccacgca gccggccacc gccgagacc agcacatcgc 120
cgaccagggt aggtcccagc ttgaagagaa agaaaacaag aagttccctg tggttaaggc 180
cgtgtcattc aagagccagg tggtcgcggg gacaaactac ttcataaagg tgcacgtcgg 240
cgacgaggac ttcgtacacc tgcgagtgtt ccaatctctc cctcatgaaa acaagccctt 300
gaccttatct aactaccaga ccaacaaagc caagcatgat gagctgacct atttctgac 360
ctgactttgg acaaggccct tcagccagaa gactgacaaa gtcacccctc gtctaccaga 420
gcgtgcactt gtgatcctaa aataagcttc atntccgggc tgtgccctt ggggtggaag 480

gggcangatt ntgcantgn ttttgcat

510

<210> 358

<211> 240

<212> DNA

<213> Homo sapiens

<400> 358

ggtcttgac tcatgagctg tccccacatt aggttaaaa acagatgcaa ttcccgacg 60
tctaaaccaa accactttca ccgtacacg accgggggta tactacggc aatgctctga 120
aatctgtgga gcaaaccaca gtttcatgcc catcgtccta gaattaattc ccctaaaaat 180
ctttgaaata gggcccgat ttaccctata gcacccctc tacctcctc agagccaaaa 240

<210> 359

<211> 340

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (320)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (331)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (334)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (335)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (338)

<223> n equals a,t,g, or c

<400> 359

ggcagagct cataaggaaa tctacccta tgtcatccag gaacttagac caactttaa 60
tgaaactggga atctccactc cggaggaaact gggccttgac aaagtgtaaa ccgcatggat 120
gggcttcccc aaggatttat tgacattgct acttgagtgt gaacagttac ctggaaatac 180
tgatgataac atattacctt atttgaacaa gttttccttt attgagtacc aagccatgta 240
atggtaactt ggactttaat aaaaaggaaa tgagtttgaa ccggaaaaaa aaaaaaaaaa 300
aaaaaaaaa aaagaaaaan aaagaaggga nagnnaanaa 340

<210> 360
<211> 501
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (147)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (148)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (192)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (242)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (244)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (364)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (455)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (465)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (474)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (491)
<223> n equals a,t,g, or c

<400> 360
cccacgcgtc cgcccacgcg tccgatgcct atcatatagt aaaaccacgc ccatgacccc 60
taacaggggc cctctcagcc ctccaatga cctccggcct agccatgtga tttcacttcc 120
actccataaa gctcctcata ctaaggnta ctaaccaaca cactaaccat ataccaatga 180
tggcgcgatg tnacacgaag aaagcacata ccaagggcac cacacaccac ctgtccaaaa 240
angncttcga tacgggataa tcctatttat tacctcagaa gtttttttct tcgcaggatt 300
ttctgagctt ttacactcca gcctagccct acccccaact aagaaggnac tggccccaac 360
aagnatcacc cgctaaatcc ctanaatcca ctccataaca ctccgtatat cgatcaggat 420
atcatcactg actcacatat ctaatagaaa aaacngaaac aatantcanc atgnttatta 480
aatttatggt ntctatttac c 501

<210> 361
<211> 393
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (357)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (359)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (364)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (367)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<400> 361
gccccattat tcctagaacc aggcgacctg cgactccttg acgttgacaa tcgagtagta 60
ctcccgattg aagcccccat tcgtataata attacatcac aagacgtctt gcactcatga 120
gctgtcccca cattaggctt aaaaacagat gcaattcccg gacgtctaaa ccaaaccact 180
ttcacgcta cagcaccggg ggtatactac ggtcaatgct ctgaaatctg tggagcaaac 240
cacagtttca tgcccatcgt cctagaatta attcccctaa aaatccttga aataggggccc 300
gtatttacct tatagcacc cctctacccc ctctagagcc aaaaaaaaaa aaaaagntnc 360
aagnttnaaa aaactgaaaa tcagaaaagn ttt 393

<210> 362
<211> 664
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (308)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (535)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (551)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (567)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (642)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (660)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (662)

<223> n equals a,t,g, or c

<400> 362

```
gagtaggtct acaagacgct acttccccta tcatagaaga gcttatcacc tttcatgac 60
acgccctcat aatcattttc cttatctgct tcctagtccct gtaggccctt ttcctaacac 120
tcacaacaaa actaactaat actaacatct cagacgctca ggaaatagaa accgtctgaa 180
ctatccctgcc cgcacatcc tagtcctcat cgcctccca tccctacgca tcccttacat 240
aacagacgag gtcaacgac cctcccttac catcaaatca attggccacc aatggtagtg 300
aacctacnaa gtacaccgac tacggcggac taatcttcaa ctctacata cttccccatt 360
attcctagaa ccaggcgacc tgcgactcct tgacgttgac aatcgagtag tactcccgat 420
tgaagcccc attcgtataa taattacatc acaagacgct ttgcactcat gagctgtccc 480
cacattaggc ttaaaaacag atgcaattcc ccgacgtcta aaccaaacca ctttnaccgg 540
tacacgaacc nggggtatta ctacggncaa tgctctgaaa tctggggagc aaaaccacag 600
gttcatgcc atcgtccctag aaataaattc ccctaaaaat cnttgaaaaa aggggcccgn 660
antt 664
```

<210> 363

<211> 595

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (466)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (467)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (485)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (515)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (552)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (557)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (559)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (568)
<223> n equals a,t,g, or c

<400> 363
gcgcaagtag gtctacaaga cgctacttcc cctatcatag aagagcttat cacctttcat 60
gatcacgccc tcataatcat tttocttata tgcttcctag tcctgtatgc ccttttccta 120
acactcacaa caaaactaac taataactaac atctcagacg ctcaggaaat agaaaccgtc 180
tgaactatcc tgcccgcctat catcctagtc ctcacgcgcc tcccatccct acgcatcctt 240
tacataacag acgaggtcaa cgatccctcc cttaccatca aatcaattgg ccaccaatgg 300
tactgaacct acgagtacac cgactacggc ggactaatct tcaactccta catacttccc 360
ccattattcc tagaaccagg cgacctgcga ctccttgacg ttgacaatc gagtagtact 420
cccgattgaa gccccattc gtataataat tacatcacaa gacgtnnttg cactcatgag 480
ctgtncccac attaaagctta aaaacagatg cattnccgga cgtctaaacc aaaccacttt 540
caccgatata cnaaccngng ggtatacnac cggccaatgc ttctgaaatc ttgtg 595

<210> 364
<211> 441
<212> DNA
<213> Homo sapiens

<400> 364
acaagacgct acttccccta tcatagaaga gcttatcacc ttctatgac acgccctcat 60
aatcattttc cttatctgct tcctagtccg gtatgccctt ttcctaacac tcacaacaaa 120
actaactaat actaacatct cagacgctca ggaaatagaa accgctctgaa ctatcctgcc 180
cgccatcatc ctagtcccca tcgccctccc atccctacgc atcctttaca taacagacga 240
gggtcaacgat cctcccttt accatcaaat caattgggcc accaatggta ctgaacctac 300
gagtacaccg actacggcgg actaatcttc aactcctaca tacttcccc attattccta 360
gaaccaggcg acctgcgact ccttgacgtt gacaatcgag tagtactccc gattgaagcc 420
cccattcgta taataattaa c 441

<210> 365
<211> 367
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (347)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (362)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (363)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (366)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (367)
<223> n equals a,t,g, or c

<400> 365
catacttccc ccattattcc tagaaccagg cgacctgcga ctcccttgacg ttgacaatcg 60
agtagtactc ccgattgaag cccccattcg tataataatt acatcacaag acgtottgca 120
ctcatgagct gtccccacat taggcttaaa aacagatgca attcccggac gtctaaacca 180
aaccactttc accgctacac gaccgggggt atactacggt caatgctctg aaatctgtgg 240
agcaaaccac agtttcatgc ccatcgtcct agaattaatt cccctaaaaa tctttgaaa 300
tagggcccggt atttacccta tagcaccccc tctaccccct ctagagncaa aaaaaaaaaa 360
annagnn 367

<210> 366
<211> 460
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (51)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (154)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (211)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (322)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (336)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (340)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (355)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (356)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (378)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (411)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (413)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (415)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (453)
<223> n equals a,t,g, or c

<400> 366
ttttcgaggc caacgatccc tcccttacca tcaaatcaat tggccaccaa nggtaccgaa 60

```
cctacgagta caccgactac ggcggaacta tcttcaactc ctacatactt cccccattat 120
tcctagaacc aggcgacctg cgactccttg acgntgacaa tcgagtagta ctcccgattg 180
aaacccccat tcggataata attacatata nggacgtctt gcactcatga gctgccccac 240
attaggctta aaaacagatg caattcccgg acgtctaaac caaaccacta tcaccgctac 300
acgaccgggg gtatactacg gncaatgctc tgaaanctgn ggagcaaacc acagnntcat 360
gcccatcggc ctagaatnaa tccccctaaa aatctttgaa atagggcccg nantnaccct 420
atagcaccce ctctaccccc tctagagcca aanaaaaaaa 460
```

<210> 367

<211> 610

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (6)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<400> 367

```
gcaagntgac acnaaccctc actaaaggga acaaaagctg gagctccacc gcggtgcggc 60
cgctctagaa ctagtggatc ccccgggctg caggaattcg gcacgaggga atggcggctt 120
ccatctctga ggcgagcgac gctatggatc ccacagtggg ttgctaagaa ggccattttc 180
aactctccac tggaggctgc tatggcgctc cctcacctgc agcagcccag ctttctactg 240
gtagcctga aagctgactc tataaataag ccctttgcac agcagtgcc aagacttggt 300
aaagtcattg aggactttcc agcaaagtct gaaccaatca ggtccttgt gactggagca 360
gctggtcaaa ttgcatattc actgctgtac agtattggaa atggatctgt ctttggtaaa 420
gatcagatgt catcgcaaca gataaagaag acgttgcctt caaagacctg ggatgtggcc 480
attcttgtgg gctccatgcc aagaagggaa ggcattggaga gaaaagattt actgaaagca 540
aatgtgaaaa tcttcaaata ccagggtgca gccttagata aatacggcaa gaagtcagtt 600
aaagggttatt 610
```

<210> 368

<211> 548

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (370)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (378)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (412)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (429)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (471)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (490)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (495)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (528)
<223> n equals a,t,g, or c

<400> 368
ggcacgagga gcactctatt tattgtactg tgaataatga tgaaggagag tggagtggcc 60
caccacctga atgcagagga aaatctctaa cttccaaggt cccaccaaca gttcagaaac 120
ctaccacagt aaatgttcca actacagaag tctcaccaac ttctcagaaa accaccacaa 180
aaaccaccac accaaatgct caaggtacag agactccatc agttcttcaa aaacacacca 240
cagaaaatgt ttcagctaca agaaccctc caactcctca gaaaccacc acagtaaatg 300
tcccagctac aatagtaca ccaacacctc agaaaccac cacattaatg ttccagttac 360
aggagtctcn tcaacacntc aagnacacc tagtaatgtt tcagttacag gncctaccac 420
ttttcggancc caccggggc aatgttcgnc accattcccg cgcgttcggc ntttcttca 480
aaaccttttn caagnccttt tgcgttagat cctgtggcat gttttgtnc cggcccttac 540
ggccaggt 548

<210> 369
<211> 538
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (12)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (423)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (466)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (521)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (531)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (538)
<223> n equals a,t,g, or c

<400> 369
gatggcgcg anccgggacg cgcgtacgcc ctgctgcttc tcctgatctg ctttaacggt 60
ggaagtggac ttcacttaca ggtcttaagc acaagaaatg aaaataagct gcttcctaaa 120
catcctcatt tagtgcggca aaagcgcgcc tggatcacccg cccccgtggc tcttcgggag 180
ggagaggatc tgtccaagaa gaatccaatt gccaaagatac attctgatct tgcagaagaa 240
agaggactca aaattactta caaatacact ggaaaaggga ttacagagcc accttttggg 300
atatttgtct ttaacaaaga tactggagaa ctgaatgtta ccagcattct tgatcgagaa 360
gaaacaccat tttttctgct aacagggtta cgctttggat gcaagaggga acaatgtaga 420
ganacccttt agagctacgc attaagggnt cttgatatcc aattgncaac ggaaccagtg 480
gttcacacag ggatgtcttt gttggggcct gttgaagagt ngaggtgcag nacatacn 538

<210> 370
<211> 538
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (401)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (492)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (500)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (536)
<223> n equals a,t,g, or c

<400> 370
gagctcagtc cccatcccga agcacagggc aggacgtcgc ggcggagtgg ggaagcgagg 60
agtcctgtggc cgggagcttg gaggctgagt ttgagaaagc tgcagaggag gttaggcacc 120
ttaagaccaa gccatcggat gaggagatgc tggtcatcta tggccactac aaacaagcaa 180
ctgtgggcga cataaataca gaacggcccg ggatgttgga cttcacgggc aaggccaagt 240
gggatgcctg gaatgagctg aaagggactt ccaaggaaga tgccatgaaa gcttacatca 300
acaaagtaga agagctaaag aaaaaatacg ggatatgaga gactggattt gggtactgtg 360
ccatgtgttt atcctaaact gagacaatgc cttgtttttt nctaataccg tggatggtgg 420
gaatccggga aaataaccag ttaaccagc tactcaaggc tgctcaccat acggtcttaa 480
cagattaggg gntaaaacgn ttactgactt ccttgagtag ttttacctga aaccantt 538

<210> 371
<211> 224
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (26)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (48)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (59)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (68)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (90)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (108)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (113)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (123)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (171)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (192)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (204)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (215)
<223> n equals a,t,g, or c

<400> 371
gctgtgcct acacaccacc ntcgcnagaa gctgtgcagc gcattgcnga gtctcaccng 60
cagtc tanca gcaatttgaa tgagaaccan gcctcagagg aggagganga gcngggggag 120
ctncggggagc tgggttatcc nagagaggaa gatgaggagg aagaggagga ngatgaagaa 180
gaggaagact angaggacag ccangctgaa gaccngagcg gaga 224

<210> 372
<211> 459
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (6)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (52)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (90)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (105)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (139)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (140)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (176)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (208)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (328)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (394)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (429)
<223> n equals a,t,g, or c

<400> 372
aactnnccca atgccgaact gggaggcccc tttaccaga tgaacggagt gnccggaaat 60
ggcatgaaca acattgacat gactggagan aagaagtcgt tgganctccc atatcccagc 120
agctttgtct ccgtctctnn acctagaaac cagaccttca cttacatggg caagtntctc 180
attgaccctc agtaccctgg tgccagcngc taccagaag gcataatcaa tattgtgagt 240
gcaggcatct tgcaaggggt cacttcccca gcttcaacca cagcctcatc cagcgtcacc 300
tctgcctccc ccaaccact ggccacanga cccctgggtg tgtgcacccat gtcccagacc 360
cagcctgacc tggaccacct gtactctccg ccancgcctc ctccctctta ttctggctgt 420
gcaggaganc tctaccagga cccttctgcg ttctgtta 459

<210> 373
<211> 422
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (27)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (47)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (60)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (66)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (70)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (78)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (87)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (105)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (267)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (331)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (359)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (393)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (401)
<223> n equals a,t,g, or c

<400> 373
ggcagacctg aggcccatcc ccaaggngcc cagccgtagc ttcaggntct ccgccagggn 60
ccctcncggn tgcttctngt ttttaanact tccagatcga tectntcctc ggtaccgtgg 120
gctttgggtc tggcctccac ggggtggcct tcaccctgaa agcagtttgc cgagaatgta 180
tgtgagccaa gttcgccgcc aagggggagg gccagttggg agcctgccga gcgggccaag 240
aaagtagagg acatgatgaa gaagctntgg ggtgacaggt gagccccggg gaatatggtg 300
ggggattcct gaaaactggg ggtagtggca ncaacgtagg ggcgtggtgt nctattcana 360
attggttcaa gccacacaagt ttgcccata atnttcctt ngggaccaac ttccaagta 420
at 422

<210> 374
<211> 342
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (4)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (17)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (20)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (292)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (301)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (305)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (325)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (332)

<223> n equals a,t,g, or c

<400> 374

```
gcanaacatc agcgcnnaacn gganataaaa agattatcca cagagcattc cagtgtatca 60
gagtatcatc cagccgatgg ctatgcgttc agtagcaaca ttacacaag aggatccac 120
ctggaccaag gggaagctgc tgttgctttt aagccaactt ctaatcgcca tattagattg 180
aaattatgaa ccaactccaa acacaacca agaaatatgc ccaatcccag ttatgacttt 240
gtttgccagg ggaaacaaca gttgagctct ccgggttcct taaggaaatga tntttttaga 300
naatnccttg aatgaatcg gaaancatgg tngaaaaagt tt 342
```

<210> 375

<211> 387

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (19)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (162)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (305)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (335)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (366)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (374)
<223> n equals a,t,g, or c

<400> 375
ggcagagctc tgcgcgcant ccgcttgact cagctcaccg agattctgtc agggggtgtt 60
tatattgaga agaacgataa gctttgtcac atggacacaa ttgactggag ggacatcgtg 120
agggaccgag atgctgagat agtgggtgaag gacaatggca gnaagctgtc ccccctgtca 180
tgaggtttgc aaggggcgat gctgggggtcc tggatcagaa gactgccaga cattgaccaa 240
gaccatctgt gcttcctcag tgtaattggt cactgctttg gggcccaacc ccaaccagtg 300
gttgncatga tgagtgtggc cgggggttgt ttcanggcc cttcaggac acagactgtt 360
tttgnntggc cggnatattca atggaca 387

<210> 376
<211> 492
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (203)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (318)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (333)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (336)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (405)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (416)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (440)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (444)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (448)

<223> n equals a,t,g, or c

<400> 376

```
ggcagcgagaa gcaaaactcaa agcatttgagg tcaaaacatc ctgattgttt tatacatggt 60
ttggatgata aattcagtaa agatacaggc ataatcctta tatgcaaata agttgatggg 120
tattttctat ggtgatttca gacaaggaac ctgcagggtca ctctggaaga tggttacatt 180
gaattgagca ccagcgatag ggncggccca atttttaaat ctccacagac gtatatggat 240
ggtttactgc attatgtatc tgtaataaag gacaactctg ggtgagtcga ataaataactt 300
ctgtcagagc tgtgagtnga gttttactct ctnttngttt taatggaatt ttctggtgtc 360
tttttgcaaa aaatttgtcc ttgattacca aaaatacttt cactnataag tcttgnocct 420
ctttcctttt ttcccatatn caanttttct tcataaaaaa aataaaacttg ccacacatgg 480
gtgggcttca tt 492
```

<210> 377

<211> 336

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (15)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (36)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (66)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (70)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (116)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (163)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (197)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (227)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (229)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (231)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (249)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (280)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (292)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (308)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<400> 377
gacngtaccg tcccaaatcc cgggtcgacc cacgntccg aagcagcagt tagcccccg 60
cccgntgtn tgtccccaga gccatggaga gagccagtct gatccagaag gccaanctgg 120
cagagcaggc cgaacgctat gaggacatgg cagccttcat ganaggcgcc gtggagaagg 180
gcgaggagtc tccctcnaag agcgaacact gctctcagta gcctatnana ncgtggtggg 240
cggccagang gctgcctgga ggggtgctgc cagtattgan cagaaaagca angaggaggg 300
gttcgganag gagaaggggc ccgangtgcg ttgaat 336

<210> 378
<211> 488
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (20)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (35)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (43)
<223> n equals a,t,g, or c

<400> 378
cngtcagtna cggtaccggn aattccccggg tgcgnaccca cgnctgtgccg ccgagaccgc 60
tgccttgctc ggaagggggc gagcggctgc tgcccaccca gaagcagccc gggggcggcc 120
aggtaactc cagccgctac aagacggagc tgtgccgcc ctttgaggaa aacgggtgcct 180
gtaagtacgg ggacaagtgc cagttcgac acggcatcca cgagctccgc agcctgacct 240
gccaccccaa gtacaagacg gagctgtgcc gcacctcca caccatcggc ttttgccct 300
acggggcccg ctgccacttc atccacaacg ctgaagagcg ccgtgccctg gccggggccc 360
gggacctctc cgctgaccgt ccccgccctc agcatagctt tagtttgctg ggtttcccag 420
tgccgctgcc accgcccgtg ccaccgggct gctggacagc ccacgtccat caaccaacc 480
cctatttt 488

<210> 379
<211> 398
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (46)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (124)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (344)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (385)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (386)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (394)
<223> n equals a,t,g, or c

<400> 379
ggcacgaggg ggcctcccgc tgtggccacc tctgccgcgg ccgcgncgcc gcgtcctacc 60
ctgcccctccg tgcctctctg ctgccgcagt cgctggcggc gccggccgcg ttcccgcacgc 120
gcantaacag ccaggagtcc aaaactactt acctggaaga ccttccacca cccctgagt 180
atgaattggc cccgtccaag ttagaagagg aagtggatga tgtctttctc attcgagctc 240
aaggactgcc ctgggtcatg gcactatggg aagatgtggc ttttaacttt tttttccaga 300
cttgcagaat ccggcaacgg ttgagtaatg ggaattacat tttntccct aaaaacaaga 360
gatgggggaa aacgtaaggg ggtnnntggc cttnaatt 398

<210> 380
<211> 455
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (421)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (442)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (447)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (448)
<223> n equals a,t,g, or c

<400> 380
atgccatggt gcggctggtc aagtgcgacg tctacccctg ccccaacaca gtggactgct 60
tcgtgtcccg cccacccgag aaaaccgtct tcaccgtctt catgctagct gcctctggca 120
tctgcatcat cctcaatgtg gccgaggtgg tgtacctcat catccgggcc tgtgcccgcc 180
gagcccagcg ccgctccaat ccaccttccc gcaagggtc gggcttcggc caccgcctct 240
cacctgaata caagcagaat gagatcaaca agctgctgag tgagcaggat ggctccctga 300
aagacatact gcgcncaacc ctggcacggg ggctgggctg gctgaaaaaa acgaccgtgc 360
tcggntgtga tgccacatac caggcaacct cccatccac cccgaccttg cctggggcga 420
nccctccttc tccctgcccg tncccanng ctcac 455

<210> 381
<211> 421
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (3)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (4)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (8)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (23)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (156)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (191)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (398)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (410)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (413)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (421)
<223> n equals a,t,g, or c

<400> 381
ntnngcgnct gcaggctgaa cantagtggga tccaaagaat tcggcagcag tgggaacgag 60
gggtgaagcg gcccaggtga ggctcgccat ggccaaacag ctgcaggccc gaagctagac 120
gggatcgact acaacccctg ggtggagttt gtgaantggc cagtgcgcat gacgtcgtga 180
acttgggccca nggcttcccg gatttcccac caccagactt tgccgtggaa gcctttcagc 240
acgctgtcag tggagacttc atgcttaacc agtacaccaa gacatttggt taccaccac 300
tggacgagga tcctggcaat ttctttgggg gagctgctgg gtcaaggata agaccggtc 360
agggatgtgc tggtgactgt tggtggnat gggggccngt ttcaaaagcn ttncaggcc 420
n 421

<210> 382
<211> 545
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (57)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (466)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (505)
<223> n equals a,t,g, or c

<400> 382
aaggnagaca caancctcac taaaggggaac aaaagctgga gctccaccgc ggtgcgnccg 60
ctctagaact agtggatccc cggggctgca ggaattcggc acgagcggcg gcggcggcgg 120
cgggccagtc ggtgtatgcc ttctcggcgc gcccgctggc cggcggggag cctgtgagcc 180
tggtctccct gcggggcaag gtactactta tcgagaatgt ggcgtccctc tgaggcacca 240
cgttccggga ctacaccag atgaacgagc tgcagcggcg cctcggaccc cggggcctgg 300
tggtgctcgg cttcccgtgc aaccagtttg ggcacacagga gaacgccaag aacgaagaga 360
ttctgaattc cctcaagtac gtccggcctg gtggtgggtt cgagcccaac ttcattgctct 420
tcgagaagtgc cgaggtgaac ggtgcggggg cgcacctctt cttcgncttc ctgcgggagg 480
ccctgccagc tcccagcgac gacgncaccg cgcttatgac cgaccccaag ctcacacact 540
ggtct 545

<210> 383
<211> 375
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (224)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (242)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (264)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (282)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (329)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (333)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (368)
<223> n equals a,t,g, or c

<400> 383
gcttttcattg ccaagtcctt ctatgacctc agtgccatca gcctggatgg ggagaaaagta 60
gatttcaata cgtctcgggg cagggccgtg ctgattgaga atgtggcttc gctctgaggc 120
acaaccaccc gggacttcac ccagctcaac gagctgcaat gccgctttcc caggcgcctg 180
gtggtccttg ggcttccctt gcaaccaatt tgggacatca gganaactgt caaaatgagg 240
anacctgaa cagttcctaat tatnttccgt cctgggggtgg gnataccaac ccccttccc 300
cctttttcca aaaattttaa ggttaaatng gcnaaaaaaa acatcctntt tttcccccta 360
ccttaaanga aaaac 375

<210> 384
<211> 530
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (178)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (354)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (361)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (382)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (386)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (415)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (426)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (436)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (515)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (527)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (529)
<223> n equals a,t,g, or c

<400> 384
gctcgtgccg tgagacacaa ccaccoggga cttcaccag ctcaacgagc tgcaatgccg 60
ctttcccagg cgcttggtgg tccttggtt cccttgcaac caatttgac atcagagcag 120

```
gagagacaga agtagcaaac cctctttcga gatgtccctc cagccccaga agtacctnca 180
gcctcacacc atctcttcag cctagcaagt tgctggaggg agtctataac ctaccaggag 240
ccagccagcc atttgatca agaaatagaa atctgcaggg tacagtggct cacacctata 300
atcccagcgc tttgggaggg taagtcttag gacaaaggca aggaagaaaa gcangaactt 360
naaaatccaa ttccttttgg gncctnaaat ttaaccttca agttcaaggg agctnaagta 420
aggcanaagg ccaaanggct ttttacttna acaacaacgg tgccaatttg gaaaggcaag 480
ggcaaggcaa aaaaccaggg ggcaagaagg aaaangggaa aaggggntnt 530
```

<210> 385

<211> 465

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (151)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (310)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (312)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (329)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (338)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (347)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (351)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (354)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (373)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (383)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (438)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (449)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (450)

<223> n equals a,t,g, or c

<400> 385

```
cacactcaag ttctttcctg ccagtgccga caggacggtc atcgattaca acggggaacg 60
cacgctggat ggttttaaga aattcctgga gagcggtggc caggatgggg caggggatga 120
tgacgatctc gaggacctgg aagaagcaga ngagccagac atggaggaag acgatgatca 180
gaaagctgtg aaagatgaac tgtaatacgc aaagccagac ccgggcgctg ccgagacccc 240
tcggggggct gcacaaccag cagcagcgca aggcttccga gccttgnggg ctcggtttga 300
aaagaagggn tngccgggaa acccagggna actctctnga agttganact ncancctag 360
aaaacgtccg ttnaaccccg ttnccttctt ctgcttttcg ggtttttgga aaagggattc 420
atttcagggc aggccaanct tgttggggnn tgttcctgaa accat 465
```

<210> 386

<211> 733

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (606)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (621)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (689)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (703)
<223> n equals a,t,g, or c

<400> 386
gcggtgatca tgggggcccc gggtcgggc aagggcaccg tgctcgtcgc catcactaca 60
cacttcgagc tgaagcacct ctccagcggg gacctgctcc gggacaacat gctgcggggc 120
acagaaattg gcgtgttagc caaggctttc attgaccaag ggaaactcat ccagatgat 180
gtcatgactc ggctggccct tcatgagctg aaaaatctca ccagtatag ctggctgttg 240
gatggttttc caaggacact tccacaggca gaagccctag atagagctta tcagatcgac 300
acagtgatta acctgaatgt gccctttgag gtcattaaac aacgccttac tgctcgtcgg 360
attcatcccg ccagtggccg agtctataac attgaattca accctcccaa aactgtgggc 420
attgatgacc tgactgggga gcctctcatt cagcgtgagg atgataaacc agagacgggt 480
atcaagagac taaaggctta tgaagaccaa acaaagccag tcctggaata ttaccagaaa 540
aaaggggtgc tggaacatt ctccggaaca gaaaccaaca agatttgcc ctatgtatat 600
gcttttctac aactaaagt ncacaaaagaa gccagaaagc ttaattact tcatgaggag 660
aaatgtgtgt aactattaat agtaagaang gcaaaccctc tantccttgg atttaaaact 720
ggttttctaa aac 733

<210> 387
<211> 180
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (6)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (43)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (61)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (123)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (133)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (151)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (153)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (159)
<223> n equals a,t,g, or c

<400> 387
nctttnatct ntgaagtgga tctacaatgg ctteagcagt gtntccagt tcctaggact 60
ntacaagaaa tctggaaaac ttgtattctt caggcttggg taatgtcagg caaaaccact 120
gtngttgcac atngcttcaa aggtgcgcag ntngtgggnc aacattgttt cccaacactt 180

<210> 388
<211> 428
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (49)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (84)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (90)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (197)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (263)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (285)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (286)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (306)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (366)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (395)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (414)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (418)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (427)

<223> n equals a,t,g, or c

<400> 388

```
gggaacagta agaccgagga ccagcgcaac gaggagaagg cgcacgtgna ggccaacaaa 60
aagatcgaga agcagctgca gaangacaan caggtctacc gggccacgca ccgcctgctg 120
ctgctgggtg ctggagaatc tggtaaaagc accattgtga agcagatgag gatcctgcat 180
gttaatgggt ttaatgnaga cagtgagaag gcaaccaaag tgcaggacat caaaaacaac 240
ctgaaaagagg cgattgaaac cantgtggcc gccatgagca acctnnngtc ccccctggga 300
gctggnaaac ccgagaaac cagttcagag tggactacat cctggagtgt tatgaacgtg 360
cctggntttt gacttccctc cggattctat gagcnggcaa ggtctgtgga ggtnaagntt 420
ctgctgna                                     428
```

<210> 389

<211> 454

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (181)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (184)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (209)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (240)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (283)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (287)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (292)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (309)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (316)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (319)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (369)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (402)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (424)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (426)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (437)

<223> n equals a,t,g, or c

<400> 389

accacgcgt cgcacgc gtccgagct caaacctc caggagccc aggagcccag 60
gggtgggaaa ctccaggaact ttgcacccat ccctggtgaa cctgtggttc ccatcctctg 120
tagcaaccgc aactttccag aagaactcaa gcctctctgc aagagcccaa tgcccaggag 180

ntanttcaga ggctggagga aatcgctgna ggacccgggg cacatgtgga aatctgtgcn 240
tacgctggcc tgtaccgat gctaagggg cttgccactg gcnggcntcc cntccgcagc 300
aggggaagnc ttttcntcnt gcagaaagg ccacccatga tattccattc cccagcagtt 360
caactaacnt ggttccattc gggaaggagc agccggggaa gnattgggtt gattggaagg 420
cttngnccca aatggtncct tccctggcaa tttt 454

<210> 390

<211> 553

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (29)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (30)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (58)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (77)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (414)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (417)

<223> n equals a,t,g, or c

<400> 390

gcaaganaaa attacctca ctaaagggnn caaaagctgg agctccaccg cgggtcgnc 60
gctctagaac tagtgntcc cccgggctgc aggaattcgg cacgagcagg gacgctaaga 120
ttgctacctg gactttcgtt gaccatgctg tcccgggtgg tactttccgc cgccgccaca 180
gcgccccctc tctgaagaat gcagccttcc taggtccagg gaccctatgt actcggaact 240

gggcttatct tgtacgcttt atccaaagaa atatatgtga ttagcgcaga gaccttcact 300
gccctatcag tactaggtgt aatggtctat ggaattaaaa aatatgggcc ctttgttgca 360
gactttgctg ataaactcaa tgagcaaaaa cttgcccac tagaagaggc gaanagngct 420
tccatccaac acatccagaa tgcaattgat acggagaagt cacaacaggc actgggtcag 480
aagcgccatt accttttttg atgtgcaaaag gaataacatt gctatggctt tggaagtac 540
ttaccggga acg 553

<210> 391

<211> 632

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (586)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (613)

<223> n equals a,t,g, or c

<400> 391

cagtttcgaa antaaccctc actaaagggg acaaaagctg gagctccacc gcggtgcggc 60
cgctctagaa ctagtggatc ccccgggctg caggaattcg gcacgaggtc ctgtgcggtc 120
acttagccaa gatgcctgag gaaaccacaga cccaagacca accgatggag gaggaggagg 180
ttgagacgtt cgcctttcag gcagaaattg cccagttgat gtcattgatc atcaatactt 240
tctactcgaa caaagagatc tttctgagag agctcatttc aaattcatca gatgcattgg 300
acaaaatccg gtatgaaagc ttgacagatc ccagtaaatt agactctggg aaagagctgc 360
atattaacct tataccgaac aaacaagatc gaactctcac tattgtggat actggaattg 420
gaatgaccaa ggctgacttg atcaataacc ttgttactat cgccaagtct gggaccaaag 480
cgttcatgga agctttgag gctggtgcag atatctctat gattggccag ttcggtgttg 540
gtttttattc tgcttatttg gttgctgaga aagtaactgt gatcancaa cataacgatg 600
atgagcagta cgnttgggag tcctcagcag gg 632

<210> 392

<211> 600

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (6)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (162)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (518)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (571)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (572)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (587)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (590)
<223> n equals a,t,g, or c

<400> 392
gctcanatct gccttctgga gactgcgccg tcctcccggg agagccagaa agaggacatg 60
gctgctgggc agcgggaagc gaggccccag gtgtcactga cattcgagga cgtggctgtg 120
ctctttacct gggatgagtg gagaaagctg gctccttctc anagaaactt gtaccgggat 180
gtgatgctgg agaactatag gaacctggtc tctactgggac tctcatttac caaaccaaaa 240
gtcatctccc tgttcagca aggagaagat ccctgggagg tggagaaaga cagttctggt 300
gtctcctctc taggatgtaa gagcacacct aaaatgacaa agtcaactca aactcaggat 360
tcatttcagg agcagataag gaaaagattg aaaagggatg aaccctggaa cttcatatca 420
gaaagatcct gcatatatga agagaaatta aagaaacagc aggacaaaaa tgaaaattta 480
caaataatth cagttgcccc tacaaaaatc cttactgnag atagaagcca taaaaatgtt 540
gaatttgccc aaaacttcta cctgaaatca nncttcatta agcaccngan aattgcctaa 600

<210> 393
<211> 531
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (63)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (267)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (352)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (394)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (395)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (400)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (451)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (464)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (471)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (476)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (512)
<223> n equals a,t,g, or c

<400> 393
tgcagctctt ttcattttgc catccttttc cagctccatg atggttctgc aggtttctgc 60
ggnccccccg gacagtggct ctgacggcgt tactgatggt gctgctcaca tctgtgggtc 120
cagggcaggg ccactccaga gaattacctt ttccagggac ggcaggaatg ctacgcgttt 180
aatggtaaca gccagaagga catcctggag gagaaagcgg gcagtgccgg aacaggatgt 240
gcagacacaa cttacggagc tgggcgngcc catggaccct gcagcgccga gttccagcct 300
agggtggaat gttttccccc tccaagtagg gggcccttgg cagcaacaca anctgcttgt 360
ttggccaagt gacggntttt taccagga gcannngtan tcctgtcaac tggattggag 420
gcacatttga ttttgcgccg ttagacatga nggggttggg ttcngtttgt naaatntggg 480
ttgggaattt taatgaaaaa ggggggcaaa gnaattttaa aggggggttt t 531

<210> 394
<211> 404
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (269)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (295)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (330)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (382)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (399)
<223> n equals a,t,g, or c

<400> 394
gggactcttaa ttctgtaata tatcaaggaa tctcgtaaaa ccgacactaa aacgtccctg 60
cctacaaatc atccggccaa attatgagtt cattgtatta tgcgaatgct ttattttcta 120
aatatccagc ctcaagttcg gttttcgcta ccggagcctt cccagaacaa acttcttggtg 180
cgtttgcttc caacccccag cgcccgggct atggagcggg ttccggcgct tccttcgccc 240
cctcgatgca gggcttgtag cccggcggng ggggcatggc gggccagagc gcggncggcg 300
tctacgcggc cggctatggg ctcgagccgn gttccttcaa catgcactgc gcgccctttg 360
agcagaaacc tctcgggggt gngcccnggn gaattccgnc aagg 404

<210> 395
<211> 499
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (10)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (89)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (90)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (93)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (95)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (98)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (112)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (115)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (119)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (120)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (125)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (149)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (172)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (211)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (223)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (225)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (299)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (426)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (442)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (443)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (445)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (457)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (470)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (477)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (484)
<223> n equals a,t,g, or c

<400> 395
ggcacgaggn gcggcggggc tggagccccc atcctcctgc ggcagatgtt cgagcctgta 60
agctgcacct tcacgtacct gctgggtggn agngngtncc gggaggccgt tntgntcggn 120
ccagnccctgg aaacagcgcc tcgggatgnc cagctgatca aggagctggg gnttcggctg 180
ctctatgctg gcgttgaga ccagggccag ncctggccac aantnaggct gtgtcacctt 240
cgttctgaat gaccacagca tggccttcac tggagatgcc ctgttgatcc gtgggtgtng 300
gcggacagac ttccagcaag gctgttgcca agaccttgta accattcggg ccatgaaaag 360
atctttcaaa atttccagga gactgtcttg atctaacctg gttcaagggt accatggggg 420
ttaaantgtt caacgtggag gnnngggagg acttttnaaa cccttggttn aaccttnaat 480
tttnaggagt ttttcaaaa 499

<210> 396
<211> 526
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (445)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (502)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (508)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (519)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (523)
<223> n equals a,t,g, or c

<400> 396
ggtccccgtc acgctgactt tccgtgcagt gctgtggtgc gaaaatgcct cgccgctcct 60
ggtagacgaa gaggaagaca aacctacagt cgcttccaaa ctctagagtt ggaaaaggaa 120
tttcttttta acccctatct gaccaggaaa agaagaatcg aggtttccca cgccctagcc 180
ctcaccgaga gacaggtaaa aatctggttc cagaacagga gaatgaaatg gaaaaaggaa 240
aacaacaagg acaaatattcc cgtttcccg caggaggtga aggacgggga aacgaaaaag 300
gaagcccaag agctggagga agacagagcc gaaggcctga caaattaact tctaccttta 360
aaatttacca cagactatta aaactaataa tcaccatag ctgtggacac cacctatatt 420
ctttggtgga aaggacctta cctgngtttc aagctacctt catgtcactg gtcttgaggg 480
tttctgggct tttagagagg antttgngg tttaaaaang ttntag 526

<210> 397
<211> 443
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (94)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (443)
<223> n equals a,t,g, or c

<400> 397
gccccacgct cccgggtgctc cttacctccc cgggtggtcgg gcgcgaatta ctggaaattg 60
gcttttcccg ttggggccga agtaccttcc ctgnggcggc gactcagcgg ggtgtcgttc 120
ggccggcgtg acgcagccgg atcggcgcca gacggaaaacc tagcggtgac tgtatctgaa 180
ttttgcagct gcagaatgtg tagtacctta aaagggttggc aacaatgagt aaaccagaat 240
taaaaggaaga caagatgctg gaggttcact ttgtgggaga tgatgatgtt cttaatcaca 300
ttctagatag agaaggagga gctaaattga agaaggagcg agcgacttt tggtcaacct 360
caaaaaaata ataaagaagc cagaatatga ttggaggaa gatgaccagg aggtcttaaa 420
agatcagaac tatgtgaaa ttn 443

<210> 398
<211> 652
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (31)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (64)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (68)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (78)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (88)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (91)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (100)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (103)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (111)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (116)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (134)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (261)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (383)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (397)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (549)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (579)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (594)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (606)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (620)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (639)
<223> n equals a,t,g, or c

<400> 398
gcccacgcgt ccgnanacgc ntccgtttgg nccaagatgg catggaatga ggaaggcgct 60
tccttgngga ctggtggnct tggcgtnct ncgggctgtn agnacctcat ntatgngcac 120
ttaccaaaag cggntgaaaa ttgtggaagt tggccccga gatggactac aaaatgaaaa 180
gaatatcgta tctactccag tgaaaatcaa gctgatagac atgctttctg aagcaggact 240
ctctgttata gaaaccacca nctttgagtc tcctaagtgg gttccccaga tgggtgacca 300
cactgaagtc ttgaagggca ttcanaagtt tcctggcatc aactaccag tcctgacccc 360

```
aaatttgaaa ggcttcgagg cancggnacc atgactnctg tgaggatgca gcactccctg 420
gcaggtcaga cctatgccgt gcccttcacg cagccagacc tgcggcgaga ggaggccgtc 480
cagcagatgg cggatgccct gcagtacctg cagaaggctc tggagacatc ttcagcaagt 540
gggtgctgnc actcaccgcc acctgatgag agggccatnc ctgtctgggc aatnccagca 600
acacancctt tgggagcaan ccccttgggg aatcccggnc tggggaaccc at 652
```

<210> 399

<211> 341

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (39)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (84)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (137)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (153)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (302)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (310)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (332)

<223> n equals a,t,g, or c

<400> 399

```
ggcacgagag gacctggata aagcacttct aaaggcctnt caggacatgt tcgacaagaa 60
aaccaaggct tccctttacc tctnactca caatgggaac atgtacacct catccctgta 120
cgggtgcctg gccctgnttc tgtcccacca cnttgcccaa gaactggctg gctccaggat 180
tggtgccttc tcttatggct ctggttttagc agcaagtttc ttttcatttc gagtatcccg 240
```

tctaaagggtg ttctgcagat ccatggaaaag cttctgggaa acgtatgcta gcagagcttc 300
tncccggtgan tcatatTTTT aagatccac tnttagctgg a 341

<210> 400
<211> 604
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (392)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (484)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (502)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (509)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (559)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (593)
<223> n equals a,t,g, or c

<400> 400
gcctccgcat aggcctgcgt cagattaaaa cactgaactg acaattaaca gcccaatata 60
tacaatcaac caacaagtca ttattaccct cactgtcaac ccaacacagg catgctcata 120
aggaaagggtt aaaaaaagta aaaggaaactc ggcaaatctt accccgcctg ttaccacaaa 180
acatcacctc tagcatcacc agtattagag gcaccgcctg cccagtgaac catgtttaac 240
ggccgcggta ccctaaccgt gcaaaggtag cataatcact tggttcctta ataggacac 300
gtatgaatgg cttcacgagg gttcagctgt ctcttacttt taaccagtga aattgacctg 360
cccgtgaaga ggcggcatga cacagcaaga cnaagaagac cctatggagc ttttaattat 420
taatgcaaac agtacctaca aaccacacagg tcctaactac caaacctgca ttaaaaaatt 480
cggntggggc gacctcggac anaaccaanc tccagcagta catgctaaac ttaccagcaa 540
acgactatat ctaattganc atacttgaca aggacaagta cctaggaaca ggnaatcttt 600
ttaa 604

<210> 401
<211> 519
<212> DNA
<213> Homo sapiens

<400> 401
cataatcact tgttccttaa ttagggacct gtatgaatgg ctccacgagg gttcagctgt 60
ctcttacttt taaccagtga aattgacctg cccgtgaaga ggcgggcata acacagcaag 120
acgagaagac cctatggagc tttaatttat taatgcaaac agtacctaac aaaccacag 180
gtcctaaact accaaacctg gcattaaaaa ttctgggttg ggcgacctcg gagcagaacc 240
caacctccga gcagtacatg ctaagacttc accagtcaaa gcgaactact atactcaatt 300
gatccaataa cttgaccaac ggaacaagtt accctaggga taacagcgca atcctattct 360
agagtccata tcaacaatag ggtttacgac ctcgatgttg gatcaggaca tcccgatggt 420
gcagccgcta ttaaagggtc gtttgttcaa cgattaaagt cctacgtgat ctgagttcag 480
accggagtaa tccaggtcgg tttctatcta ctttcaaat 519

<210> 402
<211> 265
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (23)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (213)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (218)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (233)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (255)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (264)
<223> n equals a,t,g, or c

<400> 402
gctgccagta gcatatgctt gtntcaaaga ttaagccatg catgtctaag tacgcacggc 60
cgggtacagt aaactgcgaa tggctcatta aatcagttat gggtcctttg gtcgctcgct 120
cctctcctac ttggataact gtggtaattc tagagctaata acatgccgac gggcgctgac 180
ccccttcgcg ggggggatgc gtgcatttat canatcanaa ccaacccggt canccctct 240
ccggcccccg ccggngggcg gcgnc 265

<210> 403
<211> 325
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (137)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (138)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (172)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (173)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (200)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (263)
<223> n equals a,t,g, or c

<400> 403
gcagggtactt gggtctccag tgggagctgc cctctcgaag gcaggacagc ggtggcgga 60
gatataaaga cctgaagata gtcttttctg tccaaagatg gaaaacagta ctactacat 120
ttctcgggag gagcttnnag aactacaaga ggcatttaata aaaatagata tnnacaatag 180
tgggtatgtc agtgactatn aacttcaaga cctgtttaag gaagcaagcc ttcctctgcc 240
tggctacaag gtgcgagaga ttntggagaa aattctatca gttgctgaca gcaacaaaga 300
tggcaaaatc aattttgaag agttt 325

<210> 404

<211> 540
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (473)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (489)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (526)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (537)
<223> n equals a,t,g, or c

<400> 404
ataactatat ctttcttctt ctgtctccgc ccacccacat ttttctcttt ccctttctct 60
ctttggggac cctcccccat gctcccctgc cccatcccct ttcccccctc ccgtcttctc 120
atccctccat tcccatcttt cccagcaat taccagctct ggctgggtcg ccacaacttg 180
tttgacgacg aaaacacagc ccagtttgtt catgtcagtg agagcttccc acaccctggc 240
ttcaacatga gcctcctgga gaaccacacc cgccaagcag acgaggacta cagccacgac 300
ctcatgctgc tccgcctgac agagcctgct gataccatca cagacgctgt gaaggctcggg 360
aagttgcccc ccaggaacc cgaagtgggg gagcacctgg ttggcttccg gctggggcaa 420
gcattgaacc agaagaattt cttaatttca gaagatcttc aaatggtgng ganccttcaa 480
aaatcctgnc taaaggaata aatggcgaaa aaagccaac gttcanaaag gtggacngga 540

<210> 405
<211> 319
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (94)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (296)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (298)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (301)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (309)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (313)
<223> n equals a,t,g, or c

<400> 405
ctgcnccgcc gcgggcccga gctgcgcgct gaaggccggc aagacagcga gcggtgcggg 60
cgaggtgtgt cgctgtctgt ctgagcagag cgtnggccat ctcgcgctgc gccggggggcc 120
cggggcgcgcc ctgcctgccc tgctggacga gcagcaggta aacgtgctgc tctacgacat 180
gaacggctgt tactcacgcc tcaaggagct ggtgcccacc ctgccccaga accgcaagtg 240
aagcaagggtg gagattttcc agcacgtcat cgactacatc agggacttta atttgnantt 300
naatcggant ccnaatttg 319

<210> 406
<211> 355
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (10)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (23)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (27)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (41)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (46)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (57)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (86)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (121)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (124)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (145)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (150)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (171)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (216)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (251)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (266)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (283)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (304)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (321)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<400> 406

ggcanaggtg cagctgaccc tgcgggncac acagtgccca nagggnaagt ccgtaanatg 60
ccacgtgaaa gcactacacg attccnagcc aggatgtaac tgtgcccctg cccagttccc 120
ncancttccc catgctgccca ncccngattn tggctgaaac gaccggccct ngaggacctg 180
gtcttaggtt cagaagcgaa gcatcacgtg acacantgac cggcctgaga gaatgcctct 240
ggtgaccact ntnacctggg acgcenttaa agtgggaaga gcncgtgttc aaggaccacc 300
tggngcgtga cctgtgtggg nttgntacag gtgttccagt gtgctgactg gttgt 355

<210> 407

<211> 437

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (110)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (272)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (296)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (370)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (389)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (412)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (430)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (435)
<223> n equals a,t,g, or c

<400> 407
ggcanagcac aagcccagca acaccaaggt ggacaagaga gttgagtcca aatatggtcc 60
cccatgcccc tcatgcccag cacctgaatt tcctgggggg accatcagtn ttctgtgtcc 120
ccccaaaacc caaggacact ctcatgaatc tcccggaccc ctgaaggta cgtgcgtggt 180
ggtggacgtg aagccaggga agaccccag gtccagtcca actggtacgt ggatggcgtg 240
gaggtgcata atgccaagac aaagccgcgg gnggagcagt tcaacagcac gtaccngtgg 300
ttcagcgtcc ttcaccgtcc tgcaccagga tggcttgaac ggcaaggagg ttacaagtgg 360
caaggttttn caacaaaggg cttcccgtnc ttcttgggg aaaaccttt tncaaaggcc 420
aaggggcagn cccgngg 437

<210> 408
<211> 310
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (102)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (137)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (277)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (289)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (294)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (298)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (303)
<223> n equals a,t,g, or c

<400> 408
gggggatatt gtagtggtgg tagctgctct aatttctact tctaccacat ggacgtctgg 60
ggcgaaagga ccacggtcac cgtctcctca gcctccacca anggccatc ggtcttcccc 120
ctggcgccct gctccangaa cacctccgaa aacacagcgg cctgggctg cctggtaag 180
gactacttcc ccgaaaccgg tgacggtgto ttggaactca gggggctctg accagcgggg 240
tgcacacctt cccagctgtc ctacagtcct caggaancta ctccctcanc accntggnga 300
cgntgcctcc 310

<210> 409
<211> 421
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (52)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (74)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (235)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (246)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (279)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (346)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (382)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (384)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (405)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (417)

<223> n equals a,t,g, or c

<400> 409

```
gctcagtttag gacccataacg gaaccatgga agccccagcg cagcttctct tntcctgct 60
actctggctc ccanatacca ctggagaaat actgatgacg cagtctccag ccaccctgct 120
tgtgtctcca ggggaaagag tcaccctctc ctgcagggcc gggcagagtg ttacagcaa 180
cttagcctgg tatcagcaga aacctggcca ggctcccagg ctctcatgt atggntcatc 240
caccanggcc actgatgtcc cagtcagggt cagtggcant gggctctggga cagagttcac 300
tctcaccatc agcagcctgc agtctgacga ttctgcagtc tatttntgct agcagtatat 360
tatgtggcct ggaaccttcg gncnagggac caagggggaa atcanacgaa ctggggntgc 420
a 421
```

<210> 410

<211> 448

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (9)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (342)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (361)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (384)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (421)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (424)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (443)
<223> n equals a,t,g, or c

<400> 410
gtacggcnnng ntnntttcgg gtcgaccgcg ccttccgcgg acgcgtgggt tcggacgcgt 60
gggcggaacgc gtggggccga ggctgccaag atgcttgag aagcactgag caagaaccct 120
ggctacatca aacttcgcaa gattcgagca gccagaata tctccaagac gatcgccaca 180
tcacagaatc gtatctatct cacagctgac aaccttgtag tgaacctaca ggatgaaagt 240
ttcaccaggg gaagtgcag cctcatcaag ggtaagaaat gagcctagtc accaagaact 300
ccaccccccag aggaagtggg tctgcttctc cagttttgga gnaccaccag gggccagaa 360
nagccctacc ccgcccatt atcntgcaat ggtcccaca ccgggttccc tgaaccctct 420
nngnttaagg aaaatgaaa tanccctt 448

<210> 411
<211> 141
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (67)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (95)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (101)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (120)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (138)
<223> n equals a,t,g, or c

<400> 411
gattgatctc atgtgcaaga aaatgaagca cctgtggttc ttcctcctgc tggtagcagt 60
ctcccanatg cgtcctgtcc caggtgcagc tgcangagtc ngggccagga ctggtgaagn 120
cttcggagac cctgtccntc a 141

<210> 412
<211> 473
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (403)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (461)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (471)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (472)
<223> n equals a,t,g, or c

<400> 412
ggaaaggagc cccagccctg agattcccag gtgtttccat tcggtgatca gcactgaaca 60
cagaactcac catggagttt ggactgagct gggttttcct tgttggtatt ttaaaagggtg 120
tccagtgtga agtgcagctg gtggagtctg ggggagccgt agtacagcct ggggggtccc 180
ttagactctc ctgtgaagcc tctggattca cctttgacaa ttatgccatg cactgggtcc 240
gtcaagctcc ggngaagggt ctggagtggg tctgtctcat cagtcgggat ggtcgtgaga 300
catattttgc agactctatg aagggtcggg tcaccatctc cagagacaac agcaaaaact 360
gcctgtatct ccaagtgaac agtctgagag ttgaggacac cgncttgat tactgtgcaa 420
aagatatccc ggggtcgtcg gtatggacgt ctggggtaaa nggacaccng nna 473

<210> 413
<211> 328
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (183)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (187)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (188)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (228)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (241)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (242)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (288)
<223> n equals a,t,g, or c

<400> 413
ggctctggac tccaaggcct ttccacttgg tgatcagcac tgagcacaga ggactcacca 60
tggaattggg gctgagctgg gttttccttg ttgctatattt agaaggtgtc cagtgtgagg 120
tgcaagctggt ggagctctgg ggaggcttgg tccaagccgg gggggtccct gagactctcc 180
tgngcannct ctggcttccc cctttataac catgggatga cctgggtncg ccaggctcca 240
nngaaggggc tggaatgggt ggccaccata cagtgatgaa aaactganaa atactatgtg 300
gactctgtga agggccgatt caccatct 328

<210> 414
<211> 575
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (14)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (361)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (470)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (511)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (527)
<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (539)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (544)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (573)

<223> n equals a,t,g, or c

<400> 414

```
ggccgtgaga ttcncaggag ttccacttg gtgacagca ctgaacacag accaccaacc 60
atggagtttg ggccatagctg ggttttcctt gttgctattt taaaagggtg ccactgtgag 120
gtgcagctgg tggagtctgg gggaggcttg gtacagccag ggcgggtccct gagactctcc 180
tgtacaactt ctggattcac ctttgagat tattctatga gctgggtccg ccaggctcca 240
gggaaggggc tggagtgggt aggtttcatt agaagcaaag cgcattgggtg gacaacagaa 300
tacgcccgct ctgtgaaaag gcagattcac catctcaaag agatgattcc acaggcatcg 360
nctatctggc aaatgaacag cctgaaaccg aggacacaga cattattact gtctagacat 420
gactacaggc acaccctgg ctactggggg cagggaaccc tggtcaccgn cttctctggc 480
ttccaccaag ggccatcgtc ttccccctgg ngcccttgtt ccaggancac ttccgaaanc 540
cagnggcctg ggcttgctgg gcaagggtc ttncc 575
```

<210> 415

<211> 438

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (37)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (65)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (105)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (139)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (157)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (190)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (201)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (269)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (281)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (296)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (297)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (301)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (303)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (312)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (323)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (330)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (331)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (333)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (350)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (356)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (371)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (385)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (397)
<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (422)

<223> n equals a,t,g, or c

<400> 415

```
gtccagtgtg aggtgcagct ggtggagtct gggggangct tggtagagcc tggggggtct 60
ctganactct cctgtgcagc ctctggattc accttcagta gttangacat gcattgggtc 120
cgccaagtgt caggaanaang tctggaatgg gtctcangta ttgatcctgc tggtaacaca 180
aactatccan gctccgtgaa nggccgattc atcatctcca gagaaaatga caagagctcc 240
tcgtatcttc aaaatgaatg ggctgacanc cggggaaaac ngtgtgtaat attgtnnaaa 300
nanaaattgc anttcctggt aantgggtan ntncgatctc ttggggccgn nggaancctt 360
ggttaaatgt nttcctcaag aattnccga accaagnccc caaagggttt tcccgcctaa 420
ancctttgaa aaagaaac                                     438
```

<210> 416

<211> 502

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (22)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (48)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (105)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (135)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (145)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (312)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (367)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (428)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (451)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (455)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (474)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (480)
<223> n equals a,t,g, or c

<400> 416
ggcagagca acgctggcaa tnataacaca aaatattcac agaagtnca ggacagagtc 60
accattacca gggacacatc cacgaacaca gcctacatgg atctnagcag cctgagatct 120
gaagacacgg ctgtntatta ttgtncgagg ggattttttg gggaccgtga ttactactac 180
tactactaca tggacgtctg gggcaaagg accacggtca ccgtctctc agcatccccc 240
accagcccca aggtcttccc gctgagcctc tgcagcaccc agccagatgg gaacgtggtc 300
atcgctgct tngtccaggg ctctctcccc caggagccac ttcagtgtgg acctggagcg 360
aaaggggnaca gggcgtgacc gccagaaaat tcccacccag ccaggatgcc tccgggggac 420
ctgtacanca cgagcagcca gctgaccctg ncggncacaa gtgccttagc cggnaagttn 480
cgttgacatt gccacgtgga ag 502

<210> 417
<211> 427
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (230)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (233)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (352)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (363)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (388)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (411)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (427)
<223> n equals a,t,g, or c

<400> 417
gatcatgagt atctatgtga attggtatcg gcagaaagca ggaaaagccc ctgagctcct 60
gatccacggt ggcattccagc ttacaaagtg gagtccccgg aaggttcagt ggcaggggga 120
tctgggacag atttcactct caccattagc agtctgcaac ctgaagattt tggcacttac 180
ttctgtcaac agaattacaa tgtcccgtgg aacgttcggc cagggggaacn aangtgga 240
atgaaaccga actgtggctg caccatctgt cttcatcttc ccgccatctg atgagcagtt 300
gaaatcttga actgcctctg ttgtnttgcc tgctgaataa cttctatccc anaaaaggcc 360
aanttcattg gaaggtgat aacccccncc atcggttact cccggaaaat ntcccaaacc 420
ggacgcn 427

<210> 418
<211> 308
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (126)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (132)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (182)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (254)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (262)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (270)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (282)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (297)

<223> n equals a,t,g, or c

<400> 418

gctccaactc tggcaacaca gccactttga ccatcagcgg gactcaggct atggatgagg 60
ctgactatta ctgtcaggcg tgggacagta gcgctgtggt cttcggcgga gggaccaggc 120
tgacntcct angtcagccc aaggctgccc cctcggtcac tctgttcccg ccctcctctg 180
angagcttca agccaacaag gccacactgg tgtgtctcat aaatgacttc taccgggaa 240
gccgtgacag tggncctggaa angcagatan caaccccggt cnaggcgga ttggganaaa 300
caaccaca 308

<210> 419

<211> 482

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature
<222> (370)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (441)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (462)
<223> n equals a,t,g, or c

<400> 419
tcgacccacg cgtccgcagc tgtgggcaca agaggcagca ctcaggacaa tctccagcat 60
ggcctggtct cctctcctcc tcaactctcct cgctcactgc acagggtcct gggcccagtc 120
tgtgtgacg cagccgcct cagtgtctgg ggccccaggg cagaggggtca ccatctcctg 180
cactgggagc agctccaaca tcggggcagg ttatgatgta cactggtacc agcagcttcc 240
aggaacagcc cccaaagtcc tcacttatgg taacagcaat cggccctcag gggtcctga 300
ccgattctct ggctccaagt ctggcacctc agcctccctg gccatcactg ggctccaggc 360
tgaagatgan gttgattatt actgccagtc ctatgacagc agcctgggtg gttcgggtgt 420
cggcggaaag accaagctga ncgtcctang tcagcccaaa gntgccccct cggttaactct 480
gt 482

<210> 420
<211> 462
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (85)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (86)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (100)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (438)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (446)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (452)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (456)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (459)
<223> n equals a,t,g, or c

<400> 420
gacaggattc aggacaatct ncaccatggc cggcttccct ctccctctca ccctcctcac 60
tcactgtgca nggtcctggg ccannctgt gctgactcan ccaccctcan cgtctgggac 120
ccccgggcag aggggtcacca tctcttggtc tggaagcagc tccaacatcg gaactaatta 180
tgtatactgg taccagcagc tcccaggaac ggcccccgaa gtcctcatct ataagaatga 240
tcagcggccc tcaggggtcc ctgaccgatt ctctggetcc aagtctggca cctcagcctc 300
cctggccatc ggtgggctcc ggtccgagga tgaggctgat tattactgtg catcatggga 360
tgacagcctg agtgggtccg tcttcggcgg agggaccaag ctgaccgtct tgggtcagcc 420
caaggctgcc ccctcgggna ctctgntccc gnectnctnt ga 462

<210> 421
<211> 435

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (324)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (391)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (416)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (423)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (431)
<223> n equals a,t,g, or c

<400> 421
ggaggcagag ctctgggaat ctcaccatgg cctggacccc tctcctgctc cccctcctca 60
ctttctgcac agcgggaagag aggccaggcc cctgtggtgg tcattctatga ggacaacaaa 120
cgaccctccg ggatccctga gagattctct ggctccacct cagggacatt ggccaccgtg 180
attatcagtg gggcccagggt ggacgatgat actgacttct actgtcagtc aacacacagt 240
agtaataatg gtaggtccgt atgtcttcgg aactgggacc aaggtcaccg tccttggtca 300
gccccaggcc aaccccaatg tcantctgtt cccggcctcc tctgaaggag nttcaagcca 360
aacaaaggca cactaagtgt gtctgatcag ngagttttaa ccgggaagtg tgaantggg 420
ctnggaagca nattg 435

<210> 422
<211> 334
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (101)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (122)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (135)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (145)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (310)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (323)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (324)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (326)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (331)

<223> n equals a,t,g, or c

<400> 422

gcagaactct nggtgtctca ccatggcctg gatcccteta cttctcccc tcctcactct 60
ctgcacagac tctgaggcct cccatgaatt gagacaacca ntttcggtgt cagtgtcccc 120
angacagacg gccangatca cctgntctgg agatgcattg ccagaacaat ctattttttg 180

```
gtatcaacag aagccaggcc aggccctgt attggtgatt tataaagtcc atgagaggcc 240
gtcagatgcc ctgaacgatt ctctggctcc aggtcacaga caacagtcac gttgaccatc 300
agtggagccn agggcaaaaa atnngngggg ngtg 334
```

<210> 423

<211> 597

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (26)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (61)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (81)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (115)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (117)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (118)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (132)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (168)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (174)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (176)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (189)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (199)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (201)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (204)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (209)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (217)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (230)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (257)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (259)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (272)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (277)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (296)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (333)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (344)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (393)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (412)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (413)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (414)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (427)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (457)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (492)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (511)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (513)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (550)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (552)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (580)
<223> n equals a,t,g, or c

<400> 423
nggatttngc caagctctga atcaanctca ctaaaggga caaaagggtg agctccaccg 60
nggtggcggc cgctctagaa ntactggatc ccccggttg cccgggttcg gcganannng 120
acaatntagc cnatgtaggg tctggtcccc caaatccaga tctgtcgnaa acantnttgg 180
atcaatacnt tgccatggnc ncanaaaanc atgggtncaa catggaacan gctcttgga 240
tgctcttctg gcataancnt aatatccaaa antcatnggc tgatttgccc aacttnaccc 300
ctttcccaga taagtggact gtggaagata aantcttatt tgancaagcc ttacttttc 360
atgggaaaac tttcataca atccaaccaa tgnttcaca taaatctata gnnngtctg 420
tgaaatntta ctattcttg aaaaaagacg aagactnaaa ctattgtgat ggatcgccat 480
gccccgaaa cnaaacggga cgggaagaga ncnacgatga actggaacaa gcaaatggaa 540
caatcccatn gnacttgaag ttggatccaa accaagaaan ccaaaggga gtcccct 597

<210> 424
<211> 143
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (42)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (92)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (96)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (99)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (131)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (138)
<223> n equals a,t,g, or c

<400> 424
gaacacatat aagagcaaaa tctggatcgg caggtgttac anacacaatg tatacgactg 60
gaggcccggt actacagcct cagcctgacg gnacancanc tctcccacat cgtggcggag 120
ttgaggaacc ngaaacanaa aat 143

<210> 425
<211> 323
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (7)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (291)
<223> n equals a,t,g, or c

<400> 425
gggcagngga gctctattgc caccatgagt ttctccggca agtaccaact gccagagcca 60
ggaaaacttt gaagccttca tgaaggcaat cggctctgccg gaagagctca tccagaaggg 120
gaaggataatc aaggggggtgt cggaaatcgt gcagaatggg aagcacttca agttcaccat 180
caccgctggg tccaaagtga tccaaaacga attcacggtg ggggaggaat gtgagctgga 240
gacaatgaca ggggagaaag tcaagacagt gggtcagttg gaaggtgaca ntaaactggt 300
gacaactttc aaaaacatca agt 323

<210> 426
<211> 683
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (79)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (94)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (105)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (112)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (114)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (120)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (124)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (129)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (134)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (136)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (139)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (158)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (163)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (188)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (195)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (206)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (232)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (234)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (241)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (251)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (262)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (264)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (275)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (286)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (291)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (301)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (324)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (326)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (338)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (370)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (395)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (399)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (409)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (418)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (487)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (536)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (558)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (578)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (601)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (607)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (633)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (648)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (662)
<223> n equals a,t,g, or c

<400> 426

gaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
aaaaaaaaaa aaaaaaaang ggcggccgtt ttanaggatc caagnttacg tncncgtgcn 120
tgcnacgtna tagntntntnt atagggtcac cttaaattnaa ttnactggcc gtcgttttac 180
aacgtcgnga ctggnaaaac cctggngtta cccaacttaa tcgccttgca gnanatcccc 240
ntttcgccag ntggcgtaat ancnaaaagg cccgnaccga tcgcctttcc naacagttgc 300
ncagcctgaa tggcaaattg gacnccct gttagcgngc attagcncg gcgggtgtgg 360
gggttaccn cagcgtgacc gttacanttg ccagngccnt agcgcccgnt cctttcgntt 420
tcttcccttc ctttttcgcc acgttcgccg gttttccccg taaagcttta aatggggggc 480
tcccttnagg gttccgattt agggctttac gggaccttga ccccaaaaaa ctttgnttag 540
ggggatgggt cagctagnng ggccattgcc cttgatanac gggtttttcg cccttttgac 600
nttggantcc caggtttttt aaataggggg gcntttttgt tccaaaantg gggaacaaac 660
antttaaccc ctttttttgg ggg 683

<210> 427

<211> 369

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (323)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (337)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (344)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (349)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (350)

<223> n equals a,t,g, or c

<400> 427

tgggcctgag atgcaggtaa aactgctcca gtccctgggc ctgaaatcaa ccctcatcac 60
cgatggctcc actcccatca acctcttcaa cacagccttt gggctgctgg ggatggggcc 120
cgagggtcca gccctgggc agaaaggttg gcattgggcc cagccctgga agggggatat 180
ccccccagtc ttgctcaagc ccctcaagct cctggaaaac accactttgt gcctgttctg 240
cgcttactcc tgattaatac aatgaatttt cttgggcatt ttacaatttc caacacttta 300
aaaaaaaaaa aaaaaaattc aanggggggg ccggttncca attnccccnn ttttatttct 360
tttaaaatc 369

<210> 428
<211> 299
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (79)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (121)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (295)
<223> n equals a,t,g, or c

<400> 428
ncctaatactc ggaatcctaa gtggtcgctt gaggaccggc cggaattccc gggcaccac 60
gcgtcggcgg tagctgctnt tcctggggcc cgtgggcctc atcatgtacc tcgggggcgt 120
nttcttcac aaccggcagc gctctagcac tgccatgaca gtgatggccg acctgggcga 180
gcgcatggtc agggagaacc tcaaagtgtg gatctatccc gaggggtactc gcaacgacaa 240
tggggacctg ctgcctttta agaagggcgc cttctacctg gcagtcagg cacangtgc 299

<210> 429
<211> 538
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (124)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (133)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (325)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (374)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (376)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (394)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (397)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (414)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (432)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (435)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (440)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (460)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (472)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (473)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (482)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (484)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (498)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (499)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (529)
<223> n equals a,t,g, or c

<400> 429
gaagcagaga tctgaagaca gcatgtacac agccattccc cagagtggct ctccattccc 60
aggctcagtg caggatccag gcctgcatgt gtggcgggtg gagaagctga agccggtgcc 120
tgtngcgcaa ganaaccagg gcatcttctt ctcgggggac tcctacctag tgctgcacaa 180
tggcccagaa gaggtttccc atctgcacct caacacactg ctgggagagc ggcctgtgca 240
gcaccgcgag gtaaggggca atgagtctga cctcttcatg agctacttcc cacgggggctt 300
caagtaccag gaaggtgggt tggantcagc atttcacaag acttccacag gagccccagt 360
tgccatcaag aaantntacc aggtgaaggg gaanaanaaa tccgtccaac gagngggcat 420
gaattgggaa anttnaaatn ggggttggtt acctggaatn ggcaaaaatt tnnctgggtt 480
gngnaatcaa atttganna aaagggggga ttgcctggat cgggatttnc aaggaagc 538

<210> 430
<211> 552
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (434)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (505)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (508)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (514)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (523)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (535)
<223> n equals a,t,g, or c

<400> 430
gcggacacgg acaggattga cagattgata gctctttctc gattccgtgg gtggtggtgc 60
atggccgttc ttagttggtg gagcgatttg tctggttaat tccgataacg aacgagactc 120
tgccatgcta actagttacg cgacccccga gcggtcggcg tcccccaact tcttagaggg 180
acaagtggcg ttcagccacc cgagattgag caataacagg tctgtgatgc ccttagatgt 240
ccggggctgc acgcgcgcta cactgactgg ctccagcgtgt gcctacccta cgccggcagg 300
cgcgggtaac ccgttgaacc ccattcgtga tggggatcgg ggattgcaat tattccccat 360
gaacgaggaa ttcccagtaa gtgcgggtca taagcttgcg ttgattaagt ccctgccccct 420
tcaacccttc tggnccttcg gaccactgng acttttccat ctttcttaaa cggaggctgg 480
gccttggggg gggggctggc ctggnntntg ttgncccaa ganggtgctt tcaanggtga 540
ggaacttttt tt 552

<210> 431
<211> 218
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (31)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (36)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (37)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (43)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (56)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (60)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (83)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (104)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (105)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (107)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (136)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (143)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (159)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (168)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (171)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (181)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (196)
<223> n equals a,t,g, or c

<400> 431
ccccggcct gcaggaaatt cgagcagagt ncacannag ggnaccaccg tcctgntctn 60
ccagttcggg ttgaatgcaa gtncctagccg attttcccta caannantcc agttgattac 120
aattcttcct gtacgncaga gancctgcc tttaaaagnt gccaacgntt ncctgacgag 180
ncctgcagcc acagtncggc aattcctaca agtgccaa 218

<210> 432
<211> 610
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (401)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (587)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (590)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (592)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (593)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (604)
<223> n equals a,t,g, or c

<400> 432
gcacacccct gggcgcccc gtcacccagc cgtatgcctt ggactccaag gtcaagcaaa 60
taggaggtgg aattcagagc atcacctaca cccacaacgg agacatcagc cgaaagccca 120
acacacgtaa gcagaagaac ggcttccccg ccaacttcat ccactcgctg gactcctccc 180
acatgatgct caccgcccctg cactgctaca ggaagggcct gaccttcgtc tctgtgcacg 240
actgttactg gactcacgca gctgatgtct ccgtcatgaa ccagggtgtgc cgggagcagt 300
ttgtccgctt gcacagcgag cccatcctgc aggacctgtc cagattcctg gtcaagcggg 360
tctgctctga gcccagaag atcttgaggg ccagccagct naaggagaca ctgcaggcgg 420
tgcccaagcc aggggccttc gacctggagc aggtgaagcg ttccacctac ttcttcagct 480
gacaccccggt gaggccttgc agtgtgtaaa taaagctctt ttgccacccc aaaaaaaaaa 540
aaaaaaaaact tggggggggg gcccgggacc caattgccct atagggnggn gnnttacaat 600
tcantggccg 610

<210> 433
<211> 328
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (26)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (123)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (131)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (132)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (222)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (223)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (225)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (244)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (249)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (266)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (291)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (314)

<223> n equals a,t,g, or c

<400> 433

```
caaatgtatg gaactaaca gatggtcca tatcgagatt caaagttaac ccattctgttc 60
aagaactact ttgatgggga aggaaaagtg cggatgatcg tgtatgtgaa cccaaggct 120
gangattatg nngaaaactt ncaagtcatt agatttgcgg aagtgactca agaagttgaa 180
gtagcaagac ctgtagacaa ggtaatatgt ggtttaacgc cnngnaggag atacagaaac 240
cagntcngg gtccagttgg aaatgnacca ttgggtactg acgtggtttt ncagagtttt 300
ccacctttgc cgtncatgcg aaattttt 328
```

<210> 434

<211> 535

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (76)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (77)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (274)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (351)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (405)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (454)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (491)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (504)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (512)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (517)
<223> n equals a,t,g, or c

<400> 434
gggttcgacc cacgcgtccg aggctgaaca ggacagaaag atagaagaag tcagagatgc 60
catggaaaat gaaatnngaa actctctgaa caagaattac aatttcgtcg tctcagtcaa 120
gagcaagttg acaactttac tctggatata aatactgcct atgccagact cagaggaatc 180
gaacaggctg ttcagagcca tgcagttgct gaagagggaag ccagaaaagc ccaccaactc 240
tggttttcag tggaggcatt aaagtacagc atgnaagacc tccatctggc agaaacacct 300
actatcccgc tgggtagtgg cagttgaagc ccatccaaag ccaactgttc nggataatga 360
atttcaccca agctttaacc gcagctatc ccttcagag ttccnggacc cgtgggggtg 420
ttacatggaa gagaaccctt agggcccgtt ttcncatggt ggtttccaaa aattggggccc 480
cgagggtagg nccaatggat tggnttgga ancccgnaat tagcttgta cccgt 535

<210> 435
<211> 524
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (138)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (154)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (157)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (238)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (258)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (259)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (262)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (269)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (279)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (308)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (337)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (338)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (339)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (340)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (341)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (342)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (343)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (344)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (345)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (347)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (350)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (354)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (375)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (380)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (498)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (500)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (518)

<223> n equals a,t,g, or c

<400> 435

```
ccctgctcca ccgtccccgt caccacggag gtttcgtacg ccggtgcac caagaccgtc 60
ctcatgaatc attgctccgg gtccctgcggg acatttgtca tgtactcggg ccaagcccag 120
gccctggacc acagcttncct cctgctgcaa aganganaaa accagccagc gtgagggttg 180
tcctgagctt gcccgaatgg cggctcgtt gacacacacc tacaccaca tcgaaganct 240
gccagtgcc aagaacacnnt cntgggggnt tcccgaacng gaacttccc cgggggccgg 300
ggggcttncc ccttagcatc tggaacgtag gtttttnnnn nnnnnngnnan aaanccccct 360
ttcaattgcc cttcnaaaan ttttacctcc ccccgaaact ctttaaaact cttaaactcg 420
ggttcctctt cttccaaata ttatgttcta atttttttt tcatccctgc tttcccaata 480
taaactcagg gggaaatncn aaaaaaaaaa aaatccangg gggg 524
```

<210> 436

<211> 384

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (79)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (124)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (177)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (191)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (215)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (320)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (341)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (349)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (366)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (374)

<223> n equals a,t,g, or c

<400> 436

ggcanaggat caggctggag aagtggatca caccttggtg ggacaatgca caggtggcgg 60
ttacttcacg cagttcagna ccagctcggg gtccgcggaa gaggcagccc tactggagtc 120
tcgnattcct taccctaaaga ggaagcagca gtgcctgcaa tttttctata aaatgancgg 180
ggaagtcctt ncagacagac ttcgttgtct gggtnccagga ggggatgaca gcacaggcaa 240
tgttcgcaat tgggtgaagg gcagactttt caaggagatg atgaccacaa tttggaaaat 300
tggcccatgt ggggtgcttcn aagaggaaca gaatttcggg naccttttnc caggggcaca 360
aaaggnggac ctnagagct tcaa 384

<210> 437
<211> 390
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (304)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (317)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (372)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (385)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (386)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<400> 437
cttgctcaat gacgccgtca ctgtggctct gtatcacctc tttgaggagt ttgccaaacta 60
cgaacacgtg ggcacgtgg acatcttcct cggttcctg agcttcttcg tgggtggcct 120
gggcgggggtg cttgtgggcg tggctacgg ggtcatcgca gccttcacct cccgatttac 180
ctccacatc cgggtcatcg agccgtctt cgtcttcctc tacagctaca tggcctactt 240
gtcagccgag ctcttcacc tgtcaggcat catggcgctc atagcctcag gagtggatgat 300
gcgncctat gtgggangcc aacacttcca caagttccca caacaacatc aaataatttc 360
ctggaagatg tnggagcagc gtcannaagn 390

<210> 438
<211> 234
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature

<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (24)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (47)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (82)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (114)
<223> n equals a,t,g, or c

<400> 438
nccccggact tttagtgttaa ngcncctacc ccctgcaggt accggtncgg aattcccggg 60
tttgactcca cgcgttttgc tncattaaca acataaaacc ctcatggaca cganaaaaca 120
ccctcatggt catacaccta tccccattc tctctctatc cctcaacccc gacatcatta 180
ccgggttttc ctcttaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 234

<210> 439
<211> 144
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (36)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (43)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (105)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (114)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (135)
<223> n equals a,t,g, or c

<400> 439
nnctatcgat ggggagcaag agtcttttaa ataagnccat ttnagtcctt ggtaacaag 60
ggtttaaagt ggagcgaatg cacatcacag acatgaaatt ggctnacctg cctngcttag 120
aagcccttgg tgttnaggtc aaca 144

<210> 440
<211> 411
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (69)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (86)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (132)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (209)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (230)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (246)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (276)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (340)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (346)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (390)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (404)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (410)

<223> n equals a,t,g, or c

<400> 440

```
gccaaagtcc accccaactc tgtgcacatc tgtgcggtgg ttgtggagta cgagaccaag 60
gctggccgna tcaacaaggc cgtggncacc aactggctgc gggccaagga gcctgccggg 120
gagaacggcg gncgcgcgct ggtgcccatg ttcgtgcgca agtcccagtt ccgcctgccc 180
ttcaaggcca ccacgcctgt catcatggng ggccccggca ccgggggtgtn acccttcata 240
ggcttnatcc aggagcgggc ctggctgcga cagcanggca aggaggtggg ggagacgctg 300
ctgaactacg gctgccgccg ctcggatgag gactacctgn accggnagga gctggcgagc 360
ttccacaggg acggtgcgct caccagctn aacgtggcct tctnccgggn a 411
```

<210> 441
<211> 623
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (73)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (162)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (232)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (252)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (289)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (360)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (365)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (407)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (435)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (472)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (495)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (502)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (504)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (509)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (538)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (560)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (586)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (587)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (619)
<223> n equals a,t,g, or c

<400> 441
gccagcagnc accacagcgc ctgcttcctc ggccctgaaa tcatgcccct aggtctcctg 60
tggtggggcc tancctgtt gggggctctg catgcccagg cccaggactc cacctcagac 120
ctgatccag cccacctct nagcaaggtc cctctgcagc anaacttcca cgacaaccaa 180
ttccacggga agtggtatgt ggtacgcctg gcacggaatg caattctcag anaacacaaa 240
gaccgcgaaa anatgtatgc caccatctat gagctgaagg aaacaaganc tacaatgtcc 300
ctccgcctgt ttaagaaaaa aaaatgtgac tacttggatc aggaattttg gtccaaggtn 360
gccanccggc gaattccacc ttggggacct ttaaaattgc cttggantaa ccaatttcct 420
ccttccaatt gttancacca attacaacng ccttcttttg gttttcttcc anaaaatttc 480
tccaaacaag gaatncttcc anancccccnc ttccggaaaa acaaggaatt aattccgnac 540
ttaaaggaaa aattctccgn tctcccattt cttgggcccc ctaaanncaa attcgtcttc 600
ccttttccaa ccaacattnt ttt 623

<210> 442
<211> 211
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (47)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (58)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (77)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (93)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (121)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (131)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (178)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (195)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (207)
<223> n equals a,t,g, or c

<400> 442
tgggcagcca cactgcacgc aggctgggcc gactgagggg ctcagangcc aggtctctnag 60
gcccacgcag ggcctanggt gggaagatgg cangtggggg cggcgacctg agcaccagga 120
ngctgaatga ntgtatttca ccagtagcaa atgagatgaa ccattcttct gcacacanc 180
acgatttgca aaggntgttc acggaanacc a 211

<210> 443
<211> 399
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (316)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (347)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (358)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (391)
<223> n equals a,t,g, or c

<400> 443
gtctgagtgg atggacactg cctcttagaa ctagaactta gaactttatc ttgaaaatgt 60
accactgttg cagaagctcc tcacagagta tgtgtcaggc atttttaacc tgctaaaggc 120
aagaagaagt gttcaccaca tagttgcaaa ggtcttcaac ttgccacagc caacagaaaa 180
atcaaaatga ttgaaccctt tgggaatcaa gtatattgtg gccaggccag tgtattctac 240
aaatgctttt gaggaaaatc ataaaaagac aggaagacat cataagacat ttctggatca 300
tctcaaagtg tgttgnaact gttccccaca aaaggcaaga gaattgncct ctctttgntt 360
cccatagcat tttgntgcc agcataccgg nttaaagaa 399

<210> 444
<211> 465
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<400> 444
gcgggactcc aaatgggtcg cagtcgcagc cgctctccac ggagggaacg taggcgttcc 60
cggtccacat cccgggagag agaacgcagg cgccgagaaa ggtccaggtc tcgggagaga 120
gatcgagaaa ggagccgctc gcgatccccg caccgaagac gctcccgatc tccaagacga 180
catagatcca catctccttc ccttctcga ctgaaaagaaa gaagagatga ggaaaagaaa 240
gaaacaaaaa aaacaaagag caaagaacgg cagattactg aggaagactt agagggcaaa 300
acagaggaag aaatagaaat gatgaagtta atgggatttg cctcctttga ctccacaaaa 360
ggtaagaagg tggatngctc tgtaaatgcn tatgccataa atgtctctca gaagaggaag 420
tacaggtatg catagcccc atttttgttt gaactaataa atcag 465

<210> 445
<211> 297
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<400> 445

```
ggcagtnctct ctgccaggct ggggatgatt ccaatagtaa taagaagaat gctgacctcc 60
aagtgtctaaa gcccgagccc gagctgggtt atgaagacct gaggggctca gtgaccttcc 120
actgtgccct gggccctgag gtggcaaacg tggccaaaat tctgtctggc agagagtggg 180
gaaaagacgc ggtttccagc ttgcagattt gttaagtttc tcaggcagat ttgactttc 240
agcctttcat acttgtttaa gcaactattt gtattaaatg aagtttttg aaaaaca 297
```

<210> 446

<211> 448

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (22)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (24)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (306)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (366)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (369)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (435)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (440)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (448)
<223> n equals a,t,g, or c

<400> 446
tcgacccacg cgctccgccga nnanagtttg gttttgcgga ttgccaccga tgattccaaa 60
gcagtctgtc gcctcagtg taaattcggg gccacactca aaatcagcag gcttctcttg 120
gaacggggcga gagagctgaa tattgacatc atcgggtgtca gcttccacgt gggaagtggc 180
tgcaccgatc ctggagacct tcgtgcaagc catctccgat gcccgctgtg tcttcgacat 240
ggggagctga gggttggttc aacatgtatc tgcttgatat cgggtggtggg ctttcctggg 300
atctgnagga tgtgaaactt aaatttgga gagatcacca tgttatcaac ccagccctgg 360
gacaantant tcccggggccg aattcggggcg tggacattca tagccgagcc cgggcaggat 420
aattaagttg gcttncagcn ttcaagcn 448

<210> 447
<211> 268
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (10)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (36)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (43)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (45)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (60)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (99)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (132)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (135)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (140)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (159)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (183)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (192)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (206)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (222)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (243)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (260)

<223> n equals a,t,g, or c.

<220>

<221> misc feature

<222> (262)

<223> n equals a,t,g, or c

<400> 447

```
ggcacgaggn agtgtcccag ncggttcgt gtcgcnatgg ggnanatcga gtgggccatn 60
tgggccaacg agcagggcgt ggcgtccggc ctaatcctna tcaccggggg catcgtggcc 120
acagctgggc gnttnaccn ntgg tacttt ggtgcctant ccattgtggc gggcgtgttt 180
gtntgcctgc tngagtacc ccgggnaaag aggaagaagg gntccaccat ggtgcgatgg 240
ggncagaagt acatgaccgn cntggtga 268
```

<210> 448

<211> 425

<212> DNA

<213> Homo sapiens

<400> 448

```
gacaccttca tccgtcacat cgcctgctg ggctttgaga agcgcttcgt acccagccag 60
cactatgtac atgttcctgg tgaaatggca ggacctgtcg gagaaggtgg tctaccggcg 120
cttcaccgag atctacgagt tccatctccc aagtggtttg acgggcagcg ggccgccgag 180
aaccaccagg gcacacttac cgagtactgc ggcacgctca tgagcctgcc caccaagatc 240
tcccgtgtgc cccacctcct cgacttcttc aagggtgcgc ctgatgacct caagctcccc 300
acggacaacc agacaaaaaa gccagagaca tacttgatgc ccaaagatgg caagagtacc 360
gcgacagaca tcaccggccc catcatcctg cagacgtacc gcgccattgc caactacgag 420
aagac 425
```

<210> 449

<211> 88

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (25)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (26)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (73)
<223> n equals a,t,g, or c

<400> 449
tcgttgccct gttactgtct gtggnnatgt gcatgggtcaa tttcatgatc ttatgggaact 60
ctttagaatt ngnggaaaat caccggat 88

<210> 450
<211> 214
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (8)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (26)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (31)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (49)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (53)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (68)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (75)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (89)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (91)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (102)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (108)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (113)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (114)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (141)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (154)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (193)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (204)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (205)
<223> n equals a,t,g, or c

<400> 450
atctccantg gaaccacggt gtggcngatg ntggggcctt cctgcggtnc aangagccct 60
taccgganta ngtnggatg gtgactgant nttcaaataa antacganac tgnnccaacc 120
tctctgccac ggacatccaa ntgcttgac gcanatacca gttggaagca gagtttggtg 180
gggtgtctca acnaaaagtt gagnaagca acgt 214

<210> 451
<211> 473
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (370)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (444)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<400> 451
gcggacgcgt ggggcaagac ttttgcccgg tacctttcat tccggcgtga caacaatgag 60
ctgttgctct tcatactgaa gcagttatg gcagagcagg tgacatatca gcgcaaccgc 120
tttggggccc agcaggacac tattgaggtc cctgagaagg acttggtgga taaggctcgt 180
cagatcaaca tccacaacct ctctgcattt tatgacagt agctcttcag gatgaacàag 240
ttcagccacg acctgaaaag gaaaatgatc ctgcagcagt tctgaggccc tatgccatcc 300
ataaggattc cttgggattc tggtttgggg tggtcagtgc cctctgtgct ttatggacac 360
aaaaccagan cacttgatga actcggggta ctagggtcag ggcttatagc aggatgtctg 420
gctgcacctg gcatgactgt ttgnttctcc aacctgcctt gtgcttctna cct 473

<210> 452
<211> 397
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (29)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (225)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (330)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (376)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (385)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (389)
<223> n equals a,t,g, or c

<400> 452
gggtcgaccc acgcgtccgc ccacgctcnc gcttgcttgg acatcaggta ccaactgcac 60
tcgttttggc attgcagcta aatatcagtt ggatccact gcttccattt ctgcaaaagt 120
caacaactct agcttaattg gagtaggcta tactcagact ctgaggcctg gtgtgaagct 180
tacactctct ggctctggta gatgggaaga gcattaaatg ctggnaggcc acaagggttg 240
ggctcgccct gggagttggg aggcttaatc cagctgaaag aaaccttttg ggaatgggat 300
atccagaaga tttgggcctt aatataattn ccattgtgga ccagcagcag gctttttttc 360
cccccaagaa gattgntcca aaccnangnt gatctcc 397

<210> 453
<211> 463

<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (393)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (404)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (426)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (435)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (453)
<223> n equals a,t,g, or c

<400> 453
ggcaagatgg tgttgcagac ccaggtcttc atttctctgt tgctctggat ctctggtgcc 60
tacggggaca tcgtgatgac ccagtctcca gactccctgg ctgtgtctct gggcgagagg 120
gccaccatca actgcaagtc cagccagagt gttttataca gtcceaaca taagaactac 180
ttaacttgg accagcagaa accaggacag cctcctaagc tgctccttta ctgggcatct 240
acccgggaat ccgggggtccc tgaccgattc agtggcagcg ggtctgggac agatttcact 300
ctcaccatca gcagnctgca ggctgaagat gtggcagatt attactgtca gcaatattat 360
actactccct ggacgttcgg ccactggacc aangtgaaa tcanacgaaa ctgtggctgc 420
accatntgcc tcatntttcc gccatctggt gancagttga aat 463

<210> 454
<211> 332
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (95)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (332)

<223> n equals a,t,g, or c

<400> 454

```

tggctcatct tggctgtgat ttcagtatat gatttagtgg ctgttttgtg tccgaaaggt 60
ccacttcgta tgctggttga aacagctcag gaganaaatg aaacgctttt tccagctctc 120
atttactcct caacaatggg gtggttggtg aatatggcag aaggagaccc ggaagctcaa 180
aggagagtat ccaaaaattc caagtataat gcagaaaagca cagaaaggag tcacaagaca 240
ctgttgacaga gaatgatgat ggcgggttca gtgaggaatg ggaagcccag aaggacagtc 300
atctagggcc tcatcgctct acacctgaat cn                               332
```

<210> 455

<211> 429

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (99)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (105)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (107)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (109)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (143)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (148)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (158)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (172)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (243)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (271)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (274)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (292)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (314)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (346)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (382)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (392)
<223> n equals a,t,g, or c

<400> 455
ggcnagtagt nnggggtccc aggggttggt tttaaagtag tgacttctac caacatgtcc 60
cgtgggtcca gagagcgggt ttgaccgcca cattaccant ttttnancng agggtcggct 120
ctaccaagta ggaatatgct ttnaaggnta ttaaccangg tggacttaca tnagtagctg 180
atcagcggga aagactgtgc agtaattgtc acacagaaga aagtacctga caaattattg 240
gantccagca cagtgactca cttattcaag ntanctggaa acattgggtg tntgaagacc 300
ggaatgtcag ctgncagcag atcccaggta cagagggcac gctatnaggc agctaacttg 360
gaatacaagt atggctatga gnttcctgtg gncatgcctg tgtaaaagga tttccgggtat 420
ttctcaggt 429

<210> 456
<211> 623
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (38)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (352)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (482)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (516)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (540)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (554)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (570)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (588)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (604)

<223> n equals a,t,g, or c

<400> 456

```
tggcggcttc cttcaccgc aacccgagag acgaccncc gggcccgccc cgcggaagcc 60
gccggttgcc aggccaagga gtggactagg gtcgccgggg aagcggtttg ggagagccca 120
tggtgactgc gtgagtggag cccagctgtg tggatgccc agcatggatg actacatggt 180
cctgagaatg attggggagg gctcgttcgg cagagctctt ttggttcaac atgaaagcag 240
taatcaagat gtttgccatg aaagaaataa ggcttccaag tctttctcta atacacagaa 300
ttctaggaag gaggctgttc ttttagccaa aatgaaacac cctaattattg gngccttcaa 360
agaatcattt gaaagctgca ggacacttgt atattgtgat ggaatactgt gatggacggg 420
atctaattgca aaagattaaa cagcagaaaa ggaaaagtta tttcctgaag acatgatact 480
tnaatgggtt acccaaagt gccttgagat aaaatnacat ttaccaagaa acgtgtgctn 540
cccaagagat tttnaaagtc ccaaaaaatn tttttcctta acctcaanaa ttgggaaaaa 600
gttnaaaatt ggggaaaaac ttt 623
```

<210> 457

<211> 441

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (20)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (28)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (185)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (223)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (233)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (250)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (391)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (399)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (417)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (432)

<223> n equals a,t,g, or c

<400> 457

```
atccngaaac tгнаанссгн tgtgtcanct gaaaaacccat tcgggtgtgtc tgcacttgga 60
ggetgcactt gcttgcttcc acttatgctt gttctcagaa cacaaacaaa acctgtgaag 120
agtgcctgaa aaacgtctcc tgtcttttgt gcaacactaa caagcttgtc ttggactacc 180
agttncaaaг tcttgccacc ggcttccctt tgгтааттаа ctncctgcac ctngggaatt 240
ttgttggtgn aaacttagaa gcgctgaatc atcacatgtt cggtagtccg gggaaaccctc 300
ctcctggggc attggcatct gctgcttgct tgcttgctgc aagaagaaaa aaagaaccgc 360
aaacccgгac aggaattaaa gaaaaaaggc nttgcgttna accgгaaaga aaaagcngaa 420
taccggccgg angaacггaa a 441
```

<210> 458

<211> 419

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (252)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (278)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (306)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (402)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (404)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (408)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (409)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (410)

<223> n equals a,t,g, or c

<400> 458

ncttcctnta ggagcaggag ggatgagcaa gggctctcca gcgcgacagg acatggagaa 60
agagcgagag actctgcagg cctggaagga gcgcgtgggg caggagctgg accgcgtggt 120
ggctttcttg atggagcact cccacgacca ggagcacggg ggcttcttca cgtgccttgg 180
ccgcgagggg cgggtgtatg atgacctcaa gtatgtgtgg ctgcagggga ggcaggatatg 240
gatgtattgt cngcctgtac cgcactttcg agcgcttncg ccatgctcag cttctggacg 300
cagcanaagc aggtggtgag ttcttgctgc ggtatgcccg ggtggcacct tctggcaaga 360
agtgtgcctt tgtgctgact cggtgacggc cgcccgggtca angngcannn aaccatctt 419

<210> 459

<211> 509

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (284)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (308)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (331)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (369)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (377)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (393)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (405)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (410)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (414)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (419)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (424)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (440)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (472)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (485)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (490)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (493)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (499)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (502)
<223> n equals a,t,g, or c

<400> 459
aattcggcac gagctgaagt nactgatgag tataaaaatn atgtaaaaaa cagatctgtt 60
tatattaaag gcttcccaac tgatgcaact cttgatgaca taaaagaatg gttagaagat 120
aaaggtcaag tactaaatat tcagatgaga agaacattgc ataaagcatt taagggatca 180
atthttgttg tgthtgatag cattgaatct gctaagaaat ttgtagaggc ccctggccag 240
aagtacaaag aaccagacct gctaatactt ttcaaggccg gttnccttgc caaaaaatga 300
ggaagaanca aataagtga gctaattagg nttacagggc agagcaacca agttgaggag 360
tcctgatgna ctctggngaa gttggtcctt ctnatttcgg ggttngtgcn gcctgggng 420
ttcncacttt tcctcagggn taatgtgctc ccaggccagg gtttttttag anccggcttg 480
taccngtcan tgnccctng cnggtcttg 509

<210> 460
<211> 468
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (20)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (445)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (450)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (459)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (465)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (468)
<223> n equals a,t,g, or c

<400> 460
cgcccttttg taccgggtccn gaattcccgg ggcttccctc ctcttccttt cttcgccatc 60
gtgggtgtgtt cttgactccg ctgctcgcca tgtcttctca caagactttc aggattaagc 120
gattcctggc caagaaacaa aagcaaatc gtcccattcc ccagtggatt cggatgaaaa 180
ctggaaataa aatcaggtac aactccaaaa ggagacattg gagaagaacc aagctgggtc 240
tataaggaat tgcacatgag atggcacaca tatttatgct gtctgaaggc cacgatcatg 300
ttaccatatt aagctgaaaa tgtcaccact atctggagat ttcgacgtgt tttcctctct 360
gaatctgtta tgaacacgtt gggtggctgg attcagtaat aaatatgtaa ggcctttctt 420
tttaaaaaaa aaaaaaaaaa aaaaanaaan acaaaaaana tcaanaaa 468

<210> 461
<211> 580
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (18)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (54)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (249)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (340)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (427)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (433)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (452)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (462)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (466)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (470)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (472)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (478)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (509)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (510)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (556)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (567)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (575)

<223> n equals a,t,g, or c

<400> 461

```
ggggggggng ggggggtntt aggtttgcct aaaaatattt aggtgacact atanaaggta 60
cgcttgcagg taccgggtccg gaattcccgg gtcgacccac gcgtccgtcg acccacgcgt 120
ccgcccacgc gtccgcccac gcgtccgccc acgcgtccgc tgtgtcgtaa aatgggggtc 180
ccttactgca ttatcaaggg aaaggcaaga ctgggacgtc tagtccacag gaagacctgc 240
accactgtng ccttcacaca ggtgaactcg gaaagacaaa ggcgcttttg ctaagctggg 300
ggaagctatc aggaccaatt acaatgacag atacgatgan atccgcccgt ccactggggg 360
tggaatgtt ccctgggtcc ctaattcttg ttgctccgta tcgccaact ccgaaaaagg 420
caaaagntta aanaactttg ccacttaaac tngggttaaa tnttcnctgn tnaatttncc 480
ctgttccctt aaaaataatt gaaattatnn aatttcccc tccccaaaaa aaaaaaaaaa 540
gggggggccc ttaaanattc cccccnaag gggcnaatta 580
```

<210> 462

<211> 549

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (11)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<400> 462

```
acnngnanac nancctcact aaaggaaca aaagctggag ctccaccgcg gtgcggccgc 60
tctagaacta gtggatcccc cgggctgcag gaattcggca cgaggcgccg ccacaatggt 120
gcgcatgaat gtcctggcag atgctctcaa gagtatcaac aatgccgaaa agagaggcaa 180
acgccagggtg cttattaggc cgtgctccaa agtcatcgtc cggtttctca ctgtgatgat 240
gaagcatggt tacattggcg aatttgaaat cattgatgac cacagagctg ggaaaattgt 300
tgtgaacctc acaggcaggc taaacaagtg tggggtgatc agccccagat ttgacgtgca 360
actcaaagac ctggaaaaat ggcagaataa tctgcttcca tcccgccagt ttggtttcat 420
tgtactgaca acctcagctg gcatcatgga ccatgaagaa gcaagacgaa aacacacagg 480
agggaaaaatc ctgggattct tttctaggg atgtaataca tatatttaca aataaaatgc 540
ctcatggac 549
```

<210> 463

<211> 480

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (7)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (130)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (320)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (410)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (416)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (450)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (455)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (456)
<223> n equals a,t,g, or c

<400> 463
gcctctngcg ggccctgttc aagcggaatc ctgccaaaccg gctcggctcc ggccctgatg 60
gggcagagga aatcaagcgg catgtcttct actccaccat tgactggaat aagctatacc 120
gtcgtgagan cagccacccc ttcaagccag cagtggctca gcctgatgac accttctact 180
ttgacaccga gtacacgtcc cgcacaccca aggattcccc aggcattcccc cccagcgtg 240
gggcccacatca gctgtttccgg ggcttcagct tcgtggccac cggcctgatg gaagacgacg 300
gcaagcctcg tgccccgcan gcaccctgc actcgggtgg acagcaactc catgggaaga 360
acctgggttt tagtgacggc tacgtggtaa aggagacaat tgggtgtgggn tcctantctg 420
agtgaagcg ctgtgtccac aaagggccan aacannaata atgcaatagg ggaggtaaca 480

<210> 464
<211> 220
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (17)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (18)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (30)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (42)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (58)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (63)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (67)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (72)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (105)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (129)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (147)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (157)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (170)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (213)

<223> n equals a,t,g, or c

<400> 464

```
agggacacgt cccatnnngt ggcgggcgcg ctgcggccct tngtgcaggc cacggtgncc 60
gcnaccngaa gncagcctgt ttggaccta aagcggccct tcctnagccg ggagtcgctg 120
agcggccang cctgcgatcg acttgtngtc gactccntgg gtgctcaatn tccctgcttc 180
tttttgtaa ttcccacaca gacatcaagg tgnctgattt 220
```

<210> 465

<211> 438

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (330)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (366)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (383)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (403)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (414)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (428)

<223> n equals a,t,g, or c

<400> 465

```
ataggcatga gccaccacgc caggccggga atgccctatt ctaattcaca taaagactcc 60
atgtgggcta gcaggcatgt gaaatgtctg ccagtgcac aacagctata taactaagaa 120
gttatctttt tcaactgtctt ctggtactct tcatttcact tgggtttttt agcaattcag 180
gaagctggaa tgtttgaatg ggtttttagct tgataccttc ttctttttcc catttggcag 240
ataataccac tagtctgacg gataaacacc tggacccaat cagggaaaat ctgggaaaagc 300
actgggaaaa actgtgcccg taaactgggn ttcacacagt ctcagattga tgaattgacc 360
atgacnatga gcgagatgga tgnaagaaag gttaccagat gcnccaaaat gggngatgag 420
gaaggctnaa ggggccccg                                     438
```

<210> 466

<211> 127

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (53)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (87)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (94)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (118)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (122)

<223> n equals a,t,g, or c

<400> 466

```
gcacgattca tgcaaaaagg naactaagca ctatgagatg cttgctaata gancagctgc 60
aaatgggtcac tgcattgata tttatgnttg tgcncctgat caaactggac ttttgganct 120
gnagtgt                                     127
```

<210> 467
<211> 439
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (40)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (194)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (295)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (409)
<223> n equals a,t,g, or c

<400> 467
gctggaatct ttacaggaga accacttcca agaagatgan cagtttctgg gagccgttat 60
gccaaaggctt ggcattggaa tggatacttg tgcattcct ttgaagcacg gtgggctttc 120
cttggttcaa accacagatt acatttacct gatcgtagac gacccttaca tgatgactcc 180
tgcagttgct gaantcaggc ctgtcccctg cccacacttg gcactgggca taaagcaatt 240
agggaggaag caggaaagcc ctctgctgct gttgcaactg aatacatgct ggcangataa 300
catgtgccaa tgtcctcagt gactctatgc aatgggggtc acggaatgtg acaatatgct 360
gatgctcctt ggaatcataa taaaatgacc gacaggaaaa ggataaatna tcctctgatt 420
atccaagggtt ttaaaacca 439

<210> 468
<211> 484
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (350)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (399)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (412)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (415)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (449)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (463)
<223> n equals a,t,g, or c

<400> 468
gagcaggccc gttggaagtg gttgtgacaa cccagcaat gtggagaagc ctggggccttg 60
cctggctctc tgtctcctcc catcgggagg aacagagagc caggaccaa gctccttatg 120
taagcaaccc ccagctggac ataagagatc aagatcaatg ctaaaactcca atgggttcagt 180
gactgtggtt gtcttcttca agccagctga tacctgtcat actgcaggca tctaaattag 240
aaacctgcga gtaaaactga agaaagaagg atattctaata attcttatat tgtgtaatat 300
taaggaaatct cttctcgata aaatcacaca tctaagaata aggttaaaan atatttctgt 360
tataccagaa gaaacaacaa tgctgactct ttaatgaana aataactctat antanatggg 420
cgctgaatac tgttgctttc tcacttcang aaaacataat gtntgaaaat ggacgtttca 480
ttaa 484

<210> 469
<211> 489
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (308)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (348)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (368)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (371)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (387)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (417)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (420)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (458)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (468)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (477)
<223> n equals a,t,g, or c

<400> 469
cgccctggcga ttaccggtct caccatggag cggaaagtgc ttgcgctcca ggcccgaag 60
aaaaggacca aggccaagaa ggacaaagcc caaaggaaat ctgaaactca gcaccgaggc 120
tctgctcccc actctgagag tgatctacca gagcaggaag aggagattct gggatctgat 180
gatgatgagc aagaagatcc taatgattat tgtaaaggag gttatcatct tgtgaaaatt 240
ggagatctat tcaatgggag ataccatgtg atccgaaagt taggctgggg acacttttca 300
acagtatngg ttatcatggg gtattccagt taagttttta ttggttcntg taaaaagatt 360
agtttaganga ngtctaaaag ggtttgntct agttccatgg tggacccagg ttcaacnacn 420
ctagccggtc cattaccaga atttgttgtt ggcaaaagncc cgatcctngg ggaaccnttt 480
ctttcgaa 489

<210> 470
<211> 318
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (2)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (12)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (21)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (32)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (73)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (102)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (103)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (106)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (114)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (151)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (188)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (223)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (258)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (264)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (280)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (281)
<223> n equals a,t,g, or c

<400> 470
cntcactata tntgaggagc nggtaccct gnaggtaccg gtccggaatt cccgggtcga 60
cccacgcgtc cgntatgaca acctgatcac accagccatg annggngccg gctncctgca 120
ggggaacgtc gattcttgcc agggtgacag nggagggcct ctggtcactt cgaagaacaa 180
tatctggngg ctgatagggg atacaagctg gggttctggc tgngccaaaag cttacagacc 240
aggagtgtac ggggaatgnga tggnatcac ggactggatn natcgacaaa tgagggcaga 300
cggctaattcc acatggct 318

<210> 471
<211> 455
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (306)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (417)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (426)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (431)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (440)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (446)
<223> n equals a,t,g, or c

<400> 471
ggagtataca ccatgagcaa agctcaccct cccgagttga aaaaatttat ggacaagaag 60
ttatcattga aattaaatgg tggcagacat gtccaaggaa tattgcgggg atttgatccc 120
tttatgaacc ttgtgataga tgaatgtgtg gagatggcga ctagtggaca acagaacaat 180
attggaatgg tggtaatacg aggaaatagt atcatcatgt tagaagcctt ggaacgagta 240
taaataaatgg ctgttcagca agagaaaccc atgtcctctc tccatagggc ctgttttact 300
atgatnttaa aaattaagtc atgtacattt tcatattaaa ctttttgta aataaaacttt 360
tgtaataatc aaaaaaaaaa aaaaaaaaaa aaaccaagg gggggcccg tccccantcc 420
cccctntttt nattcctttn aaaatncctt ggccc 455

<210> 472
<211> 676
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (605)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (669)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (672)
<223> n equals a,t,g, or c

<400> 472

```
gctccaggag tgggacggag ggagctggcc gggatgaagt ctgagactat gtcctgagaa 60
gaaagagtgt atcgtattgg ttgaaaagtt ggtggggtcg ggcttaagcg gaggaggggg 120
ctctctggcc cttactcggc agatggggcc ggagagagga cgggaggtgc cgggagaaca 180
tcgagggacc ggtggaggaa gggtagctgg atgagttttg attcatcatg gataatctgt 240
catcagaaga aattcaacag agagctcacc agattactga tgagtctctg gaaagtacga 300
ggagaatcct gggtttagcc attgagtctc aggatgcagg aatcaagacc atcactatgc 360
tggaatgaaca aaaggaacaa ctaaaccgca tagaagaagg cttggaccaaa ataaataagg 420
acatgagaga gacagagaag actttaacag aactcaacaa atgctgtggc ctttgtgtct 480
gccccatgtaa tagaacaagg aactttgagt ctggcaaggc ttataagaca acatggggag 540
atggtggaga aaactcacct tgcaatgtag tatctaaaca gccaggcccg gtgacaaatg 600
gtcanccttta gcaaccaaca acaggagcag ccagtgggtg atacattaaa cccataacta 660
atgatcccng anaaga 676
```

<210> 473

<211> 512

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (1)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (403)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (457)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (487)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (495)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (500)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (506)

<223> n equals a,t,g, or c

<400> 473

```
ntcgacagaa ggggtacggct gcgagaagac gacagaaggg tacggctgcg agaagacgac 60
agaagggggc agcatggcgt acccggggca tcctggcgcc ggcggcgggt actaccag 120
cgggtatgga ggggtcccg gagggcctgc gtttcccgga caaactcagg atccgctgta 180
tggttacttt gctgctgtag ctggacagga tgggcagata gatgctgatg aattgcagag 240
atgtctgaca cagtctggca ttgctggagg atacaaacct tttaacctgg agacttgccg 300
gcttatgggt tcaatgctgg atagagatat gtctggcaca atgggtttca atgaatttaa 360
agaactctgg gctgtactga atggctggag acaacacttt atnaattttt gacactgcag 420
gaatggaaca agtagacca caagaattgg ataaagnccc tgacaacaat gggatttaag 480
gtttgantcc ccaanctggn gaattnaatt gc 512
```

<210> 474

<211> 272

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (18)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (59)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (69)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (74)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (104)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (122)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (129)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (150)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (185)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (191)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (234)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (249)
<223> n equals a,t,g, or c

<400> 474
ggcacagcgg cgggccnngt cgtccggcgg ttgcggatgt cgggetgggc ggacgagcnc 60
ggcgtcgang gccnacgggc gcatctacgt ggggaacttc cgancgacgt gcgcgagaag 120
gnacttggnna ggacctgttc tacaagtacn gccgcatccg cgagatcgag ctcaagaacc 180
ggcannccctc ntcctgtcgg cttcgtgcg cttcgaggaa ccccgagat gcanaggatg 240
ctatttatng gaagaaatgg ttatgattat gg 272

<210> 475
<211> 338
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (8)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (15)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (36)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (119)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (127)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (170)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (173)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (204)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (250)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (284)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (298)

<223> n equals a,t,g,, or c

<220>

<221> misc feature

<222> (308)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (318)

<223> n equals a,t,g, or c

<400> 475

```
naatctcngt ttggnnggcc cgcccgagg tccggnccgg aactcccggg tcgacccacg 60
cgccgggcta ccattctgaa acaaatgcaa gtattacatc cagcagccag aatgctggng 120
gagctgncta aggctcaaga tatagaagca ggagatggca ccacatcagn agncatcatt 180
gctggctccc tcttagattc ttgnaccaag cttcttcaga aagggattca tccaaccatc 240
atttctgagn cattccagaa ggccctggaa aagggcattg aaancttgac tgacatgnct 300
cgacctgngg aactgagnga cagagaaact ttggtaaa 338
```

<210> 476

<211> 424

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (330)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (378)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (381)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (388)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (424)
<223> n equals a,t,g, or c

<400> 476
ttaagttcat gaattccaga gttttcaaga aaattcaagc tttgaaggct tcaccgtcta 60
agaaacggtg caattcaatc gccgccctaa aggccacttc acaggagatt gtgtcctcaa 120
ttagccagga atggaaggat gagaagcggg atttgctgac tgaaggacaa agtttttagca 180
gccttgatga agaagccctg ggatcccgcac acaggccaga cctgggccct agcactccat 240
cactgtttga agctgcttcc ttggcaacca caatttctact tcttcctata cgtcaatggg 300
cattatccac aagacaaggc ctacaatttn ttcaaaccag gtaatatattt ggtgcaattc 360
aaattaaccc ttcaaggntt nttaaccnca atcaaccatt agggcctgaa cccgtttttt 420
ttgn 424

<210> 477
<211> 228
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (6)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (111)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (153)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (157)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (165)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (207)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (223)
<223> n equals a,t,g, or c

<400> 477
ggcatncatg tcttcagaaa catccataag accaatctct gtgatcttat caccagcctt 60
ttgtgccttt nggttctttt gccaaccaaa gaactcaatg aacacttctn ntccaagctt 120
aaggcaccga ttcctattga acttggtgtt gtngtancag ccacnttaac ctctcatttt 180
ggaaaactac atgaaaatta taattcnagt attgctggac atnttccc 228

<210> 478
<211> 486
<212> DNA
<213> Homo sapiens

<400> 478
aactagtctc gaacggcccc ggggacacgc tggatctgac ctactggatt gacggcaccc 60
ggcatgtggt ctccctggag gacgtcggcc tggctgactc gcagtggaag aacgtcaccg 120
tgcaggtggc tggcgagacc tacagcttgc acgtgggctg cgacctcata gacagcttcg 180
ctctggacga gcccttctac gagcacctgc aggcggaaaa gagccgatg tacgtggcca 240
aaggctctgc cagagagagt cacttcaggg gtttgcttca gaacgtccac ctagtgtttg 300
aaaactctgt ggaagatatt ctaagcaaga aggggtgcca gcaaggccag ggaggtaggt 360
gtgttgtgaa aaatgcattt tatatactgg cttggatgga tttctattgt gacatgggtg 420
atgtgtgtgt gtgtatgtgt gtgcattcat gcttgtaaaa atttaagtaa aaatatttgg 480
gtaact 486

<210> 479
<211> 524
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (70)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (369)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (409)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (432)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (471)
<223> n equals a,t,g, or c

<400> 479
ggtgcctgct gcctcaaacc accttgtgta aaatctgcaa tacggcttct tccatgtacc 60
tgtgcctgan ctgtctgcaa taaagaagaa tttgtaaaaa aacggcacct cgctttgctt 120
ttcctccgca accatgtctg acaaaccgga tatggctgag atcgagaaat tcgataagtc 180
gaaactgaag aagacagaga cgcaagagaa aaatccactg ccttccaaag aaacgattga 240
acaggagaag caagcaggcg aatcgtaatg aggcgtgctg cgccaatatg cactgtacat 300
tccacaagca ttgccttctt attttacttc ttttagctgt ttaactttgt aagatgcaaa 360
gaggttggnt caagtttaaa tgactgtgct gcccctttca catcaaagna ctactgacaa 420
cgaaggccgc gncgtgccttt cccatctgtc tatctatctg gctggcaggg naggaaagaa 480
cttgcattgt ggtgaaggaa gaagtggggt ggaagaagtg ggggt 524

<210> 480
<211> 306
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (15)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (19)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (34)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (52)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (64)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (66)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (76)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (97)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (126)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (129)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (130)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (133)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (134)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (145)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (146)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (161)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (166)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (181)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (186)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (194)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (200)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (233)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (267)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (275)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (304)

<223> n equals a,t,g, or c

<400> 480

```
ggcagagggc ccgcnagcng ccgccctcct cctnagcctg ctgctgttcg gnttcaccct 60
agtntnaggc acaggnagcag agaagactgg cgtgtgnccc gagctccagg ctgcaccagc 120
aactgncann cannagtgcg tcctnnggaa cagcgaaatg ncccgnacaa cctccaagtg 180
nctgcnaggc gggncgtgtg cacccttctgc ctctctgccc caatgatgaa ggnggggttc 240
tgccccccagg tgnaacatta aacttttccc aggtngggcc tctgttcggg caccatgcca 300
ggtngt                                     306
```

<210> 481

<211> 473

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (364)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (411)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (418)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (431)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (445)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (459)

<223> n equals a,t,g, or c

<400> 481
gccttttagct gccatcttgc gtccccgcgt gtgtgcgcct aatctcaggt ggtccacccg 60
agacccttg agcaccaacc ctagtcccc gcgcggcccc ttattcgctc cgacaagatg 120
aaagaaacaa tcatgaacca ggaaaaactc gccaaactgc aggcacaagt gcgcattggt 180
gggaaaggaa ctgctcgag aaagaagaag gtggttcata gaacagccac agcagatgac 240
aaaaaacttc agttctcctt aaagaagtta ggggtaaaca atatctctgg tattgaagag 300
gtgaatatgt ttacaaacca aggaacagtg atccacttta acaaccctaa agttcaggca 360
tctntggcag cgaacacttt caccattaca ggccatgctg agacaaagca nctgacanaa 420
atgctacca ncatcttaaa ccagnttggt gcggatagnc tgactaagtt taa 473

<210> 482

<211> 571

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (4)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (13)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (70)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (92)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (297)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (307)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (494)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (550)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (565)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (569)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (571)
<223> n equals a,t,g, or c

<400> 482
gannacttct ctngaaacgc ccgtaccggt cttgaattcc cggctcgacc cacgcgtccg 60
gatgacggtn cggggcgccct ctcccagga cncttgggtc aagggtggagt atgcctacag 120
cgacaacagc ctggaccccg ggctttttgt agaaagcacc cgcaagggga gtgtagtggtc 180
cagagctaat agcatcggtt ccaccagtgc ctcttctgtc cccaacacag atgatgagga 240
cagtgattac caccaggagg cctacaagga gtcctacaaa gaccggcggc ggcgcgnaac 300
acacttnagg cttgagcaga agaggaggga cgccatcaag agaggctatg atgaccttca 360
gaccatcgtc cccacttgcc agcagcagga cttctccatt ggctcccaa agctcagcaa 420
agccatcgtc tacaaaagac cattgactac attcagtttt tgcacaagga gaagaaaaag 480
caggaggagg aggngtcacg ttacgcaagg atgtaccggc ctaaagatca tgaaagtga 540
ctatgagcan attgtgaagg cacantagna n 571

<210> 483
<211> 220
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (16)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (69)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (99)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (136)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (179)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (183)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (185)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (215)
<223> n equals a,t,g, or c

<400> 483
aatacaacgg gaatcnctaa ctgccttgcg tgaagggtccc ggtccggaat tcccggggtcg 60
acccacgcnt ncgggccagg atgctgaatc tgctgctgnt ggcgctggcc gtcctggcga 120
gccgcgccta cgcggnccct gccccaggcc aggcctgca gcgagtgggc atcgtcggng 180
gtnangaggc ccccaggagc aagtggccct ggcangtgag 220

<210> 484
<211> 439
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (5)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (18)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (31)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (35)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (36)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (394)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (406)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (414)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (434)
<223> n equals a,t,g, or c

<400> 484
gggaggaggag gggaaatnct tctcgggtatc nggtnnctgg agctccaccg cgggtggcggc 60
cgctctagaa ctagtggatc ccccgggctg caggaattcg gcacgagcag cccagcagcg 120
gctgaccctc tgcctgcggg gaagggagtc gccaggcggc cgtcatggcg gtgtcggaga 180
gccagctcaa gaaaatggtg tccaagtttt taacgatggc agttccaggg aactaatgaa 240
cctcactgga acaatccctg tgccttatag aggtaataca tacaatattc caatatgcct 300
atggctactg gacacatacc catataatcc ccctatctgt tttgttaagc ctactagttc 360
aatgactatt aaaacaggaa agcatgttga ttgnccaaaa aaattngggg gggnaaaaaa 420
ggggaaaatt ttanttttt 439

<210> 485

<211> 115
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (11)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (69)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (71)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (87)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (89)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (104)
<223> n equals a,t,g, or c

<400> 485
gacaaagtaa ngcatccctc ctgaccatca gacgttgatc ttgctggga aacatctgga 60
aaatggacnc nccctgtctg actacancnt ccacaaagaa tccnccctgc acctg 115

<210> 486
<211> 558
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (12)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (151)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (159)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (255)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (270)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (353)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (402)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (404)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (410)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (416)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (488)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (495)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (517)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (539)
<223> n equals a,t,g, or c

<400> 486
ntctctgngg tntagctgaa cgccgaggac cggccgaatt cccgggtcgac cccgcgttcg 60
ttcttttagc atcaaggaat ctgatgagaa agcaggaaaag atcttttcac agatgaacaa 120
tattatgagt aaaagtttgc atgatgatgg ntttactgnt ccacagatta ttgaaatgga 180
gctggatagt caggagcaag ttgttggtgc aggatcctcc tgtgacttac attcagcaat 240
ttgcagatgc agcanccaac cttacctctn cggattctga gaagtggaaac tctgtgtttc 300
ccaagcctgg gacttttggtt caagtgcttg aggctgcaaa gtttgcatag gangagaatc 360
tttgataaac tcaataactc cttgtgctga cggaattcat tngntggtan gactgnggac 420
atgccctggt gaatccttga gtgcaataaa tcaaagtaga aggccttgna taatttaaata 480
taaattanac tctgnactat gtaataaccg gaaaggnaac tgcaatcaaa ttgcttgna 540
tggaatgggg caaaatta 558

<210> 487
<211> 354
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (13)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (15)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (18)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (21)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (79)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (114)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (115)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (120)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (127)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (155)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (195)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (203)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (229)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (239)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (307)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (315)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (339)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (343)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (346)
<223> n equals a,t,g, or c

<400> 487
gctctgctca acngngangg ntcggcacia cattcagaag gaattatgtc ttcattgctgc 60
tcaaggtctc gctcagctna aggcattgtac ctacaaaggt cacaagacag gtggnactgn 120
agagcanata tgggagatcc agaaggatca acttntatac tatccattct taaaaatgtg 180
cctttcagca aatgntgagc atncaagctt agtggatgca acccatcana accactccna 240
aaatggatag ttagccaaaa tgattaagcg ttccttaaaa ttaacttgac caaggaaata 300
tcctttntca taaanctgtg actaggcata cactgtagnt gtnganaatt atgc 354

<210> 488
<211> 508
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature

<222> (134)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (236)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (242)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (258)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (275)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (288)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (298)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (300)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (314)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (349)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (375)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (387)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (400)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (407)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (412)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (413)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (424)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (429)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (434)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (440)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (441)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (450)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (456)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (458)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (462)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (469)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (475)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (478)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (485)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (493)
<223> n equals a,t,g, or c

<400> 488

```
aattcggcac gagaaagcgc atggagacta agggagctgg agtgaccctg aatgttcttg 60
aaatgacttc tgaagattta gaaaatgctc taaaagcagt catcaatgac aaaagttaca 120
aggagaacat catncgcctc tccagccttc acaaggaccg cccggtggag ccgctggacc 180
tgcccggtgt ctgggtggag tttgtgatga ggcacaaggc cgcgccacac ctgcgncctg 240
cnccccacgg acctcacntg gtaccagtac cattnccttg gccgtgantt ggtttccntn 300
ttggcccttg gtcntgaaat tggccttaat aacctttaa attttgctnt natggttacc 360
cgaaaatttt ggggnaaaaa agggccnttt aagaaaagcc caaaatnaga cnnccctttg 420
aaangggtn gganaaaagg naaaattttt accttncncg gnatttccna atttnaanag 480
atagngttta aanatttttt ttttttagg 508
```

<210> 489

<211> 425

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (3)

<223> n equals a,t,g, or c

<400> 489

```
ccnccgtggg atgttgagct gaggccggag gagaagtagc agtcgctggc agagcacaca 60
ggctgctctg ggggatgagc tgggtgcgtt aaggaacagg ccagcactgg cattcgcaag 120
cagtggggaa ggggagagat gccgaggttg tcagtatcct gactttcaga ggcctttttt 180
tgtttgtttt aatttttgc agattgatat taaaaactca tgtggaggaa ctcaaggaat 240
gtttagaaga ccaaaagtcc ccaatgacag gaacaaaagc aaccaatttt taactttctc 300
ttctcattcc tgttttcatt gatttccac atgtagtcc tttgttcagg aagtccttgg 360
ggaaattaag gatctttgaa gctctgaaat aggtgatcag gttagtggg tctgtcagct 420
gtctg 425
```

<210> 490

<211> 607

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (2)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (5)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (6)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (8)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (9)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (58)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (443)
<223> n equals a,t,g, or c

<400> 490
ancannncna caccacctca ctaaaggga caaaagctgg agctccaccg cggcgcncc 60
gctctagaac tagtggatcc cccgggctgc aggaattcgg cagagaagc cgtcaaggag 120
tagaaattgg tatgcttaga agcagattct aaaagcagtt tctcttcaga acatcttttt 180
tcataccact tgataagcat cttgaaacac catggctgta gctgcagtaa aatgggtgat 240
gtcaaagaga actatcttga aacatttatt tccagtccaa aatggagctt tatattgtgt 300
ttgtcataaa tctacgtatt ctctctacc agatgactat aattgcaacg tagagcttgc 360
tctgacttct gatggcagga caatagtatg ctaccaccct tctgtggaca ttccatatga 420
acacacaaaa cctatccctc ggncagatcc tgtgcataat aatgaagaaa cacatgatca 480
agtgtgtaaa accagattgg aagaaaaagt tgaacacctt gaggaaggac ctatgataga 540
acaacttagc aaaatgtttc ttactacta agcaccgttg gtatcctcat ggacgggtatc 600
acagatg 607

<210> 491
<211> 371
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (81)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (265)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (336)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (339)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (353)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (364)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (368)

<223> n equals a,t,g, or c

<400> 491

```
aaactgtaat tgcaatgttg aggggtgact tgctaaccag tcagttgaat tccaggcaca 60
gattgagttg cttctggagt nggcagggtc actgtgaacg cctacgtgag cctcttctac 120
accataaaagc gggcacaaagt ggtcagtcct gaaagagtcg gaagctggca tatcggccgc 180
ccaagtgaac ctgtccagtg tctgcttgcc atcctgccag aacaggccct caagcccaag 240
agccatccca ggcctgtttc agctncagct aaagcctctc tttcatctgg aagaagaggc 300
aagggggcag gagaccaggc tctagctctg gggccntent tcagcccca tcnggggaat 360
aaanttantt t 371
```

<210> 492

<211> 440

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (347)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (353)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (397)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (402)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (434)
<223> n equals a,t,g, or c

<400> 492
ttgagtcgtg ttaatgtaag aatgactcct atcattagga gtgctgctcg gaggttactc 60
acctttggga gtaatactga agagaggggt ctgcagaaag gatgtgtatg aagcttagat 120
aataatggct gtttcgtaaa ctgtttgaga cctattaatg aaaatgacta tttcttgctg 180
tttttatcca acgtctgcat tttccccctt taaagctgcg gtctcctggt tgataaaaga 240
atattggcca gtattgcaga ttttaactgg atttggtga tcctccaggg gaccagtttc 300
tgtgggcgtg tattggagca ggtttgtctt taactcttaa atggttnggt ccnaattttt 360
aaaaaggaag ggccccaagt agcccagatt taaaggngta tnccattcc caaagttcct 420
tggaacaatt accnaaagga 440

<210> 493
<211> 90
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (1)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (20)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (27)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (49)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (56)
<223> n equals a,t,g, or c

<400> 493

ngggagctgg cctggagcgn atgggcnccg tgatggatcg catggccanc ggctgncagc 60
gcatggggccc atcaatctgg agcggatcgg 90

<210> 494

<211> 218

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (28)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (30)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (46)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (47)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (53)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (78)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (87)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (95)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (135)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (163)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (167)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (192)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (216)

<223> n equals a,t,g, or c

<400> 494

```
gggtcgaccc acgcggtccg ttttttntn ttttttagn tccannttt tantcaagaa 60
ctcatacaaa attttccnga taaatgnaat ttaancctcg ttttctcct cttcttcgtc 120
ctggttaatc tgggnagtaa cgtaattcgt aactctcttt ggntgtnagc aactgacgcg 180
caaccagttc angtagatta ttcttcttca aatatngt 218
```

<210> 495

<211> 148

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (26)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (83)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (96)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (111)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (116)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (122)

<223> n equals a,t,g, or c

<400> 495

gtttttaaaaa aaaacatgtc atgtangttg tctaaaaata aaatgcattt aaactcaaaa 60
aaaaaaaaaaaa aaaaaaaaaa aanagggggg gggcgnggga aaaaaaaaaa ngaggntgag 120
gntttgaaga agggaagggg ggccccc 148

<210> 496

<211> 536

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (355)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (382)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (412)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (444)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (460)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (484)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (534)

<223> n equals a,t,g, or c

<400> 496

```
atgacccaca ctgccagaag agaattcagt attttttact tttcacactg tcttcccgat 60
gaggggaagga gccagccag aaagcactac aatcatgggc aagttcccaa ctgagtcac 120
ttgtgagtgg ataatcagga aaaatgagga atccaaaaga caaaaatcaa agaacagatg 180
gggtctgtga ctggatcttc tatcattcca attctaaatc cgacttgaat attcctggac 240
ttacaaaatg ccaagggggg gactggaagt tgtgggatat cagggtataa attatatccg 300
tgagtggggg gagggaagac cagaattccc ttggaattgt gtattgatgc cattntaagc 360
ctaaaagatc accttgtatt cncctttacct tctaaaagcc attatttatg gngttagaag 420
aagaggaaga aattcaggtc ccgnaaaaca cttttgttcn gggggggggc cgtacccaat 480
tcgncccata gtgagtcgta ttacaatcac gggccgctcg ttacacggcg gacnng 536
```

<210> 497

<211> 93

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (84)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (85)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (88)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (89)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (91)

<223> n equals a,t,g, or c

<400> 497

gtgaaacatc aataaagacc acttaatgca cgctttcaaa aaaaaaaaaa aaaaaaaaaa 60
aaaaaaaaaa aaaaaaaaaa aaannaanna nat 93

<210> 498

<211> 392

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (8)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (50)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (104)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (106)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (113)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (160)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (180)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (181)

<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (245)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (248)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (334)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (363)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (377)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<400> 498
cgggaccnaa ggaggagctg cgtggcgggcg gtggcgacat ggcggacctn ccgaggcggg 60
tgacgcgccc cctgatgatg ggcctccagg ggagctcggg cctnanggcc tgnacggtcc 120
agaggaaaag ggccgggatc gtgaccggga ggcacggcgn acaccggagc gagcgcgagn 180
ncgccgggac cgggatcgtg accgtgaccg tgaccgcgag cacaaacggg gggagcgggg 240
cattnaancg gggcagggat gaagcccga gttgggggcg gtggccagga caacgggttg 300
gaaggttttg gcaacgacag ccgagaattt tacntgaatt ttaaggcggc nacgttactt 360
gtnccggaat ttggttnttt attgaagttn cc 392

<210> 499
<211> 262
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (23)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (40)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (60)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (76)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (78)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (104)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (114)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (160)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (209)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (212)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (218)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (219)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (226)

<223> n equals a,t,g, or c

<400> 499

```
ggtaagagca acgtgctttg ggngcagaga agagggagan agcagcatct tgcctggatn 60
agccagggga cacagnanag aagcccacct ggacacaaca ccanaaaggc ttcntattcg 120
ggtgtggagt cttttcagca acccggtcca gtactgggcn gatacagcca tccaccttac 180
agatgtgtct acgtgacgct ctgccattnc anctcgggna ctatangtaa ttctcaagaa 240
agccctcatt tttataacct gg                                     262
```

<210> 500

<211> 437

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (309)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (317)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (384)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (398)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (405)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (406)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (418)

<223> n equals a,t,g, or c

<400> 500

```
accacaaaagc cccctggag catcttcccg gctggcagga ccatgccatc tctgtggaga 60
agggtgctggg gaggaagtc cttccagtgc cacatggagt gaggccctgc ccatgctggg 120
gactttgggg aggaatttgg tattctggtg gccttgctca gctctcattg agatcttttc 180
ctatcagaat gttagtgaat atacttcgca gctctttgtt cagcaataag gaatattctt 240
tcaattcttg ctcttcaagc caatttacta caccatttg tctttccaaa attcatccca 300
acggtatant tggtttnggt cctccttggg ttcaatctgt ttcctggtaa aattgaacct 360
gtccattgtg aatcctctta attncctcct cccaaaanaa ccagnncttg taaatttntg 420
gccaaactcca ggggccca 437
```

<210> 501

<211> 180

<212> DNA

<213> Homo sapiens

<220>

<221> misc feature

<222> (162)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (165)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (166)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (174)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (177)

<223> n equals a,t,g, or c

<400> 501

```
gagaactagt ctgcagtttt tttttttttt ttttttggtt ttttggctct ttcaaaggta 60
atggcccatc gatgagcatt tttaacatac tccatagtct tttcctgtgg tgtaggtct 120
ttatttttat ttttttcctg ggggctgggt ggggttgggg gncanngggg gaantgncct 180
```

<210> 502
<211> 436
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (101)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (174)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (197)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (199)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (222)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (225)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (229)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (262)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (266)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (276)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (287)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (290)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (297)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (327)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (328)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (330)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (336)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (343)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (364)
<223> n equals a,t,g, or c

<220>

<221> misc feature
<222> (385)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (390)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (436)
<223> n equals a,t,g, or c

<400> 502
tacaaactgt cctttattca gagtgagact gcggaacatt aataatttat cacgcgggga 60
gtccccagaa gccctgtgcc cacgaacccc tgtgggcgga nggagagaag cggggactcc 120
gggagcttcc tgagagggcc gtgtcttggg agcaagggtga catatccagt ccangcacgc 180
ggaacatgac tcagaantng ggaacaaaaa aaccatcccc cncngaang gggggggcca 240
ggccccataa aagcacaatg gnggcnggtg gaattngggt taaaatntcn gggttcnaaa 300
aagaccatat ttttttttcc cagttcnntn gcccncctt ttnttgtaa gggggggggt 360
taanattccc ccctccccgg gaaanaaaan tacctttctt aggaagaagg gcccccccc 420
ccttccccgc tttttn 436

<210> 503
<211> 418
<212> DNA
<213> Homo sapiens

<220>
<221> misc feature
<222> (41)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (50)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (55)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (56)
<223> n equals a,t,g, or c

<220>
<221> misc feature

<222> (57)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (61)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (62)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (72)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (94)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (97)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (106)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (109)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (110)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (111)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (112)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (116)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (125)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (126)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (127)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (130)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (179)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (210)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (212)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (221)

<223> n equals a,t,g, or c

<220>

<221> misc feature

<222> (226)

<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (237)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (239)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (245)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (253)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (256)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (257)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (289)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (337)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (350)
<223> n equals a,t,g, or c

<220>
<221> misc feature
<222> (351)
<223> n equals a,t,g, or c

THIS PAGE BLANK (USPTO)